

14769

Sample Submission Form

Amino Acid Laboratory
 University of California, Davis
 1020 Vet Med 3B
 1089 Veterinary Medicine Drive
 Davis, CA 95616
 Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
 Non-federal funds ID/Account Number
 to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/index.cfm>

Vet/Tech Contact: _____
 Company Name: _____
 Address: _____

(b) (6)

Email: _____
 Tel: _____ Fax: _____

(b) (6)

(b) (6)

Billing Contact: _____ TAX ID: _____
 Email: _____ Tel: _____

(b) (6)

(b) (6)

(b) (6)

Patient Name: _____
 Species: Feline
 Owner's Name: _____

(b) (6)

(b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
 Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 124 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

14771

Sample Submission Form

Amino Acid Laboratory
 University of California, Davis
 1020 Vet Med 3B
 1089 Veterinary Medicine Drive
 Davis, CA 95616
 Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
 Non-federal funds ID/Account Number
 to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/index.cfm>

Vet/Tech Contact: _____
 Company Name: _____
 Address: _____

(b) (6)

Email: _____
 Tel: _____ Fax: _____

(b) (6)

Billing Contact: _____ TAX ID: _____
 Email: _____ Tel: _____

(b) (6)

Patient Name: _____
 Species: Feline
 Owner's Name: _____

(b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
 Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 536 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

Congestive Heart Failure Due to Reversible Cardiomyopathy in Patients With Hyperthyroidism

GUILLERMO E. UMPIERREZ, MD,* SRIDEVI CHALLAPALLI, MD,*
CAM PATTERSON, MD†

ABSTRACT: The authors describe the clinical characteristics and response to therapy of seven patients with hyperthyroidism, dilated cardiomyopathy, and low-output cardiac failure. All patients (4 women and 3 men, age 47 ± 4 years, mean \pm standard error of the mean) were admitted with the primary diagnosis of congestive heart failure. The cause of hyperthyroidism was Graves' disease in six patients, and toxic multinodular goiter in one. On admission, the mean serum T_4 was 21 ± 1 $\mu\text{g/dL}$ and mean serum T_3 : 411 ± 77 ng/mL , and serum thyroid-stimulating hormone was suppressed (<0.03 $\mu\text{U/mL}$) in all patients. Two-dimensional echocardiogram showed biventricular or four chamber dilatation and impaired left ventricular performance. Therapy of heart failure and hyperthyroidism resulted in rapid clinical improvement. During follow-up (5 months to 9 years), left ventricular ejection fraction improved from a mean of 28% to a mean ejection fraction of 55% ($P < 0.01$). Resolution of dilated cardiomyopathy with normalization of systolic function was achieved in five patients, and improvement from severe to mild left ventricular dysfunction was observed in two patients. We conclude that some patients with hyperthyroidism may have a reversible form of dilated cardiomyopathy and "low-output failure." Assessment of thyroid hormone status in patients with heart failure might permit the identification of patients with dilated cardiomyopathy and thyrotoxicosis who are likely to have reversible cardiac dysfunction. **KEY INDEXING TERMS:** Hyperthyroidism; Dilated cardiomyopathy; Heart failure. [Am J Med Sci 1995;310(3):99-102.]

The association of hyperthyroidism and cardiovascular dysfunction is well established.¹⁻⁴ Hyperthyroidism is one of the most common causes of sustained hyperkinetic circulatory disease.^{5,6} This high-output state results from a direct effect of thyroid hormones that increase heart rate and cardiac contractility,^{7,8} and from an indirect effect of thyroid hormones on the peripheral circulation that results in increased blood volume and peripheral vasodilatation.⁸ This increase in cardiac work leads to cardiac hypertrophy and increased ejection fraction.⁹⁻¹¹ Paradoxically, in some patients with hyperthyroidism, high-output congestive heart failure develops despite increased cardiac performance.^{1,6,7} Likoff and Levine,³ in 1943, reported that among 409 cases of thyrotoxicosis, 21 patients had congestive heart failure in the absence of other forms of heart disease. Similarly, Sandler and Wilson² reported that 150 of 462 patients with thyrotoxicosis had evidence of cardiac dysfunction—auricular fibrillation, congestive heart failure, cardiomegaly, or all three. More recently, there have been several reports of a reversible cardiomyopathy in thyrotoxicosis, especially in children.^{12,13}

For several years, we recognized that some patients with hyperthyroidism may have low-output heart failure. Although they often had moderate to severe symptoms of heart failure and echocardiographic documented dilated cardiomyopathy, they usually experience a rapid improvement with conventional treatment of heart failure and hyperthyroidism. In this article, we describe the clinical characteristics and initial echocardiographic findings at presentation, and the response to therapy of seven patients with hyperthyroidism who had low-output heart failure and biventricular dilatation and hypokinesia.

Material and Methods

Seven patients, admitted with both hyperthyroidism and congestive heart failure due to dilated cardiomyopathy, served as the study population. All patients were seen by the endocrinology consult service at Grady Memorial Hospital between 1985 and 1994. The pri-

From the *Department of Medicine, Emory University School of Medicine, Atlanta, Georgia, and †Cardiovascular Biology Laboratory, Harvard School of Public Health.

Submitted October 14, 1994; accepted April 3, 1995.

Correspondence: Guillermo E. Umpierrez, MD, Assistant Professor of Medicine, Department of Medicine, Emory University School of Medicine, 69 Butler Street, Atlanta, GA 30303.

Table 1. Patient Characteristics and Admission Laboratory Values in Patients With Thyrotoxic Heart Failure

#	Age	Sex	T ₄ μg/dL	T ₃ ng/mL	T ₃ RU %	TSH uU/mL	Cause of Hyperthyroidism	Therapy of Hyperthyroidism
1	44	F	23	381	60	<0.03	Graves	RAI
2	82	F	20	196	48	<0.03	Toxic MNG	Surgery
3	28	F	26	800	69	<0.1	Graves	RAI
4	35	M	16	287	37	<0.1	Graves	RAI
5	65	F	21	434	68	0.1	Graves	RAI
6	29	M	19	271	36	<0.03	Graves	RAI
7	41	M	22	509	50	<0.01	Graves	RAI

T₄ = serum thyroxine, T₃ = triiodothyronine; TSH = thyrotropin (thyroid-stimulating hormone); MNG = multinodular goiter; RAI = radioactive iodine therapy.

mary diagnosis on admission was congestive heart failure. The diagnosis of heart failure was based on the Framingham criteria¹⁴ and confirmed by the presence of cardiomegaly and evidence of pulmonary edema on chest x-ray. Patients were excluded from the study population if they had a history of valvular heart disease, angina pectoris, myocardial infarction, alcoholism, or longstanding systemic arterial hypertension. All patients on admission had a cardiac evaluation that included a 12-lead electrocardiogram and a two-dimensional (2-D) echocardiogram to determine left ventricular performance, cardiac chamber dimensions, and valvular integrity, and to exclude regional wall motion abnormalities. Left ventricular performance was assessed by the fractional shortening of the left ventricle. Fractional shortening (%) was calculated from the diastolic septal-posterolateral axes measured at the level of the chordae tendineae of the mitral valve¹⁵ using the formula:

$$\frac{(\text{left ventricular diastolic diameter} - \text{left ventricular systolic diameter})}{(\text{left ventricular diastolic diameter} \times 100)}$$

Normal values of fractional shortening of the left ventricle in adults ranged from 27–37%. Determination of fractional shortening of the left ventricle allows rapid and noninvasive estimation of systolic ejective fraction (ejection fraction: % fractional shortening × 1.7).¹⁶

The diagnosis of hyperthyroidism was based on history and signs of hyperthyroidism, with laboratory data including elevated serum thyroxine (normal 5–12 μg/dL) and triiodothyronine (normal 70–190 ng/dL) concentrations and suppressed thyrotropin (thyroid-stimulating hormone, normal 0.4–3.5 μU/mL) levels. The response of left ventricular function to antithyroid therapy was assessed clinically and by repeat 2-D echocardiogram after thyroid function had returned to normal.

Congestive heart failure was treated with a combination of diuretics, digitalis, angiotensin-converting-enzyme inhibitors and oxygen therapy. Low dose propranolol (30–60 mg daily) was used in 4 patients. Pa-

tients were treated initially with propylthiouracil (450–800 mg daily) or methimazole (30–60 mg daily). Once clinical symptoms improved, euthyroidism was maintained for several months with lower doses of antithyroid drugs, followed by definitive therapy with radioactive iodine or surgery.

Results

The clinical characteristics and results of thyroid function tests on admission are shown in Table 1. Seven patients with hyperthyroidism—four women and three men—with a mean age of 47 ± 4 years (range 28–82 years) are reported. The primary diagnosis on admission in all patients was congestive heart failure. The diagnosis of hyperthyroidism was known before admission in two patients. In the other patients, the diagnosis was suspected on clinical grounds and confirmed by elevated thyroid hormones. Six patients had Graves' disease and one had toxic multinodular goiter. The duration of hyperthyroid symptoms was greater than 1 year in most patients (range 4 months to 6 years). On admission, the mean serum T₄ concentration was 21 ± 1 μg/dL, serum T₃ concentration: 411 ± 77 ng/dL, and T₃RU: 51 ± 5%, and serum thyroid-stimulating hormone was suppressed in all patients.

Therapy of heart failure and hyperthyroidism resulted in a rapid clinical improvement in all patients. Treatment of hyperthyroidism was started on admission in the two patients with known hyperthyroidism or as soon after the diagnosis was confirmed by laboratory studies. Symptoms of heart failure improved within 1–4 days of medical therapy, and hyperthyroidism was controlled within 2 months of antithyroid therapy. A maintenance regimen of antithyroid medication was administered for 2–6 months and was followed by definitive therapy of hyperthyroidism with radioactive iodine or surgery. I¹³¹ was given to six patients, and one patient underwent surgery for coexistent toxic multinodular goiter and primary hyperparathyroidism.

Admission electrocardiograms revealed sinus tachycardia in 4 patients, atrial flutter with 2:1 block in two patients, and atrial fibrillation in one patient. Electro-

cardiographic abnormalities resolved in all patients after treatment of hyperthyroidism. Similarly, radiologic evidence of cardiomegaly and pulmonary congestion improved in most patients during follow-up.

In Table 2, the echocardiographic findings on admission and during follow-up are shown. 2-D echocardiograms performed soon after admission had either bi-ventricular or four chamber dilatation and impaired systolic function. The left atrium diameter was 45 ± 1 mm (mean \pm standard error), the left ventricular diastolic diameter (LVDd) was 55 ± 1 mm, and the left ventricular systolic diameter (LVDs) was 46 ± 1 mm. The mean fractional shortening of the left ventricle ($(LVDd - LVDs)/LVDd$, percent) was $17 \pm 1\%$, with an estimated ejection fraction of $28 \pm 2\%$. Therapy of heart failure and hyperthyroidism resulted in rapid clinical improvement, with resolution of signs and symptoms of heart failure in all patients. Repeat 2-D echocardiogram within 6–12 months of resolution of hyperthyroidism showed resolution of dilated cardiomyopathy with normalization of systolic function in five patients, and improvement from severe to mild ventricular dysfunction in the other two patients (Figure 1). The mean left ventricular ejection fraction improved from $28 \pm 2\%$ to $55 \pm 5\%$ after resolution of hyperthyroidism. Cardiovascular drugs were discontinued in the five patients with normal left ventricular ejection fraction, while the other two patients were continued on a low dose of angiotensin-converting-enzyme inhibitors.

Discussion

The major importance of this study is the documentation of a reversible low-output heart failure due to dilated cardiomyopathy in hyperthyroidism. Admission 2-D echocardiogram showed biventricular or four chamber dilatation and markedly impaired systolic function in all patients. Complete echocardiographic resolution of dilated cardiomyopathy and normaliza-

Table 2. Echocardiographic Measurements Before and After Treatment of Hyperthyroidism

#	Before/After Treatment		
	Left Atrial Size (mm)	Fractional Shortening (%)	Ejection Fraction (%)
1	45/34	12/40	20/68
2	46/35	21/31	36/53
3	39/26	19/38	32/65
4	45/35	15/22	25/38
5	44/33	17/37	29/62
6	42/35	13/24	22/40
7	46/32	20/36	34/61
M \pm SE	$45 \pm 1/33 \pm 1$	$17 \pm 1/33 \pm 3$	$28 \pm 2/55 \pm 5$

The fractional shortening of the left ventricle was calculated by the formula $(LVDd - LVDs)/LVDd \times 100$. Left Ventricular ejection fraction: % fractional shortening $\times 1.7$.¹⁶




Figure 1. Left ventricular performance assessed by the fractional shortening of the left ventricle (%) in seven patients with dilated cardiomyopathy, congestive heart failure, and hyperthyroidism.

tion of left ventricular systolic function was achieved in 5 of 7 patients, and improvement from severe to mild cardiomyopathy was observed in 2 patients (Figure 1). The most important historical finding was the long duration of symptoms of untreated hyperthyroidism (4 months to 6 years). This is in agreement with previous reports.^{2,3} Likoff and Levine³ reported that the duration of thyrotoxicosis in patients with clinical evidence of severe heart failure was 55 months, with moderate heart failure duration was 12 months, and with no failure it was 8 months.

Left ventricular dilatation, congestive heart failure, and death have been reported in necropsy studies in humans with hyperthyroidism¹⁷⁻¹⁸ and in spontaneously hyperthyroid animals,¹⁹ in the absence of other forms of heart disease. Histologically, interstitial and perivascular fibrosis, myocardial hypertrophy, necrosis, and cell edema are reported. Ultrastructurally, increased number, size, and complexity of cardiac mitochondria have been described in animal models of hyperthyroidism.²⁰ In addition, based on recent evidence, excess thyroid hormone causes reversible replacement of the normal myosin isoenzyme concentrations.^{21,22} Although abnormalities in myosin isotypes in myocardial tissues from patients with thyrotoxic cardiomyopathy have not been described, changes in myosin concentration with different contractile properties may prove to be important in the development of thyrotoxic heart disease.

Thyrotoxicosis has long been recognized as a major cause of tachyarrhythmias.^{2,23} Atrial fibrillation is the most common arrhythmia, occurring in 7–21% of pa-

tients with hyperthyroidism. In our patients, the admission electrocardiogram revealed sinus tachycardia in four patients and supraventricular tachyarrhythmia in three patients. Correction of hyperthyroidism was followed by restoration of normal sinus rhythm in all patients. Altered cardiac excitability due to increased beta-adrenergic receptor density in myocardial tissue^{6,23,24} and mechanical distension of the left atrium appears to be important in the development of atrial fibrillation in hyperthyroidism.²⁵

This study does not address the etiology of dilated cardiomyopathy in patients with thyrotoxicosis, and the potential mechanisms underlying this process remain to be elucidated. It is possible that long-lasting tachycardia and high-output state induced by thyroid hormone may eventually lead to dilatation of the left ventricle and a progressive decline in systolic performance. Indeed, Ikram⁵ demonstrated improvement in contractile function after treatment in patients with longstanding thyrotoxicosis and overt heart failure. It is also plausible that thyroid hormone, which is known to translocate to the nucleus of cardiac cells in association with its receptor and to act as a trans-activating factor,^{26,27} may alter the expression of certain cardiac proteins (myosin heavy chain, sarcoplasmic reticulum calcium-activated adenosine triphosphatase) in cardiomyocytes,^{22,28} which may result in contractile dysfunction in patients with hyperthyroidism.

In summary, we suggest, with this report, that in addition to the "high-output congestive heart failure" commonly described in patients with hyperthyroidism, some patients may have a reversible form of dilated cardiomyopathy and "low-output congestive heart failure." Conventional treatment of hyperthyroidism usually results in rapid resolution of the clinical manifestations of heart failure, in partial or complete reversal of cardiomyopathy and in marked improvement of left ventricular systolic function. Assessment of thyroid hormone status in patients with dilated cardiomyopathy might permit the identification of patients who have thyrotoxicosis and are, therefore, likely to have reversible cardiac dysfunction.

Acknowledgment

The authors thank Lewis S. Blevins, MD, and Nelson B. Watts, MD, for critical review of the manuscript.

References

1. Woeber KA. Thyrotoxicosis and the heart. *N Engl J Med.* 1992;327:94-8.
2. Sandler G, Wilson GM. The nature and prognosis of heart disease in thyrotoxicosis: A review of 150 patients treated with ¹³¹I. *Q J Med.* 1959;28:347-69.
3. Likoff WB, Levine SA. Thyrotoxicosis as the sole cause of heart failure. *Am J Med.* 1943;206:425-34.
4. Forfar JC, Muir AL, Sawers SA, Toft AD. Abnormal left ventricular function in hyperthyroidism: Evidence for a possible reversible cardiomyopathy. *N Engl J Med.* 1982;307:1165-70.
5. Ikram H. The nature and prognosis of thyrotoxic heart disease. *Q J Med.* 1985;54:19-28.

6. DeGroot WJ, Leonard JJ. Hyperthyroidism as a high cardiac output state. *Am Heart J.* 1970;79:265-75.
7. Polikar R, Burger AG, Scherrer U, Nicod P. The thyroid and the heart. *Circulation.* 1993;87:1435-41.
8. Klein I. Thyroid hormone and the cardiovascular system. *Am J Med.* 1990;88:631-7.
9. Merillon JP, Passa PH, Chastre J, Wolf A, Gourgon R. Left ventricular function and hyperthyroidism. *Br Heart J.* 1981;46:137-43.
10. Grossman W, Robin NI, Johnson LW, Brooks HL, Selenkow HA, Dexter L. The enhanced myocardial contractility of thyrotoxicosis: Role of beta adrenergic receptor. *Ann Intern Med.* 1971;74:869-74.
11. Biondi B, Fazio S, Carella C, et al. Control of adrenergic overactivity by B-blockade improves the quality of life in patients receiving long term suppressive therapy with levothyroxine. *J Clin Endocrinol Metab.* 1994;78:1028-33.
12. Cavallo A, Casta A, Fawcett HD. Is there is a thyrotoxic cardiomyopathy in children? *J Pediatr.* 1985;107:531-6.
13. Shapiro S, Steier M, Dimich I. Congestive heart failure in neonatal thyrotoxicosis. *Clin Pediatr.* 1975;14:1155-6.
14. McKee PA, Castelli WP, McNamara PM, et al. The natural history of congestive heart failure: The Framingham Study. *N Engl J Med.* 1971;285:1441-6.
15. Folland ED, Parisi AF, Moynihan PF, et al. Assessment of left ventricular ejection fraction and volumes by real-time, two-dimensional echocardiography. *Circulation.* 1979;60:760-6.
16. Quinonez MA, Pickering E, Alexander JA. Percentage of shortening of the echocardiographic left ventricular dimension: Its use in determining ejection fraction and stroke volume. *Chest.* 1978;74:59-65.
17. Shirani J, Barron MM, Pierre-Louis ML, Roberts WC. Congestive heart failure, dilated cardiac ventricles, and sudden death in hyperthyroidism. *Am J Cardiol.* 1993;72:365-8.
18. Saphir O. Myocarditis: A general review, with an analysis of two hundred and forty cases. *Arch Pathol.* 1942;33:88-137.
19. Liu S, Peterson ME, Fox PR. Hypertrophic cardiomyopathy and hyperthyroidism in the cat. *J Am Vet Med Assoc.* 1984;185:52-7.
20. Page E, McCallister LP. Quantitative electron microscopic description of heart muscle cells-application to normal, hypertrophied and thyroxin-stimulated hearts. *Am J Cardiol.* 1973;31:172-81.
21. Chizzonite RA, Everett AW, Prior G, Zak R. Comparison of myosin heavy chains in atria and ventricles from hyperthyroid, hypothyroid, and euthyroid rabbits. *J Biol Chem.* 1984;259:15564-71.
22. Rohrer DK, Hartong R, Dillmann WH. Influence of thyroid hormone and retinoic acid on slow sarcoplasmic reticulum Ca⁺⁺ ATPase and myosin heavy chain alpha gene expression in cardiac myocytes. *J Biol Chem.* 1991;266:8638-46.
23. Olshausen KV, Bischoff S, Hahaly G, et al. Cardiac arrhythmias and heart rate in hyperthyroidism. *Am J Cardiol.* 1989;63:930-3.
24. Golf S, Lovstad R, Hansson V. B-adrenoceptor density and relative number of B-adrenoceptor subtypes in biopsies from human right atrial, left ventricular, and right ventricular myocardium. *Cardiovasc Res.* 1985;19:636-41.
25. Iwasaki T, Naka M, Hiramatsu K, et al. Echocardiographic studies on the relationship between atrial fibrillation and atrial enlargement in patients with hyperthyroidism of Graves' disease. *Cardiology.* 1989;76:10-7.
26. Izumo S, Nadal-Ginard B, Mahdavi V. All members of the MHC gene family respond to thyroid hormone in a highly tissue-specific manner. *Science.* 1986;231:597-600.
27. Izumo S, Lompre AM, Matsuoka R, et al. Myosin heavy chain messenger RNA and protein isoform transitions during cardiac hypertrophy. Interaction between hemodynamic and thyroid hormone-induced signals. *J Clin Invest.* 1987;79:970-7.
28. Klein I, Ojamaa K. Cardiovascular manifestations of endocrine disease. *J Clin Endocrinol Metab.* 1992;75:339-42.

(b) (6)

(b) (6)

Client ID:	(b) (6)	Patient ID:	(b) (6)
Client Name:	(b) (6)	Name:	(b) (6)
Spouse/Other:	(b) (6)	Breed:	Retriever, Labrador Mix
Address:	(b) (6)	Sex:	Spayed Female
	(b) (6)	Color:	Brindle
Telephone:	(b) (6)	Age:	6 Yrs. 9 Mos.
		DOB:	(b) (6)

Referring Veterinarian:	(b) (6)	Dr. Jennifer Jones
Practice:	(b) (6)	FDA
Phone:	(b) (6)	
FAX:	(b) (6)	

Dear Dr. Jones,

Please find attached the normal results of the iron panel on (b) (6). Please let me know if you have any other recommendations in (b) (6).

I would love to discuss our findings on dilated cardiomyopathy and dietary relationships in our clinic over the past year. In particular, 75% of our DCM cases in 2017 for which we have adequate dietary histories were on grain free diets. We have started to do a survey of other cardiac patients during the same time period to see if we can get an idea of what percentage of our referral population were feeding these type of diets so we can see if this is truly significant. As we have been looking harder at these cases, we have found some other interesting things. For instance, two patients that were found to be taurine deficient were being fed Zignature diets. One was on the kangaroo variety and the other the pork variety. There continues to be a lot of discussion about diets and dilated cardiomyopathy on our list serve, but I can see it degenerating into I diagnosed DCM and the dog is on this diet so therefore this is a problem, which may or may not be true evidence to support a causative role of the diet.

I am not sure what information the FDA would want on all this at this point. We are certainly concerned that there may be a wider concern than simply the kangaroo and lentil diets that we first identified as a potential problem. Feel free to give me a call if you wish to discuss this further, or if you can suggest a direction for us to go with this inquiry. My clinic number is (b) (6) and I am in the clinics Monday through Thursday, or my cell phone is (b) (6).

Sincerely,

(b) (6)

		800.218-sub 1	800.218-sub 2	800.218-sub 6	
		Case Sample	Storebought	Case sample	Label
		California Naturals Kangaroo & Lentil	California Naturals Kangaroo & Lentil	California Naturals Kangaroo & Lentil	Product Nutrient Analysis (website label)
(b) (4)	Ca	1.30%	1%	0.93%	0.83%
	Mg	0.13%	0.14%	0.15%	0.17%
	P	0.74%	0.67%	0.68%	0.71%
	Fe	30 mg/kg	30 mg/kg	31 mg/kg	305 mg/kg
	Co	0.12 mg/kg	0.14 mg/kg	.14 mg/kg	n/a
	Cu	21 mg/kg	19 mg/kg	16 mg/kg	13.61 mg/kg
	Zn	240 mg/kg	280 mg/kg	200 mg/kg	193.37 mg/kg
	Se	0.7 mg/kg	0.65 mg/kg	.68 mg/kg	0.08 mg/kg
	Ca:P	1.76:1	1.49:1	1.37:1	
	Cu:Zn	0.09:1	0.07:1	0.08:1	
(b) (4)	Tau	~0.26%	1.06 mg/g = ~0.11%	1.22 mg/g = ~0.12%	
	Cystine	2.32 mg/g = ~0.23%	2.31 mg/g = ~0.23%	2.5 mg/g = ~0.25%	
	Met	5.78 mg/g = ~0.58%	5.53 mg/g = ~0.55%	7.78 mg/g = ~0.78%	0.61%
	Met-Cys	~0.81%	~0.78%	~1.03%	0.97%
MSU	Iodine	not tested	4.04 ug/g (ppm)	1.87 ug/g (ppm)	

AAFCO	
AAFCO-Adult Maint	Issues
0.5 to 2.5%	none
0.06%	none
0.4 to 1.6 %	none
40 mg/kg	below AAFCO & Label
25 mg/kg-chicks/rats/sheep max	unlikely
7.3 mg/kg	none
80 mg/kg	none
0.35 to 2 mg/kg	label should be higher
1:1 to 2:1	none
0.09:1-not AAFCO	none
0.1% in Cats	
n/a	
0.33%	none
0.65%	none
1 ppm (min) to 11 ppm (max)	none

<http://www.californianaturalpet.com/products/1741>

to align w/ AAFCO maintenance claim

Report Details - EON-266814								
ICSR:	1053335							
Type Of Submission:	Initial							
Report Version:	FPSR.FDA.PETF.V.V1							
Type Of Report:	Both							
Reporting Type:	Voluntary							
Report Submission Date:	2016-06-06 11:15:17 EDT							
Reported Problem:	Problem Description: Presented 5/8/2016 for lethargy; on physical exam the patient was dyspneic and pleural effusion identified on cursory ultrasound and DV thoracic x-rays - given lasix and placed in oxygen. Transferred to cardiology service; evaluation including echocardiogram on 5/9/16 revealed dilated cardiomyopathy, moderate left atrial enlargement, pleural effusion and azotemia. Plasma taurine was submitted to University of Wisconsin. Lab results received 5/15/16 - plasma taurine 24nmol/ml (ref range 60-120, critical level <40). Recheck echocardiogram on 5/15/16 revealed same changes as prior and a thrombus in her left ventricle. Medications/supplements included taurine 250mg PO BID, Mirtazepine 15mg tablets (Give 1/4 tablet PO every 3d PRN), Furosemide 12.5 mg tablets (Give 1/4 tablet PO SID), Pimobendan 1.5mg tiny tabs (Give 1 tablet PO BID). Patient presented on (b) (6) for partial aortic thromboembolism and owner's elected euthanasia. Review of patient's diet history revealed that all 5 cats in household had been fed Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry for approximately 3 years. The 4 remaining cats were tested for taurine deficiency and 2/4 had whole blood levels indicating deficiency: 5/21/2016 - Whole Blood Taurine submitted at the University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016 (b) (6): 9yr male neutered domestic long hair: 196 nmol/ml (b) (6): 8y female spayed domestic short hair: 368 nmol/ml (b) (6): 9yr male neutered domestic long hair: 124 nmol/ml (b) (6): 9yr male neutered domestic long hair: 536 nmol/ml							
	Date Problem Started: 05/08/2016							
	Concurrent Medical Problem: No							
	Outcome to Date: Died Euthanized							
	Date of Death: (b) (6)							
Product Information:	Product Name: Merrick Purrfect Bistro Grain Free Real Chicken Recipe							
	Product Type: Pet Food							
	Lot Number: <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td>Lot Number:</td> <td>16025 DL1 38310 14131</td> </tr> <tr> <td>Expiration Date:</td> <td>07/26/2017</td> </tr> </table>	Lot Number:	16025 DL1 38310 14131	Expiration Date:	07/26/2017			
	Lot Number:	16025 DL1 38310 14131						
	Expiration Date:	07/26/2017						
	UPC: 2280838310							
	Package Type: BAG							
	Package Size: 5.4 kilogram							
	Purchase Date: 05/01/2016							
	Number Purchased: 1							
	Possess Unopened Product: No							
	Possess Opened Product: Yes							
	Storage Conditions: Stored in bag indoors							
Product Use Information: <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td>Description:</td> <td>Fed to cats in bowl</td> </tr> <tr> <td>Last Exposure Date:</td> <td>05/08/2016</td> </tr> <tr> <td>Time Interval between Product Use and Adverse Event:</td> <td>3 Months</td> </tr> <tr> <td>Product Use Stopped After the Onset of the</td> <td>Yes</td> </tr> </table>	Description:	Fed to cats in bowl	Last Exposure Date:	05/08/2016	Time Interval between Product Use and Adverse Event:	3 Months	Product Use Stopped After the Onset of the	Yes
Description:	Fed to cats in bowl							
Last Exposure Date:	05/08/2016							
Time Interval between Product Use and Adverse Event:	3 Months							
Product Use Stopped After the Onset of the	Yes							

FDA-CVM-FOIA-2019-1704-000010

	Adverse Event:			
	Adverse Event Abate After Product Stop:	No		
	Product Use Started Again:	No		
	Perceived Relatedness to Adverse Event:	Definitely related		
	Other Foods or Products Given to the Animal During This Time Period:	Yes		
Manufacturer/Distributor Information:	Name:	Merrick Pet Care, Inc		
	Type(s):	Manufacturer		
	Address:	P.O. Box 9800 Amarillo Texas 79105 United States		
	Contact:	Phone:	18006647387	
		Web Address:	www.merrickpetcare.com	
	Possess One or More Labels from This Product:	Yes		
Purchase Location Information:	Name:	(b) (6)		
	Address:	(b) (6) United States		
Animal Information:	Name:	(b) (6)		
	Type Of Species:	Cat		
	Type Of Breed:	Mixed (Cat)		
	Gender:	Female		
	Reproductive Status:	Neutered		
	Weight:	5.3 Kilogram		
	Age:	12 Years		
	Assessment of Prior Health:	Good		
	Number of Animals Given the Product:	5		
	Number of Animals Reacted:	3		
	Owner Information:	Owner Information provided:	Yes	
		Contact:	Name:	(b) (6)
			Phone:	(b) (6)
	Address:	(b) (6) United States		
Healthcare Professional	Practice Name:	(b) (6) OIA-2019-1704-000011		

	Information:	Contact:	Name: (b) (6)	
			Phone: (b) (6)	
			Email: (b) (6)	
		Address:	(b) (6) United States	
Sender Information:	Name:	(b) (6)		
	Address:	(b) (6) United States		
	Contact:	Phone:	(b) (6)	
		Email:	(b) (6)	
	Permission To Contact Sender:	Yes		
	Preferred Method Of Contact:	Email		
Reported to Other Parties:	Manufacturer			
Additional Documents:				

Report Details - EON-266821					
ICSR:	1053339				
Type Of Submission:	Initial				
Report Version:	FPSR.FDA.PETF.V.V1				
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)				
Reporting Type:	Voluntary				
Report Submission Date:	2016-06-06 11:44:41 EDT				
Reported Problem:	Problem Description: Another household cat diagnosed with dilated cardiomyopathy and taurine deficiency - separate report filed (FDA ICSR ID 1053335). Euthanized or (b) (6) due to aortic thromboembolism. Review of the patient's diet history revealed that all 5 cats in household had been fed Merrick Purrfect Bistro Grain Free Real Chicken Recipe Feline dry for approximately 3 years. Remaining 4 cats in household tested for taurine deficiency - whole blood samples submitted to University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results received on 5/27/16 - (b) (6) 196nmol/ml - started on taurine supplementation 250mg PO BID for 2-3 weeks. Diet was changed at the time of other cat's diagnosis (5/15/15). Patient also diagnosed with hyperthyroidism on same day as blood submitted for taurine testing - history of weight loss. An echo was not performed on this patient therefore it is unknown if he had evidence of DCM.				
	Date Problem Started: 05/27/2016				
	Concurrent Medical Problem: No				
	Outcome to Date: Not Applicable				
Product Information:	Product Name: Merrick Purrfect Bistro Grain Free Real Chicken Recipe				
	Product Type: Pet Food				
	Lot Number: <table border="1" style="display: inline-table; vertical-align: top;"> <tr> <td>Lot Number:</td> <td>16025 DL1 38310 14131</td> </tr> <tr> <td>Expiration Date:</td> <td>07/26/2017</td> </tr> </table>	Lot Number:	16025 DL1 38310 14131	Expiration Date:	07/26/2017
	Lot Number:	16025 DL1 38310 14131			
	Expiration Date:	07/26/2017			
	UPC: 2280838310				
	Package Type: BAG				
	Package Size: 5.4 kilogram				
	Number Purchased: 1				
	Possess Opened Product: Yes				
	Storage Conditions: stored in bag indoors				
	Product Use Information:	Description: fed to cats in bowl			
		Last Exposure Date: 05/15/2016			
Product Use Stopped After the Onset of the Adverse Event: Yes					
Perceived Relatedness to Adverse Event: Definitely related					
Other Foods or Products Given to the Animal During This Time Period: No					
Manufacturer/Distributor Information:	Name: Merrick Pet Care, Inc				
	Type(s): Manufacturer				
	Address: P.O. Box 9800 Amarillo Texas 79105 United States				
FDA-CVM-FOIA-2019-1704-000013					

		Contact:	Phone: 18006647387	
			Web Address: www.merrickpetcare.com	
		Possess One or More Labels from This Product:	Yes	
	Purchase Location Information:	Name:	(b) (6)	
		Address:	(b) (6) United States	
Animal Information:	Name:	(b) (6)		
	Type Of Species:	Cat		
	Type Of Breed:	Mixed (Cat)		
	Gender:	Male		
	Reproductive Status:	Neutered		
	Weight:	4.4 Kilogram		
	Age:	9 Years		
	Assessment of Prior Health:	Fair		
	Number of Animals Given the Product:	5		
	Number of Animals Reacted:	3		
	Owner Information:	Owner Information provided:	Yes	
		Contact:	Name:	(b) (6)
			Phone:	(b) (6)
Address:	(b) (6) United States			
Healthcare Professional Information:	Practice Name:	(b) (6)		
	Contact:	Name:	(b) (6)	
		Phone:	(b) (6)	
		Email:	(b) (6)	
Address:	(b) (6) United States			
Sender Information:	Name:	(b) (6)		
	Address:	(b) (6)		
		United States		
	Contact:	Phone:	(b) (6)	
		Email:	(b) (6)	
Permission To Contact Sender:	Yes			

FDA-CVM-FOIA-2019-1704-000014

	Preferred Method Of Contact:	Email
	Reported to Other Parties:	Manufacturer

Additional Documents:

Report Details - EON-266827											
ICSR:	1053345										
Type Of Submission:	Initial										
Report Version:	FPSR.FDA.PETF.V.V1										
Type Of Report:	Both										
Reporting Type:	Voluntary										
Report Submission Date:	2016-06-06 12:11:20 EDT										
Reported Problem:	Problem Description: Another household cat diagnosed with dilated cardiomyopathy and taurine deficiency - separate report filed (FDA ICSR ID 1053335). Euthanized or (b) (6) due to aortic thromboembolism. Review of the patient's diet history revealed that all 5 cats in household had been fed Merrick Purrfect Bistro Grain Free Real Chicken Recipe Feline dry for approximately 3 years. Remaining 4 cats in household tested for taurine deficiency - whole blood samples submitted to University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results received on 5/27/16 - (b) (6) 124nmol/ml - started on taurine supplementation 250mg PO BID for 2-3 weeks. Diet was changed at the time of other cat's diagnosis (5/15/15). An echo was not performed on this patient therefore it is unknown if he had evidence of DCM.										
	Date Problem Started: 05/27/2016										
	Concurrent Medical Problem: Yes										
	Pre Existing Conditions: patient is obese										
	Outcome to Date: Not Applicable										
Product Information:	Product Name: Merrick Purrfect Bistro Grain Free Real Chicken Recipe										
	Product Type: Pet Food										
	Lot Number: <table border="1" style="display: inline-table; vertical-align: top;"> <tr> <td>Lot Number:</td> <td>16025 DL1 38310 14131</td> </tr> <tr> <td>Expiration Date:</td> <td>07/26/2017</td> </tr> </table>	Lot Number:	16025 DL1 38310 14131	Expiration Date:	07/26/2017						
	Lot Number:	16025 DL1 38310 14131									
	Expiration Date:	07/26/2017									
	UPC: 2280838310										
	Package Type: BAG										
	Package Size: 5.4 kilogram										
	Purchase Date: 05/01/2016										
	Number Purchased: 1										
	Possess Unopened Product: No										
	Possess Opened Product: Yes										
	Storage Conditions: stored in bag indoors										
	Product Use Information: <table border="1" style="display: inline-table; vertical-align: top;"> <tr> <td>Description:</td> <td>fed to cats in bowl</td> </tr> <tr> <td>Last Exposure Date:</td> <td>05/15/2016</td> </tr> <tr> <td>Product Use Stopped After the Onset of the Adverse Event:</td> <td>Yes</td> </tr> <tr> <td>Perceived Relatedness to Adverse Event:</td> <td>Definitely related</td> </tr> <tr> <td>Other Foods or Products Given to the Animal During This Time Period:</td> <td>No</td> </tr> </table>	Description:	fed to cats in bowl	Last Exposure Date:	05/15/2016	Product Use Stopped After the Onset of the Adverse Event:	Yes	Perceived Relatedness to Adverse Event:	Definitely related	Other Foods or Products Given to the Animal During This Time Period:	No
	Description:	fed to cats in bowl									
Last Exposure Date:	05/15/2016										
Product Use Stopped After the Onset of the Adverse Event:	Yes										
Perceived Relatedness to Adverse Event:	Definitely related										
Other Foods or Products Given to the Animal During This Time Period:	No										
Manufacturer/Distributor Information: <table border="1" style="display: inline-table; vertical-align: top;"> <tr> <td>Name:</td> <td>Merrick Pet Care, Inc</td> </tr> <tr> <td>Type(s):</td> <td>Manufacturer</td> </tr> <tr> <td>Address:</td> <td>P.O. Box 9800 Amarillo</td> </tr> </table>	Name:	Merrick Pet Care, Inc	Type(s):	Manufacturer	Address:	P.O. Box 9800 Amarillo					
Name:	Merrick Pet Care, Inc										
Type(s):	Manufacturer										
Address:	P.O. Box 9800 Amarillo										
FDA-CVM-FOIA-2019-1704-000016											

			Texas 79105 United States
		Contact:	Phone: 18006647387 Web Address: www.merrickpetcare.com
		Possess One or More Labels from This Product:	Yes
	Purchase Location Information:	Name:	(b) (6)
		Address:	(b) (6) United States
Animal Information:	Name:	(b) (6)	
	Type Of Species:	Cat	
	Type Of Breed:	Mixed (Cat)	
	Gender:	Male	
	Reproductive Status:	Neutered	
	Weight:	9.5 Kilogram	
	Age:	9 Years	
	Assessment of Prior Health:	Good	
	Number of Animals Given the Product:	5	
	Number of Animals Reacted:	3	
	Owner Information:	Owner Information provided:	Yes
		Contact:	Name: (b) (6) Phone: (b) (6)
		Address:	(b) (6) United States
	Healthcare Professional Information:	Practice Name:	(b) (6)
		Contact:	Name: (b) (6) Phone: (b) (6) Email: (b) (6)
		Address:	(b) (6) United States
Sender Information:	Name:	(b) (6)	
	Address:	(b) (6) United States	
	Contact:	Phone:	(b) (6)

FDA-CVM-FOIA-2019-1704-000017

	Email:	(b) (6)
	Permission To Contact Sender:	Yes
	Preferred Method Of Contact:	Email

Additional Documents:

**Follow-up Case Information Uniform Data Entry Form
Vet-LIRN**

Date (mm/dd/yy) Jun 9, 2016
EON/CC Number: 266,814

PATIENT INFORMATION

Pet Name (b) (6)
 Dog Cat
Breed DSH
Age in years (if < 6 months, put 0.5) 12
Gender: M MN F FS

This form serves as a Uniform Data Entry Form to capture additional case specific information not clear from the Consumer Complaint or Medical Records in a standardized manner. Because each follow-up interview made with owners features questions tailored specifically to the case, each box of information contained in this Uniform Data Entry Form may not be completed.

HISTORY-Additional Comments from Owner

Owner's Description of What Happened: lethargic, difficulty moving, not herself

Any Health Problems Prior to the Event (e.g. allergies, surgeries): (b) (6): none; none in the past; all house cats

Sensitive GI tract (e.g. stomach upset when switching foods, eats a lot of grass) Yes

Changes to the pet's diet prior to illness Yes

Date Diet Change:

CLINICAL INFORMATION--Additional Comments from Owner on What Happened

Appetite Increased Decreased

Water Consumption Increased Decreased

Vomiting Yes

Urination Increased Decreased

Diarrhea Yes

Lethargy Yes

Duration of Diarrhea (days)

Other:

Blood in Feces Fresh,Red
 Coffee Ground
 Black,Tarry

MEDICATIONS-Taken Prior to the Event and Mentioned by Owner

List medications mentioned by owner (e.g. NSAIDs, steroids, heartworm/flea prevention, antibiotics, etc.)

List probiotics, vitamins, or supplements mentioned by owner:

**Follow-up Case Information Uniform Data Entry Form
Vet-LIRN**

EON/CC Number: 266,814

Owner: (b) (6)

Pet's Name: (b) (6)

DIET-Any other foods the owner mentions were given to the animal during this period. (check all that apply)

Commercial Dry Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name: Merrick Grain Free Bistro Chicken-started in January of 2014, print out of all pet food purchases thru Jan 2014 --> on the Chicken pretty much the entire time; owner has a listing of all purchases from pet

Commercial Wet-Canned Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name: Fancy Feast Wet food given to (b) (6) to stimulate her appetite after her illness onset, but she only licked the gravy. So for ~1 week before the other 4 house cats were tested for taurine, only (b) (6)

Commercial Wet-Pouch Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name:

Commercial-Raw Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name:

Homemade-Raw Product Use as Part of Diet: Primary Secondary Occasional

Describe Product Type:

Homemade-Cooked Product Use as Part of Diet: Primary Secondary Occasional

Describe Product Type:

Table Scraps/Human Food (as an occasional contribution to diet) Describe Product Type(s): not in past 5 years

Pet Treat Products Product Use as Part of Diet: Primary Secondary Occasional

Commercial Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Rawhides or Pig Ears Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Marrow Bones Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Chicken Jerky Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Duck Jerky Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Sweet Potato Jerky or Treats Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

**Follow-up Case Information Uniform Data Entry Form
Vet-LIRN**

EON/CC Number:

Owner:

Pet's Name:

DIET-continued-Any other foods the owner mentions were given to the animal during this period. (check all that apply)

Other Treats Product Label Name/Lot: Date first fed
How Product Administered Date last fed

ENVIRONMENTAL EXPOSURES-Environmental Exposures Mentioned by the Owner Potentially Affecting the Animal's Overall State of Health Prior to the Event . (check all that apply)

- Indoor Outdoor Indoor & Outdoor Carrion Rodents Grapes or Raisins Nuts
- Plants Trash Hunt Pet Shows Sporting Events Pet Recreation Facilities
- Livestock Poultry Reptiles Pet Birds Small Mammals Untreated Surface Water
- Anti-freeze Mushrooms Heavy Metals Ticks Urban Suburban Rural

Comments:

HOUSEHOLD-Signalment of Additional Animals Given the Product mentioned by the owner.

Animal 1 Reacted

Animal 2 Reacted

Animal 3 Reacted

Comments

Food and Drug Administration Office of Regulatory Affairs

Summary Report

For Sample Number: 958500

TD Sample Number:

Import Sample Number

This is an accurate reproduction of the original electronic record as of 08/24/2016

Sample Class: Normal Everyday Sample **Sample Origin:** Domestic **Sample Basis:** Surveillance
Sample Flag: **Sample Type:** Official **Collecting District:** NWJ-DO
Home District: **Orig C/R and Records To:** NWJ-DO **Collection PACs:** 71R801

Product Name: Poultry Prod Pet Cat Food; Not Elsewhere Classified (NEC); Packaged Food (Not Commercially Sterile)

Product Description: See Remarks Section.

Collection Reason: Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

Lab: SRL **Split Num:**0 **Date Received:** 06/29/2016 **Date Out of Lab:** 08/04/2016
District **District Conclusion** **District**
Conclusion: No Action Indicated (NAI) **Made By:** Tweedley, Karen P 08/12/2016
Disposition **Disposition** **Disposition**
Reason: NAI By Examining District **Authorized By:** Tweedley, Karen P **Authorized Date:** 08/12/2016

Performing Org	PAC	LID	PAF	Compliance No	Lab Class-Description	Laboratory Status
ACNA-N	71R801		NAR		1 - In Compliance	Completed

Lab Conclusion

Sample Narrative - Method: AccQTag AAA(Waters) Analysis - Taurine
Amt Found - 0.187% (dry matter basis)
Meets AAFCO minimum requirement of 0.10%

Sample Narrative - Method: Instrument Manual (Denver Instrumentst IR60)/ AOAC 930.15 Analysis - Moisture
Amt Found - 2.20%
Amt Declared - 11.00% max

Lab Conclusion Date **Lab Conclusion Made By**
08/04/2016 Hawes,Brian M

Food and Drug Administration Office of Regulatory Affairs

Summary Report

For Sample Number: 958501

TD Sample Number:

Import Sample Number

This is an accurate reproduction of the original electronic record as of 08/24/2016

Sample Class: Normal Everyday Sample **Sample Origin:** Domestic **Sample Basis:** Surveillance
Sample Flag: **Sample Type:** Official **Collecting District:** NWJ-DO
Home District: **Orig C/R and Records To:** NWJ-DO **Collection PACs:** 71R801

Product Name: Poultry Prod Pet Cat Food; Not Elsewhere Classified (NEC); Packaged Food (Not Commercially Sterile)

Product Description: See Remarks

Collection Reason: Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

Lab: SRL **Split Num:**0 **Date Received:** 06/29/2016 **Date Out of Lab:** 08/04/2016
District **District Conclusion** **District**
Conclusion: No Action Indicated (NAI) **Made By:** Tweedley, Karen P 08/12/2016
Disposition **Disposition** **Disposition**
Reason: NAI By Examining District **Authorized By:** Tweedley, Karen P **Authorized Date:** 08/12/2016

Performing Org	PAC	LID	PAF	Compliance No	Lab Class-Description	Laboratory Status
ACNA-N	71R801		NAR		1 - In Compliance	Completed

Lab Conclusion

Sample Narrative - Method: AccQTag AAA Analysis - Taurine
Amt Found - 0.156% (dry matter basis)
Meets AAFCO minimum requirement of 0.10%

Sample Narrative - Method: Instrument Manual (Denver Instrumentst IR60)/ AOAC 930.15 Analysis - Moisture
Amt Found - 1.99%
Amt Declared - 11.00% max

Lab Conclusion Date **Lab Conclusion Made By**

08/04/2016 Hawes,Brian M

Food and Drug Administration Office of Regulatory Affairs

Summary Report

For Sample Number: 958504

TD Sample Number:

Import Sample Number

This is an accurate reproduction of the original electronic record as of 08/24/2016

Sample Class: Normal Everyday Sample **Sample Origin:** Domestic **Sample Basis:** Surveillance
Sample Flag: **Sample Type:** Official **Collecting District:** NWJ-DO
Home District: **Orig C/R and Records To:** NWJ-DO **Collection PACs:** 71R801

Product Name: Poultry Prod Pet Cat Food; Not Elsewhere Classified (NEC); Packaged Food (Not Commercially Sterile)

Product Description: See Remarks Section.

Collection Reason: Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

Lab: SRL **Split Num:**0 **Date Received:** 06/29/2016 **Date Out of Lab:** 08/04/2016
District **District Conclusion** **District**
Conclusion: No Action Indicated (NAI) **Made By:** Ciaccia, Andrew 08/17/2016
Disposition **Disposition** **Disposition**
Reason: NAI By Home District **Authorized By:** Ciaccia, Andrew **Authorized Date:** 08/17/2016

Performing Org	PAC	LID	PAF	Compliance No	Lab Class-Description	Laboratory Status
ACNA-N	71R801		NAR		1 - In Compliance	Completed

Lab Conclusion

Sample Narrative - Method: AccQTag AAA Waters Analysis - Taurine
Amt Found - 0.176% (dry matter basis)
Meets AAFCO minimum requirement of 0.10%

Sample Narrative - Method: Instrument Manual (Denver Instrumentst IR60)/AOAC 930.15 Analysis - Moisture
Amt Found - 2.79%
Amt Declared - 11.00% max

Lab Conclusion Date **Lab Conclusion Made By**
08/04/2016 Hawes,Brian M

OR Final Report

A. Study Identification:

Study Number: 800.180

Study Director: Renate Reimschuessel, VMD PhD

Division: Vet-LIRN

Division Code: HFV – 500

Other Investigators:

Jennifer Jones	Vet-LIRN
Sarah Nemser	Vet-LIRN
Olgica Ceric	Vet-LIRN
Jake Guag	Vet-LIRN
David Rostein	OS&C

B. Descriptive Title of Study:

Investigation into the death of one cat and low blood taurine levels of two other cats after consumer Merrick Purrfect Bistro Grain Free Chicken Recipe cat food

Name and Address of Testing Facilities:

Mod II
Vet-LIRN and DAFM
Center for Veterinary Medicine
Office of Research
8401 Muirkirk Road
Laurel, MD 20708

C. Starting and Completion Date: Starting Date: 6/7/2016
Ending Date: 6/13/2016
Final Report Submitted Date: 9/2/2016

Case Summary

Complaint: June 7, 2016 Vet-LIRN received consumer complaints, EON-226814, EON-226821, and EON-226827, reporting the death of one cat and low blood taurine levels of 2 others after consuming Merrick Purrfect Bistro Grain Free Real Chicken Recipe food for cats.

Signalment: (b) (6)-12 year old female spayed (FS) domestic shorthair (DSH); (b) (6)-9 year old male castrated (MC) domestic longhair (DLH); (b) (6)-8 year old FS DSH; (b) (6)-9 year old MC DLH; (b) (6)-9 year old MC DLH

Signs: congestive heart failure, dilated cardiomyopathy (DCM), an aortic thromboembolism, low plasma taurine levels, weight loss, hair loss

Medical Records: Medical records were received and reviewed.

(b) (6): 12 yo FS DSH-euthanized

Presenting complaint 5/8/2016: lethargy starting (b) (6), indoor only → recheck (b) (6) b/c very weak, PD, difficulty walking, inappetant → (b) (6), feed Merrick Purrfect Bistro Grain Free Real chicken recipe ~3 yr, prior to this fed Dick Van Pattons Indoor Dry-Chicken & Salmon; vet called Merrick → bag purchased 2 weeks prior per owner. Vet doesn't think a single bag/lot issue because takes several months to develop → urinating & defecating outside litterbox, soft stool, no change to appetite, ambulating well → (b) (6) dyspnea beginning (b) (6), recheck → by (b) (6) weak hindlimbs → by (b) (6) 1 hindlimb worse than other → by (b) (6), recheck b/c dragging right hindlimb → euthanized, vet spoke w/ Merrick QA team and the taurine in that lot # was sufficient

PE 5/8: BCS 7/9, T 93, HR 150, RR 60, muffled heart sounds, inc respiratory effort, dull lung sounds

-5.9: heart sounds slightly muffled, RR 30 w/ slight effort

-5.10: cat PD, inappetant, RR ~28

-5.25: tachypneic, mild inc resp effort, RR 48, faint referred upper airway noise

(b) (6) tachypneic, moderate dyspnea, RR 48-60, non-weight bearing Right hindlimb, Right HL: no femoral pulse, cold paw pad; T 94.8, laterally recumbent

Labs: 5.9.2016 **BGA:** Hct 40, Na 146.3 (146.2-156.2), K 4.99 (3.41-4.71), Cl 107.8 (117-125.3), Ca 1.17 (1.16-1.35), Mg 1.08 (0.33-0.49), Glu 156 (72-132), Lac 9.7 (0.7-1.9), BUN 67 (20-33), Ct 5.3 (1.1-3.5)

-5.9 Renal panel: ALP 11 (14-111), ALT 140 (12-130), BUN 74 (16-36), Cl 100 (112-129), Na 138 (150-165), Ct 1.4 (0.8-2.4)

-5.15 BGA: Hct 43, Na 145.3, K 3.33, Cl 104.5, BUN 61, Ct 3.1

-5.25 BGA: Hct 34, Na 147.3, K 6.65, Cl 115.1, BUN 31, Ct 1.3

5.9 **PCV/TS:** 42/6.8

-5.15 PCV/TS: 39/7.8

-5.25 PCV/TS: 32/6

5.9 **Plasma taurine:** 24 (60-120, critical <40)

5/8 Cursory US: mild-moderate pleural effusion, R>L

5/8 Rads: cardiac silhouette difficult to visualize, pleural effusion, moderate inc opacity area caudal to left cranial lung lobe

5/9 Echocardiogram: mod LA enlargement, LV enlarged, mild RA & RV enlargement, trivial MR &

TR, dec aortic & pulmonary flow, moderate volume Pleural effusion → DCM
 -5.15: small volume PE, no pericardial effusion, large mass in LV-thrombus
 -5.25: thrombus unchanged
 (b) (6) small vol PE, large LV thrombus

Treatments: Lasix, thoracocentesis: 5/8 (25mL), 5/9 (120 mL), 5/25 (160 mL), O2, pimobendan, taurine, mirtazapine, MaxCal, buprenex, telazol, acepromazine, butorphanol, beuthanasia

Name	Clinical Signs	Lab work Abnormalities	Taurine Level (300-600)**	Outcome	Comments
(b) (6)	Chronic weight & hair loss, polyphagia, BCS 1.5-2/5, alopecia	ALP 174 (hi) ALT 243 (hi) TT4 20.2 (hi)	196	Supplement taurine, treat hyperthyroidism	Lives above garage
(b) (6)	Copious otic debris-AU, resorptive tooth lesion (UR PM3)	TP 9.1 (hi) Glob 6.5 (hi) Gamma Globulins 3.6 (hi)	368	Treat ears; Potential Diagnosis of: Lymphoma, myeloma, or chronic inflammatory disease	Lives above garage
(b) (6)	Moderate tartar, some matted hair	nsf	124	Supplement taurine	Lives above garage
(b) (6)	Significant gingivitis, heavy tartar PM3's, loose canine tooth (UR)	nsf	536	None	Lives in house because (b) (6) is aggressive toward him

**Taurine Level >200 associated with No Risk of DCM.

nsf = No significant findings

Owner Interview: An owner interview was completed in order to understand the feeding history and the impact of any potential environmental exposures. The owner also sent a copy of her cat food purchase history. According to the document, Merrick Grain Free Bistro Chicken Adult cat food was most frequently purchased.

Presenting complaint: (b) (6) had lethargy, difficulty moving, not acting like herself. Indoor cats

Prior MHx: none significant for all cats (5) in household

Diet: Merrick Grain Free Bistro Chicken-started in January of 2014, --> on the Chicken variety pretty much the entire time; owner has a listing of all purchases from pet store and she will send us a copy tomorrow; fed prior to the Merrick food was Dick Van Pattens Natural Balance-2 indoor cat formulas-owner thinks the chicken & salmon type but not 100% sure, she can find out if needed: owner mentions the consistency changed when the company was purchased by Del Monte which prompted her to switch foods to Merrick; After the illness onset for (b) (6), Fancy Feast Wet food fed to stimulate her appetite, but she only licked the gravy. So for ~1 week before the other 4 house cats were tested for taurine, only (b) (6) got the Fancy Feast leftovers (solid chunks) not consumed by (b) (6)

Response: Vet-LIRN collected medical records for review and completed an owner interview for more information on the feeding history and potential environmental exposures. The Office of Regulatory Affairs (ORA) local district office sampled and tested product.

Results: Three regulatory samples were collected. The measured taurine content for each was 0.187%, 0.156%, and 0.176%, on a dry matter basis.

Conclusion: The medical records showed three of five cats in a household had low blood taurine levels. One cat, (b) (6), had low blood taurine and dilated cardiomyopathy (DCM). DCM can be caused by low dietary and blood taurine levels. She was euthanized due to a thromboembolism, a complication arising from DCM. (b) (6) lived above the garage with three other cats: (b) (6), (b) (6) and (b) (6). (b) (6) also had low blood taurine levels (<200 nMol/mL), which increase the risk for developing DCM. Neither cat had an echocardiogram, and it is unclear if they had any evidence of DCM. One cat, (b) (6), who lived separately from the other cats, had the highest blood taurine level. (b) (6) received a supplemental wet food for approximately one week prior to the blood taurine level check. The dietary interview and purchase history indicate the chicken variety cat food was fed most frequently. Over time, if the food were deficient in taurine, the cats could develop low blood taurine and thus DCM. If all cats ate the same diet deficient in taurine, you would expect all cats to have low blood taurine levels. (b) (6) normal blood taurine level could be due to individual variation. (b) (6) had the highest blood taurine level of all the cats. His blood taurine level prior to eating the supplemental wet food is unknown. However, if he were taurine deficient, it is possible for the supplemental wet food to improve his blood taurine level. This could explain why (b) (6) had the highest blood taurine level of all the cats.

According to AAFCO, cat food must contain a minimum of 0.10% taurine on a dry matter basis. All three cat food samples tested by ORA are in compliance. Because taurine deficiency develops over time, the cats would have had to consume taurine deficient product over a period of months to years. It is unknown if prior lots or varieties of the food contained adequate taurine levels. It is unlikely this lot of food caused the cats' taurine deficiency.

Supplemental Information

01-800.180-EON-multiple	(b) (6) CC: Consumer complaints
02-800.180-EON-multiple	(b) (6) MedRec: Medical records
03-800.180-EON-multiple	(b) (6) Interview: Owner interview
04-800.180-EON-multiple	(b) (6) Feed: Purchase history
05-800.180-EON-multiple	(b) (6) Results: District testing results
06-800.180-EON-multiple	(b) (6) Summary: Vet-LIRN summary document

SIGNATURES

Deputy Director OR Date

Director OR Date

Study Director Date

Vet-LIRN Case Summary Document

Vet-LIRN Case Number:	800.180
EON/CC #:	EON-226814-226821-226827
Vet-LIRN Initiation Date:	6.7.2016
MedRec: Requested:	6.7.2016
MedRec: Received:	6.7.2016
MedRec: Significant finding:	(b) (6)-DCM,
Vet-LIRN Tests (planned):	MRx, owner interview, ORS to sample
Vet-LIRN Test Results:	ORS sample-in compliance
Result Interpretation:	
IF NFA, justification:	Completed MRx, Interview

COMPLAINT:

#1-for (b) (6)-12 yo FS Mixed Breed Feline: Presented 5/8/2016 for lethargy; on physical exam the patient was dyspneic and pleural effusion identified on cursory ultrasound and DV thoracic x-rays – given lasix and placed in oxygen. Transferred to cardiology service; evaluation including echocardiogram on 5/9/16 revealed dilated cardiomyopathy, moderate left atrial enlargement, pleural effusion and azotemia. Plasma taurine was submitted to University of Wisconsin. Lab results received 5/15/16 - plasma taurine 24nmol/ml (ref range 60-120, critical level <40). Recheck echocardiogram on 5/15/16 revealed same changes as prior and a thrombus in her left ventricle. Medications/supplements included taurine 250mg PO BID, Mirtazepine 15mg tablets (Give 1/4 tablet PO every 3d PRN), Furosemide 12.5 mg tablets (Give ¼ tablet PO SID), Pimobendan 1.5mg tiny tabs (Give 1 tablet PO BID). Patient presented on (b) (6) for partial aortic thromboembolism and owner's elected euthanasia. Review of patient's diet history revealed that all 5 cats in household had been fed Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry for approximately 3 years. The 4 remaining cats were tested for taurine deficiency and 2/4 had whole blood levels indicating deficiency: 5/21/2016 - Whole Blood Taurine submitted at the University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016 - (b) (6): 9yr male neutered domestic long hair: 196 nmol/ml (b) (6): 8y female spayed domestic short hair: 368 nmol/ml (b) (6): 9yr male neutered domestic long hair: 124 nmol/ml (b) (6): 9yr male neutered domestic long hair: 536 nmol/ml

For #2 for (b) (6)-9yo MC Mixed Breed feline: Another household cat diagnosed with dilated cardiomyopathy and taurine deficiency - separate report filed (FDA ICSR ID 1053335). Euthanized on (b) (6) due to aortic thromboembolism. Review of the patient's diet history revealed that all 5 cats in household had been fed Merrick Purrfect Bistro Grain Free Real Chicken Recipe Feline dry for approximately 3 years. Remaining 4 cats in household tested for taurine deficiency - whole blood samples submitted to University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results received on 5/27/16 (b) (6) 196nmol/ml - started on taurine supplementation 250mg PO BID for 2-3 weeks. Diet was changed at the time of other cat's diagnosis (5/15/15). Patient also diagnosed with hyperthyroidism on same day as blood submitted for taurine testing - history of weight loss. An echo was not performed on this patient therefore it is unknown if he had evidence of DCM.

#3 (b) (6)-9yo MC Mixed breed feline-OBESE- for Another household cat diagnosed with dilated cardiomyopathy and taurine deficiency - separate report filed (FDA ICSR ID 1053335). Euthanized on (b) (6) due to aortic thromboembolism. Review of the patient's diet history revealed that all 5 cats in household had been fed Merrick Purrfect Bistro Grain Free Real Chicken Recipe Feline dry for approximately 3 years. Remaining 4 cats in household tested for taurine deficiency - whole blood samples submitted to University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results received on 5/27/16 - (b) (6) 124nmol/ml - started on taurine supplementation 250mg PO BID for 2-3 weeks. Diet was changed at the time of other cat's diagnosis (5/15/15). An echo was not performed on this patient therefore it is unknown if he had evidence of DCM.

Signalments: 5 cats in household

- (b) (6)-deceased-12 yo FS Feline Mix, DCM and Aortic Thromboembolism
- (b) (6)-9 yo MC DLH, no echocardiogram, hyperthyroidism, low plasma taurine
- (b) (6)-8 yo FS DSH, plasma taurine wnl
- (b) (6)-9 yo MC DLH, no echocardiogram, low plasma taurine
- (b) (6)-9 yo MC DLH, plasma taurine wnl

Signs: 5 cat household: congestive heart failure in 1 cat with dilated cardiomyopathy and aortic thromboembolism, low plasma taurine → euthanized; other 4 living cats unknown if DCM present because no echocardiogram performed; 2 of 4 living cats had low plasma taurine and supplementation begun; 1 cat with low taurine also diagnosed w/ hyperthyroidism and had a history of weight loss

Food: Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry for approximately 3 years

Owner: (b) (6)

Vet: (b) (6)

Vet-LIRN PLAN OF ACTION: MRx and owner interview, ORA regulatory sampling based on interview results

FINAL CONCLUSION: pending

Follow-up: email vet permission to contact owner and requesting MRx

6.8.2016

JJ-Vet calling owner to let them know we'll be calling (were ok with it when the vet submitted the report). Vet sent MRx.

MRx summary:

(b) (6)-8 yo FS DSH:

Presenting complaint 5/21/2016: taurine check. Lives in a room above the garage with 3 other cats, indoor only, no Rx → recheck 6/1 protein electrophoresis, ddx: LSA vs myeloma vs chronic inflam

PE: FORL right upper PM3, mild tartar, copious black otic debris-AU that is mildly pruritic on cleaning

Labs: 5.21.2016

CBC: nsf

Chem: TP 9.1 (5.2-8.8), Glob 6.5 (2.3-5.3)

6.1 **Taurine:** 368 (300-600, no risk if >200)
Protein Electrophoresis: TP 8.2 (5.2-8.8), Glob 5.3 (2.3-5.3), alpha 0.3 (norm),
alpha 2 0.7 (norm), beta 0.6 (0.3-0.9), Gamma 3.6 (0.3-2.5)

Treatments: tresaderm

(b) (6) -9 yo MC DLH

Presenting complaint 5/21/2016: one of 4 cats kept in finished room above garage, chronic weight loss past few months & taurine check, was losing hair >1 yr but told related to anxiety, voracious appetite, indoor only, no Rx

PE: slightly feisty, alopecia caudal dorsum/ventrum/lateral thighs, BCS 1.5-2/5

Labs: 5.21.2016

CBC: nsf

Chem: ALP 174 (6-102), ALT 243 (10-100)

T4: 20.2 (0.8-4)

Taurine: 196 (300-600, no risk if >200)

Treatments: Tapazole (methimazole), taurine

(b) (6): 9 yo MC DLH

Presenting complaint 5/21/2016: lives in the owner's house, not above the garage b/c (b) (6) is aggressive to him, no Rx, Diet: Merrick;

PE: right upper canine loose, significant gingivitis, heavy tartar PM3's

Labs: 5.21.2016

CBC: nsf

Chem: nsf

Taurine: 536 (300-600, no risk if >200)

(b) (6): 9 yo MC DLH

Presenting complaint 5/21/2016: taurine check, one of 4 cats living above garage in a room; 2 dogs in house, Diet: Merrick; no Rx

PE: some matted hair, BCS 4/5, moderate tartar overall

Labs: 5.21.2016

CBC: MCHC 30.8 (30-38), Plt 188 (200-500)-clumped

Chem: Glob 5.3 (2.3-5.3)

Taurine: 124 (300-600, no risk if >200)

Treatments: Taurine

(b) (6): 12 yo FS DSH-euthanized

Presenting complaint 5/8/2016: lethargy starting 5/7, indoor only → recheck 5/15 b/c very weak, PD, difficulty walking, inappetent → 5/19, feed Merrick Purrfect Bistro Grain Free Real chicken recipe ~3 yr, prior to this fed Dick Van Pattons Indoor Dry-Chicken & Salmon; vet called Merrick → bag purchased 2 weeks prior per owner. Vet doesn't think a single bag/lot issue because takes several months to develop → urinating & defecating outside litterbox, soft stool, no change to appetite, ambulating well → 5/25 dyspnea beginning 5/24, recheck → by 5/26 weak hindlimbs → by 5/29 1 hindlimb worse than other → by (b) (6) recheck b/c dragging right hindlimb → euthanized, vet spoke w/ Merrick QA team and the taurine in that lot # was sufficient

PE 5/8: BCS 7/9, T 93, HR 150, RR 60, muffled heart sounds, inc respiratory effort, dull lung sounds

-5.9: heart sounds slightly muffled, RR 30 w/ slight effort

-5.10: cat PD, inappetent, RR ~28

-5.25: tachypneic, mild inc resp effort, RR 48, faint referred upper airway noise

(b) (6): tachypneic, moderate dyspnea, RR 48-60, non-weight bearing Right hindlimb, Right HL: no femoral pulse, cold paw pad; T 94.8, laterally recumbent

Labs: 5.9.2016 **BGA:** Hct 40, Na 146.3 (146.2-156.2), K 4.99 (3.41-4.71), Cl 107.8 (117-125.3), Ca 1.17 (1.16-1.35), Mg 1.08 (0.33-0.49), Glu 156 (72-132), Lac 9.7 (0.7-1.9), BUN 67 (20-33), Ct 5.3 (1.1-3.5)
-5.9 Renal panel: ALP 11 (14-111), ALT 140 (12-130), BUN 74 (16-36), Cl 100 (112-129), Na 138 (150-165), Ct 1.4 (0.8-2.4)
-5.15 BGA: Hct 43, Na 145.3, K 3.33, Cl 104.5, BUN 61, Ct 3.1
-5.25 BGA: Hct 34, Na 147.3, K 6.65, Cl 115.1, BUN 31, Ct 1.3
5.9 **PCV/TS:** 42/6.8
-5.15 PCV/TS: 39/7.8
-5.25 PCV/TS: 32/6
5.9 **Plasma taurine:** 24 (60-120, critical <40)

5/8 Cursory US: mild-moderate pleural effusion, R>L

5/8 Rads: cardiac silhouette difficult to visualize, pleural effusion, moderate inc opacity area caudal to left cranial lung lobe

5/9 Echocardiogram: mod LA enlargement, LV enlarged, mild RA & RV enlargement, trivial MR & TR, dec aortic & pulmonary flow, moderate volume Pleural effusion → DCM

-5.15: small volume PE, no pericardial effusion, large mass in LV-thrombus

-5.25: thrombus unchanged

-6.1: small vol PE, large LV thrombus

Treatments: Lasix, thoracocentesis: 5/8 (25mL), 5/9 (120 mL), 5/25 (160 mL), O2, pimobendan, taurine, mirtazapine, MaxCal, buprenex, telazol, acepromazine, butorphanol, beuthanasia

Thoughts: since (b) (6) lives in house with owners, could he be getting more supplements of taurine (e.g. table scraps of meat) vs the other 4 cats living above the garage. The clinician commented that if food was the source of the taurine deficiency, it was interesting the array of presentations/levels in the 5 cats.

Left msg for owners to arrange interview.

JJ-Owner sent follow-up email & voicemail. Will email to arrange interview.

6.9.2016

JJ-Owner interview:

Presenting complaint: (b) (6) had lethargy, difficulty moving, not acting like herself. Indoor cats

Prior MHx: none significant for all cats (5) in household

Diet: Merrick Grain Free Bistro Chicken-started in January of 2014, --> on the Chicken variety pretty much the entire time; **owner has a listing of all purchases from pet store and she will send us a copy tomorrow**; fed prior to the Merrick food was Dick Van Pattens Natural Balance-2 indoor cat formulas- owner thinks the chicken & salmon type but not 100% sure, she can find out if needed: owner mentions the consistency changed when the company was purchased by Del Monte which prompted her to switch foods to Merrick;

After the illness onset for (b) (6), Fancy Feast Wet food fed to stimulate her appetite, but she only licked the gravy. So for ~1 week before the other 4 house cats were tested for taurine, only (b) (6) got the Fancy Feast leftovers (solid chunks) not consumed by (b) (6). **(Could this explain his higher blood taurine than the other housemates???)**

Owner will send the list of pet food purchases from Merrick tomorrow. Will forward to group.

6.13.2016

JJ-Received owner's receipts from her food purchase history. Forwarded to DR on 6/10. NFA-completed interview and MRx review.

Final conclusion: Based on the medical records, 3 of the 5 cats in the household had low blood taurine levels. One cat, (b) (6), had documented low blood taurine and was euthanized due to a thromboembolism, a complication arising from dilated cardiomyopathy (DCM). DCM can be caused by low dietary and blood taurine levels. (b) (6) lived above the garage with 3 other cats: (b) (6). (b) (6) also had low blood taurine levels (<200 nMol/mL) associated with risk for DCM. Neither cat had echocardiograms to confirm DCM. It is unclear why (b) (6) had taurine levels within reference range, but may be due to individual variation. One cat, (b) (6), who lived separately from the other cats, had the highest blood taurine level. (b) (6) received a supplemental wet food for approximately 1 week prior to the taurine level check. If the dry food regular diet were deficient in taurine, it is possible the supplemental wet food could have improved his taurine levels. This may explain why (b) (6) had the highest blood taurine levels of all the cats, which were within normal range. The dietary interview and purchase history indicate the Chicken variety was most often fed to the cats. Over time, if the food were deficient in taurine, the cats could develop low blood taurine and thus DCM. The ORS product sampling and taurine testing will provide more information on the food taurine content.

7.21.2016

JJ-Checked w/ DR. Testing results from ORA still pending.

7/27/2016

Taurine results received

7/28/16

JJ-DAF reviewing the taurine results.

8.24.2016

JJ-DAF reviewed the results:

OK Everyone. The product appears to be a dry extruded product, for which the AAFCO Cat Food Nutrient Profiles content for taurine is 0.10% on a dry matter basis. Clearly all three samples were analyzed to contain more than that amount of taurine. On a dry matter basis the concentration of taurine in the samples was analyzed to be:

FACTS #	Amount Taurine Found	%Moisture	%Dry Matter	Amount
	Taurine on a Dry Matter Basis			
958500	0.183g/100g \approx 0.18%	2.20%	$100 - 2.20 = 97.80\%$	
	$0.183/0.9780 = 0.187\%$			
958501	0.153g/100g \approx 0.15%	1.99%	$100 - 1.99 = 98.01\%$	
	$0.153/0.9801 = 0.156\%$			
958504	0.171g/100g \approx 0.17%	2.79%	$100 - 2.79 = 97.21\%$	
	$0.171/0.9721 = 0.176\%$			

All of the Dry Matter Taurine percentages are above 0.10%. IF any of the samples were canned cat food, they would not be in compliance with the AAFCO Cat Food Nutrient Profiles for the recommended minimum taurine content and IF the label indicated the product was formulated to meet the AAFCO Cat Food Nutrient Profiles the product would be misbranded.

The answer to the question of consequence/causation of the taurine content in the product from which these three samples originated to the cats in the consumer complaint is that this(ese) lot(s) of product are not indicated to be causative. However, dilated cardiomyopathy from taurine deficiency occurs over a long period of exposure to a deficient diet (months to a year or more), so, if these cats were eating the Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry for the 3 years indicated in the complaint, it is possible that the product was deficient for some long interval of time during that three year period and that a return to "normal" taurine levels in the diet were insufficient to correct the problem in the three cats that developed low blood taurine and the two with dilated cardiomyopathy. Treatment for dilated cardiomyopathy caused by taurine deficiency takes higher daily doses of taurine for several months than normal dietary amounts and is not completely curative.

Recommendations for regulatory steps to consider: [REDACTED] (b) (5)

NFA.

8.25.2016

JJ-ORA final results received from DR. Filed.

3 subs taurine content: 0.176%, 0.187%, 0.156% on dry matter basis → in compliance w/ 0.10% minimum set by AAFCO

10/14/16

OC-received FOIA request related to the case, preparing documentation. Deadline: 10/26/16.

[REDACTED] (b) (6)

16879

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/aal/aal.html>

Vet/Tech Contact: _____ (b) (6) / Contact: (b) (6) Date: 6-20-17

Company Name: _____ (b) (6)

Address: _____ (b) (6)
_____ (b) (6)

Email: _____ (b) (6)

Tel: _____ (b) (6) Fax: _____ (b) (6)

Billing Contact: _____ (b) (6) TAX ID: _____

Email: _____ (b) (6) Tel: _____ (b) (6)

Patient Name: _____ (b) (6)

Species: Canine

Owner's Name _____ (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)
Plasma: _____ Whole Blood: 276 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

(b) (4)



(b) (6)

PET OWNER: (b) (6)
 SPECIES: Canine
 BREED:
 GENDER: Male
 AGE: 8 Years
 PATIENT ID: (b) (6)














(b) (6)
 (b) (6)
 ACCOUNT #: (b) (6)
 ATTENDING VET: (b) (6)

LAB ID: 2600050841
 ORDER ID: 105299928
 COLLECTION DATE: 6/18/17
 DATE OF RECEIPT: 6/19/17
 DATE OF RESULT: 6/19/17

(b) (4) HealthChek™ - Standard CBC

Hematology

6/19/17 (Order Received)
 6/19/17 5:44 PM (Last Updated)

TEST	RESULT	REFERENCE VALUE							
RBC	4.98	5.39 - 8.7 M/ μ L	L 						
Hematocrit	35.0	38.3 - 56.5 %	L 						
Hemoglobin	12.2	13.4 - 20.7 g/dL	L 						
MCV	70	59 - 76 fL							
MCH	24.5	21.9 - 26.1 pg							
MCHC	34.9	32.6 - 39.2 g/dL							
% Reticulocyte	2.3	%							
Reticulocyte	115	10 - 110 K/ μ L	H 						
Reticulocyte Comment	<p>A reticulocyte count of greater than 110 K/μL of blood is considered evidence of bone marrow response to an increased peripheral demand. Depending on the degree of anemia, a reticulocyte count <110 K/μL may indicate an inadequate bone marrow response. Serial monitoring of the erythrogram and reticulocyte count may be useful to evaluate bone marrow responsiveness over time.</p> <p>The following chart may be used as a guideline to determine appropriateness of regenerative response.</p> <p>Degree of bone marrow response (K/μL):</p> <table border="0"> <tr> <td>Mild</td> <td>110-150</td> </tr> <tr> <td>Moderate</td> <td>150-300</td> </tr> <tr> <td>Marked</td> <td>>300</td> </tr> </table>			Mild	110-150	Moderate	150-300	Marked	>300
Mild	110-150								
Moderate	150-300								
Marked	>300								
WBC	9.1	4.9 - 17.6 K/ μ L							
% Neutrophil	58.6	%							
% Lymphocyte	29.8	%							
% Monocyte	8.7	%							
% Eosinophil	2.9	%							
% Basophil	0.0	%							
Neutrophil	5.333	2.94 - 12.67 K/ μ L							
Lymphocyte	2.712	1.06 - 4.95 K/ μ L							
Monocyte	0.792	0.13 - 1.15 K/ μ L							
Eosinophil	0.264	0.07 - 1.49 K/ μ L							
Basophil	0	0 - 0.1 K/ μ L							

(b) (4)



(b) (6)

PET OWNER: (b) (6)

DATE OF RESULT: 6/19/17

LAB ID: 2600050841

Hematology (continued)

TEST	RESULT	REFERENCE VALUE	
Platelet	208	143 - 448 K/ μ L	

Chemistry

6/19/17 (Order Received)
6/19/17 5:44 PM (Last Updated)

TEST	RESULT	REFERENCE VALUE	
Glucose	84	63 - 114 mg/dL	
(b) (4) SDMA	^a 14	0 - 14 μ g/dL	
Creatinine	0.5	0.5 - 1.5 mg/dL	
BUN	16	9 - 31 mg/dL	
BUN:Creatinine Ratio	32.0		
Phosphorus	5.6	2.5 - 6.1 mg/dL	
Calcium	11.3	8.4 - 11.8 mg/dL	
Sodium	154	142 - 152 mmol/L	H
Potassium	5.4	4.0 - 5.4 mmol/L	
Na:K Ratio	29	28 - 37	
Chloride	107	108 - 119 mmol/L	L
TCO2 (Bicarbonate)	30	13 - 27 mmol/L	H
Anion Gap	22	11 - 26 mmol/L	
Total Protein	6.5	5.5 - 7.5 g/dL	
Albumin	4.1	2.7 - 3.9 g/dL	H
Globulin	2.4	2.4 - 4.0 g/dL	
Alb:Glob Ratio	1.7	0.7 - 1.5	H
ALT	57	18 - 121 U/L	
AST	83	16 - 55 U/L	H
ALP	84	5 - 160 U/L	
GGT	4	0 - 13 U/L	
Bilirubin - Total	<0.1	0.0 - 0.3 mg/dL	
Bilirubin - Unconjugated	0.0	0.0 - 0.2 mg/dL	
Bilirubin - Conjugated	0.1	0.0 - 0.1 mg/dL	
Cholesterol	159	131 - 345 mg/dL	

(b) (4)



(b) (6)

PET OWNER: (b) (6)

DATE OF RESULT: 6/19/17

LAB ID: 2600050841

Chemistry (continued)

TEST	RESULT	REFERENCE VALUE	H
Creatine Kinase	1,162	10 - 200 U/L	<input type="checkbox"/>
Hemolysis Index	^b 1+		<input type="checkbox"/>
Lipemia Index	^c N		<input type="checkbox"/>

^a BOTH SDMA AND CREATININE ARE WITHIN THE REFERENCE INTERVAL which indicates kidney function is likely good. Evaluate a complete urinalysis and confirm there is no other evidence of kidney disease.

^b Index of N, 1+, 2+ exhibits no significant effect on chemistry values.

^c Index of N, 1+, 2+ exhibits no significant effect on chemistry values.

Case Description

Patient Information

Patient name: (b) (6)

Species: Canine

Breed: Mix

Age: 8yr MN

Case Information

Referrer: (b) (6)

Created: 06/19/2017 4:38:15pm

Modified: 06/19/2017 4:52:56pm

Clinic: (b) (6)

Clinician: (b) (6)

Modality: CR

Patient ID: (b) (6)

Sex: M

Description: THORAX II VIEWS

History

history of chronic cough-worsening, not responsive to cough tabs or sid lasix, still eating and dirinking well

Physical Findings

4/6 murmur, clear lungs, soft not painful abdomen

Report

Radiographic Findings

3 thoracic radiographs made to June 19, 2017. Previous radiographs from the referring veterinarian are on the server but cannot be evaluated for comparison due to image transfer artifact.

There is moderate generalized cardiomegaly. The heart measures approximately 11.5 VHS. Peripheral pulmonary vessels are normal in diameter. There is dorsal displacement of the carina by the enlarged heart. The lungs are normal. There is no evidence of cardiogenic pulmonary edema or free pleural fluid. The trachea is normal in diameter.

Conclusion

1. Cardiomegaly. Valvular endocardiosis and insufficiency as most likely. Concurrent pulmonary hypertension is not excluded. There is no evidence of congestive heart failure.
2. Irritation/compression of the carina/principal bronchi by the enlarged heart could be contributing to the cough.
3. Allergic/inflammatory bronchitis or infectious tracheobronchitis may be present without radiographic changes and could also be an underlying cause for chronic cough.

Consider cardiology consult for echocardiography. Consider treatment for bronchitis.

(b) (6)

06/19/2017 4:52:56pm

Radiology Report powered by Remedy View

170
x
with history
by

(b) (6)

(b) (6)
(b) (6)

Patient Information

Patient: (b) (6) Age: 8 years Referring Veterinarian: (b) (6)
Patient Number: (b) (6) Weight:(kg) 6.20 Cardiologist: (b) (6)
(Cardiology)
Breed: Shih Tzu Sex: M Client Number: (b) (6)
Exam Date: 06/19/2017 17:26 BSA: 0.34

History: (b) (6) was presented for evaluation of lethargy and progressive coughing. (b) (6) has had a progressive cough that started approximately 6 months ago. He was started on a cough suppressant (unknown dose/type) and Lasix (5 mg PO q24h) with no improvement. His respiratory rate/effort are normal at home. He has not had any episodes of weakness/collapse but has been lethargic over the past few weeks.

(b) (6) is on a limited ingredient diet with kangaroo as the sole protein source. He had a negative heartworm test in February and receives Sentinel monthly (1/2 of a 26-50# tablet?).

Physical Examination: T 100.8 P 128 R 30
Grade 3-4/6 left apical systolic murmur with radiation to the right base. Fair femoral pulses. Regular rhythm. Normal lung sounds. Eupneic. Normal abdominal palpation. PLNs WNL. Excessive bruising noted after jugular venipuncture (none from the first venipuncture). No other bruising/petechiation noted. Thin body condition. MM light pink/slightly tacky. CRT < 2 sec. Fundic exam: Abnormal. Suspect partial retinal detachment on nasal aspect of retina OS with hyperreflectivity around the edge. No abnormalities identified OD.

Diagnostic Tests: Blood Pressure: 102 mmHg (#3 cuff, left forelimb)
Thoracic Radiographs: Generalized cardiomegaly (progressive as compared to previous films). No evidence of cardiogenic edema. Mild diffuse bronchointerstitial pattern.
CBC: Mild non/pre-regenerative anemia.
Chemistry Profile: Mild hypernatremia (154 mmol/L) with mild hypochloremia (107 mmol/L). Mild elevation in bicarb.
Taurine level pending
Echocardiogram: See below. Sinus rhythm on ECG.

Echocardiographic Report

2D ECHO

LA Systolic Diameter LX	3.3 cm	Aortic Root Diameter	1.2 cm
-------------------------	--------	----------------------	--------

DOPPLER

AV Peak Velocity	60.7 cm/s	PV Peak Gradient	1.1 mmHg
AV Peak Gradient	1.5 mmHg	TR Peak Velocity	292 cm/s
MR Peak Velocity	545 cm/s	TR Peak Gradient	34.2 mmHg
PV Peak Velocity	51.8 cm/s		

M-MODE

LV Diastolic Diameter MM	4.3 cm	LVPW Diastolic Thickness MM	0.58 cm
LV Systolic Diameter MM	4 cm	LVPW Systolic Thickness MM	0.64 cm
LV Fractional Shortening MM	7.1 %	LVPW Percent Thickening MM	0.091
LV Diastolic Volume Cube	79 cm ³	IVS to PW Ratio MM	1
LV Systolic Volume Cube	63.2 cm ³	LV Mass MM	70.3 g
LV Ejection Fraction Cube	0.2	LV Mass Normalized MM	206 g/m ²
IVS Diastolic Thickness MM	0.58 cm	RV Diastolic Diameter MM	0.3 cm
IVS Systolic Thickness MM	0.63 cm	MV E Point Septal Separation	1.6 cm
IVS Percent Thickening MM	0.068		

Left Ventricle: Severe dilation (normalized LVIDd 1.5) with marked reduction in contractility (normalized LVIDs 2.25).

Left Atrium: Moderate dilation

Right Ventricle: Normal

Right Atrium: Normal

Mitral Valve: Mildly thickened valve leaflets. 3+ mitral regurgitation.

Aortic Valve: Normal

Tricuspid Valve: Mildly thickened valve leaflets. 1+ tricuspid regurgitation.

Pulmonic Valve: Normal valve morphology. Mild pulmonic insufficiency.

Aorta: Normal

Pericardium: Normal

Diagnosis

Dilated cardiomyopathy- This is a disease characterized by weakening of the heart muscle and dilation of the heart chambers. As the disease progresses, it can lead to congestive heart failure (fluid in the lungs causing shortness of breath and cough). Abnormal heart rhythms are common and can result in sudden death. Most commonly this is an inherited disease, though it can occur secondary to a deficiency in an amino acid called taurine. This is a relatively uncommon disease in small breed dogs so we generally recommend checking taurine levels in these cases.

Abnormal fundic exam - suspect partial retinal detachment of left eye - this change is not what I typically expect with taurine deficiency, but I strongly recommend having (b) (6) evaluated by a veterinary ophthalmologist to better characterize this change. As we discussed, it may be possible that this change is potentially linked with the heart disease and learning more about this may help us get a better idea of a reason for both conditions. Also, it may be important to initiate treatment to help protect his vision and decrease risk of further changes in the eye that could also eventually cause pain. I do not believe that he is in pain today from this.

Recommendations

Give all medications as directed:

Pimobendan (Vetmedin) 2.5 mg tablets- Give 1 tablet by mouth in the morning and 1/2 of a tablet in the evening. Give at 12 hour intervals.

This is a drug that is approved for the treatment of congestive heart failure secondary to dilated cardiomyopathy or chronic valve disease (endocardiosis). However, two studies of dogs with dilated cardiomyopathy, one in Doberman Pinschers (PROTECT) and one in Irish Wolfhounds, have shown a delay in the onset of heart failure in preclinical dogs treated with Vetmedin compared to placebo. Recently, another study (EPIC) has shown significant prolongation of the asymptomatic period in animals with progressive disease and heart enlargement from chronic valve disease, prior to the onset of congestive heart failure, as well. This is off-label use of this medication. In our experience, side effects are uncommon, but it is important that you advise us if you feel your pet is having any potential adverse effects from this medication. The reported potential side effects listed for this medication are increased heart rate, vomiting, diarrhea, inappetence, uneasiness, incoordination, convulsions, increased drinking and increase urinating.

Enalapril (Enacard, Vasotec) 2.5 mg tablets- Give 1/2 of a tablet by mouth every 12 hours.

This medication is a strong drug that dilates blood vessels, permitting the heart to pump blood more efficiently. It can lower blood pressure (hypotension) and cause changes in kidney function and electrolyte values. If your pet develops weakness or depression, decrease the drug dose by 1/2 and call. A kidney panel and blood pressure should be reevaluated 7-10 days after beginning this medication.

Taurine 500mg - Give 500mg orally every 12 hours. We will discontinue this if his taurine level comes back in the normal range.

You may discontinue the furosemide. The Cough-Tabs can be used as needed, especially at night, for the cough.

One thing that can be very helpful for home monitoring is checking sleeping or resting respiratory rates. A recent study showed that even pets with severe heart disease rarely have resting respiratory rates greater than 30 breaths per minute unless they are starting to decompensate for that disease. Elevated respiratory rates at home may be even more sensitive than chest radiographs at picking up early decompensation. Count your pet's respiratory rate when he/she is at rest or sleeping (not within 20 minutes of being active). If his/her respiratory rate is greater than 30 breaths per minute, recheck again in a couple of hours. If persistently elevated above this level, call.

With advanced heart disease, our biggest dietary concerns are adequate calorie content and low sodium content. We aim for less than 100mg sodium per 100 kilocalories (kcal) in patients with significant structural heart disease. We do not advise protein restriction unless there is concurrent kidney disease (i.e. kidney diets are not advised unless there is concurrent kidney disease). You had already investigated the sodium content of the kangaroo diet and it is within this range, but we may recommend a diet change if his taurine level were to come back abnormal.

We will make further recommendations for follow-up after we receive the taurine level and after we see what the ophthalmologist determines with his eye. He should have a kidney panel and blood pressure rechecked after 7-10 days on the enalapril. This can be scheduled as a technician appointment. We will also want to hear about how his cough responds to the new cardiac medications when he comes for this recheck.

(b) (6) (Cardiology)

(Electronically Signed)

Final Date: 19 June 2017 19:42

Amended: 19 June 2017 19:44

(b) (6)

Notes to our clients

-Please bring all medications to your pet's scheduled appointments.

-We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).

-Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.

-If an emergency arises with your pet, (b) (6) is a 24 hour facility.

(b) (6)

(b) (6)

Client ID:	(b) (6)	Patient ID:	(b) (6)
Client Name:	(b) (6)	Name:	(b) (6)
Spouse/Other:	(b) (6)	Breed:	Shih Tzu Mix
Address:	(b) (6)	Sex:	Neutered Male
Telephone:	(b) (6)	Color:	White
		Age:	8 Yrs. 6 Mos.
		DOB:	(b) (6)

Referring Veterinarian: (b) (6)
Practice: (b) (6)
Phone:
FAX:

Cardiology Reevaluation

Reevaluation of:

Dilated cardiomyopathy, Cough

Owner notes concern over significant increase in cough. They have been traveling frequently due to a recent death in the family and (b) (6) is not tolerating the car rides like he has in the past. Typically, (b) (6) is very calm in the car and will fall asleep right away for the trip but recently he is very anxious with respirations up into the 80s. His normal resting respirations have been averaging in the low 20s (22) when not coughing and not traveling. Owner describes the cough as being forceful but non-productive (does not sound congested). She does report an improved cough with the first two days of Vetmedin but it quickly returned to baseline and has continued to progress. Last night he had a bad night and was unable to sleep comfortably. No change in the cough with the cough tabs prescribed by (b) (6)' regular vet. Owner also reports that there was no improvement in the cough when (b) (6) was receiving his furosemide. (b) (6) has also had three urinary accidents in the house since starting the new medications.

Physical Exam:

	6/19/2017	7/6/2017
	1:44 PM	11:38 AM
Vital Sign	ER	038
Weight	6.2 kilograms	6.08 kilograms
Temp	100.8	99.6
HR	128	164
RR	30	56
RQ		Panting

Grade 4/6 left apical systolic murmur with radiation to the right base. Adequate femoral pulses. Normal lung sounds. Eupneic. Normal abdominal palpation. PLNs WNL. MM pink/moist. CRT < 2 sec.

Diagnostics:

Chest radiographs: Unchanged cardiomegaly. No evidence of cardiogenic edema.
Renal panel: clinically unremarkable
Blood pressure: 138 mmHg (#3cm cuff, left forelimb)

Diagnosis:

Dilated cardiomyopathy
Cough - progressive
Urinary accidents - if (b) (6) continues to have accidents in the house a urinalysis would be recommended to rule out a urinary tract infection. This can be done through your regular vet.

Recommendations:

Please give the following medications as directed:

ITEM DESCRIPTION	DIRECTIONS
Enalapril 2.5mg tablets	Give 1/2 tablet by mouth every 12 hours
Vetmedin 2.5mg tablets	Give 1 tablet by mouth once daily in the mornings and 1/2 tablet by mouth once daily in the evenings
Cough tabs	Give 1/2 tablet by mouth every 12 hours as previously prescribed by your regular vet

ADD:

Doxycycline 100mg tablets Give 1/2 tablet by mouth once every 12 hours. Give with food or flush with water after medicating to avoid esophageal irritation.

(b) (6) does not have evidence of active heart failure on his x-rays. I suspect his cough is secondary to underlying primary airway disease. As we discussed, Lasix can have some anti-inflammatory effects which may account for the increase in cough after stopping this drug. Unfortunately, work up for primary airway disease usually involves general anesthesia to obtain an airway sample. (b) (6) is not a good candidate for this given the severity of his heart disease. We can try empirical therapy starting with doxycycline as it is an antibiotic that targets many respiratory pathogens but also has anti-inflammatory effects.

If his cough does not improve over the weekend, please call us on Monday. If you notice any respiratory concerns, you can always bring (b) (6) in through the emergency department. We will make further recommendations based on how he responds to the doxycycline.

(b) (6)

*****Notes to our clients*****

-Please bring all medications to your pet's scheduled appointments.

-We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS OUTSIDE OF (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays, and weekends) MAY BE ASSOCIATED WITH AN AFTER HOURS FILLING FEE.

-Check out www.goodrx.com and enter your local zip code to search for the best prices on your medications at your local pharmacies.

-If an emergency arises with your pet, (b) (6) is only a phone call away. (b) (6) is a 24 hour facility and the emergency veterinarians can always reach the cardiologist on-call.

-Please schedule your recommended recheck as soon as possible. Our schedule tends to book up quite quickly and we want to make sure that we see your pet in a timely manner.

Vet-LIRN Case Summary Document

Vet-LIRN Case Number:	
EON/CC #:	EON-350158
Owner LAST Name:	(b) (6)
Vet LAST Name:	(b) (6)
Vet-LIRN Initiation Date:	3/28/2018
MedRec: Requested:	Received with Complaint
MedRec: Received:	
MedRec: Significant finding:	
Vet-LIRN Tests (planned):	
Vet-LIRN Test Results:	
Result Interpretation:	
IF NFA, justification:	

COMPLAINT Narrative: At the time of diagnosis (b) (6), (b) (6) was a 13 year old female spayed Labrador retriever who had been maintained on a Zignature Kangaroo formula. She presented with a history of a progressive cough which, prior to presentation, became productive and she coughed up a small volume of pink foam (possible pulmonary edema). On examination she had a 2/6 left apical systolic heart murmur and on echo diagnosed with advanced dilated cardiomyopathy with severe left ventricular dilation, moderate to severe left ventricular systolic dysfunction, and moderate to severe left atrial dilation. Thoracic radiographs were suspicious for early congestive heart failure. A whole blood taurine level was submitted and was low at 168. She was treatment with furosemide, benazepril, pimobendan, spironolactone, taurine and l-carnitine and her diet was changed to Royal Canin Early Cardiac. At her recheck in 2/26/18, (b) (6) heart had improved significantly with now mild dilated cardiomyopathy with normalized left atrial dimensions, mild left ventricular dilation and low normal left ventricular systolic function. The furosemide was able to be discontinued at this time.

Signalment: (b) (6) -13 yr FS Lab

Signs: productive, progressive cough

Food Product: Zignature Kangaroo Formula

Plan:

- MRx
- Open product for Tau, Cysteine, Methionine, +/- Beta-Alanine

MRx summary:

Presenting complaint 10/31/2017: developed a cough on 10/25, Rads and labwork at the vet showed ALP 440, GGT 30, mildly low Lymph, cardiomegaly → treated with hydroxyzine, doxycycline, hydrocodone → stopped drugs Monday b/c cough worsened → to ER on (b) (6) after coughing up pink tinged foam; no lethargy, continues to eat and drink; UTD on vaccines and HWP, no drugs → treat with Lasix, benazepril, vetmedin, spironolactone, Tau, L-carnitine and **vet recommended a diet change** → recheck 2/26/18: intermittent cough, related to excitement, change diet to RC Early Cardiac → on

recheck improved → suspect Tau responsive DCM-mild, suspect cough secondary to bronchial or primary respiratory disease

PE (b) (6): LS-OU, HR 132, mild periodontal disease, Gr II/VI, left apical protosystolic murmur, questionable mild inc bronchovesicular sounds bilaterally, SC mass left ventrum;

PE 2/26: Gr III/VI pansystolic, PMI MV, reg rhythm with S3 gallop, HR 130, BCS 6/9, hepatomegaly

Labs: (b) (6) BP 100 (based on Echo)
-2/26:155 mg Hg, direct measurement
11/3: Tau-blood: 168 (200-350)

Rads 10/27: generalized cardiomegaly, left atrial enlargement, slight left auricular bulge, increased sternal contact & rounded heart, dorsal tracheal deviation, prominent pulmonary vasculature with questionably mild inc interstitial opacity in caudal-dorsal lungs, suggesting early CHF/PE

(b) (6) **Echo:** severe LV hypertrophy, mild-mod MV regurgitation, mod-sev LA dilation, mild TV regurg, mild RV & RA dilation, mod-sev lower systolic function values
-2/26: mild LV dilation, mild MV regurg, normal LA, mild TV regurg, normal RV & RA, low normal systolic functional indices of LV

(b) (6) **ECG:** normal sinus rhythm

An article about beta-alanine: <https://academic.oup.com/alcalc/article/36/1/29/138000>

If Tau & Cys/Met are normal, we may need to reconsider other MOA's causing this, unrelated to the food.

I emailed the vet to request the full MRx and see if lot/best by information available for the leftover food.

4/4/2018

JJ-Vet sent the full MRx available and does not have any leftover food. We will purchase the food for testing. A dog from a previous case without food (800.218- (b) (6) Cocker Spaniel with Low Tau and also eating Zignature Essentials Kangaroo.

MRx added to above summary.

Vet-LIRN Case Summary Document

Vet-LIRN Case Number:	800.261
EON/CC #:	EON-350158
Owner LAST Name:	(b) (6)
Vet LAST Name:	(b) (6)
Vet-LIRN Initiation Date:	3/28/2018
MedRec: Requested:	Received with Complaint
MedRec: Received:	
MedRec: Significant finding:	
Vet-LIRN Tests (planned):	<ul style="list-style-type: none"> • MSU <ul style="list-style-type: none"> ○ Iodine • (b) (4) <ul style="list-style-type: none"> ○ Cys-Met-Tau
Vet-LIRN Test Results:	
Result Interpretation:	
IF NFA, justification:	

COMPLAINT Narrative: At the time of diagnosis (10/31/17), (b) (6) was a 13 year old female spayed Labrador retriever who had been maintained on a Zignature Kangaroo formula. She presented with a history of a progressive cough which, prior to presentation, became productive and she coughed up a small volume of pink foam (possible pulmonary edema). On examination she had a 2/6 left apical systolic heart murmur and on echo diagnosed with advanced dilated cardiomyopathy with severe left ventricular dilation, moderate to severe left ventricular systolic dysfunction, and moderate to severe left atrial dilation. Thoracic radiographs were suspicious for early congestive heart failure. A whole blood taurine level was submitted and was low at 168. She was treatment with furosemide, benazepril, pimobendan, spironolactone, taurine and l-carnitine and her diet was changed to Royal Canin Early Cardiac. At her recheck in 2/26/18, (b) (6) heart had improved significantly with now mild dilated cardiomyopathy with normalized left atrial dimensions, mild left ventricular dilation and low normal left ventricular systolic function. The furosemide was able to be discontinued at this time.

Signalment: (b) (6)-13 yr FS Lab

Signs: productive, progressive cough

Food Product: Zignature Kangaroo Formula

Plan:

- MRx
- Open product for Tau, Cysteine, Methionine, +/- Beta-Alanine

MRx summary:

Presenting complaint 10/27 to rDVM: developed a cough on 10/25, cough for 3-4 days, not lethargic, normal eating/drinking, no vomiting or diarrhea, worse when lying down, dog didn't cough while in clinic except for a tracheal cough when pulling on the leash → treated with hydroxyzine, doxycycline, hydrocodone → stopped all 3 drugs Monday b/c cough worsened → to ER on (b) (6) after coughing up

pink tinged foam; no lethargy, continues to eat and drink; UTD on vaccines and HWP, no drugs → treat with Lasix, benazepril, vetmedin, spironolactone, Tau, L-carnitine and **vet recommended a diet change** → labwork done 11/14 → to rDVM 11/16: doing well → recheck 2/26/18: intermittent cough, related to excitement, change diet to RC Early Cardiac → on recheck improved → suspect Tau responsive DCM-mild, suspect cough secondary to bronchial or primary respiratory disease → recheck 3/13: resting RR 16 rpm, minimal coughing only when excited, since switching to cardiac food BMs are dense and tenesmus, owner is weaning dog off lasix

PE 10/27 @ rDVM: numerous lipomatous & dermal masses, no audible murmur or arrhythmia, shallow breathing

PE (b) (6) @ specialist: LS-OU, HR 100 bpm, mild periodontal disease, Gr II/VI, left apical protosystolic murmur, questionable mild inc bronchovesicular sounds bilaterally, SC mass left ventrum, mildly tense cranial abdominal palpation

PE 11/16 @ (b) (6): mild underbite, H/L wnl

PE 2/26: Gr III/VI pansystolic, PMI MV, reg rhythm with S3 gallop, HR 130, BCS 6/9, hepatomegaly

PE 3/13: T 99.9F, RR 56, HR 124 bpm, Gr III/VI murmur, rest nsf

Labs: 10/27 CBC: Lym 1.01 (1.05-5.1)

-3/13: Lym 1044 (1060-4950), Plt 615 (143-448), Plt inc on direct

10/27 Chem: ALP 440 (23-212), GGT 30 (0-11), rest nsf

-11/14: Glu 51 (70-143), Glob 4.7 (2.5-4.5), ALP 621, GGT 31

-3/13: Na:K 27, ALP 2243 (5-180), GGT 117 (0-13)

(b) (6) BP 100 (based on Echo)

-2/26: 155 mg Hg, direct measurement

-3/13: 130-140 mmHg, direct measurement

11/3 Tau-blood: 168 (200-350)

3/13 UA: 1.010, pH 5

3/13 TT4: 0.8 (1-4)

Rads 10/27: generalized cardiomegaly, left atrial enlargement, slight left auricular bulge, increased sternal contact & rounded heart, dorsal tracheal deviation, prominent pulmonary vasculature with questionably mild inc interstitial opacity in caudal-dorsal lungs, suggesting early CHF/PE

(b) (6) **Echo:** severe LV hypertrophy, mild-mod MV regurgitation, mod-sev LA dilation,

mild TV regurg, mild RV & RA dilation, mod-sev lower systolic function values

-2/26: mild LV dilation, mild MV regurg, normal LA, mild TV regurg, normal RV & RA, low normal systolic functional indices of LV

(b) (6) **ECG:** normal sinus rhythm

Prior MHx: 7/2017: doing well at home-**occasionally coughs**, several SQ masses, no murmur or cough on tracheal palpation; 10/23/2017-vaccines, doing well per O, no murmur ausculted, not been getting HWP consistently,

An article about beta-alanine: <https://academic.oup.com/ajcal/article/36/1/29/138000>

If Tau & Cys/Met are normal, we may need to reconsider other MOA's causing this, unrelated to the food.

I emailed the vet to request the full MRx and see if lot/best by information available for the leftover food.

4/4/2018

JJ-Vet sent the full MRx available and does not have any leftover food. We will purchase the food for testing. A dog from a previous case without food (800.218 (b) (6) Cocker Spaniel with Low Tau and also eating Zignature Essentials Kangaroo.

MRx added to above summary.

4/11/2018

JJ-JG received the sample. I prepared the lab submission forms and will aliquot the sample today for testing.

Vet-LIRN Case Summary Document

Vet-LIRN Case Number:	800.261
EON/CC #:	EON-350158
Owner LAST Name:	(b) (6)
Vet LAST Name:	(b) (6)
Vet-LIRN Initiation Date:	3/28/2018
MedRec: Requested:	Received with Complaint
MedRec: Received:	
MedRec: Significant finding:	
Vet-LIRN Tests (planned):	<ul style="list-style-type: none"> • MSU <ul style="list-style-type: none"> ○ Iodine • (b) (4) <ul style="list-style-type: none"> ○ Cys-Met-Tau
Vet-LIRN Test Results:	<ul style="list-style-type: none"> • Iodine < 10 ppm-no suspicion of exogenous thyroid tissue • Tau
Result Interpretation:	
IF NFA, justification:	

COMPLAINT Narrative: At the time of diagnosis (b) (6), (b) (6) was a 13 year old female spayed Labrador retriever who had been maintained on a Zignature Kangaroo formula. She presented with a history of a progressive cough which, prior to presentation, became productive and she coughed up a small volume of pink foam (possible pulmonary edema). On examination she had a 2/6 left apical systolic heart murmur and on echo diagnosed with advanced dilated cardiomyopathy with severe left ventricular dilation, moderate to severe left ventricular systolic dysfunction, and moderate to severe left atrial dilation. Thoracic radiographs were suspicious for early congestive heart failure. A whole blood taurine level was submitted and was low at 168. She was treatment with furosemide, benazepril, pimobendan, spironolactone, taurine and l-carnitine and her diet was changed to Royal Canin Early Cardiac. At her recheck in 2/26/18, (b) (6) heart had improved significantly with now mild dilated cardiomyopathy with normalized left atrial dimensions, mild left ventricular dilation and low normal left ventricular systolic function. The furosemide was able to be discontinued at this time.

Signalment: (b) (6)-13 yr FS Lab

Signs: productive, progressive cough

Food Product: Zignature Kangaroo Formula

Plan:

- MRx
- Open product for Tau, Cysteine, Methionine, +/- Beta-Alanine

MRx summary:

Presenting complaint 10/27 to rDVM: developed a cough on 10/25, cough for 3-4 days, not lethargic, normal eating/drinking, no vomiting or diarrhea, worse when lying down, dog didn't cough while in clinic except for a tracheal cough when pulling on the leash → treated with hydroxyzine, doxycycline, hydrocodone → stopped all 3 drugs Monday b/c cough worsened → to ER on (b) (6) after coughing up pink tinged foam; no lethargy, continues to eat and drink; UTD on vaccines and HWP, no drugs → treat with Lasix, benazepril, vetmedin, spironolactone, Tau, L-carnitine and **vet recommended a diet change** → labwork done 11/14 → to rDVM 11/16: doing well → recheck 2/26/18: intermittent cough, related to excitement, change diet to RC Early Cardiac → on recheck improved → suspect Tau responsive DCM-mild, suspect cough secondary to bronchial or primary respiratory disease → recheck 3/13: resting RR 16 rpm, minimal coughing only when excited, since switching to cardiac food BMs are dense and tenesmus, owner is weaning dog off lasix

PE 10/27 @ rDVM: numerous lipomatous & dermal masses, no audible murmur or arrhythmia, shallow breathing

PE (b) (6) @ specialist: LS-OU, HR 100 bpm, mild periodontal disease, Gr II/VI, left apical protosystolic murmur, questionable mild inc bronchovesicular sounds bilaterally, SC mass left ventrum, mildly tense cranial abdominal palpation

PE 11/16 @ rDVM: mild underbite, H/L wnl

PE 2/26: Gr III/VI pansystolic, PMI MV, reg rhythm with S3 gallop, HR 130, BCS 6/9, hepatomegaly

PE 3/13: T 99.9F, RR 56, HR 124 bpm, Gr III/VI murmur, rest nsf

Labs: 10/27 CBC: Lym 1.01 (1.05-5.1)
-3/13: Lym 1044 (1060-4950), Plt 615 (143-448), Plt inc on direct
10/27 Chem: ALP 440 (23-212), GGT 30 (0-11), rest nsf
-11/14: Glu 51 (70-143), Glob 4.7 (2.5-4.5), ALP 621, GGT 31
-3/13: Na:K 27, ALP 2243 (5-180), GGT 117 (0-13)
(b) (6) BP 100 (based on Echo)
-2/26: 155 mg Hg, direct measurement
-3/13: 130-140 mmHg, direct measurement
11/3 Tau-blood: 168 (200-350)
3/13 UA: 1.010, pH 5
3/13 TT4: 0.8 (1-4)

Rads 10/27: generalized cardiomegaly, left atrial enlargement, slight left auricular bulge, increased sternal contact & rounded heart, dorsal tracheal deviation, prominent pulmonary vasculature with questionably mild inc interstitial opacity in caudal-dorsal lungs, suggesting early CHF/PE

(b) (6) **Echo:** severe LV hypertrophy, mild-mod MV regurgitation, mod-sev LA dilation, mild TV regurg, mild RV & RA dilation, mod-sev lower systolic function values
-2/26: mild LV dilation, mild MV regurg, normal LA, mild TV regurg, normal RV & RA, low normal systolic functional indices of LV

(b) (6) **ECG:** normal sinus rhythm

Prior MHx: 7/2017: doing well at home-**occasionally coughs**, several SQ masses, no murmur or cough on tracheal palpation; 10/23/2017-vaccines, doing well per O, no murmur ausculted, not been getting HWP consistently,

An article about beta-alanine: <https://academic.oup.com/alcal/article/36/1/29/138000>

If Tau & Cys/Met are normal, we may need to reconsider other MOA's causing this, unrelated to the food.

I emailed the vet to request the full MRx and see if lot/best by information available for the leftover food.

4/4/2018

JJ-Vet sent the full MRx available and does not have any leftover food. We will purchase the food for testing. A dog from a previous case without food (800.218 (b) (6)), Cocker Spaniel with Low Tau and also eating Zignature Essentials Kangaroo.

MRx added to above summary.

4/10/18

JG – Received the sample. Treat-sub1 (Zignature, Kangaroo formula)

4/11/2018

JJ-JG received the sample. I prepared the lab submission forms and will aliquot the sample today for testing.

4/12/2018

JJ-I prepared the samples and sent them to MSU for iodine screening and (b) (6) for Tau/Cys/Met screening.

5/4/2018

JJ-The MSU iodine results were < 10 ppm and not suspicious for exogenous thyroid tissue.

The (b) (4) results came back for Taurine, Cystine, and Methionine.

- Taurine = 45.5 mg/100g = 0.0455g/100g = 0.046% As Is Basis
If we assume a max of 10% moisture per the label (= 90% DMB),
then $0.0455 / 0.90 = 0.05\%$ DMB, which is less than the AAFCO minimum for cats eating extruded foods (0.1% DMB.)
- Cystine = 293 mg/100g = 0.293 g/100g = 0.29% As Is Basis
If we assume a max of 10% moisture per the label (= 90% DMB), then $0.293 / 0.90 = 0.33\%$ DMB
- Methionine = 358mg/100g = 0.358 g/100g = 0.36% As Is Basis
If we assume a max of 10% moisture per the label (= 90% DMB),
then $0.358 / 0.90 = 0.4\%$ DMB, which is greater than the AAFCO minimum for growth & reproduction of 0.35% DMB.
The Methionine-cystine % = $0.4\% + 0.33\% = 0.73\%$ DMB, which is greater than the AAFCO minimum for growth & reproduction of 0.7% DMB.

BLUF: Taurine was low based on the AAFCO minimum for feline extruded foods.

Patient Demographics

(b) (6)		Study Date: 11/01/2017			
Patient ID: (b) (6)		Accession #:		Alt ID:	
DOB:	Age:	Gender:	Ht:	Wt: 67lb 4oz	BSA:
Institution: CVCA (b) (6)					
Referring Physician:					
Physician of Record:		Performed By:			
Comments:					

Adult Echo: Measurements and Calculations

2D

LVIDd (2D)	6.23 cm	LVAd (A4C)	34.40 cm ²	IVSd (2D)	0.932 cm
LVPWd (2D)	0.791 cm	LVAs (A4C)	25.70 cm ²	RVIDd/LVIDd	0.139
EDV (2D-Teich)	196 ml	EDV (A4C)	141 ml	RVIDd (2D)	0.866 cm
EDV (2D-Cubed)	242 ml	ESV (A4C)	88.8 ml	LA Area	24.1 cm ²
A4Cd		LV Mass (Cubed)	239 g	LA Dimen (2D)	4.2 cm
LV Vol	141 ml				
LV Length	6.89 cm				
LV Area	34.4 cm ²				
A4Cs		IVS/LVPW (2D)	1.18	LA/Ao (2D)	1.75
LV Vol	88.8 ml				
LV Length	6.13 cm				
LV Area	25.7 cm ²				
LVLd (A4C)	6.9 cm	SV (A4C)	52.2 ml	AoR Diam (2D)	2.4 cm
LVLs (A4C)	6.1 cm	EF (A4C)	37.0 %		

MMode

IVSd (MM)	0.966 cm	SV (MM-Teich)	78.0 ml	LVPW % (MM)	21.1 %
LVIDd (MM)	6.30 cm	FS (MM-Teich)	19.4 %	RVIDd (MM)	0.322 cm
LVPWd (MM)	0.859 cm	EF (MM-Teich)	38.8 %	LA Dimen (MM)	3.7 cm
IVSs (MM)	1.11 cm	EDV (MM-Cubed)	250 ml	AoR Diam (MM)	2.3 cm
LVIDs (MM)	5.08 cm	ESV (MM-Cubed)	131 ml	LA/Ao (MM)	1.61
LVPWs (MM)	1.04 cm	SV (MM-Cubed)	119 ml	MV D-E Exc Dist	1.4 cm
IVS/LVPW (MM)	1.12	EF (MM-Cubed)	47.6 %	MV D-E Slope	43.6 cm/s

EDV (MM-Teich)	201 ml	FS (MM-Cubed)	19.4 %	MV E-F Slope	19.1 cm/s
ESV (MM-Teich)	123 ml	IVS % (MM)	14.9 %	MV EPSS	1.4 cm

Doppler

LVOT Vmax		MV Peak A Vel		Lat A` Vel	10.7 cm/s
Max PG	7 mmHg	Vel	75.2 cm/s		
Vmax	134 cm/s	PG	2 mmHg		
RVOT Vmax		MV E/A	1.6	E`/A` Lateral	1.2
Max PG	2 mmHg				
Vmax	77.1 cm/s				
MR Vmax		Lat E` Vel	12.7 cm/s	TR Vmax	
Max PG	100 mmHg			Max PG	40 mmHg
Vmax	501 cm/s			Vmax	315 cm/s
MV Peak E Vel		E/Lat E`	9.8		
Vel	1.24 m/s				
PG	6 mmHg				

Other Measurements

Dimensions: 2D LAX

LA lax (2D) 5.41 cm

Dimensions: Diameters

LVID/Ao (2D) 2.60

EF & Volume: Simpson's


Sphericity Id 1.1







Dimensions: Diameters


LVEDDN 2.31

LVID/Ao (2D) 2.60

Images







Signature

Signature:
Name(Print):

Date:

Patient Chart for (b) (6)
Date: 03-14-18, Time: 5:05p

Client: (b) (6)
Page: 2

11-25-13	64.00
09-16-13	69.00
07-10-13	59.30
07-25-12	68.30
01-09-12	71.00

MEDICAL HISTORY

Date	By	Code	Description	Qty (Variance)	Photo
03-13-18	JO2	(b) (4)	(b) (4) Requisition	(b) (6)	
		865	Senior Screen		
			Attachments\8546	(b) (6), (b) (6)	3.14.18 Lab.pdf
		C153	Office Visit - Recheck		
		P163	Blood Pressure		
			CHECK-IN Patient check-in		
			SMT: 11-14-17 at 10:51a: recheck blood chemistry proile w/ electrolytes O wants AB		
			CMO: 02-12-18 at 4:59p: called o to r/s - ok with JO		
			AB2: 02-27-18 at 11:15a: exam, BP, sr screen per last CVCA report		
			SMT: 03-12-18 at 2:09p: LMOM		

Age: 12y Weight: 71.50 Temp: 99.90 Respiration: 56.00 Pulse: 124.00
CRT: pink 1-2 secs.

SUBJECTIVE SECTION

exam, BP, sr screen per last CVCA report

resting resp around 16 per mom, doing well at home, eating/drinking normal, bathroom normal, minimal coughing only when excited, since o switched to cardlac food BMs are very dense and sometimes has trouble passing stool, no vomiting, no other concerns, per o is weaning p off lasix

OBJECTIVE SECTION

ABNORMALITIES

Oral Cavity
mm pink

Cardiovascular
III/VI murmur as previously described

Respiratory
Respiratory rate normal; lungs eupneic

Lymphatic
All palpable LN's WNL

Other
euhydrated, BAR

PLAN SECTION

NOTES

BP 130-140 (LHL, size 5 cuff)

Patient Chart for (b) (6)
Date: 03-14-18, Time: 5:05p

Client: (b) (6)
Page: 3

Date	By	Code	Description	Qty (Variance)	Photo
			Senior screen to (b) (4) UA free catch		
			Disc firm stools and mild tenesmus- adv can trial metamucil but rec confirm with cardiologist ok to add in.		

121217 B

(b) (6)

CVCA CONSULTATION REQUEST FORM

Date: (b) (6)

Client Id #: (b) (6) Client Name: (b) (6)

Address: (b) (6) City: (b) (6) State: (b) (6) Zip: (b) (6)

Telephone:

Cellular: : (b) (6)
Cellular: : (b) (6)

Animal Name: (b) (6) Species: Canine Breed: Labrador Retriever

Color: Yellow Sex: spayed female Weight: 0Kg.

Date of Birth: (b) (6) Age: 13 Yrs. 0 Mos.

Referring Veterinary Hospital: No Vet

Doctor's Name: No Vet

Referring Veterinary Hospital Phone #: (b) (6)

(b) (6) Requesting Consult: (b) (6)

Relevant History / Physical Findings:

Cough started last Wednesday. Radiographs and blood work were performed. Radiographs revealed suspected cardiomegaly. Blood work showed mild ALP and GGT elevations. The owner made cardio-consultation on Friday however her cough got worse with pink tinged foam so (b) (6) was brought to (b) (6) for a cardiology consultation. (b) (6) has been a healthy dog with no current medications. She is up to date on vaccination and heartworm preventative.

Current Medications:

Hydroxyzine, Doxycycline, and hydrocodone, which was stopped because her coughing got worse with those medications.

Radiographs performed at:

- RDVM
- (b) (6)

Consulting Cardiologist:

(b) (6)	(b) (6)
(b) (6)	

CVCA, Cardiac Care for Pets

(b) (6)
Phone: (b) (6)
Email: (b) (6)@cvcavets.com
www.cvcavets.com



Client: (b) (6)
Co-owner:
Patient name (b) (6)
Species: Canine
Breed: Labrador Retriever
Sex: FS
Age: 13 years and 5 months old
Weight: 33.18kg. / 73.15 lbs

(b) (6)
Email:

Cardiac Evaluation Report Exam Date: 02/26/2018

Diagnosis

- Mild, improved dilated cardiomyopathy - suspect taurine-responsive
- Mild, improved mitral and very mild tricuspid valve regurgitation as cause of heart murmur
- Normal, improved left atrial chamber dilation
- Mild, improved eccentric left ventricular chamber dilation
- Low normal, improved left ventricular contractility/heart muscle function
- Cough - suspect bronchial/primary respiratory disease

Medications

- Decrease Lasix/Furosemide 40 mg tablets - Give 1 and 1/2 tablets twice daily for 1 week then decrease to 1 tablet twice daily for 1 week then decrease to 1/2 tablet twice a day for 1 week then discontinue. Please call if you note an increase respiratory rate while decreasing the Lasix. If there is an increase in cough (but normal respiratory rate), we will consider adding in a bronchodilator.
- Continue Benazapril 10 mg tablets - Give 1 and 1/2 tablets twice daily
- Continue Vetmedin/Pimobendan 7.5 mg EZ tablets - Give 1 tablet twice daily.
- Continue Spironolactone 25 mg tablets - Give 1 tablet twice daily.
- Continue Taurine 1500 mg twice daily.
- Continue L-carnitine 1500 mg three times daily.
- You may purchase the taurine and L-carnitine at any health food or nutrition store or www.puritanspride.com. You may also obtain the L-carnitine in bulk powder form from North Carolina State University by calling 919-513-6325.
- Continue with monthly heartworm and flea/tick control as prescribed by (b) (6)

Please allow 24-48 hours for CVCA to process prescription refill requests.

Refill all medications indefinitely unless directed by CVCA or your primary care veterinarian.

- **Please check all medications and dosages on your discharge report against the pharmacy labels.**

Please Note

- Please see our website www.cvcavets.com for more information about (b) (6) dilated cardiomyopathy.

Nutrition Recommendations:

- Continue the Royal Canin Early Cardiac diet.
- Consider fish oil supplements (omega-3 fatty acids). Her dose is approximately EPA 1220 mg and DHA 760 mg total per day. Please start at 1/2 the dose for one week, then increase to the full dose if tolerating well thereafter. Please avoid Cod liver oil and flax seed as well as products with Vit A and/or D.

For more information about fish oils, please visit --<http://vet.tufts.edu/heartsmart/diet/important-nutrients-for-pets-with-heart-disease/>

- In addition to the supplements approved by Tuft's Veterinary Nutrition Service, other reputable brands include Welactin and Nordic Naturals. (b) (6) may have additional brand recommendations.

Activity Recommendations

- Continue normal activity as she wants and is able to do. Please allow (b) (6) to take more breaks and rest during activity.
- Please avoid exercise in the hot/humid weather.

At Home Monitoring:

- In order to monitor for the development of early congestive heart failure in the out-patient setting, we recommend monitoring your pet's resting respiratory rate several times a week. Normal resting respiratory rates should be less than 30 breaths per minute. Consider using a respiratory rate monitoring application to track (b) (6) respiratory rate - Cardalis or BI Pharma have reliable phone applications. Please contact us if you note a persistent or progressive increase.

Future Anesthesia/Fluid Recommendations

- We expect (b) (6) to tolerate carefully monitored general anesthesia with normal preoperative bloodwork and a balanced anesthetic regimen. During anesthesia, we recommend careful monitoring of ECG, BP and pulse ox and 1/2 usual surgical fluid rate (ie: 2-4 ml/kg/hr). Carefully monitor for several hours post-operatively for signs of respiratory congestion and consider chest radiographs if these signs occur. There is some risk associated with all anesthetic events.
- Avoid medications with tachycardia as a side effect, such as ketamine, telazol and glycopyrrolate. Cleared for low dose atropine if needed for intraprocedure bradycardia. Avoid medications that significantly alter blood pressure such as acepromazine and Domitor.
- (b) (6) should not receive corticosteroids (prednisone) in the future please contact CVCA for recommendations, if corticosteroids are indicated.

Reevaluation

- Recheck with (b) (6) in the next 2-4 weeks and every 6 months for wellness care as directed, close auscultation, blood pressure and complete lab tests including blood and urine testing (CBC/Chemistry/Urinalysis/Thyroid evaluation). Please forward these results when available.
- Please recheck with CVCA in 6 months for a follow up consultation/examination, blood pressure, and echocardiogram. Please contact us or schedule an earlier appointment if (b) (6) has any problems or symptoms indicative of worsening heart disease or if recommended by (b) (6)

We thank you for trusting in CVCA to care for (b) (6) today. Please do not hesitate to call us with any questions or concerns.

Sincerely,

(b) (6)

Visit Summary

Heart Rate: 130

BP: 155 mmHg

Cuff Size/Location: 6 cuff/LF

History: Recheck DCM, suspected early CHF; doing well; RRR - 16 bpm, increased Lasix in January due to increased cough; cough seems to be intermittent and related to excitement; good appetite; 3 kg weight gain since 10/2017; walks 30-45 minutes per day - slow pace, at times winded but recovers very quickly.

(b) (6) developed a cough last Wednesday (10/25/17). Radiographs and blood work were performed by (b) (6). The lab work (which is unavailable for review) reportedly showed an elevated ALP 440 and GGT 30 and mild lymphopenia. Thoracic radiographs were performed which revealed cardiomegaly. (b) (6) was treated with hydroxyzine 50mg BID, doxycycline 200mg AM and 100mg PM, and hydrocodone 5mg q8-12h. All medications were stopped on Monday as her cough had worsened and she was presented to the (b) (6) for a cardiac evaluation as her coughing had worsened and she had brought up a small volume of pink-tinged foam after a coughing fit. During this time there has been no evidence of lethargy and she continues to eat and drink normally at home.

PPHx: None

Meds: None

Other: UTD on vaccinations, On HW preventative

Diet: changed from Zignature (Kangaroo) to Royal Canin Early Cardiac

Physical Exam Findings: 3/6 pansystolic murmur, PMI - mitral valve, regular rhythm with S3 gallop; LUNGS - clear all fields, panting, normal effort; SI. overweight body condition (BCS - 6/9); Pink mm; PP - SS; PLN - WNL; ABD - hepatomegaly; BAR

Echocardiographic Findings

Mild left ventricular eccentric dilation - significant improvement compared to previous exam; mild, improved centrally located mitral regurgitant jet, normal, improved left atrial dimensions on 2D imaging and on M-mode imaging, mild, low velocity eccentric low velocity tricuspid regurgitation, subjectively normal right ventricular and right atrial dimensions, normal left and right ventricular outflow velocities, low normal, improved indices of systolic function (FS% and EF% by modified Simpson's, normal EPSS, normal transmitral inflow velocities and E:A wave ratio on spectral Doppler tracings, normal TDI E':A' ratio of the lateral mitral annulus, no masses, effusions or heartworms observed.

Comments

Dear (b) (6),

Thank you for sending (b) (6) to see us with (b) (6) today. I am quite pleased with (b) (6) exam today. She has had remarkable improvement in her echocardiogram with the cardiac medications, change in diet and supplementation with Taurine and L-carnitine. Her risk for congestive heart failure at this point is very low so we will be weaning (b) (6) off the Lasix/furosemide while (b) (6) monitors (b) (6) respiratory rate. Her current cough is likely due to respiratory disease and if the cough progresses/worsens, we will consider adding in a bronchodilator, such as Theophylline. Right now, with the marked improvement, (b) (6) long-term prognosis has improved considerably. I suspect we will be able to further discontinue cardiac medications if her heart remains stable. We will continue to closely monitor (b) (6) heart disease via serial echocardiography and institute further therapy when progression is noted. While on this course of medication, it is important to monitor the chemistry profiles and blood pressures. Hopefully, (b) (6) will continue to do so well - she's a sweetie!

We appreciate your continued referrals and the trust you place in CVCA to co-manage your cardiac patients. We look forward to working with you on this case and others. In an effort to continue to improve CVCA's service to both you and your clients, please visit our website at www.cvcavets.com and complete our online referring veterinarian survey.

Sincerely,

(b) (6)

Patient Demographics

(b) (6)		Study Date: 02/26/2018			
Patient ID:	(b) (6)	Accession #:	Alt ID:		
DOB:	Age:	Gender:	Ht:	Wt: 73lb 0oz	BSA:
Institution: Philips Medical					
Referring Physician:					
Physician of Record:				Performed By: (b) (6)	
Comments:					

Adult Echo: Measurements and Calculations

2D

LVIDd (2D)	5.01 cm	LVAd (A4C)	21.30 cm ²	IVSd (2D)	1.24 cm
LVPWd (2D)	1.20 cm	LVAs (A4C)	13.90 cm ²	RVIDd/LVIDd	0.139
EDV (2D-Teich)	119 ml	EDV (A4C)	61.9 ml	RVIDd (2D)	0.695 cm
EDV (2D-Cubed)	126 ml	ESV (A4C)	33.3 ml	LA Area	15.8 cm ²
A4Cd		LV Mass (Cubed)	186 g	LA Dimen (2D)	2.9 cm
LV Vol	61.9 ml				
LV Length	5.90 cm				
LV Area	21.3 cm ²				
A4Cs		IVS/LVPW (2D)	1.03	LA/Ao (2D)	1.21
LV Vol	33.3 ml				
LV Length	4.79 cm				
LV Area	13.9 cm ²				
LVLd (A4C)	5.9 cm	SV (A4C)	28.6 ml	AoR Diam (2D)	2.4 cm
LVLs (A4C)	4.8 cm	EF (A4C)	46.2 %	HR - AV	82 bpm

MMode

IVSd (MM)	1.09 cm	SV (MM-Teich)	52.1 ml	LVPW % (MM)	40.9 %
LVIDd (MM)	4.96 cm	FS (MM-Teich)	22.4 %	RVIDd (MM)	0.806 cm
LVPWd (MM)	0.965 cm	EF (MM-Teich)	44.9 %	LA Dimen (MM)	3.1 cm
IVSs (MM)	1.58 cm	EDV (MM-Cubed)	122 ml	AoR Diam (MM)	2.4 cm
LVIDs (MM)	3.85 cm	ESV (MM-Cubed)	57.1 ml	LA/Ao (MM)	1.29
LVPWs (MM)	1.36 cm	SV (MM-Cubed)	64.9 ml	MV D-E Slope	25.7 cm/s
IVS/LVPW (MM)	1.13	EF (MM-Cubed)	53.2 %	MV E-F Slope	13.6 cm/s

EDV (MM-Teich)	116 ml	FS (MM-Cubed)	22.4 %	MV EPSS	0.3 cm
ESV (MM-Teich)	63.9 ml	IVS % (MM)	45.0 %		

Doppler

LVOT Vmax		MV E/A	1.6	E`/A` Medial	1.3
Max PG	18 mmHg				
Vmax	211 cm/s				
RVOT Vmax		Med E` Vel	5.71 cm/s	TR Vmax	
Max PG	3 mmHg			Max PG	6 mmHg
Vmax	91.2 cm/s			Vmax	125 cm/s
MV Peak E Vel		E/Med E`	8.5		
Vel	0.488 m/s				
PG	1 mmHg				
MV Peak A Vel		Med A` Vel	4.54 cm/s		
Vel	30.8 cm/s				
PG	0 mmHg				

Other Measurements

Dimensions: Diameters

LVID/Ao (2D)	2.09
EDVI	57.4 ml/m ²
ESVI	30.9 ml/m ²


EF & Volume: Simpson's


Sphericity Id	1.2
---------------	-----

Dimensions: Diameters

LVEDDN	1.77
LVID/Ao (2D)	2.09

Images





Signature

Signature:
Name(Print):

Date:

CVCA, Cardiac Care for Pets

(b) (6)
Phone: (b) (6)
Email: (b) (6)@cvcavets.com
www.cvcavets.com



Client: (b) (6)
Co-owner:
Patient name: (b) (6)
Species: Canine
Breed: Labrador Retriever
Sex: FS
Age: 13 years and 5 months old
Weight: 33.18kg. / 73.15 lbs

(b) (6)

Cardiac Evaluation Report Exam Date: 10/31/2017

Diagnosis

- Advanced dilated cardiomyopathy - ruleout idiopathic vs. taurine-responsive
- Mild to moderate mitral valve regurgitation as cause of heart murmur
- Trace tricuspid valve regurgitation
- Moderate to severe left atrial chamber dilation
- Severe eccentric left ventricular chamber dilation
- Moderate to severe decrease in contractility/heart muscle function
- Mild left ventricular wall thinning
- Mild right atrial and right ventricular chamber dilation
- Progressive cough - rule out: early left sided congestive heart failure vs. mainstem bronchial compression

Medications

- Begin Lasix/Furosemide 40 mg tablets - Give 1 tablet twice daily.
 - > For mild increases in respiratory rate/effort, you may give an additional dose of Lasix.
 - > If you are consistently giving an additional dose of Lasix, please contact our office so we may help adjust medications long-term.
 - > We may increase this dose in the future based on at home monitoring of breathing and recheck blood work.
- Begin Benazapril 10 mg tablets - Give 1 tablet twice daily for 4 days then increase to 1 and 1/2 tablet twice daily thereafter.
- Begin Vetmedin/Pimobendan 5mg tablets - Give 1 and 1/2 tablets twice daily. Will switch to 7.5 mg EZ tablets at 1 tablet twice daily. The 7.5mg tablet will be compounded through (b) (6), please call them to set up shipping and billing (b) (6)
- Please call if you notice a decrease in appetite, vomiting, lethargy, weakness or any other signs of illness while beginning/adjusting the medications.
- Continue with monthly heartworm and flea/tick control as prescribed by (b) (6)

In 2 weeks, if (b) (6) is eating and feeling well:

- Begin Spironolactone 25 mg tablets - Give 1 tablet once daily for 4 days then increase to 1 tablet twice daily thereafter.

< Begin Taurine 1500 mg twice daily.
< Begin L-carnitine 1500 mg three times daily.
< You may purchase the taurine and L-carnitine at any health food or nutrition store on www.puritanspride.com. You may also obtain the L-carnitine in bulk powder form from North Carolina State University by calling 919-513-6325.

Please allow 24-48 hours for CVCA to process prescription refill requests.

Refill all medications indefinitely unless directed by CVCA or your primary care veterinarian.

< **Please check all medications and dosages on your discharge report against the pharmacy labels.**

Please Note

< Please see our website www.cvcavets.com for more information about (b) (6) dilated cardiomyopathy.

Nutrition Recommendations:

< (b) (6) is on a specialized diet which could be contributing to taurine deficiency. Please change her to a new diet, as her housemate is on a novel protein diet - consider prescription diets such as Royal Canin or Science Diet. Please discuss diet options with (b) (6)

< In patients with early/mild heart failure, CVCA recommends feeding a diet with less than 80 mg of sodium per 100 kCal of food (50-80 mg/100 kCal). In patients with refractory heart failure signs, further sodium restriction may be beneficial.

< For more information about sodium content of various foods, please visit:

- o Dog: http://vet.tufts.edu/wp-content/uploads/reduced_sodium_diet_for_dogs.pdf
- o Treats: http://vet.tufts.edu/wp-content/uploads/treats_for_dogs_with_heart_disease.pdf

< CVCA recommends avoiding kidney diets unless (b) (6) has kidney disease that warrants protein restriction.

< Diet changes should be done gradually (ie. over ~1 month) to avoid GI upset and avoided until (b) (6) is stable and eating well on the cardiac medications, usually about 2 weeks after starting or adjusting therapy.

< If you are interested in a consultation with a veterinary nutritionist, please visit -<http://vetnutrition.tufts.edu/make-an-appointment/>

< CVCA recommends fish oil supplements (omega-3 fatty acids) in many dogs with cardiac disease. Her dose should be approximately EPA 1220 mg and DHA 760 mg total per day. Please start at 1/2 the dose for one week, then increase to the full dose if tolerating well thereafter. Please avoid Cod liver oil and flax seed as well as products with Vit A and/or D.

For more information about fish oils, please visit --<http://vet.tufts.edu/heartsmart/diet/important-nutrients-for-pets-with-heart-disease/>

< In addition to the supplements approved by Tuft's Veterinary Nutrition Service, other reputable brands include Welactin and Nordic Naturals. (b) (6) may have additional brand recommendations.

Activity Recommendations

< Keep (b) (6) very quiet for the next 3-4 days with only brief leash walks to eliminate.

< Once her coughing has resolved, (b) (6) may gradually resume activity as she wants and is able to do. Please allow (b) (6) to take more breaks and rest during activity.

< Please try avoid burst type activity, as this increases the arrhythmia risk and avoid exercise in the hot/humid weather.

< Please try to warm (b) (6) up for 5-10 minutes with walking prior to moderate activity and take more rests during more vigorous activity.

At Home Monitoring

< Monitor for signs of cough, respiratory difficulty, exercise intolerance, abdominal swelling, weakness, lethargy, etc. If you note any of these symptoms, please notify CVCA or (b) (6) as these symptoms may indicate recurrent congestive heart failure. If you note an increase in cough, respiratory rate or effort, please feel free to give an additional dose of Lasix/Furosemide, while contacting CVCA.

< In order to monitor for the development of early congestive heart failure in the out-patient setting, we recommend monitoring your pet's resting respiratory rate several times a week. Normal resting respiratory rates should be less than 30 breaths per minute. Consider using a respiratory rate monitoring application to track (b) (6) respiratory rate - Cardalis or BI Pharma have reliable phone applications. Please contact us if you note a persistent or progressive increase.

< In addition, (b) (6) is sadly at increased risk for sudden cardiac death due to her cardiac disease. Dobermans are particularly at risk for development of severe, sudden malignant arrhythmias that sadly may result in sudden death. However, we hope to minimize these risks with our treatment plan.

Future Anesthesia/Fluid Recommendations

- Avoid intravenous or subcutaneous fluid therapy in the future, if possible. If fluid therapy is indicated, please contact CVCA.
- (b) (6) should not receive corticosteroids (prednisone) in the future please contact CVCA for recommendations, if corticosteroids are indicated.
- Avoid elective anesthesia, as (b) (6) is at high risk for complications due to the degree of cardiac disease. If anesthesia is necessary in the future, please contact CVCA for recommendations for monitoring and anesthetics.

Reevaluation

- Please recheck with (b) (6) in the next day or two to obtain taurine levels. Please forward these results when available.
- Please recheck with (b) (6) in 2 weeks for a follow up examination and blood chemistry profile with electrolytes and as recommended by (b) (6) Please forward these results when available.
- Please recheck with (b) (6) every 4-6 months for a follow up examination and blood chemistry profile with electrolytes and as recommended by (b) (6) Please forward these results when available.
- Please recheck with CVCA in 5 months for a follow up consultation/examination, blood pressure, and echocardiogram. Please contact us or schedule an earlier appointment if (b) (6) has any problems or symptoms indicative of worsening heart disease or if recommended by (b) (6)

Visit Summary

Heart Rate: 132 bpm

BP: 100mmHg (based on MR gradient)

History:

(b) (6) developed a cough last Wednesday (10/25/17). Radiographs and blood work were performed by (b) (6). The lab work (which is unavailable for review) reportedly showed an elevated ALP 440 and GGT 30 and mild lymphopenia. Thoracic radiographs were performed which revealed cardiomegaly. (b) (6) was treated with hydroxyzine 50mg BID, doxycycline 200mg AM and 100mg PM, and hydrocodone 5mg q8-12h. All medications were stopped on Monday as her cough had worsened and she was presented to the (b) (6) for a cardiac evaluation as her coughing had worsened and she had brought up a small volume of pink-tinged foam after a coughing fit. During this time there has been no evidence of lethargy and she continues to eat and drink normally at home.

PPHx: None

Meds: None

Other: UTD on vaccinations, On HW preventative

Diet: Zignature (Kangaroo)

Physical Exam Findings:

BAR, sweet but nervous

OP/EENT: Pink, moist mucous membranes, CRT <2s, mild periodontal disease, LS OU, clear AU, No nasal or ocular discharge, no cough on tracheal palpation

PLN: WNL

H/L: Grade 2/6 left apical protosystolic heart murmur, regular rhythm, strong synchronous femoral pulses, RR: 36 breaths/min, questionable mild increase in bronchovesicular sounds bilaterally, no crackles or wheezes ausculted, eupneic

Abd: Soft non-painful abdominal palpation, no palpable masses or fluid wave

MS/Neuro: BCS 5/9, Amb x 4, Mentally alert and appropriate

Integ: Normal turgor, subcutaneous mass left ventrum

Other Diagnostics:

10/27/17 pDVM CXR: Generalized cardiomegaly characterized by widening of the cardiac silhouette and loss of the caudal cardiac waist consistent with left atrial enlargement. Slight left auricular bulge. Increased sternal contact and rounding of the right heart on the VD radiograph. Dorsal deviation of the trachea. Prominent pulmonary vasculature with a questionable mild increase in interstitial opacity in the caudodorsal lung fields which may suggest early congestive heart failure/pulmonary edema.

Echocardiographic Findings

Severe left ventricular eccentric hypertrophy with apical rounding and increased sphericity, mild-moderate centrally

FDA-CVM-FOIA-2019-1704-000071

located mitral regurgitant jet, moderate-severe secondary left atrial dilation on 2D imaging and moderately-severely increased LA:Ao ratio on M-mode imaging, mild eccentric low velocity tricuspid regurgitation with mildly elevated estimated right ventricular pressures consistent with mild pulmonary hypertension, mild right ventricular and right atrial dilation, normal left and right ventricular outflow velocities, moderately to severely depressed indices of systolic function (FS% and EF% by modified Simpson's - LVDI 144ml/m², LVSI 90ml/m²), increased EPSS, elevated transmitral inflow velocities and E:A wave ratio on spectral Doppler tracings, normal TDI E':A' ratio of the lateral mitral annulus, no masses, effusions or heartworms observed.
ECG during echocardiogram: Normal sinus rhythm. No ventricular ectopy noted.

Comments

Dear (b) (6)

Thank you for sending (b) (6) to see us with (b) (6) today. Sadly, (b) (6) has dilated cardiomyopathy with moderate to severe systolic dysfunction and moderate to severe left atrial dilation. This places her at a high risk of developing congestive heart failure and with the progression in her cough I am concerned that we may be dealing with congestive heart failure at this time. We have begun therapy to control congestive heart failure, support cardiac function, slow down the progression of the heart disease and improve survival. We are now seeing more dogs on specialized diets that are developing taurine deficiency and we have discussed submission of taurine levels to evaluate whether this may be a contributing factor to (b) (6) condition. (b) (6) is interested in pursuing this test at your clinic, taurine levels should be drawn and placed in a heparinized tube (green top) and should be frozen and submitted to (b) (4) (who sends it to UC Davis). It will be interesting to see if this is a contributing factor to (b) (6) condition.

We will continue to closely monitor (b) (6) heart disease via serial echocardiography and institute further therapy when progression is noted. While on this course of medication, it is important to monitor the chemistry profiles and blood pressures. Dogs with dilated cardiomyopathy are at a higher risk of developing ventricular arrhythmias. None were noted today; however, it will be important to monitor for arrhythmias periodically in the future. Unfortunately, the prognosis is guarded after the onset of congestive heart failure, and we discussed with the (b) (6) family that the average survival is ~ 6-12 months.^{1,2} Survival time is highly individually variable depending on response to therapy.

We appreciate your continued referrals and the trust you place in CVCA to co-manage your cardiac patients. We look forward to working with you on this case and others. In an effort to continue to improve CVCA's service to both you and your clients, please visit our website at www.cvcavets.com and complete our online referring veterinarian survey.

Sincerely,

(b) (6)

(b) (6)

Case Summary:

(b) (6) a 13 Yrs. 0 Mos. old, spayed female, Labrador Retriever presented on (b) (6) to the (b) (6) for a coughing.

History: (b) (6) started coughing last Wednesday. She was brought to a primary veterinarian. Radiographs and blood work were performed. Radiographs revealed suspected cardiomegaly. Blood work showed mild ALP and GGT elevations. Prescribed hydroxyzine, doxycycline, and hydrocodone, which was stopped on Monday because her coughing got worse with those medications. The owner made an appointment with a CVCA on Friday(11-1-2017). However her cough got worse with pink tinged foam so (b) (6) was brought to (b) (6) for a cardiology consultation. (b) (6) has been a healthy dog with no current medications. She is up to date on vaccination and heartworm preventative.

CBC (10-27-2017) WNL
Chem (10-27-2017) ALP 440, GGT 30, other values were WNL
OVA & Parasites (7-17-2017) Negative

Physical Exam:

	(b) (6)
	1:47 PM
Vital Sign	656
Weight	30.5 kilograms
Temp	100.5
HR	100
Resp	42
Muc_Me	Pink/Healthy
mb	
CRT	<2 sec
Mentation	QAR
Pain	0 - No visible Pain
Scale	

BCS: 5/9

EENT: MM- pink, mild calculus and gingivitis, CRT <2 sec. Oral exam- no significant findings (NSF), Lenticular sclerosis on OU, throat -NSF.

Hydration appears: within normal limits (WNL)

Peripheral lymph nodes: Palpate WNL.

Airway: RR= 30 BPM, no upper respiratory noise, airway not compromised.

Respiration: RR= 24 RPM, Eupneic with no crackles or wheezes. Bilateral breath sounds ausculted, normal bronchovesicular sounds.

Cardiovascular: HR = 100 BPM, Heart auscults with NSF. No murmurs noted. Femoral pulses are adequate and synchronous.

Abdomen: Mildly tensed cranial abdomen on palpation, no organomegaly was noticed,

Neurologic: Alert and responsive. Ambulatory with no CP deficits noted. Full neurologic examination was not performed.

Integument: Hair coat has NSF. A 3cm x 3 cm soft subcutaneous mass was palpated on left caudal abdomen.

Musculoskeletal: Musculature is WNL.No obvious lameness or gait disturbance.

Urogenital: WNL

Rectal: Normal stool was palpated on rectal examination.

Initial Diagnostics:

Echocardiogram

Differential Diagnosis:

Coughs -R/O heart vs lung

Client Communication:

Plan:

Please call if you have any questions or concerns.

Thank you,

(b) (6)

(b) (6)

10/31/2017 Initial PE

(b) (6)

(b) (6)

(b) (6)	Name:	Species:	Breed:	Color:	DOB:	SEX:
Date: 7/10/17	(b) (6)	Canine	Lab Retriever	Yellow	(b) (6)	F(s)
	72 lbs	Imrab 3 TE, 3 W, Tap #1580			(b) (6)	
	SATAP HW	Lyme	Echin	K9		
	Fecal to	(b) (6)	CAN NPS			
	DW @ home,	occasionally	coughs.	Several SQ masses - no	changes noted per	5.
	PE: BAR.	H/L NSF - no audible	murmur,	no cough elicited	on tracheal palp.	Abd NSF
	on palp.	Min tartar.	Several	SQ masses - palpate	upomatous	P: Due 10/12/17
	Lyme	HL vaccine	(b) (6)			
					#305581	226 ⁵⁰ 226 ⁵⁰ (b) (6)
	7/12/17	Disp: Advantix II Navy	(>55 lbs) (#1)	+ 2 free doses	Heartgard	Brown Lpk (#1)
					#305994	286 ⁷⁴ 286 ⁷⁴ (b) (6)
	10/23/17	Disp: Advantix II Navy	(>55 lb) Lpk	#1		(b) (6)
		H/G to share w/ Bon				
	10/27/17	65.5 lbs	✓ cough = x3-4d,	not lethargic,	eld 15,	no v/d, worse when laying down
	T 100.7°	PE: BAR.	Numerous upomatous & dermal masses.	MM pink, euhdrated,	min tartar.	Abd nonpainful on palp.
		H/L - no audible	murmur or arrhythmias,	no abn lung	sounds.	Breathing very shallow - thorax doesn't
		expand fully normally -	not diffuse, no	visible abd component	to breathing	P: CRP 140, ALP 440, GGT 30,
		lymphs very s/d,	NSF rest			
		P: Lat thorax xray.	heart appears sl large,			

(b) (6)

(b) (6)	Name:	Species:	Breed:	Color:	DOB:	SEX:
	(b) (6)	Canine	Lab Ret	Yellow	(b) (6)	F(S)
<p>Date: 10/27/17</p> <p>CONT: <u>Some ↑ opacity in ct-dorsal lung fields</u></p>	<p>Trachea JSF. VHS N P3</p> <p>VD thorax xray: R side of heart rounded, lung pattern less prominent on this view.</p> <p>Reviewed xrays w/ 5-disc bronchitis, lung fibrosis all age, cardiac, open.</p> <p>Disc R side of heart looks ↑, disc HWDC (at test 3 mos ago & on HUP), DEM/heart not abnrd), pericardial effusion (heart not abnrd), etc</p> <p>Rec await lab results, consider tx for allergic bronchitis, but re cardiac workup of not improving.</p> <p>Dog did not cough while in clinic except for tracheal cough when pulling on leash.</p> <p>called 5-disc GHP1, mild ↑ ALP - not likely a concern at this time.</p> <p>Rec med below, call in 3-5 d update.</p> <p>Disp: Hydroxyzine 50mg #20, 1 PO BID</p> <p>Hydrocodone 5mg #20, 1 PO Q8-12 PRN</p> <p>Doxycycline 100mg #40, 2 PO BID, 1 PO PM</p> <p>#(b) (6)</p>					

11/10/17 09:05:38

(b) (3) (B)

->

(b) (4) I Page 001

(b) (4)

(b) (6)

(b) (6)

Account: (b) (6)

Owner: (b) (6)
Patient: (b) (6)
Species: CANINE
Breed: LABRADOR_RETRIE
Age: 11Y
Gender: FS

Requisition #: (b) (6)
Accession #: (b) (6)
Order recv'd: 11/03/2017
Ordered by: (b) (6)
Reported: 11/10/2017

TAURINE (WHOLE BLOOD)

Test		Result		
TAURINE	168	(200 - 350)	L	<input type="checkbox"/>
Testing performed at University of California, Davis				

(b) (6)

11/10/2017

FINAL REPORT

PAGE 1 OF 1

CVCA, Cardiac Care for Pets

(b) (6)

Phone: (b) (6) Fax: (b) (6)
Email: (b) (6)@cvcavets.com
www.cvcavets.com



Client: (b) (6)
Co-owner:
Patient name: (b) (6)
Species: Canine
Breed: Labrador Retriever
Sex: FS
Age: 13 years and 5 months old
Weight: 33.18kg. / 73.15 lbs

Primary Care Veterinarian: (b) (6)
Primary Care Hospital: (b) (6)
Phone: (b) (6) ext: (b) (6)
Fax: (b) (6)
Email:

Cardiac Evaluation Report Exam Date: 10/31/2017

Diagnosis

- Advanced dilated cardiomyopathy - ruleout idiopathic vs. taurine-responsive
- Mild to moderate mitral valve regurgitation as cause of heart murmur
- Trace tricuspid valve regurgitation
- Moderate to severe left atrial chamber dilation
- Severe eccentric left ventricular chamber dilation
- Moderate to severe decrease in contractility/heart muscle function
- Mild left ventricular wall thinning
- Mild right atrial and right ventricular chamber dilation
- Progressive cough - rule out: early left sided congestive heart failure vs. mainstem bronchial compression

Medications

- Begin Lasix/Furosemide 40 mg tablets - Give 1 tablet twice daily.
 - > For mild increases in respiratory rate/effort, you may give an additional dose of Lasix.
 - > If you are consistently giving an additional dose of Lasix, please contact our office so we may help adjust medications long-term.
 - > We may increase this dose in the future based on at home monitoring of breathing and recheck blood work.
- Begin Benazapril 10 mg tablets - Give 1 tablet twice daily for 4 days then increase to 1 and 1/2 tablet twice daily thereafter.
- Begin Vetmedin/Pimobendan 5mg tablets - Give 1 and 1/2 tablets twice daily. Will switch to 7.5 mg EZ tablets at 1 tablet twice daily. The 7.5mg tablet will be compounded through (b) (6), please call them to set up shipping and billing (b) (6)
- Please call if you notice a decrease in appetite, vomiting, lethargy, weakness or any other signs of illness while beginning/adjusting the medications.
- Continue with monthly heartworm and flea/tick control as prescribed by (b) (6)

In 2 weeks, if (b) (6) is eating and feeling well:

- Begin Spironolactone 25 mg tablets - Give 1 tablet once daily for 4 days then increase to 1 tablet twice daily thereafter.

⟨ Begin Taurine 1500 mg twice daily.
⟨ Begin L-carnitine 1500 mg three times daily.
⟨ You may purchase the taurine and L-carnitine at any health food or nutrition store or www.puritanspride.com. You may also obtain the L-carnitine in bulk powder form from North Carolina State University by calling 919-513-6325.

Please allow 24-48 hours for CVCA to process prescription refill requests.

Refill all medications indefinitely unless directed by CVCA or your primary care veterinarian.

⟨ **Please check all medications and dosages on your discharge report against the pharmacy labels.**

Please Note

⟨ Please see our website www.cvcavets.com for more information about (b) (6) dilated cardiomyopathy.

Nutrition Recommendations:

⟨ (b) (6) is on a specialized diet which could be contributing to taurine deficiency. Please change her to a new diet, as her housemate is on a novel protein diet - consider prescription diets such as Royal Canin or Science Diet. Please discuss diet options with (b) (5)

⟨ In patients with early/mild heart failure, CVCA recommends feeding a diet with less than 80 mg of sodium per 100 kCal of food (50-80 mg/100 kCal). In patients with refractory heart failure signs, further sodium restriction may be beneficial.

⟨ For more information about sodium content of various foods, please visit:

- Dog: http://vet.tufts.edu/wp-content/uploads/reduced_sodium_diet_for_dogs.pdf
- Treats: http://vet.tufts.edu/wp-content/uploads/treats_for_dogs_with_heart_disease.pdf

⟨ CVCA recommends avoiding kidney diets unless (b) (6) has kidney disease that warrants protein restriction.

⟨ Diet changes should be done gradually (ie. over ~1 month) to avoid GI upset and avoided until (b) (6) is stable and eating well on the cardiac medications, usually about 2 weeks after starting or adjusting therapy.

⟨ If you are interested in a consultation with a veterinary nutritionist, please visit -- <http://vetnutrition.tufts.edu/make-an-appointment/>

⟨ CVCA recommends fish oil supplements (omega-3 fatty acids) in many dogs with cardiac disease. Her dose should be approximately EPA 1220 mg and DHA 760 mg total per day. Please start at 1/2 the dose for one week, then increase to the full dose if tolerating well thereafter. Please avoid Cod liver oil and flax seed as well as products with Vit A and/or D.

For more information about fish oils, please visit -- <http://vet.tufts.edu/heartsmart/diet/important-nutrients-for-pets-with-heart-disease/>

⟨ In addition to the supplements approved by Tuft's Veterinary Nutrition Service, other reputable brands include Welactin and Nordic Naturals. (b) (6) may have additional brand recommendations.

Activity Recommendations:

⟨ Keep (b) (6) very quiet for the next 3-4 days with only brief leash walks to eliminate.

⟨ Once her coughing has resolved, (b) (6) may gradually resume activity as she wants and is able to do. Please allow Lucy to take more breaks and rest during activity.

⟨ Please try avoid burst type activity, as this increases the arrhythmia risk and avoid exercise in the hot/humid weather.

⟨ Please try to warm (b) (6) up for 5-10 minutes with walking prior to moderate activity and take more rests during more vigorous activity.

At Home Monitoring:

⟨ Monitor for signs of cough, respiratory difficulty, exercise intolerance, abdominal swelling, weakness, lethargy, etc. If you note any of these symptoms, please notify CVCA or (b) (6) as these symptoms may indicate recurrent congestive heart failure. If you note an increase in cough, respiratory rate or effort, please feel free to give an additional dose of Lasix/Furosemide, while contacting CVCA.

⟨ In order to monitor for the development of early congestive heart failure in the out-patient setting, we recommend monitoring your pet's resting respiratory rate several times a week. Normal resting respiratory rates should be less than 30 breaths per minute. Consider using a respiratory rate monitoring application to track (b) (6) respiratory rate - Cardalis or BI Pharma have reliable phone applications. Please contact us if you note a persistent or progressive increase.

⟨ In addition, (b) (6) is sadly at increased risk for sudden cardiac death due to her cardiac disease. Dobermans are particularly at risk for development of severe, sudden malignant arrhythmias that sadly may result in sudden death. However, we hope to minimize these risks with our treatment plan.

Future Anesthesia/Fluid Recommendations:

- Avoid intravenous or subcutaneous fluid therapy in the future, if possible. If fluid therapy is indicated, please contact CVCA.
- (b) (6) should not receive corticosteroids (prednisone) in the future please contact CVCA for recommendations, if corticosteroids are indicated.
- Avoid elective anesthesia, as (b) (6) is at high risk for complications due to the degree of cardiac disease. If anesthesia is necessary in the future, please contact CVCA for recommendations for monitoring and anesthetics.

Reevaluation

- Please recheck with (b) (6) in the next day or two to obtain taurine levels. Please forward these results when available.
- Please recheck with (b) (6) in 2 weeks for a follow up examination and blood chemistry profile with electrolytes and as recommended by (b) (6) Please forward these results when available.
- Please recheck with (b) (6) every 4-6 months for a follow up examination and blood chemistry profile with electrolytes and as recommended by (b) (6) Please forward these results when available.
- Please recheck with CVCA in 5 months for a follow up consultation/examination, blood pressure, and echocardiogram. Please contact us or schedule an earlier appointment if (b) (6) has any problems or symptoms indicative of worsening heart disease or if recommended by (b) (6)

Visit Summary

Heart Rate: 132 bpm

BP: 100mmHg (based on MR gradient)

History:

(b) (6) developed a cough last Wednesday (10/25/17). Radiographs and blood work were performed by (b) (6). The lab work (which is unavailable for review) reportedly showed an elevated ALP 440 and GGT 30 and mild lymphopenia. Thoracic radiographs were performed which revealed cardiomegaly. (b) (6) was treated with hydroxyzine 50mg BID, doxycycline 200mg AM and 100mg PM, and hydrocodone 5mg q8-12h. All medications were stopped on Monday as her cough had worsened and she was presented to the (b) (6) for a cardiac evaluation as her coughing had worsened and she had brought up a small volume of pink-tinged foam after a coughing fit. During this time there has been no evidence of lethargy and she continues to eat and drink normally at home.

PPHx: None

Meds: None

Other: UTD on vaccinations, On HW preventative

Diet: Zignature (Kangaroo)

Physical Exam Findings:

BAR, sweet but nervous

OP/EENT: Pink, moist mucous membranes, CRT <2s, mild periodontal disease, LS OU, clear AU, No nasal or ocular discharge, no cough on tracheal palpation

PLN: WNL

H/L: Grade 2/6 left apical protosystolic heart murmur, regular rhythm, strong synchronous femoral pulses, RR: 36 breaths/min, questionable mild increase in bronchovesicular sounds bilaterally, no crackles or wheezes ausculted, eupneic

Abd: Soft non-painful abdominal palpation, no palpable masses or fluid wave

MS/Neuro: BCS 5/9, Amb x 4, Mentally alert and appropriate

Integ: Normal turgor, subcutaneous mass left ventrum

Other Diagnostics:

10/27/17 pDVM CXR: Generalized cardiomegaly characterized by widening of the cardiac silhouette and loss of the caudal cardiac waist consistent with left atrial enlargement. Slight left auricular bulge. Increased sternal contact and rounding of the right heart on the VD radiograph. Dorsal deviation of the trachea. Prominent pulmonary vasculature with a questionable mild increase in interstitial opacity in the caudodorsal lung fields which may suggest early congestive heart failure/pulmonary edema.

Echocardiographic Findings

Severe left ventricular eccentric hypertrophy with apical rounding and increased sphericity, mild-moderate centrally

FDA-CVM-FOIA-2019-1704-000080

located mitral regurgitant jet, moderate-severe secondary left atrial dilation on 2D imaging and moderately-severely increased LA:Ao ratio on M-mode imaging, mild eccentric low velocity tricuspid regurgitation with mildly elevated estimated right ventricular pressures consistent with mild pulmonary hypertension, mild right ventricular and right atrial dilation, normal left and right ventricular outflow velocities, moderately to severely depressed indices of systolic function (FS% and EF% by modified Simpson's - LVDI 144ml/m², LVSI 90ml/m²), increased EPSS, elevated transmitral inflow velocities and E:A wave ratio on spectral Doppler tracings, normal TDI E':A' ratio of the lateral mitral annulus, no masses, effusions or heartworms observed.
ECG during echocardiogram: Normal sinus rhythm. No ventricular ectopy noted.

Comments

Dear (b) (6),

Thank you for sending (b) (6) to see us with (b) (6) today. Sadly (b) (6) has dilated cardiomyopathy with moderate to severe systolic dysfunction and moderate to severe left atrial dilation. This places her at a high risk of developing congestive heart failure and with the progression in her cough I am concerned that we may be dealing with congestive heart failure at this time. We have begun therapy to control congestive heart failure, support cardiac function, slow down the progression of the heart disease and improve survival. We are now seeing more dogs on specialized diets that are developing taurine deficiency and we have discussed submission of taurine levels to evaluate whether this may be a contributing factor to (b) (6) condition. (b) (6) is interested in pursuing this test at your clinic, taurine levels should be drawn and placed in a heparinized tube (green top) and should be frozen and submitted to (b) (4) (who sends it to UC Davis). It will be interesting to see if this is a contributing factor to (b) (6) condition.

We will continue to closely monitor (b) (6) heart disease via serial echocardiography and institute further therapy when progression is noted. While on this course of medication, it is important to monitor the chemistry profiles and blood pressures. Dogs with dilated cardiomyopathy are at a higher risk of developing ventricular arrhythmias. None were noted today; however, it will be important to monitor for arrhythmias periodically in the future. Unfortunately, the prognosis is guarded after the onset of congestive heart failure, and we discussed with the (b) (6) family that the average survival is ~ 6-12 months.^{1,2} Survival time is highly individually variable depending on response to therapy.

We appreciate your continued referrals and the trust you place in CVCA to co-manage your cardiac patients. We look forward to working with you on this case and others. In an effort to continue to improve CVCA's service to both you and your clients, please visit our website at www.cvcavets.com and complete our online referring veterinarian survey.

Sincerely,

(b) (6)

CVCA, Cardiac Care for Pets

(b) (6)
Phone: (b) (6) Fax: (b) (6)
Email: (b) (6)@cvcavets.com
www.cvcavets.com



Client: (b) (6)
Co-owner:
Patient name: (b) (6)
Species: Canine
Breed: Labrador Retriever
Sex: FS
Age: 13 years and 5 months old
Weight: 33.18kg. / 73.15 lbs

Primary Care Veterinarian: (b) (6)
Primary Care Hospital: (b) (6)
Phone: (b) (6) ext: (b) (6)
Fax: (b) (6)
Email:

Cardiac Evaluation Report Exam Date: 02/26/2018

Diagnosis

- Mild, improved dilated cardiomyopathy - suspect taurine-responsive
- Mild, improved mitral and very mild tricuspid valve regurgitation as cause of heart murmur
- Normal, improved left atrial chamber dilation
- Mild, improved eccentric left ventricular chamber dilation
- Low normal, improved left ventricular contractility/heart muscle function
- Cough - suspect bronchial/primary respiratory disease

Medications

- Decrease Lasix/Furosemide 40 mg tablets - Give 1 and 1/2 tablets twice daily for 1 week then decrease to 1 tablet twice daily for 1 week then decrease to 1/2 tablet twice a day for 1 week then discontinue. Please call if you note an increase respiratory rate while decreasing the Lasix. If there is an increase in cough (but normal respiratory rate), we will consider adding in a bronchodilator.
- Continue Benazapril 10 mg tablets - Give 1 and 1/2 tablets twice daily
- Continue Vetmedin/Pimobendan 7.5 mg EZ tablets - Give 1 tablet twice daily.
- Continue Spironolactone 25 mg tablets - Give 1 tablet twice daily.
- Continue Taurine 1500 mg twice daily.
- Continue L-carnitine 1500 mg three times daily.
- You may purchase the taurine and L-carnitine at any health food or nutrition store or www.puritanspride.com. You may also obtain the L-carnitine in bulk powder form from North Carolina State University by calling 919-513-6325.
- Continue with monthly heartworm and flea/tick control as prescribed by (b) (6)

Please allow 24-48 hours for CVCA to process prescription refill requests.

Refill all medications indefinitely unless directed by CVCA or your primary care veterinarian.

- **Please check all medications and dosages on your discharge report against the pharmacy labels.**

Please Note

- Please see our website www.cvcavets.com for more information about (b) (6) dilated cardiomyopathy.

Nutrition Recommendations:

- Continue the Royal Canin Early Cardiac diet.
 - Consider fish oil supplements (omega-3 fatty acids). Her dose is approximately EPA 1220 mg and DHA 760 mg total per day. Please start at 1/2 the dose for one week, then increase to the full dose if tolerating well thereafter. Please avoid Cod liver oil and flax seed as well as products with Vit A and/or D.
- For more information about fish oils, please visit -- <http://vet.tufts.edu/heartsmart/diet/important-nutrients-for-pets-with-heart-disease/>
- In addition to the supplements approved by Tuft's Veterinary Nutrition Service, other reputable brands include Welactin and Nordic Naturals. (b) (6) may have additional brand recommendations.

Activity Recommendations:

- Continue normal activity as she wants and is able to do. Please allow (b) (6) to take more breaks and rest during activity.
- Please avoid exercise in the hot/humid weather.

At Home Monitoring:

- In order to monitor for the development of early congestive heart failure in the out-patient setting, we recommend monitoring your pet's resting respiratory rate several times a week. Normal resting respiratory rates should be less than 30 breaths per minute. Consider using a respiratory rate monitoring application to track (b) (6) respiratory rate - Cardalis or BI Pharma have reliable phone applications. Please contact us if you note a persistent or progressive increase.

Future Anesthesia/Fluid Recommendations:

- We expect (b) (6) to tolerate carefully monitored general anesthesia with normal preoperative bloodwork and a balanced anesthetic regimen. During anesthesia, we recommend careful monitoring of ECG, BP and pulse ox and 1/2 usual surgical fluid rate (ie: 2-4 ml/kg/hr). Carefully monitor for several hours post-operatively for signs of respiratory congestion and consider chest radiographs if these signs occur. There is some risk associated with all anesthetic events.
- Avoid medications with tachycardia as a side effect, such as ketamine, telazol and glycopyrrolate. Cleared for low dose atropine if needed for intraprocedure bradycardia. Avoid medications that significantly alter blood pressure such as acepromazine and Domitor.
- (b) (6) should not receive corticosteroids (prednisone) in the future please contact CVCA for recommendations, if corticosteroids are indicated.

Reevaluation

- Recheck with (b) (6) in the next 2-4 weeks and every 6 months for wellness care as directed, close auscultation, blood pressure and complete lab tests including blood and urine testing (CBC/Chemistry/Urinalysis/Thyroid evaluation). Please forward these results when available.
- Please recheck with CVCA in 6 months for a follow up consultation/examination, blood pressure, and echocardiogram. Please contact us or schedule an earlier appointment if (b) (6) has any problems or symptoms indicative of worsening heart disease or if recommended by (b) (6)

We thank you for trusting in CVCA to care for (b) (6) today. Please do not hesitate to call us with any questions or concerns.

Sincerely,

(b) (6)

Visit Summary

Heart Rate: 130 **BP:** 155 mmHg **Cuff Size/Location:** 6 cuff/LF

History: Recheck DCM, suspected early CHF; doing well; RRR - 16 bpm, increased Lasix in January due to increased cough; cough seems to be intermittent and related to excitement; good appetite; 3 kg weight gain since 10/2017; walks 30-45 minutes per day - slow pace, at times winded but recovers very quickly.

(b) (6) developed a cough last Wednesday (10/25/17). Radiographs and blood work were performed by (b) (6). The lab work (which is unavailable for review) reportedly showed an elevated ALP 440 and GGT 30 and mild lymphopenia. Thoracic radiographs were performed which revealed cardiomegaly. (b) (6) was treated with hydroxyzine 50mg BID, doxycycline 200mg AM and 100mg PM, and hydrocodone 5mg q8-12h. All medications were stopped on Monday as her cough had worsened and she was presented to the (b) (6) for a cardiac evaluation as her coughing had worsened and she had brought up a small volume of pink-tinged foam after a coughing fit. During this time there has been no evidence of lethargy and she continues to eat and drink normally at home.

PPHx: None

Meds: None

Other: UTD on vaccinations, On HW preventative

Diet: changed from Zignature (Kangaroo) to Royal Canin Early Cardiac

Physical Exam Findings: 3/6 pansystolic murmur, PMI - mitral valve, regular rhythm with S3 gallop; LUNGS - clear all fields, panting, normal effort; Sl. overweight body condition (BCS - 6/9); Pink mm; PP - SS; PLN - WNL; ABD - hepatomegaly; BAR

Echocardiographic Findings

Mild left ventricular eccentric dilation - significant improvement compared to previous exam; mild, improved centrally located mitral regurgitant jet, normal, improved left atrial dimensions on 2D imaging and on M-mode imaging, mild, low velocity eccentric low velocity tricuspid regurgitation, subjectively normal right ventricular and right atrial dimensions, normal left and right ventricular outflow velocities, low normal, improved indices of systolic function (FS% and EF% by modified Simpson's, normal EPSS, normal transmitral inflow velocities and E:A wave ratio on spectral Doppler tracings, normal TDI E':A' ratio of the lateral mitral annulus, no masses, effusions or heartworms observed.

Comments

Dear (b) (6),

Thank you for sending (b) (6) to see us with (b) (6) today. I am quite pleased with (b) (6) exam today. She has had remarkable improvement in her echocardiogram with the cardiac medications, change in diet and supplementation with Taurine and L-carnitine. Her risk for congestive heart failure at this point is very low so we will be weaning (b) (6) off the Lasix/furosemide while (b) (6) monitors (b) (6) respiratory rate. Her current cough is likely due to respiratory disease and if the cough progresses/worsens, we will consider adding in a bronchodilator, such as Theophylline. Right now, with the marked improvement, (b) (6) long-term prognosis has improved considerably. I suspect we will be able to further discontinue cardiac medications if her heart remains stable. We will continue to closely monitor (b) (6) heart disease via serial echocardiography and institute further therapy when progression is noted. While on this course of medication, it is important to monitor the chemistry profiles and blood pressures. Hopefully, (b) (6) will continue to do so well - she's a sweetie!

We appreciate your continued referrals and the trust you place in CVCA to co-manage your cardiac patients. We look forward to working with you on this case and others. In an effort to continue to improve CVCA's service to both you and your clients, please visit our website at www.cvcavets.com and complete our online referring veterinarian survey.

Sincerely,

(b) (6)

11/10/17 09:05:38

(b) (4)

->

(b) (4) Page 001

(b) (4)

(b) (6)

Owner: (b) (6)
Patient: (b) (6)
Species: CANINE
Breed: LABRADOR_RETRIE
Age: 11Y
Gender: FS

(b) (6)

Account: (b) (6)

Requisition #: (b) (6)
Accession #: (b) (6)
Order recv'd: 11/03/2017
Ordered by: (b) (6)
Reported: 11/10/2017

(b) (6)

TAURINE (WHOLE BLOOD)			
Test	Result		
TAURINE	168	(200 - 350)	L
Testing performed at University of California, Davis			

(b) (6)

11/10/2017

FINAL REPORT

PAGE 1 OF 1

Vet-LIRN Final Case Report

A. Case Identification:

Case Number: 800.218

Vet-LIRN Director: Renate Reimschuessel, VMD, PhD

Program: Vet-LIRN

Division Code: HFV – 500

Other Investigators:

Jennifer Jones, DVM	Vet-LIRN
Sarah Nemser, MS	Vet-LIRN
Olgica Ceric, DVM, PhD	Vet-LIRN
Jake Guag, MPH	Vet-LIRN
David Rotstein, DVM, MPH	OS&C CERT
Lee Anne Palmer, VMD, MPH	OS&C DVPS
Lauren Carey, DVM	OS&C DVPS

B. Descriptive Title of Case:

Investigation of two dogs with dilated cardiomyopathy after consuming California Natural Venison and Green Lentil food and California Natural Kangaroo and Lentil dog foods.

Address of Vet-LIRN Program Office:

Mod II
Center for Veterinary Medicine
Office of Research
8401 Muirkirk Road
Laurel, MD 20708

C. Initiation and Completion Date:

Initiation Date: 7/13/2017
Completion Date: 8/22/2017
Final Report Submission Date: 11/1/2017

Case Summary

Complaint: July 13, 2017, Vet-LIRN received consumer complaint, EON-323515, reporting dilated cardiomyopathy in two dogs after consuming California Natural Venison and Green Lentil food and California Natural Kangaroo and Lentil dog foods.

Signalment:

- (b)(6) 7 yr MC Miniature Schnauzer
- (b)(6) 2 yr MC Miniature Schnauzer-deceased

Signs: syncopal episodes, dyspnea, cough, heart failure

Medical Records: Vet-LIRN collected and reviewed medical records.

Name	Clinical Signs	Physical Exam	Lab Work	Significant Medical History
(b)(6)	syncopal episodes, hyporexia	P 130 bpm, mild increased breath sounds-all lung fields	suspected DCM, taurine & carnitine normal; negative infectious disease & nutritional disease testing	
(b)(6)	dyspnea, cough, inappetance, regurgitation,	P 160 bpm, R 64 rpm, pale pink mm, Gr I-II/VI left apical systolic murmur; hypokinetic, synchronous femoral pulse, jugular venous distention	P 11.7, BG 225, ALT 147, AST 1006, CK 35,930; Toxic NP, Plt 97; hepatomegaly, biventricular heart failure, cardiogenic edema; <u>Necropsy:</u> Suspect primary non-cardiogenic etiology	coffee brown urine with clumps after strenuous activity & hot outside-resolves with 24-36 hours; Crystalluria

Owner Interview: Vet-LIRN did not conduct an owner interview. However, the veterinarian mentioned:

- The owner alternated feedings between the two products
- The owner did not feed anchovies, sardines, or seafood in February or chronically
- The two dogs were from genetically different lineages
- (b)(6) had clinical signs at the time (b)(6) was treated but didn't present with CHF for several months

Response: Vet-LIRN collected medical records for review and leftover open product (Kangaroo flavor) for taurine, carnitine, and fumonisin testing.

Results: The food tested negative for fumonisin. The food taurine level (0.26% estimated Dry Matter Basis) was above the minimum level in cats (no AAFCO minimum for dogs). The food carnitine level is 0.0077% estimated on a Dry Matter Basis. There is no AAFCO carnitine minimum for dogs or cats. It is unclear whether or not the food carnitine is low, normal, or high.

Conclusion: Dilated cardiomyopathy (DCM) can be caused by a variety of etiologies including, genetic² (breed related), toxic^{3,4} (fumonisin, acrolein⁵, domoic acid, doxorubicin, lily of valley, digitalis, ionophores, sicklepod, gossypol, white snake root, ethyl alcohol, foxglove, buttercups), infectious (Bartonellosis, *Trypanosoma cruzi*), and nutritional deficiency¹ (e.g. taurine, protein restricted diets with stones, carnitine deficiency). The two genetically unrelated dogs were fed the same foods and began to experience clinical signs approximately the same time. The medical records indicate infectious disease and nutritional deficiency are unlikely etiologies. (b)(6) records indicated elevated liver enzymes and CK values, which could support a hepatotoxic and myotoxic (cardio +/- muscle) exposure. Because (b)(6) presented six months after (b)(6), it is unknown if (b)(6) also had elevated liver enzymes when (b)(6) was ill. The history also suggested no exposure to doxorubicin or domoic acid. Vet-LIRN tested the leftover bag of food from (b)(6) illness time (June 2017), but not from January, when both dogs were initially ill. A test for acrolein was not available.

The cause of the two dogs' DCM is unclear, but is likely an environmental toxin exposure. Based on the dogs' blood taurine/carnitine levels and the dry dog food test results, it is unlikely that Fumonisin, taurine, or carnitine levels in the food caused the dogs' illness.

References:

1. Sanderson SL. Taurine and Carnitine in Canine Myopathy. *Vet Clin Small Anim* 36 (2006) 1325–1343.
2. Borde D, Calvert CA, Darien BJ, Guerrero J, and Wall M. Acquired Heart and Blood Vessel Disorders in Dogs. *Merck Veterinary Manual*. Found at: <http://www.merckvetmanual.com/dog-owners/heart-and-blood-vessel-disorders-of-dogs/acquired-heart-and-blood-vessel-disorders-in-dogs>
3. Valberg SJ. Toxic Myopathies in Ruminants and Pigs. *Merck Veterinary Manual*. Found at: <http://www.merckvetmanual.com/musculoskeletal-system/myopathies-in-ruminants-and-pigs/toxic-myopathies-in-ruminants-and-pigs>
4. Garland T. Overview of Gossypol Poisoning. *Merck Veterinary Manual*. Found at: <http://www.merckvetmanual.com/toxicology/gossypol-poisoning/overview-of-gossypol-poisoning>
5. Ismahil MA, Hamid T, Haberzettl P, Gu Y, Chandrasekar B, Srivastava S, Bhatnagar A, and Prabhu SD. Chronic oral exposure to the aldehyde pollutant acrolein induces dilated cardiomyopathy. *Am J Physiol Heart Circ Physiol* 301: H2050–H2060, 2011.

Supplemental Information:

01-800.218-EON-323515-(b)(6)-CC: Consumer Complaint
 02-800.218-EON-323515-(b)(6)-MedRec: Medical Records
 03-800.218-EON-323515-(b)(6)-Results: Testing Results
 04-800.218-EON-323515-(b)(6)-Summary: Vet-LIRN Summary

SIGNATURES

Mary E. Allen -S Digitally signed by Mary E. Allen -S
DN: c=US, o=U.S. Government, ou=HHS, ou=FDA, ou=People,
cn=Mary E. Allen -S, 0.9.2342.19200300.100.1.1=1300365061
Date: 2017.11.17 14:21:13 -05'00'

Deputy Director OR

Date

John
Graham -S Digitally signed by John Graham -S
DN: c=US, o=U.S. Government, ou=HHS,
ou=FDA, ou=People, cn=John Graham -
S,
0.9.2342.19200300.100.1.1=2001387754
Date: 2017.11.17 18:07:25 -05'00'

Director OR

Date

(b) (6)

Digitally signed by Renate Reimschuessel -S
DN: c=US, o=U.S. Government, ou=HHS, ou=FDA,
ou=People, 0.9.2342.19200300.100.1.1=1300140413,
cn=Renate Reimschuessel -S
Date: 2017.11.20 14:27:01 -05'00'

Vet-LIRN Director

Date

		(b) (4)	6979064	6979065	6979066	6979067	6979068	6979069	6979070	6979071	6979072	6979073	6979074
		Client ID	800.218-sub 1	800.218-sub 2	800.218-sub 4	800.218-sub 5	800.218-sub 6	800.238-sub 1	800.238-sub 1	800.238-sub 3	800.238-sub 3	800.219-sub 5	800.219-sub 6
		Sample Description	dog food	dog food	dog food	dog food	dog food	dog treat piece 1	dog treat piece 2	dog treat piece 1	dog treat piece 2	dog treat	dog treat
Component	Unit												
Chloride	%							1.39	0.589	0.70	0.33	0.078	1.15
Taurine	mg/g			1.06	1.84	1.08	1.22						
Methionine	mg/g		5.78	5.53	4.76	6.20	7.78						
Cystine	mg/g		2.32	2.31	3.15	3.20	2.50						

(b) (4)

Report Number: 1894242-0

Report Date: 15-Aug-2017

Report Status: Final

Certificate of Analysis

Food and Drug Administration - CVM - Invoice Denise Durham

8401 Muirkirk Rd.
Laurel Maryland 20708 United States

Sample Name:	800.218	(b) (4) Sample:	6406524
Project ID	FDA_CVM-20170804-0007	Receipt Date	04-Aug-2017
PO Number	HHSF223201610005I/HHSF22301002T	Receipt Condition	Ambient temperature
Sample Serving Size	100 g	Login Date	04-Aug-2017
		Online Order	20

Analysis	Result
L-Carnitine *	
L-Carnitine	69900 ppb
Taurine	
Taurine	231 mg/Serving Size

Method References	Testing Location
-------------------	------------------

L-Carnitine (CARNITNE_S)	(b) (4)
STAREY ET AL.: JOURNAL OF AOAC INTERNATIONAL VOL. 91, NO.1, 2008. (Modified).	

Taurine (TAUR_LC_S)	(b) (4)
R. Schuster, "Determination of Amino Acids in Biological, Pharmaceutical, Plant and Food Samples by Automated Precolumn Derivatization and HPLC", Journal of Chromatography., 1988, 431, 271-284, Henderson, J.W., Ricker, R.D. Bidlingmeyer, B.A., Woodward, C., "Rapid, Accurate, Sensitive, and Reproducible HPLC Analysis of Amino Acids, Amino Acid Analysis Using Zorbax Eclipse-AAA columns and the Agilent 1100 HPLC," Agilent Publication, 2000, and Barkholt and Jensen, , "Amino Acid Analysis: Determination of Cysteine plus Half-Cystine in Proteins after Hydrochloric Acid Hydrolysis with a Disulfide Compound as Additive," Analytical Biochemistry, 177, 318-322 (1989).	

Testing Location(s)	(b) (4)
---------------------	---------

(b) (4)

(b) (4)



These results apply only to the items tested. This certificate of analysis shall not be reproduced, except in its entirety, without the written approval of (b) (4)

* This analysis is not ISO accredited.

UPS/FedEx Package Information Form

Sender (address, tel #, fax #, e-mail):

Jake Guag, M.P.H, C.P.H.
Biologist
U.S. Food & Drug Administration
Center for Veterinary Medicine
Office of Research
Vet-LIRN
8401 Muirkirk Road.
Laurel, Maryland 20708
Tel: 240-402-0917
email: Jake.Guag@fda.hhs.gov

Recipient (name, address, tel. #, fax #, e-mail):

Attn: Dr. Darcy Adin
North Carolina State University
NC State Veterinary Hospital
1060 William Moore Drive
Raleigh, NC 27607
919-513-6032
Email: dbadin@ncsu.edu

Service (Standard, Priority Overnight, etc.):

Priority overnight

Date:

01/10/2018

Special Needs (Dry Ice, Hazardous Chemicals, etc.). Dry ice shipments must list the weight of the dry ice separately	
Weight of Dry Ice (kg, lbs)	NO
Total Weight of Package (kg, lbs.):	1.7 lbs
Dimensions of Package (L*W*H):	12x12x10 inches
Value:	/
Content Description	Food collection kit

800.218

We are collecting dog food-weight 0.36 kg. In a plastic tupperware container--~5" x 5" x 2" (0.36kg = ~0.8 lb)

No hazardous materials.

Room temperature.



Taurine Deficiency Induced Dilated Cardiomyopathy in Golden Retrievers

Taurine Deficiency Induced Dilated Cardiomyopathy in Golden Retrievers

by Janet Olson, DVM, DACVIM (Cardiology)

Dilated Cardiomyopathy (DCM) is becoming more prevalent in golden retrievers. Dr. Joshua Stern, DVM, PhD, DACVIM (Cardiology) at UC Davis, starting seeing a pattern and recognized that many cases were due to dietary taurine deficiency in golden retrievers fed grain free diets. Here is what we know so far:

Background

Taurine is an amino acid that is found in high concentrations in heart and muscle. Among its many functions, it aids in normal contractile function. Evidence shows that taurine helps mediate calcium channel transports and modulates calcium sensitivity of the myofibrils.

Taurine deficiency as a cause of dilated cardiomyopathy (DCM) is not a new issue. Taurine deficiency in cats was characterized by Pion et al in the late 1980s. Taurine deficiency has since been characterized as a cause of acquired DCM in dogs as well.

Currently identified diets of concern in these golden retrievers

According to Dr. Stern, the majority of cases they are seeing at UC Davis are from grain free diets that are high in legumes, like acana pork and squash singles.

What can we do? Some Guidelines.

- **ASK:** Make sure to ask your clients (whether they own golden retrievers or not) what diets they are currently or previously have fed their dogs
- **INFORM:** Inform your clients of his issue
- **ACT:** If they are currently on, or have been on grain free diets in the past, submit baseline WHOLE blood taurine levels and AFTER submitting the WHOLE blood taurine levels, switch diets if indicated. Temporary taurine supplementation may be necessary. If levels are low, take baseline chest films, if cardiomegaly noted on the radiographs, an echocardiogram is indicated to complete your baseline evaluation. Additional therapy may be indicated.
- **GET MORE INFORMATION:** Dr. Stern has compiled many of the studies in the following links: <https://www.dropbox.com/.../AAB1sDvLZe6gE3httPskz9-0a...>
[Taurine Deficient DCM in Dogs](#)

Medical Record Review:

(b) (6)

Presenting complaint (b) (6): dyspnea, cough of 3 week duration-wheezing type more frequent at night → rDVM, treated w/ prednisone and doxycycline for kennel cough → (b) (6) inappetance, vomiting → (b) (6) dyspneic and recheck, hospitalized and treated for pneumonia, regurgitated → (b) (6) treated as outpatient, (b) (6) as syring feeding, dog regurgitated and had marked dyspnea → ER → refer to NCSU → (b) (6) put on mechanical ventilator → (b) (6) euthanized

PE (b) (6): P 160 bpm, R 64 rpm, pale pink mm, Gr I-II/VI left apical systolic murmur, femoral pulse hypokinetic but synchronous, jugular venous distention

Labwork:

(b) (6)

Labs: unremarkable (unclear what was done)

(b) (6)

Big 4: Glu 135, Azo 15-20

(b) (6) vBGA: Lact 2.4, rest wnl

(b) (6) Chem: P 6.2, K 4.9, Na 140, TP 4.2

(b) (6) Chem: BG 225, BUN 29, P 11.7, K 3.3, Cl 95, Na 144

(b) (6) Chem: BG 136, P 4.6, CK 13,621, K 4.3, Na 151, Cl 109, AST 577

(b) (6) Chem: BG 165, BUN 37, P 8.1, ALT 147, AST 1006, CK 35,930, Na 135, K 3.8, Cl 90

(b) (6)

CBC: WBC 9.4, NP 7.9, Band .18, Plt 157

-2/4: WBC 9.9, NP 8, Band .7

-2/5: WBC 6.8, TP 6.9, NP 4.2, Ban .54, Toxic NP-mild, Plt 97

(b) (6)

BP sys: 90

(b) (6)

UA post Lasix: 1.011

Cardiac troponin 0.79

BAP GM-pending

Vector borne panel: pending

Taurine/Carnitine: pending

(b) (6)

Coag: PT 9.1, PTT 14, Dimer 189, Fib 539, INR 1.09

Urine Creat: 27.9

Urine Na: pending

ECG: suspected atrial tachycardia

Rads (b) (6): concern for aspiration pneumonia

(b) (6) cardiomegaly, severe diffuse mixed interstitial to alveolar pattern most severe caudo-dorsally, hepatomegaly, dec abdominal serosal contrast

(b) (6) severe generalized cardiomegaly with biventricular heart failure, improved vs rDVM rads

(b) (6) worsening cardiogenic pulmonary edema, cannot exclude lung induced injury

(b) (6) +/- pneumonia

(b) (6) post ultrafiltration, improved cardiogenic edema, hypovolemia, residual interstitial to patchy alveolar

(b) (6) improved CHF with possible concern for bronchopneumonia, suspected hiatal hernia

(b) (6) markedly progressive alveolar pattern with significantly worse cardiogenic edema

tFAST (b) (6): severe cardiomegaly with ventricular hypocontractility

Echo (b) (6): dcm vs. myocarditis vs pacing induce vs. other (severely dilated & hypocontractile left & right ventricles, severely dilated left and right atria)

Necropsy: Lung-severe diffuse alveolar injury with marked fibrin deposition (hyaline) and marked alveolar histiocytosis and multifocal type II pneumocyte hyperplasia; mod to marked diffuse pulmonary edema; mild cardiomegaly with mild mitral valve endocardiosis and mild left ventricular hypertrophy and left atrial dilation; thorax with mild pleural effusion; Suspect primary non-cardiogenic etiology but if clinical cardiac dysfunction then functional cardiac abnormalities cannot be ruled out

Prior MHx: coffee brown urine including clumping after strenuous activity when it is hot outside and resolves with 24-36 hours; also Crystalluria

(b) (6)

Presented 6/22/2017: episodes of collapse, first occurred mid February, fall 6 seconds without losing consciousness → immediately return to normal → 2 weeks later again collapse, then on → 6/3 post 2 hour hike collapsed again; panting more than usual; good appetite for treats but reluctant to eat food since February; → recheck 7/10, doing better, no collapsing episodes except a stumbling moment when excited, respiratory rate normal, diet changed to Hill's

6/22 PE: P 130 bpm, R pant, mild increased breath sounds in all lung fields

-7/10: T 99.7F, P 136 bpm, R 36 rpm, equivocal mild dehydration <5%, Gr II/VI left apical systolic murmur

Labs: 6/22 **Big 4:** BG 64 (recheck 79), BUN 15-26

BP-sys: 130 mmHg

-7/10: 110 mmHg

EKG: left ventricular enlargement suggested

UA: 1.019

Taurine & Carnitine: normal (no values)

Vector borne panel (PCR and IFA): normal

BAP GM

Troponin 1

T4

Toxoplasma/Neospora

Chagas

Complete AA: no significant abnormalities, consulting with UC Davis

-7/10 **Renal Panel:** K 4.0

6/22 Rads: left sided congestive heart failure

-7/10: moderate left sided cardiomegaly without heart failure, moderate hepatomegaly

6/22 Echo: mitral valve endocardiosis with left atrial enlargement and heart failure, decreased left ventricular systolic function, suspected DCM

		800.218-sub 1	800.218-sub 2	800.218-sub 6	
		Case Sample	Storebought	Case sample	Label
		California Naturals Kangaroo & Lentil	California Naturals Kangaroo & Lentil	California Naturals Kangaroo & Lentil	Product Nutrient Analysis (website label)
(b) (4)	Ca	1.30%	1%	0.93%	0.83%
	Mg	0.13%	0.14%	0.15%	0.17%
	P	0.74%	0.67%	0.68%	0.71%
	Fe	30 mg/kg	30 mg/kg	31 mg/kg	305 mg/kg
	Co	0.12 mg/kg	0.14 mg/kg	.14 mg/kg	n/a
	Cu	21 mg/kg	19 mg/kg	16 mg/kg	13.61 mg/kg
	Zn	240 mg/kg	280 mg/kg	200 mg/kg	193.37 mg/kg
	Se	0.7 mg/kg	0.65 mg/kg	.68 mg/kg	0.08 mg/kg
	Ca:P	1.76:1	1.49:1	1.37:1	
	Cu:Zn	0.09:1	0.07:1	0.08:1	
(b) (4)	Tau	~0.26%	1.06 mg/g = ~0.11%	1.22 mg/g = ~0.12%	
	Cystine	2.32 mg/g = ~0.23%	2.31 mg/g = ~0.23%	2.5 mg/g = ~0.25%	
	Met	5.78 mg/g = ~0.58%	5.53 mg/g = ~0.55%	7.78 mg/g = ~0.78%	0.61%
	Met-Cys	~0.81%	~0.78%	~1.03%	0.97%

AAFCO	
AAFCO-Adult Maint	Issues
0.5 to 2.5%	none
0.06%	none
0.4 to 1.6 %	none
40 mg/kg	below AAFCO & Label
25 mg/kg-chicks/rats/sheep max	unlikely
7.3 mg/kg	none
80 mg/kg	none
0.35 to 2 mg/kg	label should be higher
1:1 to 2:1	none
0.09:1-not AAFCO	none
0.1% in Cats	
n/a	
0.33%	none
0.65%	none

<http://www.californianaturalpet.com/products/1741>

to align w/ AAFCO maintenance claim

		800.218-sub 4		
		Case Sample		
		Fromm Heartland Gold Grain Free Large Breed Adult	Product Typical Analysis (website label)	AAFCO Growth & Maint
(b) (4)	Ca	1.20%	1.14%	1.2 to 1.8%
	Mg	0.14%	0.17%	0.06%
	P	1%	1.08%	1 to 1.6%
	Fe	30 mg/kg	258.26 mg/kg	88 mg/kg
	Co	0.37 mg/kg	n/a	25 mg/kg-chicks/rats/sheep max
	Cu	25 mg/kg	25.83 mg/kg	12.4 mg/kg
	Zn	170 mg/kg	217.37 mg/kg	100 mg/kg
	Se	0.85 mg/kg	n/a	0.35 to 2 mg/kg
	Ca:P	1.2:1		1:1 to 2:1
	Cu:Zn	0.15:1		0.09:1-not AAFCO
	(b) (4)	Tau	1.84 mg/g = ~0.18%	n/a
Cystine		3.15 mg/g = ~0.32%	n/a	n/a
Met		4.75 mg/g = ~0.48%	n/a	0.35%
Met-Cys		~0.79%	n/a	0.70%
MSU	Iodione	1.58 ug/g (ppm)		

Issues	https://frommfamily.com/products/dog/gold/dry/heartland-gold-large-breed-adult/tyr
label should be higher to align w/ AAFCO growth claim	
none	
none	
below AAFCO & Label	
unlikely	
none	
none	
none	
none	
none	
none	
none	

[jical-analysis/](#)

UPS CampusShip: View/Print Label

1. Ensure there are no other shipping or tracking labels attached to your package.
2. **Fold the printed label at the solid line below.** Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup




Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages.

Hand the package to any UPS driver in your area.

Take your package to any location of The UPS Store®, UPS Access Point(TM) location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

FDA-CVM-FOIA-2019-1704-000101

FOLD HERE

<p>DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSIT 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607</p> <p>SHIP TO: JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MUIRKIRK RD LAUREL MD 20708</p>	<p>3 LBS</p> <p>RS</p> <p>DWT: 12,12,10</p>	<p>1 OF 1</p>	<p>MD 207 9-48</p> 	<p>UPS NEXT DAY AIR</p> <p>1</p> <p>TRACKING #: 1Z A44 20T 84 9169 1688</p> 	<p>BILLING: P/P DESC: Sample RETURN SERVICE</p> <p>GENTER: CVM can numbers: 6993016</p> <p>CS 20.0.27. WINTNV50 93.04.10/2017</p> 
--	---	----------------------	---	--	---

UPS CampusShip: View/Print Label

1. Ensure there are no other shipping or tracking labels attached to your package.
2. **Fold the printed label at the solid line below.** Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.




Customers without a Daily Pickup

Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages.

Hand the package to any UPS driver in your area.

Take your package to any location of The UPS Store®, UPS Access Point(TM) location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

FOLD HERE

<p>DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSIT 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607</p> <p>SHIP TO: JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MUIRKIRK RD LAUREL MD 20708</p>	<p>3 LBS</p> <p>RS</p> <p>DWT: 12,12,10</p>	<p>1 OF 1</p>	<p>MD 207 9-48</p> 	<p>UPS NEXT DAY AIR</p> <p>1</p> <p>TRACKING #: 1Z A44 20T 84 9187 0743</p> 	<p>BILLING: P/P DESC: Sample RETURN SERVICE</p> <p>GENTER: CVM can numbers: 6993016</p> <p>CS 20.0.27. WINTNV50 93.04.10/2017</p> 
--	---	----------------------	---	--	---

UPS/FedEx Package Information Form

Sender (address, tel #, fax #, e-mail):

Jake Guag, M.P.H, C.P.H.
Biologist
U.S. Food & Drug Administration
Center for Veterinary Medicine
Office of Research
Vet-LIRN
8401 Muirkirk Road.
Laurel, Maryland 20708
Tel: 240-402-0917
email: Jake.Guag@fda.hhs.gov

Recipient (name, address, tel. #, fax #, e-mail):

Attn: Dr. Darcy Adin
North Carolina State University
NC State Veterinary Hospital
1060 William Moore Drive
Raleigh, NC 27607
919-513-6032
Email: dbadin@ncsu.edu

Service (Standard, Priority Overnight, etc.):

Priority overnight

Date:

01/10/2018

Special Needs (Dry Ice, Hazardous Chemicals, etc.). Dry ice shipments must list the weight of the dry ice separately	
Weight of Dry Ice (kg, lbs)	NO
Total Weight of Package (kg, lbs.):	1.7 lbs
Dimensions of Package (L*W*H):	12x12x10 inches
Value:	/
Content Description	Food collection kit

800.218

We are collecting dog food-weight 0.36 kg. In a plastic tupperware container~5" x 5" x 2" (0.36kg = ~0.8 lb)

No hazardous materials.

Room temperature.



Network Procedures for Shipping Vet-LIRN Samples

Introduction

The purpose of this Network Procedure is to provide general information on shipping for Vet-LIRN samples. There are 5 different kinds of samples that will be covered including:


- Room Temperature samples (non-histological)
- Histological Samples
- Frozen Samples
- Urine Samples
- Exempt Patient Specimen



Including inventory and any paperwork provided by Vet-LIRN in shipment sealed in a plastic bag.


Room Temperature

Non-Histological



- Room Temperature
- Secondary packaging
- Provide cushion as needed (eg. Bubblewrap)

Histological




- Room Temperature
- Place in secondary container
- Provide cushion as needed (eg. Bubblewrap)
- Must have Exempt Animal Specimen sticker on packaging

Frozen Tissues



- Ice packs frozen for 24 hours
- Secondary packaging
- Provide cushion as needed (eg. Bubblewrap)

Urine



Exempt Patient Specimen

If there is a minimal likelihood that the sample contains a pathogen, then the packaging may be marked as "Exempt Patient Specimen". Examples include, but are not limited to:

- Serum sent for antibody testing
- Tissues sent in 10% formalin (higher than 10% formalin requires further marking, UN 3334)
- Samples to be tested for therapeutic drug monitoring or toxins
- Environmental samples not expected to contain a pathogen
- Dried blood spots placed on absorbent filter paper



Guag, Jake

From: Guag, Jake
Sent: Wednesday, January 17, 2018 9:13 AM
To: 'dbadin@ncsu.edu'
Cc: Jones, Jennifer L
Subject: FDA (Vet-LIRN) shipped sample collection kit

Dear Dr. Adin,

We shipped a food sample collection kit to your place this morning. Its tracking number is 1ZA4420T0194648732 with UPS. It is expected to arrive on tomorrow (Jan 18, 2018).

Thank you
Jake Guag

Jake Guag, MPH , CPH
Biologist
U.S. Food & Drug Administration
Center for Veterinary Medicine
Office of Research
Vet-LIRN
8401 Muirkirk Road.
Laurel, Maryland 20708
tel: 1-240-402-0917
email: Jake.Guag@fda.hhs.gov

UPS CampusShip: View/Print Label

- 1. **Ensure there are no other shipping or tracking labels attached to your package.** Select the Print button on the print dialog box that appears. Note: If your browser does not support this function select Print from the File menu to print the label.
- 2. **Fold the printed label at the solid line below.** Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup

Take your package to any location of The UPS Store®, UPS Access Point(TM) location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages. Hand the package to any UPS driver in your area.

UPS Access Point™
 BEST PAWN, INC.
 13919 BALTIMORE AVE
 LAUREL ,MD 20707

UPS Access Point™
 THE UPS STORE
 14625 BALTIMORE AVE
 LAUREL ,MD 20707

UPS Access Point™
 INTERNATIONALFOODMARKETOFBELTS
 11118 BALTIMORE AVE
 BELTSVILLE ,MD 20705

FOLD HERE

<p>JAKE GUAC 1-240-402-0917 FDA CVM/HA/Mod2 8401 MURKIRK RD LAUREL, MD 20708</p> <p>SHIP TO: DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSITY NC STATE VETERINARY HOSPITAL 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607-4065</p>	<p>2 LBS</p> <p>1 OF 1</p> <p>DWT: 12.12.10</p>	<p>NC 276 9-07</p> 	<p>UPS NEXT DAY AIR</p> <p>1</p> <p>TRACKING #: 1Z A44 20T 01 9464 8732</p> 	<p>BILLING: P/P</p> <p>CENTER: CVM can numbers: 6993016</p> <p>CS 20 0.27. WNTNVS0 93.0A.19/2017</p> 
---	---	---	--	---

UPS CampusShip: View/Print Label

- 1. Ensure there are no other shipping or tracking labels attached to your package.
- 2. Fold the printed label at the solid line below. Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup




Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages.

Hand the package to any UPS driver in your area.

Take your package to any location of The UPS Store®, UPS Access Point(TM) location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

Return Label to Vet-CRN (FHIA)

FOLD HERE

<p>DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSIT 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607</p> <p>SHIP TO: JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MUIRKIRK RD LAUREL MD 20708</p>	<p>3 LBS 1 OF 1</p> <p>RS DWT: 12,12,10</p> <p>MD 207 9-48</p> 	<p>UPS NEXT DAY AIR</p> <p>TRACKING #: 1Z A44 20T 84 9187 0743 1</p>		<p>BILLING: P/P DESC: Sample RETURN SERVICE</p> <p>CENTER: CVM can numbers: 6993016</p> <p>CS 20.0.27. WNTNVS0 93.04.10/2017</p> 
--	--	--	--	--

UPS/FedEx Package Information Form

Sender (address, tel #, fax #, e-mail):

Jake Guag, M.P.H, C.P.H.
Biologist
U.S. Food & Drug Administration
Center for Veterinary Medicine
Office of Research
Vet-LIRN
8401 Muirkirk Road.
Laurel, Maryland 20708
Tel: 240-402-0917
email: Jake.Guag@fda.hhs.gov

Recipient (name, address, tel. #, fax #, e-mail):

Attn: Dr. Darcy Adin
North Carolina State University
NC State Veterinary Hospital
1060 William Moore Drive
Raleigh, NC 27607
919-513-6032

Service (Standard, Priority Overnight, etc.):

Priority overnight

Date:

01/10/2018

Special Needs (Dry Ice, Hazardous Chemicals, etc.). Dry ice shipments must list the weight of the dry ice separately	
Weight of Dry Ice (kg, lbs)	NO
Total Weight of Package (kg, lbs.):	1.7 lbs
Dimensions of Package (L*W*H):	12x12x10 inches
Value:	/
Content Description	Food collection kit

800.218

We are collecting dog food-weight 0.36 kg. In a plastic tupperware container~5" x 5" x 2" (0.36kg = ~0.8 lb)

No hazardous materials.

Room temperature.

UPS CampusShip: View/Print Label

- 1. Ensure there are no other shipping or tracking labels attached to your package.
- 2. Fold the printed label at the solid line below. Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup




Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages.

Hand the package to any UPS driver in your area.

Take your package to any location of The UPS Store®, UPS Access Point(TM) location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

Return Label to Vet-LIAN (FDA)

FOLD HERE

<p>DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSIT 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607</p> <p>SHIP TO: JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MUIRKIRK RD LAUREL, MD 20708</p>	<p>3 LBS</p> <p>RS</p> <p>DWT: 12,12,10</p> <p>1 OF 1</p>	<p>MD 207 9-48</p> 	<p>UPS NEXT DAY AIR</p> <p>1</p> <p>TRACKING #: 1Z A44 20T 84 9169 1688</p> 	<p>BILLING: P/P DESC: Sample RETURN SERVICE</p> <p>CENTER: CVM can numbers: 6993016</p> <p>CS 20.0.27. WRTNVE50 93.0A.10/2017</p> 
---	--	---	---	---

UPS CampusShip: View/Print Label

- 1. **Ensure there are no other shipping or tracking labels attached to your package.** Select the Print button on the print dialog box that appears. Note: If your browser does not support this function select Print from the File menu to print the label.
- 2. **Fold the printed label at the solid line below.** Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup

Take your package to any location of The UPS Store®, UPS Access Point(TM) location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.




Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages. Hand the package to any UPS driver in your area.

UPS Access Point™
 BEST PAWN, INC.
 13919 BALTIMORE AVE
 LAUREL ,MD 20707

UPS Access Point™
 THE UPS STORE
 14625 BALTIMORE AVE
 LAUREL ,MD 20707

UPS Access Point™
 INTERNATIONALFOODMARKETOFBELTS
 11118 BALTIMORE AVE
 BELTSVILLE ,MD 20705

FOLD HERE

<p>JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MURKIRK RD LAUREL MD 20708</p> <p>SHIP TO: DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSITY NC STATE VETERINARY HOSPITAL 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607</p>	<p>2 LBS</p> <p>DWT: 12,12,10</p> <p>1 OF 1</p>	<p>NC 276 9-07</p> 	<p>UPS NEXT DAY AIR</p> <p>TRACKING #: 1Z A44 20T 01 9190 2873</p> <p>1</p> 	<p>BILLING: P/P</p> <p>CENTER: CVM can numbers: 6993016</p> <p>CS 20.0.27. WNTNVS0 93.0A.10/2017</p> 
--	---	---	--	---



Network Procedures for Shipping Vet-LIRN Samples

Introduction

The purpose of this Network Procedure is to provide general information on shipping for Vet-LIRN samples. There are 5 different kinds of samples that will be covered including:


- Room Temperature samples (non-histological)
- Histological Samples
- Frozen Samples
- Urine Samples
- Exempt Patient Specimen



Including inventory and any paperwork provided by Vet-LIRN in shipment sealed in a plastic bag.


Room Temperature

Non-Histological



- Room Temperature
- Secondary packaging
- Provide cushion as needed (eg. Bubblewrap)

Histological



- Room Temperature
- Place in secondary container
- Provide cushion as needed (eg. Bubblewrap)
- Must have Exempt Animal Specimen sticker on packaging


Frozen

Tissues



- Ice packs frozen for 24 hours
- Secondary packaging
- Provide cushion as needed (eg. Bubblewrap)

Urine



Exempt Patient Specimen

If there is a minimal likelihood that the sample contains a pathogen, then the packaging may be marked as "Exempt Patient Specimen". Examples include, but are not limited to:

- Serum sent for antibody testing
- Tissues sent in 10% formalin (higher than 10% formalin requires further marking, UN 3334)
- Samples to be tested for therapeutic drug monitoring or toxins
- Environmental samples not expected to contain a pathogen
- Dried blood spots placed on absorbent filter paper



UPS CampusShip: View/Print Label

1. **Ensure there are no other shipping or tracking labels attached to your package.** Select the Print button on the print dialog box that appears. Note: If your browser does not support this function select Print from the File menu to print the label.

2. **Fold the printed label at the solid line below.** Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup

Take your package to any location of The UPS Store®, UPS Access Point™ location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages.

Hand the package to any UPS driver in your area.

UPS Access Point™
BEST PAWN, INC.
13919 BALTIMORE AVE
LAUREL ,MD 20707

UPS Access Point™
THE UPS STORE
14625 BALTIMORE AVE
LAUREL ,MD 20707

UPS Access Point™
INTERNATIONALFOODMARKETOFBELTS
11118 BALTIMORE AVE
BELTSVILLE ,MD 20705

FOLD HERE

<p>JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MURKIRK RD LAUREL MD 20708</p> <p>SHIP TO: DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSITY NC STATE VETERINARY HOSPITAL 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607</p>	<p>2 LBS</p> <p>DWT: 12,12,10</p>	<p>1 OF 1</p>	<p>NC 276 9-07</p> 	<p>UPS NEXT DAY AIR</p> <p>1</p> <p>TRACKING #: 1Z A44 20T 01 9190 2873</p>		<p>BILLING: P/P</p> <p>CENTER: CVM Call numbers: 6993016</p> <p>GS 20.0.27. WINTNV50 P3.0A.10/2017</p> 
--	--	----------------------	---	---	--	--

UPS CampusShip: View/Print Label

1. **Ensure there are no other shipping or tracking labels attached to your package.** Select the Print button on the print dialog box that appears. Note: If your browser does not support this function select Print from the File menu to print the label.

2. **Fold the printed label at the solid line below.** Place the label in a UPS Shipping Pouch. If you do not have a pouch, affix the folded label using clear plastic shipping tape over the entire label.

3. GETTING YOUR SHIPMENT TO UPS

Customers with a Daily Pickup

Your driver will pickup your shipment(s) as usual.

Customers without a Daily Pickup

Take your package to any location of The UPS Store®, UPS Access Point™ location, UPS Drop Box, UPS Customer Center, Staples® or Authorized Shipping Outlet near you. Items sent via UPS Return Services(SM) (including via Ground) are also accepted at Drop Boxes. To find the location nearest you, please visit the Resources area of CampusShip and select UPS Locations.

Schedule a same day or future day Pickup to have a UPS driver pickup all your CampusShip packages.

Hand the package to any UPS driver in your area.

UPS Access Point™
BEST PAWN, INC.
13919 BALTIMORE AVE
LAUREL ,MD 20707

UPS Access Point™
THE UPS STORE
14625 BALTIMORE AVE
LAUREL ,MD 20707

UPS Access Point™
INTERNATIONALFOODMARKETOFBELTS
11118 BALTIMORE AVE
BELTSVILLE ,MD 20705

FOLD HERE

<p>JAKE GUAG 1-240-402-0917 FDA-CVM-HQ-MOD2 8401MURKIRK RD LAUREL MD 20708</p> <p>SHIP TO: DR. DARCY ADIN 919-513-6032 NORTH CAROLINA STATE UNIVERSITY NC STATE VETERINARY HOSPITAL 1060 WILLIAM MOORE DRIVE RALEIGH NC 27607-4065</p>	<p>2 LBS</p> <p>DWT: 12,12,10</p> <p>1 OF 1</p>	<p>NC 276 9-07</p> 	<p>UPS NEXT DAY AIR</p> <p>TRACKING #: 1Z A44 20T 01 9464 8732</p> <p>1</p> 	<p>BILLING: P/P</p> <p>CENTER: CVM Call numbers: 6993016</p> <p>GS 20.0.27. WINTNV50 93.0A.10/2017</p> 
--	---	---	--	--

CALIFORNIA
NATURAL

GRAIN FREE
LIMITED
INGREDIENT DIET

For Dogs With Food Sensitivities

Kangaroo & Red Lentils Recipe
ADULT DOGS 1+ YEARS

NET WT. 13.25 LBS

EXP. 01/04/2013
17251005 RF PZ

GRAIN FREE
LIMITED INGREDIENT DIET

Kangaroo &



FDA-CVM-FOIA-2019-1704-000114

MADE IN OUR
NEBRASKA PLANT

Natural dog food with added vitamins,
minerals, and other trace nutrients

CALIFORNIA

ADULT + YEARS



GRAIN FREE LIMITED INGREDIENT DIET

Kangaroo & Red Lentils Recipe

WHY LESS IS MORE

The more ingredients a food has, the higher the likelihood it may include an ingredient your dog may be sensitive to. CALIFORNIA NATURAL™ grain-free dog food recipes use fewer ingredients and zero grains to help dogs avoid the dietary triggers commonly found in many other foods.

HIGH-QUALITY KANGAROO
for dogs in need of a novel protein source

FLAXSEED
for nourishing omega-3 fatty acids and fiber



RED & GREEN LENTILS
for a unique carbohydrate alternative

SUNFLOWER OIL
for healthy skin and coat

PEAS & PEA FIBER
for a nutritious grain replacement and fiber



Kangaroo is the #1 ingredient



No white potatoes, corn, soy, dairy or eggs

INGREDIENTS:

Kangaroo, Red Lentils, Green Lentils, Peas, Sunflower Oil (preserved with mixed Tocopherols), Flaxseed, Pea Fiber, Dicalcium Phosphate, Natural Flavors, Calcium Carbonate, Salt, DL-Methionine, Minerals (Zinc Proteinate, Iron Proteinate, Copper Proteinate, Manganese Proteinate, Calcium Iodate), Vitamins (Betaine Hydrochloride, Vitamin A Supplement, Niacin Supplement, Calcium Pantothenate, Beta Carotene, Vitamin B12 Supplement, Vitamin D3 Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Biotin, Folic Acid), Vitamin E Supplement, Rosemary Extract

GUARANTEED ANALYSIS:

Crude Protein (Min) 21%
Crude Fat (Min) 11%
Crude Fiber (Max) 6.5%
Moisture (Max) 10%
Linoleic Acid (an Omega-6 Fatty Acid) (Min) 13%
Iron (Min) 100 mg/kg
Zinc (Min) 140 mg/kg
Vitamin E (Min) 300 IU/kg
Omega-3 Fatty Acids* (Min) 1%
*Not recognized as an essential nutrient by the AAFCO Dog Food Nutrient Profiles

FEEDING GUIDELINES: Every cup of our food is nutrient dense so your dog gets the right amount of calories in a complete and nutritious meal. Adjust amount fed to obtain or maintain your dog's optimal weight.

To help maintain freshness, seal bag tightly and store in a cool, dry location.

Suggested Amounts to Feed Per Day (Cups)

WEIGHT (lbs.)	FEEDING AMOUNT (Cups/day)
10	7/8
20	1 3/8
30	1 7/8
40	2 1/4
50	2 5/8
60	3
70	3 3/8
80	3 5/8
90	3 7/8
100	4 1/4
120	4 3/4
140	5 1/4



Approximate Kibble Size

CALIFORNIA NATURAL™ Limited Ingredient Diet Grain Free Kangaroo & Red Lentils Recipe Dog Food is formulated to meet the nutritional levels established by the AAFCO Dog Food Nutrient Profiles for adult maintenance.

TRANSITIONING TO OUR FOOD: If feeding CALIFORNIA NATURAL™ Dog Food for the first time or changing recipes, we suggest you blend increasing amounts of the new recipe with your old dog food for six days.



CALORIE CONTENT:

Metabolizable Energy - Calculated
3,487 kcal/kg
434 kcal/cup
1 cup = 4.38 oz. (124 g)

MARS petcare

1348053

PROOF OF PURCHASE CALIFORNIA NATURAL™ Limited Ingredient Diet Grain Free Kangaroo & Red Lentils Recipe Dog Food



MADE IN OUR NEBRASKA PLANT WITH THE WORLD'S FINEST INGREDIENTS

Manufactured by: Natura Pet Products, Inc. 2779 Rademakers Way Fremont, NE 68025 USA ©/™ Trademarks © Mars, Incorporated.

Image on Front: Rasheed, a friendly and fearless dog through the desert heat. Joshua Tree National Park in California.

QUALITY STANDARDS AS HIGH AS YOURS

Good health starts with good food - that's why only the best ingredients go into every batch of CALIFORNIA NATURAL™ dog food and why we make every bag of kibble ourselves in our Nebraska plant. It's also why our independent ISO quality checks a day based on industry-recognized standards and procedures.

QUALITY CHECKS PER DAY

100% Satisfaction Guarantee: If you are not completely satisfied with this product, return the unused portion for any reason you choose within the dotted bag seal on this bag to your retailer for a full refund. For more information, visit our Care Department at www.californianatural.com



Kangaroo is the #1 ingredient

INGREDIENTS:

Kangaroo, Red Lentils, Green Lentils, Peas, Sunflower Oil (preserved with mixed Tocopherols), Flaxseed, Pea Fiber, Dicalcium Phosphate, Natural Flavors, Calcium Carbonate, Salt, DL-Methionine, Minerals (Zinc Proteinate, Iron Proteinate, Copper Proteinate, Manganese Proteinate, Calcium Iodate), Vitamins (Betaine Hydrochloride, Vitamin A Supplement, Niacin Supplement, Calcium Pantothenate, Beta Carotene, Vitamin B12 Supplement, Vitamin D3 Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Biotin, Folic Acid), Vitamin E Supplement, Rosemary Extract

GUARANTEED ANALYSIS:

Crude Protein (Min) 21%
Crude Fat (Min) 11%
Crude Fiber (Max) 6.5%
Moisture (Max) 10%

FEEDING GUIDELINE
gets the right amount of amount fed to obtain or m

To help maintain freshness
Suggested Amounts to Feed Per Day (Cups)

WEIGHT (lbs.)	FEEDING AMOUNT (cups/day)
10	7/8
20	1 3/8
30	1 7/8
40	2 1/4
50	2 5/8
60	3
70	3 3/8
80	3 5/8
90	3 7/8
100	4 1/4

EDACVM-FOIA-2019-1704-000116

5+ YEARS

...cups

NET WT. 13

EXP: 01/04/19 13:28
17251005 RFT F2

GRAIN FREE

LIMITED INGREDIENT DIET

Kanqaroo &

ONLY
INGRI
* ADDED VIT
OTHER TRA
MATURE

Red Lentils Recipe

ADULT 1+ YEARS



LIMITED **GRAIN FREE** DIET

Kangaroo & Red Lentils Recipe
ADULT 1+ YEARS

WHY LESS IS MORE

The more ingredients a food has, the higher the likelihood it may include an ingredient your dog may be sensitive to. CALIFORNIA NATURAL™ grain-free dog food recipes use fewer ingredients and zero grains to help dogs avoid the dietary triggers commonly found in many other foods.

HIGH-QUALITY KANGAROO
for dogs in need of a novel protein source

FLAXSEED
for nourishing Omega-3 fatty acids and fiber

RED & GREEN LENTILS
for a unique carbohydrate alternative



ONLY 5 KEY INGREDIENTS

+ ADDED VITAMINS, MINERALS, OTHER TRACE NUTRIENTS & NATURAL FLAVORS

SUNFLOWER OIL
for healthy skin and coat

PEAS & PEA FIBER
for a nutritious grain replacement and fiber



Kangaroo is the #1 ingredient



No white potatoes, corn, soy, dairy or eggs

INGREDIENTS:

Kangaroo, Red Lentils, Green Lentils, Peas, Sunflower Oil (preserved with mixed Tocopherols), Flaxseed, Pea Fiber, Dicalcium Phosphate, Natural Flavors, Calcium Carbonate, Salt, DL-Methionine, Minerals (Zinc Proteinate, Iron Proteinate, Copper Proteinate, Manganese Proteinate, Calcium Iodate), Vitamins (Sialine Hydrochloride, Vitamin A Supplement, Niacin Supplement, Calcium Pantothenate, Beta Carotene, Vitamin B12 Supplement, Vitamin D3 Supplement, Riboflavin Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Biotin, Folic Acid), Vitamin E Supplement, Rosemary Extract

GUARANTEED ANALYSIS:

Crude Protein (Min)	21%
Crude Fat (Min)	11%
Crude Fiber (Max)	6.5%
Moisture (Max)	10%
Linoleic Acid	
(a) Omega-6 Fatty Acid (Min)	1.9%
Iron (Min)	100 mg/kg
Zinc (Min)	140 mg/kg
Vitamin E (Min)	300 IU/kg
Omega-3 Fatty Acids* (Min)	1%

*Not recognized as an essential nutrient by the AAFCO Dog Food Nutrient Profiles

FEEDING GUIDELINES: Every cup of our food is nutrient dense so your dog gets the right amount of calories in a complete and nutritious meal. Adjust amount fed to obtain or maintain your dog's optimal weight.

To help maintain freshness, seal bag tightly and store in a cool, dry location.

Suggested Amounts to Feed Per Day (Cups)

WEIGHT (lbs.)	FEEDING AMOUNT (cups/day)
10	7/8
20	1 3/8
30	1 7/8
40	2 1/4
50	2 5/8
60	3
70	3 3/8
80	3 5/8
90	3 7/8
100	4 1/4
120	4 3/4
140	5 1/4



Approximate Kibble Size

CALIFORNIA NATURAL™ Limited Ingredient Diet Grain Free Kangaroo & Red Lentils Recipe Dog Food is formulated to meet the nutritional levels established by the AAFCO Dog Food Nutrient Profiles for adult maintenance.

TRANSITIONING TO OUR FOOD: If feeding CALIFORNIA NATURAL™ Dog Food for the first time or changing recipes, we suggest you blend increasing amounts of the new recipe with your old dog food for six days.



QUALITY STANDARDS AS HIGH AS YOURS

Good health starts with good food - that's why only the best ingredients go into every batch of CALIFORNIA NATURAL™ dog food and why we make every bag of kibble ourselves in our Nebraska plant. It's also why our plant conducts 600 quality checks a day based on industry-recognized quality and food safety standards.

600 QUALITY CHECKS PER DAY

CALORIE CONTENT:
Metabolizable Energy - Calculated
3,487 kcal/kg
434 kcal/cup
1 cup = 4.38 oz. (124 g)



MADE IN OUR NEBRASKA PLANT WITH THE WORLD'S FINEST INGREDIENTS

Manufactured by:
Natura Pet Products, Inc.
2779 Rademakers Way
Fremont, NE 68025 USA
©/TM Trademarks
© Mars, Incorporated.

Image on Front:
Rasheed, a fast, powerful and fearless saluki courses through the desert near Joshua Tree National Park in California.

100% Satisfaction Guaranteed:

If for any reason you are dissatisfied with this product, return the unused portion along with the information contained within the dotted panel on or refund. Should you wish to contact our Consumer Care Department, please call 1-800-532-7261 or visit us at www.californianaturalpet.com

1348053
PROOF OF PURCHASE
CALIFORNIA NATURAL™
Limited Ingredient Diet Grain Free Kangaroo & Red Lentils Recipe Dog Food

7 51485 3994 1 6

FOR PROOF OF PURCHASE
CUT ALONG DASHED LINE

Ver - LTRU
D - # 2469

Kangaroo &

ONLY 5 KEY INGREDIENTS

LIMITED INGREDIENT DIET

NATURAL

dog food with added vitamins, other trace nutrients

MADE IN OUR NEBRASKA PLANT

Kangaroo & Red Lentils Recipe

ADULT 1+ YEARS

For Dogs With Food Sensitivities

INGREDIENT DIET

LIMITED

GRAIN FREE



CALIFORNIA NATURAL

GRAIN FREE LIMITED INGREDIENT DIET

		800.261-sub 1				
		Case Sample				
		Fromm Heartland Gold Grain Free Large Breed Adult	Assumptions: max moisture of 10% (label)	Product Typical Analysis (website label)	AAFCO Growth & Maint	Issues
(b) (4)	Tau	45.5 mg/100g	0.05% DMB	n/a	0.1% in Cats	LOW
	Cystine	293 mg/100g	0.33% DMB	n/a	n/a	
	Met	358 mg/100g	0.4% DMB	n/a	0.35%	none
	Met-Cys	0.33 + 0.4	0.73% DMB	n/a	0.70%	none
MSU	Iodine	4.2 mg/kg	4.67% DMB		1 to 11 ppm	none

(b) (4)

Report Number: 2119443-0

Report Date: 01-May-2018

Report Status: Final

Certificate of Analysis

Food and Drug Administration - CVM

8401 Muirkirk Rd.
Laurel Maryland 20708 United States

Sample Name:	1-dog food	(b) (4) Sample:	7192972
Project ID	FDA_CVM-20180413-0004	Receipt Date	13-Apr-2018
PO Number	HHSF223201610005I HHSF22301003T	Receipt Condition	Ambient temperature
Sample Serving Size		Login Date	13-Apr-2018
Description	800.261-sub	Online Order	20

Analysis	Result
Cystine and Methionine *	
Cystine	293 mg/100g
Methionine	358 mg/100g
Taurine	
Taurine	45.5 mg/100g

Method References	Testing Location
-------------------	------------------

Cystine and Methionine (AAAC_S)	(b) (4)
--	---------

Official Methods of Analysis of AOAC INTERNATIONAL, Method 982.30 E(a/b)

Taurine (TAUR_LC_S)	(b) (4)
----------------------------	---------

Official Methods of Analysis of AOAC INTERNATIONAL, Method 999.12, AOAC International Gaithersburg, MD, USA, (Modified)

R. Schuster, "Determination of Amino Acids in Biological, Pharmaceutical, Plant and Food Samples by Automated Precolumn Derivatization and HPLC", *Journal of Chromatography*, 431:271-284, (1988) (Modified)

Henderson, J.W., Ricker, R.D. Bidlingmeyer, B.A., Woodward, C., "Rapid, Accurate, Sensitive, and Reproducible HPLC Analysis of Amino Acids, Amino Acid Analysis Using Zorbax Eclipse-AAA columns and the Agilent 1100 HPLC," Agilent Publication, 2000 (Modified)

Henderson, J.W., Books, A., "Improved Amino Acid Methods using Agilent Zorbax Eclipse Plus C18 Columns for a Variety of Agilent LC Instrumentation and Separation Goals," Agilent Application Note 5990-4547, (2010).

Testing Location(s)	Released on Behalf of
---------------------	-----------------------

(b) (4)	(b) (4)
---------	---------

(b) (4)	
---------	---

(b) (5)

These results apply only to the items tested. This certificate of analysis shall not be reproduced, except in its entirety, without the written approval of (b) (6)

* This analysis is not ISO accredited.

Printed: 01-May-2018 2:31 pm

Page 1 of 1

Signature

KANGAROO FORMULA

EXCEEDING EXPECTATIONS
IN PET NUTRITION

LIMITED INGREDIENT FORMULA

Zignature® is formulated to serve your canine companions needs by delivering the highest quality, well balanced diets. Our limited ingredient formula uses Kangaroo Protein, No Potatoes, No Grains, No Chicken, No Eggs, No Glutens, No Tapioca, No Corn and No Soy. The health of your pet is our highest priority. By eliminating unnecessary ingredients, we are able to bring you a formula closer to how nature intended it (with added vitamins and minerals). With Zignature®, you can be assured you are giving your canine companions the best food for life.

PEAS, CHICKPEAS, FLAXSEED & ALFALFA MEAL

Zignature® uses only the highest quality nutrient dense legumes (Peas & Chickpeas), Flaxseed & Alfalfa Meal and appropriate nutrition. Along with regular exercise and proper feeding, Zignature® can help your dog reach and maintain an ideal weight.

ANIMAL PROTEIN FIRST

To give your canine companions the optimum health and best diet, Zignature® understands your canine companions are carnivores first, omnivores second and thrive with a good portion of their diet coming from animal or fish protein. At Zignature®, we ensure our first two ingredients are from Kangaroo proteins for healthy growth and maintenance to give your canine companions everything they need for a long and happy life with you.



INGREDIENTS

Kangaroo, Kangaroo Meal, Peas, Chickpeas, Pea Flour, Sunflower Oil (preserved with Citric Acid), Flaxseed, Red Lentils, Green Lentils, Dehydrated Alfalfa Meal, Pea Protein, Natural Flavors, Salt, Minerals (Zinc Proteinate, Iron Proteinate, Copper Proteinate, Manganese Proteinate, Cobalt Proteinate, Selenium Yeast), Choline Chloride, Potassium Chloride, Calcium Carbonate, Vitamins (Vitamin A Acetate, Vitamin D3 Supplement, Vitamin E Supplement, Niacin, d-Calcium Pantothenate, Thiamine Mononitrate, Pyridoxine Hydrochloride, Riboflavin Supplement, Folic Acid, Biotin, Vitamin B12 Supplement), Lactic Acid, Calcium Iodate, Preserved With Mixed Tocopherols.

AAFCO NUTRITIONAL ADEQUACY STATEMENT

Zignature® Kangaroo Formula for dogs is formulated to meet the nutritional levels established by the Association of American Feed Control Officials (AAFCO) Dog Food Nutrient Profiles for all life stages including growth of large size dogs (70 lbs. or more as an adult).

DAILY FEEDING GUIDELINES

WEIGHT OF DOG (lbs)	SERVING SIZE (Adult)*
10 or less	¼ - ¾ cup
10 - 20	¾ - 1½ cups
20 - 30	1½ - 2 cups
30 - 40	2 - 2½ cups
40 - 60	2½ - 3 cups
60 - 80	3 - 3¾ cups
80 - 100	3 - 3¾ cups

PUPPIES Feed up to 2 times per pound of body weight over adult levels.

PREGNANT FEMALES Minimal increase needed during first 6 weeks. Weeks 7, 8 & 9 increase up to 25%.

LACTATING FEMALES Immediately after whelping, nutritional demand increases by 50% compared to maintenance levels. During peak lactation (around 4 - 5 weeks), increase feeding up to 300% of regular intake. Feed in 3 equal meals per day.

When switching your dog's diet, we recommend that it should be done gradually over 7 days, increasing the amount each day with the present diet as sudden change in diet may result in digestive disturbances.

Consult your veterinarian regularly. Keep fresh water available. To maintain freshness, keep package sealed and store in a cool, dry place.

Signature®

WWW.ZIGNATURE.COM

Manufactured for: **PETS GLOBAL, INC.**
10000 INDUSTRIAL DRIVE, VALENCIA, CA 92683, USA
TEL: 1-888-897-7207



GUARANTEED ANALYSIS

Crude Protein	26.0% (min)
Crude Fat	14.0% (min)
Crude Fiber	4.5% (max)
Moisture	10.0% (max)
Calcium	1.2% (min)
Phosphorus	1.0% (min)
Omega 6 Fatty Acids*	3.0% (min)
Omega 3 Fatty Acids*	0.6% (min)

*Not recognized as an essential nutrient by the AAFCO Dog Food Nutrient Profiles.

CALORIE CONTENT

Metabolizable Energy (calculated)
3,888 kcals per kg
425 kcals per cup

BEST USED BY

KANGAROO FORMULA

100% SATISFACTION GUARANTEE
All Zignature's products are 100% Satisfaction Guaranteed. If you and your pet are not completely satisfied with this or any Zignature's products, simply return the unused portion, along with the original receipt to your Zignature's retail store for a complete refund or replacement.

ANIMAL PROTEIN FIRST

To give your canine companions the optimum health and best diet, Zignature® understands your canine companions are carnivores first, omnivores second and third with a good portion of their diet coming from animal or fish protein. At Zignature®, we ensure our first two ingredients are from Kangaroo proteins for healthy growth and maintenance to give your canine companions everything they need for a long and happy life with you.

INGREDIENTS

Kangaroo, Kangaroo Meal, Peas, Chickpeas, Pea Flour, Sunflower Oil (preserved with Citric Acid), Flaxseed, Red Lentils, Green Lentils, Dehydrated Alfalfa Meal, Pea Protein, Natural Flavors, Salt, Minerals (Zinc Proteinate, Iron Proteinate, Copper Proteinate, Manganese Proteinate, Cobalt Proteinate, Selenium Yeast), Choline Chloride, Potassium Chloride, Calcium Carbonate, Vitamins (Vitamin A Acetate, Vitamin D3 Supplement, Vitamin E Supplement, Niacin, d-Calcium Pantothenate, Thiamine Mononitrate, Pyridoxine Hydrochloride, Riboflavin Supplement, Folic Acid, Biotin, Vitamin B12 Supplement), Lactic Acid, Calcium Iodate, Preserved With Mixed Tocopherols.

Zignature® Kangaroo Formula for dogs is formulated to meet the nutritional levels established by the Association of American Feed Control Officials (AAFCO) Dog Food Nutrient Profiles for all life stages including growth of large size dogs (70 lbs. or more as an adult).

AAFCO NUTRITIONAL ADEQUACY STATEMENT

DAILY FEEDING GUIDELINES

WEIGHT OF DOG (lbs) **SERVING SIZE (Adult)***

10 or less	1/4 - 3/4 cup
10 - 20	3/4 - 1 1/4 cups
20 - 30	1 1/4 - 1 1/2 cups
30 - 40	1 1/2 - 2 cups

PUPPIES Feed up to 2 times per pound of body weight over adult levels.

PREGNANT FEMALES Minimal increase needed during first 6 weeks. Weeks 7, 8 & 9 increase up to 25%

LACTATING FEMALES Immediately after whelping, nutritional demand increases by 50% compared to maintenance levels. During peak lactation (around 4 - 5

Dehydrated Alfalfa Meal

Kangaroo

Sunflower

GUARANTEED

Crude Protein
Crude Fat
Crude Fiber
Moisture
Calcium
Phosphorus
Omega 6 Fatty Acids*
Omega 3 Fatty Acids*

*Not recognized as an essential nutrient in the AAFCO Dog Food Nutrient Profile

CALORIE CO

Metabolizable Energy

3,888 kcals per
425 kcals per

BEST USED

FDA-CVM-FOIA-2019-1704-000123



Taurine and Carnitine in Canine Cardiomyopathy

Sherry Lynn Sanderson, DVM, PhD

Department of Physiology and Pharmacology, University of Georgia, College of Veterinary Medicine, 501 DW Brooks Drive, Athens, GA 30602, USA

Dilated cardiomyopathy (DCM) is one of the most common acquired cardiovascular diseases in dogs [1–4]. Although few studies of the prevalence of DCM in the overall population of dogs have been reported, estimates range from 0.5% to 1.1% [5,6]. Only degenerative valvular disease and, in some regions of the world, heartworm infection are more common causes of cardiac morbidity and mortality in dogs. DCM is seen most commonly in large and giant breeds of dogs, although its frequency seems to be increasing in medium-sized breeds, such as the English and American cocker spaniels [4–8]. It has been reported rarely in small and miniature breeds of dogs [9].

DCM is particularly challenging to veterinarians because the cause is often unknown and can vary among dog breeds [10]. Because most cases of DCM in dogs are classified as idiopathic, most therapies can be classified as “Band-Aid therapies” that palliate the effects of this disease for a short duration but do little to address the primary disease process. Therefore, DCM is almost always a progressive disease, and most dogs will eventually succumb to their disease. Survival times in dogs with DCM are variable and can be influenced by several factors, including breed. However, the prognosis for survival of dogs with DCM remains poor, with reported survival rates of 17.5% at 1 year and 7.5% at 2 years [11–13]. Until recently, reported cases of DCM reversal in dogs were very rare.

With advancements in echocardiology, diagnostic capabilities in canine cardiology have improved dramatically over the past 2 decades. Therapeutic advances have made surprisingly little progress. Symptomatic treatment is the standard care and outcome remains poor.

Recently, more promising therapies for dogs with DCM have resulted from a clearer understanding of the importance of biochemistry and nutrition in managing this disease. Nutrition is now widely accepted as an important adjunct to medical therapy in dogs with DCM.

E-mail address: sanderso@vet.uga.edu

The importance of nutrition in managing DCM has changed dramatically in the past 10 to 15 years. Historically, dietary sodium restriction was the most common nutritional recommendation for dogs with DCM. The importance of other nutrients in the origin and management of this disease was largely unknown. More recently, widely accepted beliefs about the role nutrient deficiencies could play in DCM have been proven false, further enhancing the ability to direct therapy at an underlying cause rather than just the symptoms.

This article focuses on two nutrients, taurine and carnitine, that play an important role in the cause and treatment of DCM in some dogs. Known risk factors for developing deficiencies of these nutrients are discussed, along with the use of taurine and carnitine for treating DCM in dogs.

TAURINE

What is Taurine?

Taurine is a sulfur-containing amino acid. Unlike most other amino acids, taurine is not incorporated into proteins but rather is one of the most abundant free amino acids in the body. Taurine is found in highest tissue concentrations in cardiac muscle, skeletal muscle, the central nervous system, and platelets [14].

Other than conjugation of bile acids and detoxification of xenobiotics through conjugation and excretion in bile, the function of taurine in mammals is not well understood but is highly diverse [14,15]. Since the mid-1970s, taurine has been known to be essential for normal retinal function in cats [16]. In addition, clinical and experimental evidence collected in the late 1980s documented that taurine is essential for normal myocardial function [17–20].

Taurine is involved with numerous metabolic processes, including antioxidation, retinal photoreceptor activity, development of the nervous systems, stabilization of neural membranes, reduction in platelet aggregation, and reproduction [15,16,21–26]. Although the importance of taurine for normal myocardial function is also well recognized, the mechanisms underlying its effect on the heart remain unknown. Much of the available evidence supports the theory that taurine's major effect on cellular function in the heart is modulating tissue calcium concentrations and availability [14,27,28]. In addition, taurine may inactivate free radicals and protect the heart by changing cellular osmolality [29]. Taurine may also have an effect on osmoregulation in the myocardium. Taurine is a small but highly charged osmotically active molecule, and experts have proposed that alterations in cellular osmolality induced by changes in intracellular taurine concentration are a protective mechanism in nervous tissue and myocardium [29]. Other proposed mechanisms specifically related to myocardial function include *N*-methylation of cell membrane phospholipids [30], direct effects on contractile proteins [31,32], and interactions with the renin–angiotensin–aldosterone system [33]. Taurine is a natural antagonist of angiotension II.

Is Taurine an Essential Amino Acid in Dogs?

Taurine is an essential amino acid in cats, and it is well known that taurine deficiency can cause DCM, retinal degeneration, and reproductive anomalies in this species [18]. However, taurine is not considered an essential amino acid in dogs. One explanation for the differences in taurine requirements between cats and dogs is that the activity of cysteine sulfinic acid decarboxylase (the rate-limiting enzyme in the synthesis of taurine from cysteine and methionine) is higher in dogs than cats [34]. However, the difference in activity of this enzyme between dogs and cats does not fully explain the difference in requirements. The activity of this enzyme in humans is even lower than in cats, and taurine is not considered an essential amino acid in healthy adult humans. Therefore, cats and dogs may have additional differences that may explain why taurine is an essential amino acid in cats and not in dogs.

A study in dogs conducted in the 1980s at the University of California at Davis showed that feeding taurine-free diets or diets found to be taurine-depleting in cats [35] did not result in taurine depletion when fed to a group of eight healthy beagles [36]. In addition, results of an early clinical study in dogs, also conducted at this University soon after the relationship between taurine deficiency and DCM was discovered in cats, were unrewarding. These studies showed that dogs could not become taurine-depleted from diet alone, and that taurine did not play a considerable role in the development of DCM in dogs.

Emergence of Taurine Deficiency in Dogs with Dilated Cardiomyopathy

The belief that taurine deficiency could not cause DCM in dogs was challenged in 1989 when taurine deficiency was linked to DCM in foxes [37]. This study reopened taurine's possible role in DCM in dogs, and a collaborative study between the University of California at Davis and the Animal Medical Center in New York City was initiated [38]. In this study, plasma taurine levels were evaluated in dogs with DCM and in those with chronic degenerative mitral valve disease. Surprisingly, results of this study showed that plasma taurine concentration was low in 17% of 75 dogs with DCM, and this deficiency occurred in breeds not commonly afflicted with DCM, such as American cocker spaniels and golden retrievers. However, because the plasma taurine concentration in breeds more commonly affected with DCM were within the reference range, experts concluded that taurine deficiency was unlikely to play an important role in the etiopathogenesis or therapy of DCM in dogs.

Multicenter Spaniel Trial (MUST) Study

Anecdotal reports emerged regarding supplementing American cocker spaniels diagnosed with DCM with taurine; however, initial reports of taurine supplementation were unrewarding. When Kittelson and colleagues [8] gave taurine and L-carnitine supplements to two American cocker spaniels with DCM, both dogs experienced response. These findings initiated the Multicenter Spaniel Trial (MUST) study. In this study, baseline plasma taurine concentrations and echocardiograms were collected in 11 American cocker spaniels diagnosed

with DCM. All dogs were found to have low plasma taurine concentrations at baseline (<50 nmol/mL). After baseline information was collected, dogs were randomly assigned to receive supplementation with both taurine (500 mg by mouth every 8 hours) and *L*-carnitine (1000 mg by mouth every 8 hours) or a placebo for 4 months, and echocardiograms were reevaluated after 2 and 4 months of therapy. The group supplemented with both taurine and carnitine showed significant echocardiographic improvement, whereas dogs receiving the placebo did not.

After this initial 4-month period, dogs that had received the placebo initially received supplements of both taurine and carnitine, and subsequently showed echocardiographic improvement after 2 to 4 months of therapy. The magnitude of echocardiographic improvement in the American cocker spaniels was not as dramatic as that seen after taurine supplementation in cats with taurine deficiency DCM. Nonetheless, after 4 months of supplementation, the improvement in myocardial function in each dog was significant enough to allow discontinuation of cardiovascular drug therapy. Improvements were seen in not only cardiovascular function but also survival times. The mean survival time for dogs in this study was 28.3 ± 19.1 months, compared with an average life expectancy for dogs treated with conventional drug therapy of approximately 6 months. Based on results from this study, the current recommendation is to supplement American cocker spaniels diagnosed with DCM with both taurine and carnitine at the doses mentioned earlier.

University of Minnesota Study in Urolith-forming Dogs Diagnosed with Dilated Cardiomyopathy

Around the same time the MUST study was initiated, a separate clinical study was initiated at the University of Minnesota. The population of dogs studied consisted of those with either cystine or urate urolithiasis that developed DCM after long-term consumption of a protein-restricted diet that was being used to manage their stone disease (Sherry L. Sanderson, DVM, PhD, unpublished data, 1998). Dogs in group 1 underwent only conventional drug therapy for their heart disease, whereas those in group 2 underwent taurine and/or carnitine supplementation in addition to conventional drug therapy as needed. Dogs in group 1 that were in Modified New York Heart Association (MNYHA) functional class I and II heart failure received enalapril (0.25 mg/kg by mouth every 12 hours) and digoxin (0.01–0.02 mg/kg by mouth divided twice a day), and dogs in MNYHA functional class III and IV received furosemide (dose varied depending on severity of heart disease) in addition to enalapril and digoxin. The population of dogs in group 1 ($N = 6$) consisted of five English bulldogs (four with cystine urolithiasis, one with urate urolithiasis) and one Dalmatian with urate urolithiasis. The population of dogs in Group 2 ($N = 8$) consisted of five English bulldogs (three with cystine urolithiasis, two with urate urolithiasis), two Dalmatians with urate urolithiasis, and one miniature Dachshund with cystine urolithiasis. Because when this study was initiated experts believed that dogs with DCM did not have low plasma taurine

concentrations, none of the dogs in group 1 had these concentrations evaluated at baseline. Plasma taurine concentrations evaluated before supplementation in seven of eight dogs in group 2 ranged from 2 nmol/mL to 45 nmol/mL (mean, 20.9 nmol/mL). These results were below the reference range of 41 nmol/mL to 97 nmol/mL that the investigators established from healthy adult beagles. Echocardiography was performed at baseline and once every 2 months. Details from this study will be published later, but a few interesting and important results were noted:

1. The average life expectancy for dogs in group 1 was 10.5 months, and all dogs were euthanized because of progressive congestive heart failure that became refractory to therapy. The average life expectancy for dogs in group 2 was 47.1 months, and only three of eight dogs were euthanized because of progressive congestive heart failure. In addition, three of five dogs that did not succumb to their heart disease received only taurine and/or carnitine supplementation and no conventional drug therapy for the management of their heart disease.
2. DCM reversed in three of eight dogs in group 2. DCM returned in one dog after the owner discontinued taurine and carnitine supplementation on their own, and in an additional dog when the dose of carnitine was reduced because of diarrhea associated with carnitine supplementation.
3. Dogs consuming a protein restricted diet long term could develop taurine deficiency, in contrast to results from previous studies that concluded that a diet could not induce taurine deficiency in dogs. This finding provided an impetus for further examining the effects on plasma and whole blood taurine levels in healthy adult dogs consuming a protein restricted diet long term.

Diet-Induced Taurine Deficiency in Healthy Adult Dogs

Previous reports indicated that dogs could not develop diet-induced taurine deficiency, even when fed a diet devoid of taurine. However, based on the finding of University of Minnesota study that dogs developed low plasma taurine levels after consuming a protein-restricted diet long-term, a more controlled study was undertaken to determine the cause of this problem and evaluate the effects of long-term taurine deficiency on cardiac function in healthy adult dogs [39].

This study involved 17 healthy adult beagles. Baseline plasma and whole blood taurine levels were evaluated, and echocardiography was performed to assess cardiac function. Once baseline data was collected, dogs were fed one of three protein-restricted diets for 48 months. All three diets had similar levels of protein; one diet was also low in fat, a second was high in fat, and a third was high in fat and supplemented with *L*-carnitine at 200 mg/kg of diet. All diets contained methionine and cystine concentrations at or above recommended minimum requirements established by the Association of American Feed Control Officials (AAFCO) [40]. After diet assignment, plasma taurine and whole blood taurine concentrations and echocardiography were evaluated every 6 months.

All three dietary treatments caused a significant decrease in whole blood taurine concentration compared with baseline concentrations. Dogs in the high-fat

group also experienced a significant decrease in plasma taurine concentration. This study was the first to show that diet could induce taurine deficiency in healthy adult dogs, in contrast to previous studies.

Another important observation was that one dog with taurine deficiency developed DCM, and that taurine supplementation resulted in almost complete reversal of the disease. This study was also the first to clearly document in dogs that taurine deficiency preceded DCM, and that taurine supplementation resulted in substantially improved cardiac function, similar to cats.

Why Did Dogs Develop Taurine Deficiency While Consuming a Protein-Restricted Diet?

The exact mechanism for this problem is unknown. However, this study showed that the AAFCO recommended minimum requirements for amino acids may need to be modified in dogs consuming a protein-restricted diet long-term. Many therapeutic diets for dogs are now supplemented with taurine.

Additional Examples of Diet-Induced Taurine Deficiency in Dogs

Soybean-based diets

Taurine deficiency was identified in two unrelated dogs fed a tofu-based diet [41]. Although the diet was low in protein, it met the National Research Council's published requirements for protein and other nutrients in dogs [42]. The authors attributed taurine deficiency to the fact that the primary protein source was soybean curd, which is low in sulfur-containing amino acids and devoid of taurine compared with meat proteins [43]. In addition, soybean curd has been shown to accelerate the loss of bile acids in cats [44].

Lamb meal and rice diets

Taurine deficiency was also identified in 12 Newfoundlands consuming two different commercially available lamb meal and rice diets [41]. Echocardiography was performed in six of the dogs, and none were diagnosed with DCM. The taurine deficiency was reversed when the diet was either changed or when the lamb meal and rice diets were supplemented with methionine. This study did not identify the exact mechanism for the development of taurine deficiency in the dogs consuming the lamb meal and rice diets.

In a study by Fascetti and colleagues [45], DCM and taurine deficiency were identified in 12 large and giant-breed dogs consuming commercially available diets that contained lamb meal, rice, or both as primary ingredients. All dogs received supplements of with taurine (1000–3000 mg by mouth every 24 hours), and significant echocardiographic improvement occurred in 9 of the 12 dogs that underwent an echocardiogram repeated after taurine supplementation. The authors hypothesized that taurine deficiency caused DCM and was caused by inadequate or unavailable dietary sulfur amino acids, which are essential precursors of taurine synthesis.

In a similar report, five related golden retrievers were diagnosed with taurine deficiency and DCM [46]. Three of five dogs were consuming lamb meal and rice or lamb and rice diets. All showed significant improvement after taurine

supplementation (500 mg by mouth every 12 hours), and all five dogs survived for more than 3 years. The authors attribute the DCM to a suspected autosomal recessive mode of inheritance; however, the potential role diet played in the development of taurine deficiency warrants mentioning.

Potential Causes of Taurine Deficiency in Dogs Consuming Lamb Meal and Rice or Lamb and Rice Diets

Torres and colleagues [47] compared the effects of consuming a lamb meal and rice-based diet with effects of consuming a poultry by-product-based diet in 12 beagles aged 5 to 5.5 months. Although the differences in plasma and whole blood taurine concentrations did not differ among diet groups, dogs consuming the lamb meal and rice-based diet excreted less taurine in their urine than dogs consuming the poultry by-product-based diet. When the lamb meal and rice diet was supplemented with methionine, urinary taurine excretion increased by 54%. Because taurine homeostasis in dogs is achieved primarily through regulating renal taurine excretion, the amount of taurine excreted in urine is a sensitive indicator of the adequacy of either taurine synthesis or absorption of dietary precursor amino acids. The authors concluded that reduced bioavailability of sulfur amino acids in the lamb meal and rice diet is a likely cause of taurine deficiency. This finding is supported by the increase in urine taurine concentrations after supplementation with methionine. Johnson and colleagues [48] showed that ileal digestibility of amino acids in dogs depends on the raw material sources and the temperature used to process feeds and provides a mechanism for these specific dietary effects.

A second potential, although related, cause of taurine deficiency in dogs consuming lamb meal and rice diets was proposed [49,50]. When dietary protein is low in quality, undigested protein reaches the colon, where it serves as a substrate for bacterial growth. Some bacteria produce cholytaurine hydrolase, an enzyme that causes release of taurine from taurocholic and other bile acids that are normally conserved in the enterohepatic circulation, resulting in increased fecal loss of taurine. Studies in dogs [49] and cats [50] have found that diets containing rice bran and whole rice products provide a source of moderately fermentable fiber and high amounts of fat. These fermentable fibers may increase the number of bacteria in the colon and result in a greater loss of taurine in the feces similar to the mechanism for undigested protein. The fat content of the diet can also affect taurine metabolism through altering intestinal bacteria and subsequent changes in the excretion of bile acids.

How Should Samples be Collected to Evaluate Plasma and Whole Blood Taurine Concentrations?

Fasting versus postprandial blood samples

Although fasting has no effect on plasma taurine concentrations in humans [51], food deprivation causes a small but significant reduction in plasma taurine concentrations in cats [52]. In a study by Torres and colleagues [47], plasma taurine concentrations were significantly reduced in food-restricted dogs compared with ad libitum-fed dogs. Whole blood taurine concentrations were

also reduced, although the whole blood taurine results were not statistically significant between the two groups. Because of the potential for food intake to affect plasma and whole blood taurine concentrations in dogs, withholding food, but not water, is recommended for 8 hours before sampling.

Anticoagulant used for plasma sample collection

Paired analysis of samples comparing taurine concentrations in plasma collected in lithium heparin with those collected in sodium citrate showed that plasma taurine concentrations are higher when lithium heparin is used as the anticoagulant [38]. Because most studies have used heparinized plasma samples to evaluate plasma taurine levels in dogs, these are recommended rather than sodium citrate plasma samples.

Plasma taurine sample collection

Heparinized, nonhemolyzed blood samples should be obtained and stored on ice until they are processed. After centrifuging, the plasma should be separated immediately from the cellular components, and a small amount of plasma should be left above the buffy coat to prevent contamination of the plasma with cells. Hemolysis and platelet or white blood cell contamination falsely elevates plasma taurine concentrations. Samples should be frozen until analyzed for plasma taurine concentrations.

Whole blood taurine sample collection

Heparinized whole blood should be frozen until samples can be analyzed. Because the red blood cells are lysed before analysis, hemolyzed samples do not adversely affect whole blood taurine analysis.

Plasma and whole blood taurine samples can be sent to the Department of Molecular Biosciences at the School of Veterinary Medicine, University of California, Davis, for analysis.

Which is Better: Plasma Taurine Concentrations or Whole Blood Taurine Concentrations

Earlier studies evaluating the relationship between taurine deficiency and DCM in dogs relied primarily on plasma taurine concentrations to predict tissue taurine concentrations. Studies conducted in dogs by this author showed findings similar to those reported in cats [53]. Relying on plasma taurine concentrations alone does not reliably assess tissue taurine concentrations in dogs. Simultaneously evaluating plasma and whole blood taurine concentrations predicts skeletal and cardiac muscle taurine concentrations better than evaluating either test alone. Therefore, when evaluating taurine status in dogs with DCM, plasma and whole blood taurine concentrations should be assessed simultaneously.

Reference Ranges for Plasma and Whole Blood Taurine Concentrations in Dogs

The reference range used in earlier studies evaluating plasma and whole blood taurine concentrations in dogs was extrapolated from the reference range use in

cats. However, reference ranges for plasma and whole blood taurine concentrations in dogs were published recently (Table 1).

Delaney and colleagues [49] have also suggested that plasma taurine concentrations less than 40 nmol/mL are critically low, as are whole blood taurine concentrations less than 150 nmol/mL. In addition, Sanderson and colleagues [53] found that low plasma taurine concentrations can exist without the presence of DCM.

Therefore, results showed that the onset of clinical signs in dogs, just as in cats, was variable when taurine concentrations declined markedly below the normal range [18].

Which Dogs Diagnosed with Dilated Cardiomyopathy Should Receive Taurine Supplementation?

Evaluation of plasma and whole blood taurine concentrations is recommended for all dogs diagnosed with DCM. An association between taurine deficiency and DCM was found in various breeds of dogs, including American cocker spaniels, Newfoundlands, golden retrievers, Labrador retrievers, Dalmatians, English bulldogs, and Portuguese water dogs. Taurine supplementation is highly recommended in any of these breeds that develop DCM.

Not all dogs with DCM will show dramatic improvement with taurine supplementation. However, even if plasma and whole blood taurine concentrations are within the reference range, giving taurine supplements to dogs diagnosed with DCM may still have some benefits. Because taurine is extremely safe and inexpensive, the risks and costs of supplementation are minimal, even if dogs have normal levels of plasma and whole blood. Proposed mechanisms for the beneficial actions of taurine on the myocardium include modulating tissue calcium concentrations and availability in the heart; inactivating free radicals and protecting the heart through altering cellular osmolality; osmoregulating the myocardium; directly affecting contractile proteins; and serving as a natural antagonist of angiotension II. Dogs with DCM that do not have taurine deficiency may still benefit from some of these proposed mechanisms of action for taurine.

Table 1
Normal concentrations of taurine in dogs

Plasma (nmol/mL)	Whole blood (nmol/mL)
41–97 ^a	155–347 ^a
72.8–81.2 ^b	255.8–276.2 ^b

^aReference range established from 18 healthy adult beagles consuming a canned commercial maintenance diet. *Data from* Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L-carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am J Vet Res* 2001;62:1616–23.

^bReference range established from 131 healthy adult dogs of various breeds consuming a variety of commercial adult maintenance diets. *Data from* Delaney SJ, Kass PH, Rogers QR, et al. Plasma and whole blood taurine in normal dogs of varying size fed commercially prepared food. *J Anim Physiol* 2003;87:236–44.

Recommended dose for taurine supplementation

This author has successfully used doses of 500 to 1000 mg of taurine administered orally two to three times per day for small dogs (<25 kg), and 1 to 2 g of taurine administered orally two to three times per day for large dogs (25–40 kg). These doses have been shown to normalize plasma and whole blood taurine levels in taurine-deficient dogs. Many other doses for taurine are reported in the literature. Whether a smaller or less frequent dose of taurine than what this author recommends can be used successfully remains to be determined. If doses are used that differ from those this author recommends, plasma and whole blood taurine concentrations must be reevaluated after taurine supplementation is initiated to determine if the dose being given is effective and appropriate. Another important point is that echocardiographic improvement in myocardial function is not usually documented before 2 months of supplementation, and often no improvement is documented before 4 months of supplementation. However, the dogs may feel better clinically and be more active before improvement in cardiac function is documented. Owners must not withdraw taurine supplementation prematurely before deciding if their dogs benefit.

Where Can Taurine be Purchased?

Taurine can be purchased through several retail outlets. If taurine is purchased through a health food store, consumers must look for a product that contains a USP certification symbol on the label. This symbol ensures that what is listed on the label is exactly what is found in the product.

LEVOCARNITINE (L-CARNITINE)

What is L-Carnitine?

L-carnitine (β -hydroxy- γ -trimethylaminobutyric acid) is a small water-soluble molecule with a molecular weight of 160. In dogs, carnitine is obtained either from dietary protein or endogenous synthesis in the liver using the essential precursor amino acids lysine and methionine. Synthesis also requires iron, vitamin C, and vitamin B₆ as cofactors [54]. Although carnitine is classified as an amino acid derivative, it is not an α -amino acid and the amino group is not free. Therefore carnitine is not used for protein synthesis [55].

Carnitine is found in the body either as free carnitine, short-chain acyl carnitine, or long-chain acylcarnitine. Acylcarnitine is carnitine bound to a fatty acid. Total carnitine is the sum of all the individual carnitine fractions. The free carnitine fraction is normally higher than either the short-chain acylcarnitine fraction or the long-chain acylcarnitine fraction.

Cardiac and skeletal muscles are significant storage sites, containing 95% to 98% of the carnitine in the body [56], and carnitine is concentrated in these tissues through an active membrane transport mechanism. The heart is unable to synthesize carnitine and depends on transport of carnitine from the circulation into cardiac muscle, which results in up to a 100 \times gradient between extracellular and intracellular concentrations.

Only the *L*-form of carnitine exists naturally in the body. The *D*-form competitively inhibits the actions of the *L*-form, thereby inhibiting carnitine enzyme systems. In addition, mammals are unable to convert *D*-carnitine to *L*-carnitine, and therefore this discussion focuses on *L*-carnitine.

Why is *L*-Carnitine Important for Normal Myocardial Function?

The normal heart obtains approximately 60% of its total energy production from oxidation of long-chain fatty acids [57]. Long-chain fatty acids in the cytosol of myocardial cells combine with coenzyme A (CoA) as the first step toward beta oxidation. However, long-chain fatty acids must be transported across the inner mitochondrial membrane to generate energy, and the inner mitochondrial membrane is normally impermeable to such bulky polar molecules. Therefore, transport is accomplished through a “carnitine shuttle.” In the carnitine shuttle, the activated fatty acid in the cytosol reacts with carnitine to form a more permeable molecule. This reaction occurs on the outer surface of the inner mitochondrial membrane and is catalyzed by the enzyme carnitine acyltransferase I. The newly formed long-chain acyl-carnitine ester molecule is permeable to the inner mitochondrial membrane and is transported across this membrane, where the enzyme acyltransferase II converts the long-chain acyl-carnitine back to free carnitine and the long-chain fatty acid. Therefore, carnitine functions as a cofactor of several important enzymes necessary for transport of long-chain fatty acids from the cytosol into the mitochondrial matrix [58,59]. Once inside the mitochondria, fatty acids undergo beta oxidation to generate energy [60].

Another important function of carnitine is its buffering capacity, which modulates the intramitochondrial acyl-CoA:CoA ratio [58]. This process is important because acyl-CoA is the activated form of fatty acids used for beta oxidation and lipid synthesis. However, buildup of acyl-CoA derivatives in the mitochondria results in decreased free CoA, which inhibits oxidative metabolism. Acyl-CoA derivatives also act as detergents at high concentrations. Carnitine also facilitates removal of accumulating short- and medium-chain organic acids from the mitochondria. Therefore carnitine also has a role in detoxification in the mitochondria.

What Causes *L*-Carnitine Deficiency?

Carnitine deficiency can be a primary or secondary disorder. Primary carnitine deficiencies may arise from genetic defects in synthesis, renal transport, intestinal absorption, transmembrane uptake mechanisms, or excessive degradation of carnitine [61]. In humans, primary carnitine deficiencies have been associated with cardiomyopathies that are usually not present at birth but take 3 to 4 years to develop. *L*-carnitine therapy can prevent and reverse cardiac dysfunction in some patients.

Secondary carnitine deficiencies are believed to be much more common in humans and can have many causes [61]. In humans, carnitine deficiency can result from inborn errors of metabolism or develop in patients undergoing long-term total parenteral nutrition, vegetarians, and infants fed formulas not

supplemented with carnitine. Carnitine deficiencies are recognized in dogs, but the incidence is not known.

What are the Consequences of *L*-Carnitine Deficiency?

Carnitine deficiency has been shown to cause or be associated with DCM in humans [62–64], hamsters [65,66], and dogs [36,67–69]. More widespread studies have not been undertaken in dogs because carnitine status is difficult to thoroughly assess.

What Types of Carnitine Deficiency Exist in Dogs?

Carnitine deficiency in dogs is classified as either (1) plasma carnitine deficiency, characterized by low concentrations of free plasma carnitine; (2) systemic carnitine deficiency, characterized by low concentrations of free plasma and tissue carnitine; or (3) myopathic carnitine deficiency, characterized by low free myocardial carnitine concentrations in the presence of normal and sometimes elevated plasma carnitine concentrations. Plasma carnitine deficiency alone is not a well-documented state and is included to account for the fact that plasma carnitine, but not tissue carnitine sampling, is often pursued in veterinary medicine.

For example, if plasma carnitine concentration is used to assess carnitine status of a dog, it can help diagnose carnitine deficiency when it is low. However, if plasma carnitine concentration is normal, it does not rule out the possibility of the myopathic form of carnitine deficiency, and the myopathic form of carnitine deficiency is estimated to occur in 17% to 60% of dogs with DCM. Evaluating cardiac muscle carnitine concentrations requires a fluoroscopy-guided endomyocardial biopsy, which is not practical to perform in most private practice situations and is not without risk. Therefore, diagnosing and determining the incidence of myopathic carnitine deficiency in dogs with cardiac disease remains elusive, but may be an underdiagnosed cause of DCM in dogs.

L-Carnitine Deficiency and Associated Myocardial Disease States in Dogs

Carnitine deficiency was associated with DCM in dogs in a limited number of clinical reports [8,9,68–70]. The first reported case of carnitine deficiency was in a family of boxers [69]. The sire, dam, and two littermates were diagnosed with DCM. One offspring had a low plasma carnitine concentration and low myocardial carnitine concentration at DCM diagnosis. After undergoing treatment with high-dose *L*-carnitine (220 mg/kg/d orally), this dog's fractional shortening (FS) increased from 18% to 28%. This dog's littermate had low myocardial and normal plasma carnitine concentrations and responded similarly to high-dose *L*-carnitine supplementation, with its FS increasing from 2% to 24%. The latter dog experienced a decline in myocardial function after *L*-carnitine therapy was withdrawn. Both parents of these littermates had normal plasma and low myocardial carnitine concentrations. Unfortunately, both parents died soon after beginning *L*-carnitine supplementation.

Costa and Labuc [70] presented another case report of two boxers with DCM. One was treated with 250 mg/kg/d of *L*-carnitine orally, and the other was not treated. The myocardial concentration of carnitine was found to be low in the dog that did not receive supplementation and elevated in the dog that did.

Concurrent supplementation with carnitine and taurine has shown benefit in American cocker spaniels with DCM [8]. An unpublished study by this author in 1998 showed beneficial effects from carnitine supplementation in urolith-forming dogs diagnosed with DCM while consuming a protein-restricted diet (Sherry Lynn Sanderson, DVM, PhD, unpublished material). Both studies showed dramatic improvement in myocardial function and survival times in dogs that received supplementation.

Which Came First: Carnitine Deficiency or Dilated Cardiomyopathy?

A common argument made against the role of carnitine deficiency in dogs diagnosed with DCM is that if carnitine deficiency is diagnosed after the onset of DCM, whether carnitine deficiency caused the DCM or DCM caused the carnitine deficiency is unclear. When myocardial cells are damaged, as may occur with DCM, carnitine can leak out of the cells, resulting in low myocardial carnitine levels. In this situation, the DCM caused the carnitine deficiency. Most published studies linking carnitine deficiency to DCM in dogs have shown this scenario when carnitine deficiency was diagnosed after the onset of DCM.

In an unpublished study conducted at the University of Minnesota, this author documented carnitine deficiency before the onset of DCM in three dogs (Sherry Lynn Sanderson, DVM, PhD, unpublished material, 1998). Therefore, the association of carnitine deficiency with DCM at diagnosis may not always imply a cause-and-effect relationship. However, this study indicates that carnitine deficiency can cause DCM in dogs.

Which Dogs with Dilated Cardiomyopathy Should Receive Carnitine Supplementation?

The importance of carnitine supplementation in the treatment and survival times of some dogs with DCM should not be overlooked. In the first reported study linking carnitine deficiency to DCM in boxers, two of four dogs experienced good response to carnitine supplementation [69]. Considering the generally poor prognosis of this disease in boxers, carnitine supplementation provides owners one additional option for treating this disease, and has made a dramatic difference in the survival times and quality of life of some dogs.

The importance of carnitine supplementation in American cocker spaniels with DCM and urolith-forming dogs with DCM should also not be overlooked. Although a few anecdotal reports exist in which American cocker spaniels with DCM experienced good response to taurine supplementation alone, most cases have shown response to combined supplementation with taurine and carnitine. In the above study by this author, a miniature Dachshund diagnosed with carnitine deficiency before the onset of DCM underwent treatment

only with carnitine supplementation, and its heart disease reversed. Although DCM in many dogs is not associated with carnitine deficiency, carnitine and taurine supplementation offer the most promising hope for improved quality of life and survival times in dogs that experience response.

How is Carnitine Deficiency Diagnosed?

Because performing endomyocardial biopsies is impractical for most clinicians in private practice, most screening for carnitine deficiency relies solely on plasma carnitine levels. The method for plasma carnitine sample collection is almost identical to that used for plasma taurine sample collection. Fasting, heparinized, nonhemolyzed blood samples should be obtained and stored on ice until they are processed. The plasma should be immediately separated from the cellular components ideally in a cold-centrifuge, and a small amount of plasma should be left above the buffy coat to prevent contamination of the plasma with cells. Samples should be frozen immediately until analyzed for plasma carnitine concentrations.

What is the Recommended Dose for Carnitine Supplementation in Dogs?

The doses of carnitine being administered may contribute to the lack of favorable results with carnitine supplementation that some investigators observed. The recommended doses for carnitine supplementation in dogs with DCM vary widely in the literature. Although most authors recommend a carnitine dose of 50 to 100 mg/kg orally every 8 hours, the effective dose may depend on the form of carnitine deficiency. In a limited number of cases studied at the University of Minnesota, where pre- and post-carnitine supplemented plasma and cardiac muscle carnitine levels were obtained, this author's clinical impression was that the effective therapeutic dose in dogs with systemic carnitine deficiency was much lower than the effective dose in dogs with myopathic carnitine deficiency.

Some experts speculate that the myopathic form of carnitine deficiency may be caused by a carnitine transport defect in the heart, and much higher plasma levels of carnitine seem to be needed to overcome this defect and achieve normal concentrations of carnitine in the heart than for the systemic form of carnitine deficiency. Based on this work, the dose of carnitine recommended by this author for systemic carnitine deficiency is 100 mg/kg orally every 8 hours. However, if the myopathic form of carnitine deficiency is present or suspected, the author recommends starting carnitine supplementation at 200 mg/kg orally every 8 hours to maximize the chances that carnitine supplementation will improve myocardial function.

Carnitine is a very safe substance. Diarrhea was the only adverse effect of high doses of carnitine, reported in approximately two thirds of dogs. If diarrhea occurs, the highest dose of carnitine that the dog will tolerate without causing diarrhea should be administered. Therefore, like taurine, *L*-carnitine is a safe substance to administer, and, except for the expense, few drawbacks exist to supplementing a dog with DCM with carnitine (carnitine is much more expensive than taurine). Another important point is that the time it takes for

improvement in myocardial function to occur is very similar to that for taurine supplementation. Echocardiographic improvement in myocardial function is not usually documented before 2 months of supplementation with carnitine, and often improvement is not documented for up to 4 months. However, dogs may feel better clinically and be more active before improvement in cardiac function is documented. Owners must not withdraw carnitine supplementation prematurely before determining whether their dogs benefit.

Where Can *L*-Carnitine Be Purchased?

Although *L*-carnitine can be purchased from health food stores, this source is extremely expensive. Purity of the sample is also of great importance. Therefore, only products that contain the USP certification seal should be purchased from health food stores.

L-Carnitine can also be purchased less expensively in bulk. Bulk carnitine can be purchased from Ajinomotousa, Inc (500 Frank W Burr Boulevard; Park Central West; Teaneck, New Jersey). At last check, the company required a minimum purchase of 10 kg at one time. However, the individual expense can be reduced if several owners split an order. If carnitine is purchased in bulk, owners must measure out the carnitine they are giving to their dogs. One teaspoon of carnitine is equivalent to 2 g of carnitine. Therefore, fractions of a teaspoon can be administered if necessary. Owners must be sure to purchase *L*-carnitine, not *D*- or the *DL*- isomers, because *D*-carnitine interferes with *L*-carnitine use.

Which Dogs with Dilated Cardiomyopathy Should be Supplemented With Carnitine?

Carnitine supplementation should be recommended for boxers, American cocker spaniels, and dogs with cystine or urate urolithiasis that are diagnosed with DCM. Even if carnitine deficiency did not cause DCM, supplementing dogs with carnitine does not hurt them, and supplementation may be beneficial even if carnitine deficiency is not present. The major drawback to supplementing dogs with carnitine is the expense and occasional gastrointestinal upset.

What are the Reference Ranges for Carnitine Concentrations in Dogs?

The reference ranges for carnitine concentrations in dogs are listed in Table 2 [69].

SUMMARY

Some newer more promising therapies for dogs with DCM do not involve drugs but rather nutritional supplements. Two of the more common nutritional supplements administered to dogs with DCM are taurine and carnitine. Deficiencies of these nutrients have been shown to cause DCM in dogs, and some breeds have been shown to experience dramatic improvement in myocardial function after supplementation with one or both nutrients. Although most dogs diagnosed with DCM do not have a documented taurine or carnitine deficiency, they may still benefit from supplementation. Both nutrients are very

Table 2
Normal concentrations of carnitine in dogs

Carnitine fraction	Plasma carnitine (nmol/mL)	Cardiac muscle carnitine (nmol/mg of NCP)
Free	8.36	4.11
Esterified	0.7	0.4
Total	12.38	5.13

Abbreviation: NCP, noncollagenous protein.

safe to administer to dogs. For some owners, the high cost of carnitine is the only deterrent to giving their dogs supplements of both nutrients.

References

- [1] Cobb MA. Idiopathic dilated cardiomyopathy: advances in etiology, pathogenesis and management. *J Small Anim Pract* 1992;33:112-8.
- [2] Keene BW. Canine cardiomyopathy. In: Kirk RW, editor. *Current veterinary therapy X. Small animal practice*. Philadelphia: WB Saunders Co; 1989. p. 240-51.
- [3] Sisson DD, Thomas WP, Keene BW. Primary myocardial disease in the dog. In: Ettinger SJ, Feldman EC, editors. *Textbook of veterinary internal medicine. Diseases of the dog and cat*. 5th edition. Philadelphia: WB Saunders Co; 2000. p. 874-95.
- [4] Buchanan JW. Causes and prevalence of cardiovascular disease. In: Kirk RW, Bonagura JD, editors. *Kirk's current veterinary therapy XI*. Philadelphia: WB Saunders Co; 1992. p. 647-55.
- [5] Sisson D, Thomas WP. Myocardial diseases of dogs and cats. In: Ettinger S, editor. *Textbook of veterinary internal medicine*. 4th edition. Philadelphia: WB Saunders; 1995. p. 995-1032.
- [6] Fioretti M, Delli CE. Epidemiological survey of dilatative cardiomyopathy in dogs [abstract]. *Veterinaria* 1988;2:81.
- [7] Gooding JP, Robinson WF, Wyburn RS, et al. A cardiomyopathy in the English cocker spaniel: a clinico-pathological investigation. *J Small Anim Pract* 1982;23:133-48.
- [8] Kittleson MD, Keene B, Pion PD, et al. Results of the Multicenter Spaniel Trial (MUST): taurine and carnitine responsive dilated cardiomyopathy in American cocker spaniels with decreased plasma taurine concentration. *J Vet Intern Med* 1997;11:204-11.
- [9] Sanderson S, Osborne C, Ogburn P, et al. Canine cystinuria associated with carnitinuria and carnitine deficiency [abstract]. *J Vet Intern Med* 1995;9:212.
- [10] Calvert CA. Dilated congestive cardiomyopathy in Doberman pinchers. *Compend Contin Educ Pract Vet* 1986;8:417-30.
- [11] Monnet E, Orton C, Salman M, et al. Idiopathic dilated cardiomyopathy in dogs: survival and prognostic indicators. *J Vet Intern Med* 1995;9:12-7.
- [12] Tidholm Am Svensson H, Sylven C. Survival and prognostic factors in 189 dogs with dilated cardiomyopathy. *J Am Anim Hosp Assoc* 1997;33:364-8.
- [13] Tidholm A, Johsson L. A retrospective study of canine dilated cardiomyopathy (189 cases). *J Am Anim Hosp Assoc* 1997;33:544-50.
- [14] Tenaglia A, Cody R. Evidence for a taurine deficient cardiomyopathy. *Am J Cardiol* 1988;62:136-9.
- [15] Hayes KC. Taurine requirement in primates. *Nutr Rev* 1985;43:65-70.
- [16] Hayes KC, Carey RE, Schmidt SY. Retinal degeneration associated with taurine deficiency in the cat. *Science* 1975;188:949-51.
- [17] Huxtable RJ. From heart to hypothesis: a mechanism for the calcium modulatory actions of taurine. *Adv Exp Med Biol* 1987;217:371-87.

- [18] Pion PD, Kittleson MD, Rogers QR, et al. Myocardial failure in cats associated with low plasma taurine: a reversible cardiomyopathy. *Science* 1987;237:764-8.
- [19] Schaffer SW, Seyed Mozaffari M, Kramer J, et al. Effect of taurine depletion and treatment on cardiac contractility and metabolism. *Prog Clin Biol Res* 1985;179:167-75.
- [20] Takihara K, Azuma J, Awata N, et al. Beneficial effect of taurine in rabbits with congestive heart failure. *Am Heart J* 1986;112:1278-84.
- [21] Franconi F, Bennardini F, Mattana A, et al. Taurine levels in plasma and platelets in insulin dependent and non insulin dependent diabetes mellitus: correlation with platelet aggregation. *Adv Exp Med Biol* 1994;359:419-23.
- [22] Green TR, Fellman JH. Effect of photolytically generated riboflavin radicals and oxygen on hypotaurine antioxidant free radical scavenging activity. *Adv Exp Med Biol* 1994;359:19-29.
- [23] Rebel G, Petegnief V, Lleu P, et al. New data on the regulation of taurine uptake in cultured nervous cells. *Adv Exp Med Biol* 1994;359:225-32.
- [24] Schmidt SY. Biochemical and functional abnormalities in retinas of taurine deficient cats. *Fed Proc* 1980;39:2706-8.
- [25] Sturman JA, Hayes KC. The biology of taurine in nutrition and developments. *Adv Nutr Res* 1980;3:231-99.
- [26] Sturman JA. Dietary taurine and feline reproduction and development. *J Nutr* 1991;121:S166-70.
- [27] Huxtable RJ, Chubb J, Asari J. Physiological and experimental regulation of taurine content in the heart. *Fed Proc* 1980;39:2685-90.
- [28] Schaffer SW, Kramer J, Chovan JP. Regulation of calcium homeostasis in the heart by taurine. *Fed Proc* 1980;39:2691-4.
- [29] Huxtable RJ. Physiological actions of taurine. *Physiol Rev* 1992;72:101-63.
- [30] Hamaguchi T, Azuma J, Schaffer S. Interaction of taurine with methionine: inhibition of myocardial phospholipids methyltransferase. *J Cardiovasc Pharmacol* 1991;18:224-30.
- [31] Lake N. Loss of cardiac myofibrils: mechanism of contractile deficits induced by taurine deficiency. *Am J Physiol* 1993;264(4 Part 2):H1323-6.
- [32] Steele DS, Smith GL, Miller DJ. The effects of taurine on Ca^{2++} uptake by the sarcoplasmic reticulum and Ca^{2++} sensitivity of chemically skinned rat heart. *J Physiol* 1990;422:499-511.
- [33] Gentile S, Bologna E, Terracina D, et al. Taurine induced diuresis and natriuresis in cirrhotic patients with ascites. *Life Sci* 1994;54:1585-93.
- [34] Jacobsen JG, Thomas LL, Smith LH Jr. Properties and distribution of mammalian L cysteine sulfinic carboxylases. *Biochim Biophys Acta* 1964;85:113-6.
- [35] Pion PD, Kittleson MD, Thomas WP, et al. Clinical findings in cats with dilated cardiomyopathy and relationship to finding taurine deficiency. *J Am Vet Med Assoc* 1992;201:267-74.
- [36] Pion PD, Sanderson SL, Kittleson MD. The effectiveness of taurine and levocarnitine in dogs with heart disease. *Vet Clin North Am Small Anim Pract* 1998;28:1495-514.
- [37] Moise NS. Cardiomyopathy in the fox and association with low dietary taurine. In: *Proceedings of the Seventh American College of Veterinary Internal Medicine Forum*. 1989. p. 834-5.
- [38] Kramer GA, Kittleson MD, Fox PR, et al. Plasma taurine concentration in normal dogs and dogs with heart disease. *J Vet Intern Med* 1995;9:253-8.
- [39] Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein restricted diets. *Am J Vet Res* 2001;62:1616-23.
- [40] Association of American Feed Control Officials, Inc. Model bill and regulation. AAFCO Official Publication; 2003.
- [41] Backus RC, Cohen G, Pion PD, et al. Taurine deficiency in Newfoundlands fed commercially available complete and balanced diets. *J Am Vet Med Assoc* 2003;223:1130-6.

- [42] National Research Council. Nutrient requirements of dogs, revised 1985. Washington (DC): National Academy Press; 1985.
- [43] Spitze AR, Wong DL, Rogers QR, et al. Taurine concentrations in animal feed ingredients; cooking influences taurine content. *J Anim Physiol* 2003;87:251-62.
- [44] Kim SW, Morris JG, Rogers QR. Dietary soybean protein decreases plasma taurine in cats. *J Nutr* 1995;125:2831-7.
- [45] Fascetti AJ, Reed JR, Rogers QR, et al. Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997-2001). *J Am Vet Med Assoc* 2003;112:1137-41.
- [46] Belanger MC, Quillet M, Queney G, et al. Taurine deficient dilated cardiomyopathy in a family of Golden Retrievers. *J Am Anim Hosp Assoc* 2005;41:284-91.
- [47] Torres CL, Backus RC, Fascetti AJ, et al. Taurine status in normal dogs fed a commercial diet associated with taurine deficiency and dilated cardiomyopathy. *J Anim Physiol* 2003;87:359-72.
- [48] Johnson ML, Parsons CM, Fahey GC, et al. Effects of species raw material source, ash content and processing temperature on amino acid digestibility of animal by-product meals by cecectomized roosters and ileally cannulated dogs. *J Anim Sci* 1998;76:1112-22.
- [49] Delaney SJ, Kass PH, Rogers QR, et al. Plasma and whole blood taurine in normal dogs of varying size fed commercially prepared food. *J Anim Physiol* 2003;87:236-44.
- [50] Stratton Phelps M, Backus RB, Rogers QR, et al. Dietary rice bran decreases plasma and whole blood taurine in cats. *J Nutr* 2002;132:1745S-7S.
- [51] Trautwein EA, Hayes KC. Taurine concentration in plasma and whole blood in humans: estimation of error from intra- and interindividual variation and sampling technique. *Am J Clin Nutr* 1990;52:758-64.
- [52] Pion PD, Lewis J, Greene K, et al. Effects of meal feeding and food deprivation on plasma and whole blood taurine concentration in cats. *J Nutr* 1991;121:S177-8.
- [53] Sanderson SS, Osborne C, Gross K, et al. Reliability of canine plasma and whole blood taurine concentrations as indicators of cardiac and skeletal muscle taurine concentrations. [abstract]. *J Vet Intern Med* 1998;12:224.
- [54] Bremer J. Carnitine metabolism and function. *Physiol Rev* 1983;63:1420-80.
- [55] Leibowitz BE. Carnitine. *Adv Res Press* 1987;2:1-13.
- [56] Rebouche CJ, Engel AG. Kinetic compartmental analysis of carnitine metabolism in the dog. *Arch Biochem Biophys* 1983;220:60-70.
- [57] Neely JR, Morgan HA. Relationship between carbohydrate metabolism and energy balance of heart muscle. *Annu Rev Physiol* 1974;36:413-59.
- [58] Stumpt DA, Parker WD Jr, Angelini C. Carnitine deficiency, organic acidemias, and Reye's syndrome. *Neurology* 1985;35:1041-5.
- [59] Gilbert EF. Carnitine deficiency. *Pathology* 1985;17:161-9.
- [60] Mayes PA. Oxidation of fatty acids: ketogenesis. In: Murray RK, Granner DK, Mayes PA, et al, editors. *Harper's biochemistry*. 24th edition. Norwalk (CT): Appleton & Lange; 1996. p. 224-35.
- [61] Paulson DJ. Carnitine deficiency induced cardiomyopathy. *Mol Cell Biochem* 1998;180:33-41.
- [62] Periera RR, Scholte HR, Luyt Houwen IEM, et al. Cardiomyopathy associated with carnitine loss in kidneys and small intestines. *Eur J Pediatr* 1988;148:193-7.
- [63] Pierpont MEM. Carnitine and myocardial function. In: Carter AL, editor. *Current concepts in carnitine research*. 1st edition. Boca Raton (FL): CRC Press; 1992. p. 197-213.
- [64] Paulson DJ, Sanjak M, Shug AL. Carnitine deficiency and the diabetic heart. In: Carter AL, editor. *Current concepts in carnitine research*. 1st edition. Boca Raton (FL): CRC Press; 1992. p. 215-30.
- [65] Whitmar JT. Energy metabolism and mechanical function in perfused hearts of Syrian hamsters with dilated or hypertrophic cardiomyopathy. *J Mol Cell Cardiol* 1986;18:307-17.

- [66] Whitmar JT. L Carnitine treatment improves cardiac performance and restores high energy phosphate pools in cardiomyopathic Syrian hamsters. *Circ Res* 1987;61:396-408.
- [67] Sanderson S, Ogburn P, Osborne C. Heart disease management Indications for nondrug therapies. *Vet Forum* 1996;13:36-43.
- [68] McEntee K, Clercx C, Snaps F, et al. Clinical, electrocardiographic, and echocardiographic improvements after L carnitine supplementation in a cardiomyopathic Labrador. *Canine Pract* 1995;20:12-5.
- [69] Keene BW, Panciera DP, Atkins CE, et al. Myocardial L carnitine deficiency in a family of dogs with dilated cardiomyopathy. *J Am Vet Med Assoc* 1991;198:647-50.
- [70] Costa ND, Labuc RH. Case report: efficacy of oral carnitine therapy for dilated cardiomyopathy in boxer dogs. *J Nutr* 1995;124(supp):2687S-92S.

SEVERE REVERSIBLE DILATED CARDIOMYOPATHY AND HYPERTHYROIDISM: CASE REPORT AND REVIEW OF THE LITERATURE

Cristina Boccalandro, MD,¹ Fernando Boccalandro, MD,² Philip Orlander, MD,¹ and Chik Fong Wei, MD²

ABSTRACT

Objective: To describe a case of a 46-year-old woman with Graves' disease and reversible low-output congestive heart failure and present a comparative analysis of 23 similar cases reported in the literature.

Methods: A detailed case report is presented. In addition, a review of the pertinent literature published between 1960 and 2002 was performed to identify similar cases of dilated cardiomyopathy and thyrotoxicosis and to assess the findings in these patients.

Results: A 46-year-old woman without primary heart disease was admitted to the hospital with Graves' thyrotoxicosis and severe low-output congestive heart failure. Her left ventricular ejection fraction (LVEF) at the time of initial assessment was less than 20%, and her condition was categorized as New York Heart Association (NYHA) functional class III. Nineteen months after she was treated for hyperthyroidism, her LVEF was 49% and her status was NYHA class I. A severe hypotensive episode occurred when β -adrenergic blockade therapy was initiated. The group of 23 similar cases from the literature plus our currently described patient had a mean age of 45 years, a male-to-female ratio of 1:1.2, Graves' disease as the principal cause, and LVEF improvement from 29% to 58%.

Conclusion: Dilated cardiomyopathy is an unusual manifestation of hyperthyroidism with unclear cause. Clinicians should be aware of this entity because it is treatable and hypotension can occur if β -adrenergic blockade treatment is initiated. (*Endocr Pract* 2003;9:140-146)

(b) (6)

(b) (4)

Submitted for publication February 15, 2002

Accepted for publication August 7, 2002

From the ¹Division of Endocrinology and Metabolism and ²Division of Cardiology, University of Texas at Houston, Houston, Texas.

Address correspondence and reprint requests to Dr. Fernando Boccalandro, Division of Cardiology, University of Texas at Houston, MSB 1246, Houston, TX 77030.

© 2003 AACE.

(b) (4)

(b) (4)

(b) (4)

(b) (4)

(b) (4)

(b) (4)

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

14768

Sample Submission Form

Amino Acid Laboratory
 University of California, Davis
 1020 Vet Med 3B
 1089 Veterinary Medicine Drive
 Davis, CA 95616
 Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
 Non-federal funds ID/Account Number
 to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/Index.cfm>

Vet/Tech Contact: _____ (b) (6)
 Company Name: _____ (b) (6)
 Address: _____ (b) (6)

 Email: _____ (b) (6)
 Tel: _____ (b) (6) Fax: _____ (b) (6)

Billing Contact: Finance Dept. TAX ID: _____
 Email: _____ (b) (6) Tel: _____ (b) (6)

Patient Name: _____ (b) (6) _____ (b) (6)
 Species: Feline
 Owner's Name: _____ (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
 Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 196 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150



REVIEW

Open Access



Taurine: the appeal of a safe amino acid for skeletal muscle disorders

Annamaria De Luca*, Sabata Pierno and Diana Conte Camerino

Abstract

Taurine is a natural amino acid present as free form in many mammalian tissues and in particular in skeletal muscle. Taurine exerts many physiological functions, including membrane stabilization, osmoregulation and cytoprotective effects, antioxidant and anti-inflammatory actions as well as modulation of intracellular calcium concentration and ion channel function. In addition taurine may control muscle metabolism and gene expression, through yet unclear mechanisms. This review summarizes the effects of taurine on specific muscle targets and pathways as well as its therapeutic potential to restore skeletal muscle function and performance in various pathological conditions. Evidences support the link between alteration of intracellular taurine level in skeletal muscle and different pathophysiological conditions, such as disuse-induced muscle atrophy, muscular dystrophy and/or senescence, reinforcing the interest towards its exogenous supplementation. In addition, taurine treatment can be beneficial to reduce sarcolemmal hyper-excitability in myotonia-related syndromes. Although further studies are necessary to fill the gaps between animals and humans, the benefit of the amino acid appears to be due to its multiple actions on cellular functions while toxicity seems relatively low. Human clinical trials using taurine in various pathologies such as diabetes, cardiovascular and neurological disorders have been performed and may represent a guide-line for designing specific studies in patients of neuromuscular diseases.

Keywords: Taurine skeletal muscle, Inherited muscle disorders, Disuse muscle atrophy, Development and aging, Skeletal muscle performance


Background

Taurine (2-aminoethane-sulfonic acid) is a sulfur-containing amino acid which is not used for protein synthesis and is therefore the most abundant free amino acid in mammalian tissues, with the exception of human liver in which aspartate is the most abundant one [1, 2]. The intracellular concentration of taurine ranges between 5 and 20 $\mu\text{mol/g}$ wet weight in many tissues, especially in excitable ones, such as brain, heart and skeletal muscle [1, 3, 4]. Endogenous synthesis occurs in the liver via the cysteine sulfinic acid pathway. The metabolic reaction consists in a first oxidation of the sulfhydryl group of cysteine to cysteine sulfinic acid by the enzyme cysteine dioxygenase. Cysteine sulfinic acid is then decarboxylated to hypotaurine by the cysteine sulfinic acid decarboxylase.

Taurine is obtained by a yet unclear spontaneous or enzymatic oxidation (by hypotaurine dehydrogenase) of hypotaurine (Fig. 1). The endogenous synthesis of taurine is highly variable between individuals also in relation to nutritional state, to the amount of protein intake and to cysteine availability [1, 5]. In turn the availability of cysteine is highly dependent on the metabolic equilibrium between homocysteine and methionine, via folic acid, vitamin B12 and the efficiency of the enzyme methyltetrahydrofolate reductase. In addition, a certain amount of taurine has to be introduced with food, mostly in carnivores and, to a minor extent, in omnivores [1]. The importance of the two sources vary quite a lot between species, with some, like felines and foxes, being highly dependent on diet acquisition of taurine, as they are unable to synthesize it. These species are also particularly susceptible to deficient states, developing severe pathophysiological conditions, such as dilated cardiomyopathy, retinal degeneration and reproduction defects

*Correspondence: annamaria.deluca@uniba.it
Sezione di Farmacologia, Dipartimento di Farmacia-Scienze del Farmaco,
Università degli Studi di Bari "Aldo Moro", Bari, Italy





[3, 6]. These evidences first outlined the key role of taurine for mammalian tissue functions and helped to better understand the link between tissue distress in retaining proper taurine concentration and various pathophysiological conditions.

In fact, even in species able to synthesize taurine, the tissue-specific synthesis is relatively low, with liver being the main source according to the higher expression of enzymes as cysteine dioxygenase. Importantly, the activity of this latter enzyme strictly depends upon cysteine availability, so that the exact amount of taurine being endogenously synthesized is difficult to predict [7]. However, the high intracellular concentration is guaranteed by the presence of a specific active transporter that concentrates taurine inside the cells against gradients. The taurine transporter (TauT; encoded by the SLC6A6 gene) is a sodium and chloride ion-dependent transporter ubiquitously expressed in mammalian tissues. The concentration of taurine is 100-fold less in the plasma (20–100 μM) than in the tissues, suggesting that it is indeed required for modulating key cellular functions. Due to the high tissue concentration, taurine also works as an osmolyte. Its cellular efflux via volume-dependent or volume-independent pathways works to osmotically balance the excessive production of metabolic by-products. Both uptake systems and efflux pathways are tightly regulated at transcriptional and post-transcriptional level, leading to an accurate control of taurine intracellular levels [8].

Since its discovery in ox bile in 1827, several physiological functions have been described for the amino acid, ranging from the classical role of conjugating agent for

bile acids, to wider actions as osmotic pressure regulator, modulator of calcium homeostasis and signaling and, more recently, as an endogenous anti-oxidant and anti-inflammatory compound in various tissues. The mechanism by which taurine exerts all these different functions is still unclear. Some of the taurine actions in central nervous system (CNS), seem to occur via specific binding sites or receptors, i.e. in thalamus taurine modulates neuronal firing via activation of extra-synaptic gamma-amino butyric acid (GABA) receptor isoforms $\alpha 4\beta 2\delta$ with a greater affinity than GABA [9–12]. Such high affinity binding sites have not been evidenced in other tissues.


Skeletal muscle is one of the tissues able to concentrate the largest amount of body's taurine, via the TauT activity. Pioneer studies of Ryan Huxtable anticipated that the high taurine level is needed to maintain an appropriate calcium homeostasis, likely by ensuring a correct calcium re-uptake by the sarcoplasmic reticulum [13]. Similar actions were also described in heart, with taurine exerting complex modulation of calcium homeostasis in relation to external concentration of the cation with beneficial effects in contrasting arrhythmias or heart failure [1, 3, 4].

Transgenic mice lacking TauT gene have been generated by two separate groups [6, 14–16]. In line with a key role of taurine for maintaining proper physiological functions, the drastic reduction in content consequent to TauT deletion is associated to a variety of disorders in various tissues, such as eye, kidney, heart, nociceptive system and skeletal muscle [14–17]. These conditions resemble those occurring when taurine tissue content is

altered by pathophysiological states or by inhibitors of the taurine transporter. In spite the pre-clinical research has disclosed many conditions in which taurine supplementation may be beneficial, the therapeutic use of taurine is very limited. Taurine is commonly known for its claimed effects as energizer and anti-fatigue compound and it is present in many energy soft drinks as well as in supplement cocktails for athletes. The toxicity of taurine in this context is considered relatively low with respect to other active ingredients; actually it may also be protective against cardiovascular action of caffeine [18]. Such a protection may again result from multiple taurine actions, i.e. an antihypertensive effect via vasodilatation (by reducing adrenergic and angiotensin II actions as well as calcium-induced vasospasm) along with a reduced risk of cardiac arrhythmias via modulation of ion channels and ionic homeostasis [18]. However a certain caution is important especially when taurine is used in children and/or in association with drugs, alcohol or other food supplements [19–23]. Apart for its nutraceutical role,

taurine may exert clear pharmacological actions by modulating signaling pathways and targets or via restoration of its altered tissue levels. No systematic toxicity studies have been performed to assess the toxicological parameters for taurine; however human trials have used taurine up to 10 g/daily without overt signs of toxicity. This may also depend on the direct relationship between taurine plasma level and its excretion rate by the kidney [19].

An extensive revision of all the actions of taurine in various tissues and the wide potential usefulness of its supplementation is out of the scope of this review. However, a general overview is provided in Fig. 2. As far as inherited or acquired pathophysiological conditions of skeletal muscle are concerned, the pre-clinical findings allow to distinguish effects related to exogenous pharmacological action of taurine on rather specific targets, such as in myotonic syndromes, to conditions that may be accompanied by changes in intercellular taurine content or change in calcium homeostasis, in which a taurine supplementation may be helpful to restore altered levels.



The present review is aimed at providing the state-of-art of taurine research in skeletal muscle, with particular attention to its potential therapeutic application as orphan drug in inherited rare muscle disorders, as well as in pathophysiological conditions such as aging, malnutrition and/or muscle disuse.

Skeletal muscle ion channels as specific targets of taurine: the potential action of taurine as anti-myotonic drug

Taurine and skeletal muscle chloride channels CLC-1

In CNS, taurine has been long claimed to act as an “inhibitory” amino acid and neurotransmitter [1]. Neuronal synthesis of taurine and metabotropic taurine receptors have been described in specific areas of CNS, where taurine acts in a glycine or GABA-like manner, by enhancing hyperpolarizing chloride-mediated conductance in nervous cells [9, 11, 12]. Pre-clinical evidences were provided of a beneficial effect of taurine in controlling/preventing seizure discharges and neurotoxicity [1, 12, 24]. The ability of taurine to act as inhibitory amino acid raised attention to its possible effect as potential membrane stabilizer in skeletal muscle. We investigated about the actions of the amino acid on voltage-gated chloride channels CLC-1 that account for the macroscopic chloride conductance (gCl) of skeletal muscle. Resting gCl accounts for about 70–90% to the total membrane conductance of sarcolemma and plays a pivotal role in maintaining the sarcolemmal electrical stability by shunting the depolarization-driven potassium accumulation in transverse tubules. Thus the large gCl allows repolarization and muscle relaxation.

Loss-of-function mutations of CLC-1 are responsible of myotonic syndromes with either autosomal dominant (Thomsen disease) or recessive pattern of inheritance (Becker’s Myotonia Congenita). The resulting decrease of gCl is responsible for the pathological hyperexcitability and for the delayed relaxation, spasms and stiffness typical of the disease in both patients and myotonic animals [25–27].

Our research has shown that taurine, acutely applied *in vitro*, exerts a concentration-dependent increase of gCl in rat extensor digitorum longus (EDL) myofibers, and in parallel reduces membrane excitability [28, 29]. The effective concentrations are in the millimolar range, likely in relation to the high intracellular level of the amino acid [28, 29]. A pre-clinical evaluation of the potential anti-myotonic activity of taurine has been performed. We found that taurine does not antagonize the myotonic discharges in rats made myotonic by administration of anthracene-9-carboxylic acid, a direct chloride channel blocker, nor does it restore gCl lowered *in vitro* by the same agent. However, when rats are made myotonic

by a chronic exposure to 20,25 diazacholesterol, which reduces gCl indirectly by modifying lipid membrane composition, taurine antagonizes the electromyographic signs of myotonia if administered *in vivo*, while its acute *in vitro* application contrasts both the reduced gCl and the high frequency firing of single myofibers [30]. These results suggested that taurine can contrast myotonia if chloride channels are available for a direct modulation, implying its direct action at channel level or on a site nearby. A series of taurine analogues were tested on gCl of rat EDL myofibers to investigate the structure–activity relationship (SAR) between taurine and chloride channels. The results provided a pharmacological evidence of the presence of a specific low-affinity taurine binding site able to modulate chloride channel function and/or kinetic [31]. In particular, an increased distance between the two charged heads of taurine and/or a more distributed positive charge for the replacement of the amino group with aza-cyclo moieties lead to a decreased potency in enhancing gCl [31]. The direct action of taurine on skeletal muscle chloride channel was further confirmed by two microelectrode voltage-clamp recordings of chloride currents sustained by human CLC-1 channel heterologously expressed in *Xenopus* oocytes. In these conditions, the *in vitro* application of 20 mM taurine enhanced by 100% the chloride currents and shifted channel activation toward more negative potentials, an effect that likely accounts for the increase in resting gCl observed in native fibers [32–34]. This direct modulation adds to other possible homeostatic and modulatory roles that the high intracellular taurine has on chloride channels. However, as anticipated, the acute modulation of gCl may require fully or partly functional chloride channels, questioning about the real efficacy of taurine in CLC-1 related myotonic syndromes, especially for those mutations that seriously affect channel expression and protein level. Taurine has been tested in patients with myotonic dystrophy with encouraging results. In particular acute parenteral administrations of taurine allowed to reduce membrane excitability evaluated in relation to potassium plasma concentration after potassium-enriched infusion, suggesting again an action on membrane ionic conductance. Accordingly, a double-blind oral administration of taurine led to a long-term control of myotonic symptoms estimated as reduction of electromyographic (EMG) discharges and potassium induced-hyperexcitability [35–37]. Even taking into account the possible bias deriving from these small sized trials, the effects of taurine in myotonic dystrophy patients suggest alternative modality for decreasing membrane excitability. In fact, myotonic dystrophy type 1 (DM1) or Steinardt syndrome, is caused by expansion of a CTG trinucleotide repeat in the non-coding region of DM protein kinase with abnormalities

in mRNA metabolism and alternative splicing of certain genes. In DM1 patients, the abnormal inclusion of alternative exons 6B and/or 7A and retention of intron 2 of CLC-1 channel gene (*CLCN1*) gene have been observed. These aberrant-splicing, which may also occur in myotonic dystrophy type 2 (DM2) patients, leads to premature termination codons, with a consistent decrease of the mRNA of *CLCN1*, of CLC-1 protein and consequently of gCl [38, 39]. Therefore, the possible modulatory action of taurine on other skeletal muscle ion channels has to be taken into account.

Taurine and Nav1.4 voltage gated sodium channels

It is feasible to hypothesize a modulation by taurine of the skeletal muscle isoform of voltage-gated sodium channel (Nav1.4), involved in the generation and propagation of action potential and main target of symptomatic anti-myotonic drugs [37, 40]. The effect of taurine on sodium channels of native muscle fibers has been investigated in our laboratories by cell-attached patch clamp recordings. Taurine has a dual effect. In particular taurine enhances the sodium transients elicited by depolarizing test pulses close to the threshold for channel activation (test pulse to $-70/-50$ mV), an effect that is likely related to the observed shift of the activation curve towards more negative potentials. However, taurine reduces sodium currents at more depolarized test pulse potentials, with a 50% inhibition of the maximal peak sodium current observed at 10 mM taurine. In parallel, a left-shift of the steady-state inactivation curve has been observed, indicating the ability of taurine to stabilize the blocked channels in the inactivated state [34, 41 Desaphy and Conte Camerino, unpublished observation]. This peculiar effect of taurine on Nav1.4 channel is similar to what has been observed on cardiac sodium currents [42, 43] and underlines a complex action of the amino acid on sodium channel gating and kinetic. Our extensive structure–activity relationship studies of inhibitors of Nav1.4 channel allow to predict that the anesthetic-like action of taurine is mediated by the amino group, a main pharmacophore moiety in sodium channel blockers [44–47]. The dual ability of taurine to open chloride channels and to block sodium channels envisages a greater therapeutic action of the amino acid in myotonic states related to gain-of-function mutations of sodium channels, such as Sodium Channel Myotonia and Paramyotonia Congenita. The verification that taurine is able to compensate mutation-related biophysical alterations of Nav1.4 channels will be helpful at this regard, and is part of future projects of our laboratory. For the moment, the action of taurine on sodium channels can account for the antimyotonic effect in conditions where chloride channels are defective or dysfunctional [35, 36]. In line with this, the mechanism of

taurine action on Nav1.4 sodium channels deserves to be further investigated since it may better support its pharmacological potential and its clinical use in hyperexcitability muscle disorders (Table 1).

Role of proper taurine intramuscular level for excitation–contraction coupling and muscle performance

The ability of skeletal muscle to concentrate taurine against gradient pushed toward a better understanding of its physiological role. Adult rats were chronically treated with guanidinoethane sulfonate (GES), an inhibitor of taurine transporter (TauT) to induce a reduction of taurine content in skeletal muscle. We found that a 50% reduction of taurine in EDL muscle leads to a marked decrease in gCl, and to a parallel enhancement of sarcolemmal excitability, disclosing the ability of taurine level to exert a physiological control on chloride channel function and sarcolemmal stability [48]. The mechanism underlying this effect is not clear yet, but we cannot rule out the ability of taurine to modulate CLC-1 channel function via a fine-tuning of a calcium-dependent phosphorylation-signaling pathway, as discussed below. In line with the described ability of taurine to control calcium homeostasis in both skeletal muscle and cardiac tissue [1, 4], we found a marked alteration of mechanical threshold, i.e. the voltage at which muscle fiber contracts in response to depolarizing voltage steps, in taurine-depleted EDL myofibers. Mechanical threshold depends on the kinetic of calcium release from and reuptake by sarcoplasmic reticulum, also in relation to basal cytosolic calcium concentrations. Taurine depleted EDL muscle fibers contract at more negative potentials with respect to normal ones, implying an impact of GES treatment on calcium handling [48, 49]. Both the decrease in gCl and the shift of mechanical threshold toward negative potentials were rapidly reverted by in vitro application of millimolar concentration of taurine. Actually, depleted muscles showed a higher than normal sensitivity to exogenous taurine with respect to normal ones [48], further corroborating the link between the observed alterations and the taurine level. The contractile properties and fatigability of EDL muscles depleted of taurine by a GES treatment were investigated by Bakker's group. It was found that the treatment with GES decreases muscle taurine levels to <40% of controls and decreases the peak twitch force of EDL muscles by 20%. Also, GES-treated muscles develop a lower force in force–frequency relationship and show a slower time to fatigue, likely in relation to the lower metabolic demands of the weaker muscles [50]. Primary information about the long-term effect of taurine in skeletal muscle and, consequently, of potential usefulness of its exogenous administration

Table 1 Involvement and therapeutic potential of taurine in physio-pathological conditions and diseases of skeletal muscle

Condition	Change in Taurine content / TauT	Pathogenetic mechanisms related to changes in taurine content	General symptoms	Taurine targets	Therapeutic Potential of Taurine
Post-natal development	Age-dependent increase in TauT expression and intracellular content	Delayed development and delayed acquisition of specific phenotypic properties; metabolic dysfunction	Species-specific (due to different sensitivity to taurine deficiency)	Mitochondria; ion channels; calcium homeostasis and calcium dependent gene expression	Taurine supplementation in formula for pre-term born infants; to ensure a proper skeletal muscle phenotype differentiation
Aging	Decrease in Taurine content; no information on TauT expression	Metabolic distress; calcium dependent dysfunction; reduced regenerating ability; reduced activity of free-oxygen radicals scavengers	Sarcopenia; atrophy, weakness and fatigue degeneration, altered excitation-contraction coupling, impaired performance	Ion channels; Calcium homeostasis; oxidative stress and atrophy	To counteract the decrease in taurine content and the consequent reduction in chloride channel function and the alteration in calcium ion homeostasis; to ameliorate performance and muscle strength
Ischemia and reperfusion injury	Decrease due to a compensatory taurine efflux	Insufficient vaso-dilation in relation to muscle work; metabolic distress; oxidative stress	Hyperkalemia, muscle dysfunction; ROS-induced inflammation and damage	Metabolic-sensitive channels; mitochondria	To counteract hyper-kalemia by inhibiting K_{ATP} and KCa^{2+} channels; to prevent ischemia-induced taurine loss
Myotonic syndromes and periodic paralyses	Unknown	Primary inherited channelopathies due to loss-of-function mutations of ClC-1 chloride channel or gain-of-function mutations of Nav1.4 sodium channel	Hyperexcitability and impaired muscle relaxation	ClC-1 chloride channel; Nav1.4 sodium channel	To reduce membrane hyper-excitability through: opening of chloride channel and increase in gCl mediated by both short and long term actions; modulation of generation and propagation of action potential, by blocking sodium channel with a local-anesthetic like mechanism
Disuse	Slow-to-fast decrease in taurine content; no change in TauT expression	Myofiber phenotype transition in postural muscle; atrophy	Atrophy, change in metabolism, slow-to-fast transition; weakness	Ion channel function and expression; calcium homeostasis	To counteract disuse-induced taurine loss; to counteract myofiber transition; potential counteraction of atrophy
Duchenne muscular dystrophy and related myopathies	Change in content related to pathology phase; possible reduction of TauT expression	Alteration of calcium homeostasis; calcium-related degeneration; oxidative stress and inflammation	Progressive muscle degeneration and weakness; muscle fiber loss and fibrosis; sarcolemmal instability; altered calcium homeostasis; inflammation and oxidative stress	Chloride channel and voltage-insensitive calcium permeable channels (Leak/TRP-like); SERCA; mitochondria	To ameliorate muscle performance; to counteract taurine loss and to modulate calcium availability for contraction; to counteract contraction-induced ischemia. To contrast degeneration-induced decrease in gCl; adjuvant therapy in combination with glucocorticoids

The table summarizes the main role of taurine in various conditions of skeletal muscle, indicating evidences in relation to changes in tissue content and potential site of taurine action. Please refer to text for more detailed information and specific references.

TauT taurine transport system, *SERCA* sarco/endoplasmic reticulum calcium ATPase, *gCl* macroscopic chloride conductance, *TRP* transient receptor potential channels, *ROS* reactive oxygen species, *KATP* ATP-dependent potassium channels, *KCa* calcium activated potassium channels.

derives from studies on mice in which the TauT was genetically knocked out [6, 14–16]. TauT knockout mice (TauT^{-/-}) show more than 90% decrease in taurine content in both muscle and heart and are characterized by a marked decrease in exercise performance in exhaustive training models. Although the force of isolated muscle has not been measured in these TauT^{-/-} mice, clear abnormalities of muscle structure have been found, including signs of atrophy and muscle necrosis. Additionally, the muscles of TauT^{-/-} mice have a shift of metabolism toward the glycolytic pathway, especially in condition of exercise; this has been related to a dysfunction in mitochondrial function and in fatty acid oxidative pathways [51]. In parallel, taurine deficiency leads to cardiomyopathy characterized by remodeling of ventricular cardiomyocytes, ultrastructural damages of myofibril and mitochondria, and overexpression of markers of heart failure, such as atrial natriuretic peptide, brain natriuretic peptide and beta-myosin heavy chain [15, 16].

It is therefore evident that taurine is essential to maintain muscle performance and excitation–contraction coupling; however the mechanism for these actions is still unclear. An *in vitro* study of Berg and Bakker clearly demonstrated the ability of taurine to increase the accumulation of calcium into sarcoplasmic reticulum (SR) in isolated skinned myofibers by 35%, an effect that accounts for the greater depolarization-induced contraction of fiber exposed to 20 mM taurine. This in spite taurine slightly reduces the sensitivity of contractile apparatus to calcium [52]. Interestingly, a recent study demonstrated that a prolonged exposure to 10–20 mM taurine increases the rate of calcium uptake in both type I and type II human myofibers; an action within the SR lumen has been proposed. An increase in contractile sensitivity to calcium was also observed but exclusively in type I fibers [53]. These results reinforce the original data of Huxtable and Bressler about the ability of taurine to stimulate calcium uptake by vesicles of SR [13]. Recent insight into the role of taurine in skeletal muscle has been obtained by the group of Hayes, who supplemented rats with taurine and evaluated the outcome on various functional parameters [54]. Taurine supplementation significantly increases the amino acid content in skeletal muscle, without any adaptive change in TauT activity; in parallel an increase in force and a greater resistance and recovery after fatigue have been observed. These changes were paralleled by an increase in calsequestrin1, the calcium binding protein that works to maintain high amounts of calcium in the cisterna of SR. This suggests that taurine supplemented muscle can store a greater quantity of calcium with a consequent greater calcium availability for contraction. However, the involvement of sarco/endoplasmic reticulum calcium-ATPase (SERCA) remains

to be better clarified. A decrease in markers of oxidative stress was also found, indicating that taurine may help to control activity-related oxidative stress [48]. In support to this view, a recent report by Silva et al. showed that a daily treatment of rats with 300 mg/kg taurine for 2 weeks protects muscles against *in vivo* eccentric exercise damage, such as downhill running [55]. In particular taurine reduced protein carbonylation or oxidized thiols, without increasing the expression of endogenous anti-oxidant pathways, such as superoxide dismutase or catalase [55]. Sugiura et al. similarly found that taurine administration before strenuous exercise reduces muscle DNA damage likely via down-regulation of inducible nitric oxide synthase (iNOS) and consequent reduction of nitrosative inflammation [56]. The protective effects of taurine supplementation are due to a long term modulatory effect, likely in relation to its muscle uptake and intracellular levels. In fact acute *in vitro* application of physiological concentrations of taurine to isolated mouse soleus muscle, does not increase muscle contractile performance in term of force, fatigue resistance and recovery and does not exert any synergistic action when associated with caffeine [57]. Despite the authors suggesting a lack of ergogenic benefit by acute taurine, it is important to underline that slow twitch soleus muscle is characterized by high intracellular taurine content [58, 59], predicting its lower dependency on extracellular concentrations. Accordingly, we have shown that a chronic treatment with taurine to dystrophic mice leads to a minor increase of its intracellular content in soleus muscle than in fast twitch muscles [59].

Although taurine supplementation enhances exercise performance, its efflux during exercise and/or ischemia, with consequent decrease in tissue concentration, can also occur [60, 61]. Whether the loss of taurine is a marker of tissue damage or rather a cytoprotective mechanism against ischemic insult, is still matter of debate [60, 62, 63]. The protective effect of taurine efflux in the above conditions can be related to the need to osmotically balance, along with water movement, the increase of by-products of metabolism in the myofibers [1, 14]. However a role in the mechanism to contrast fatigue can be envisaged. In fact, taurine exerts an inhibitory control on channels that couple the metabolic state of the myofiber with membrane excitability, such as the ATP-dependent potassium (KATP) channels and calcium-activated potassium channels [64, 65]. Taurine blocks skeletal muscle KATP channel by binding the channel complex nearby the sulphonylurea receptor [64]. During ischemia–reperfusion injury, the opening of KATP are involved in the cytoprotective effect of the preconditioning mechanisms, by preventing the influx of calcium ions and preserving the ATP

content of the muscle. The efflux of taurine during exercise and/or ischemia may be required to relieve a basal inhibitory effect and to enhance the potassium efflux and membrane repolarization via the specific channels activated by ATP depletion and/or intracellular calcium accumulation. This would exert a protective action against exercise-induced fatigue or impairment in muscle performance related to ischemia–reperfusion injury [64, 65]. Accordingly, the depletion of taurine induced by GES in rat skeletal muscle significantly increases the macroscopic resting potassium conductance of about 80% [48].

Intracellular taurine can also be conjugated in mitochondria of extra-hepatic tissues to 5-taurinomethyl uridine that is present in tRNA and modulates the synthesis of mitochondrial proteins. Consequently, the fatigue and the enhanced oxidative stress observed in myopathic states by taurine depletion can also be due to respiratory chain inefficiency [4, 51, 66]. A representative scheme of the taurine actions in striated myofibers is shown in Fig. 3.

Taurine as potential therapeutic muscular agent from birth to elderly

The role of taurine for post-natal development of various organs depends upon the species-specific ability to endogenously synthesize the amino acid. Cats, that critically depend on exogenous taurine intake, develop serious impairments during post-natal development if not fed with taurine. Although less compelling for humans, prematurely born infants are believed to lack the enzymes that convert cystathionine to cysteine, and may, therefore, become taurine-deficient if not breast-fed. In fact taurine is present in mother’s milk and evidences are available about potential usefulness of taurine addition in the formula especially for pre-term births [67, 68]. The actual necessity or benefit of this practice has never been rigorously studied, and as such, taurine has yet to be proven to be important during fetal development, perhaps via epigenetic and/or organogenesis related mechanisms. Recent focus has been addressed to the potential benefit of taurine supplementation in mice during gestational period, especially when mothers are exposed to




Fig. 3 Representative scheme of the action of taurine in skeletal muscle fiber. It is shown the TauT carrier which works to maintain high intracellular taurine level, along with the actions that taurine exerts on membrane channels, sarcoplasmic reticulum, mitochondria and possibly gene expression. Putative binding sites for taurine are shown (1) on CLC-1 channel and (2) as local anesthetic drug binding site. Arrows indicate a general stimulating action while dotted lines are for inhibitory effects or yet undefined pathways. A pathway for taurine efflux under stress conditions (ischemia, osmotic stress, etc.) likely via the volume-sensitive organic anion channel (VSOAC) is also shown.

low-protein diet, a condition mimicking the low weight at birth and related to the risk of developing dysmetabolic states later on [69]. In these conditions taurine protects pancreas by decreasing islet sensitivity to cytokines and shows to have an impact on gene expression and “reprogramming” in various tissues, including skeletal muscle [70–72].

In support of the pivotal role of adequate taurine level for skeletal muscle development, we demonstrated that taurine muscle level increases during the first month of rat post-natal life [73]. This increase matches the acquisition of phenotype-specific contractile properties. In particular in rat fast-twitch EDL muscle it occurs in parallel with the post-natal increase in muscle gCl and of ClC-1 channels expression; i.e. during the acquisition of the mature profile [39, 73–75]. Adult levels are likely to be attained later, since a proton nuclear magnetic resonance (H-NMR) study showed an increase in taurine in different rat skeletal muscles from 6 to 18 weeks of age [76]. Accordingly, an age dependent increase of taurine as well as of other amino acids, has been found in muscle of metabolically healthy children (age range 1–15) with respect to adults [77].

In agreement with an active role of taurine for muscle phenotype acquisition, supplementation of mothers during pregnancy and lactation as well as of new-born rats results in a higher content of the amino acid in skeletal muscle, accompanied by a more rapid development of gCl [73]. Whether such an increase is due to a modulatory action of taurine on ClC-1 channel or to an effect on its gene expression is not known yet. Importantly, a profound alteration in gene expression has been described in liver and skeletal muscle of pups that were exposed prenatally to low protein diet, while the addition of taurine to mothers via drinking water during gestation leads to a marked protection [71, 72]. Focusing on skeletal muscle, the rescuing effect of taurine did occur for genes involved in oxidative phosphorylation and in the tricarboxylic acid cycle that were markedly down-regulated in skeletal muscle by the low protein diet. Importantly, plasma taurine concentration has been suggested to be a marker of fetal well-being and a prerequisite for normal fetal development [78]. In line with the important role of taurine for skeletal muscle development, the TauT expression increases during myogenesis and its gene has consensus site for myocyte enhancing factor 2 (MEF2), being therefore under strict control of myogenic program [79]. Also, taurine has been shown to stimulate myofiber differentiation *in vitro* [80]. Although the mechanism through which taurine may control gene expression during development is not clear yet, it appears to be a necessary factor in myogenesis, and perhaps in mitochondrial biogenesis, with key role for tissue development (Table 1).

Another condition that may benefit from taurine supplementation is aging. Age-related sarcopenia is accompanied by profound changes in hormonal and metabolic profile of skeletal muscle. An important alteration in the content of various amino acids occurs in human muscle specimen with age, as a result of age-related increase in proteolysis; in parallel a marked decrease in taurine content has been observed [81].

Besides sarcopenia, skeletal muscle of aged rats develops features that are overlapping those observed in taurine depleted muscles, i.e. a marked decrease in gCl and a change in calcium homeostasis with a shift of mechanical threshold towards more negative potentials [82, 83]. We found by high-performance liquid chromatography (HPLC) determination that muscle taurine concentration is in fact significantly decreased in muscle of aged rats; however the levels can be restored to adult values upon the exogenous administration of taurine for 3 months (1 g/kg in drinking water) [84]. Importantly, the taurine administration counteracts the decrease in gCl and the alteration in excitation–contraction coupling of aged rat EDL muscle, supporting the key role of the amino acid in the alterations observed and the potential beneficial role of its supplementation in elderly subjects (Table 1). In the EDL muscle of aged rats supplemented with taurine an almost complete recovery of the pharmacological sensitivity of gCl to either direct and indirect channel modulators, such as the enantiomers of *p*-chloro-phenoxy propionic acid and the phorbol esters, respectively, was observed. The effect of these latter, along with the amelioration of mechanical threshold observed, discloses the ability of taurine to modulate gCl by reducing the phosphorylation state of the chloride channel brought about by calcium and phospholipid-dependent protein kinase C [83, 84]. This offers a unifying mechanism for physiological taurine action via calcium homeostasis and modulation of calcium-dependent signaling pathways.

In line with the above observations, TauT^{-/-} mice show accelerated senescence, with greater muscular damage and endoplasmic reticulum stress due to accumulation of misfolded proteins. A central role of calcium mishandling has been proposed, along with the interest in maintaining adequate taurine level for contrasting aging-related muscle impairments [85].

Taurine and muscular dystrophy

The alteration of calcium homeostasis is a hallmark of muscles affected by inherited muscular dystrophy, such as in mice with X chromosome-linked muscular dystrophy (mdx), the most widely used model for Duchenne muscular dystrophy (DMD). It is believed that the absence of dystrophin, a protein with a key role for sarcolemmal integrity and mechano-transduction, leads to

sarcolemmal tears and to overactivity of voltage-insensitive cationic channels which enhance passive calcium entry, especially during work load [86–88]. This in turn leads to both the alteration of excitation–contraction coupling and to the activation of degenerative pathways [88, 89]. We have found that the EDL muscles of dystrophic mdx animals undergoing chronic exercise protocols, have features resembling taurine depleted ones, i.e. a reduction of gCl and a negative rheobase voltage for mechanical activation [89, 90]. Dystrophic muscle may have a reduced ability in retaining intracellular taurine; in fact we observed a trend of a lower than normal taurine muscle concentration in parallel with markedly high levels in plasma [89]. Accordingly, other authors found that taurine levels fluctuate in mdx muscles in relation to the disease phase, with compensatory increases being observed after acute degenerative period and glucocorticoid treatment [91, 92]. In this frame, taurine seems to be a useful marker of the dystrophic state of mdx mice when monitored by H1-magnetic resonance spectroscopy both in vivo and ex vivo, although technical problems may still limit the accurate peak resolution for quantitative evaluation [91–95]. In our experiments, the in vitro application of millimolar taurine concentrations fully restored the alteration of mechanical threshold observed in these animals [89]. Interestingly, similar results have been obtained upon chronic taurine treatment in exercised mdx mice. The in vivo treatment also significantly contrasted the decrease in gCl and lead to a significant increase of mouse strength in vivo, due to an interesting anabolic action of the amino acid in the dystrophic animals [90]. As previously mentioned, $TauT^{-/-}$ mice are characterized by a marked 80% decrease in exercise performance and increased fatigability, a feature that is classically observed in the mdx phenotype [6, 14, 90, 96]. The role of taurine in muscular dystrophy is also under study in Hayes' laboratory, where a lower expression of $TauT$ in mdx mouse muscle has been demonstrated, which is not influenced by exogenous taurine administration [97], supporting the difficulty of dystrophic muscle to retain taurine. Exercise protocols may differently modulate intramuscular taurine concentration, ranging from no change to phenotype-dependent decrease, likely in relation to the exercise type; however taurine supplementation can enhance exercise performance [60, 61]. Due to the impaired mechano-transduction of dystrophic myofibers, it would be of interest to evaluate whether the exercise protocol in mdx mice can lead to a further distress in taurine concentration and in $TauT$ expression; this is currently ongoing in our laboratory.

Based on first encouraging results, we tested the possible advantage to combine taurine with α -methylprednisolone, a glucocorticoids currently in use

in dystrophic patients [58]. A synergistic action of the two drugs in enhancing mouse strength and in restoring calcium homeostasis was observed, with a normalization of mechanical threshold and a reduction of the overactivity of the cation channels likely involved in abnormal calcium entry [58, 86, 98]. The treatment was also associated with a significant increase in taurine content in fast-twitch limb muscles, suggesting that dystrophic muscle maintains the ability to uptake taurine if adequately supplemented [58]. The synergistic action observed corroborates a potential interest of taurine as adjuvant therapy in steroid-treated patients. This is also supported by the evidence that glucocorticoids exert an inhibitory action of renal taurine re-uptake, then leading to hypotaurinemia, which in turn may have long-term negative effects on cardiovascular function [5].

Importantly, the taurine treatment to mdx mice significantly reduces the high plasma level of lactate dehydrogenase, an index of metabolic distress, and it is worth to underline that a marked increase in plasma lactate actually occurs in $TauT^{-/-}$ mice [6]. Therefore taurine can also play a role in metabolism in dystrophic muscle, similarly to what observed in exercise-challenged $TauT^{-/-}$ mice [51].

Increasing evidences suggest a link between calcium homeostasis, oxidative stress and mitochondrial distress in muscular dystrophy, leading to reconcile all these taurine actions under few main mechanisms, although not fully clear yet [99, 100]. As already mentioned, taurine supplementation contrasts the exercise-induced increase in oxidative markers, without enhancing the level of endogenous anti-oxidant [55]. Other evidences support that the sulfonic amino acid is actually incapable of scavenging the common oxidants, namely, superoxide, hydrogen peroxide and hydroxyl radical, which instead are the main products of enhanced NADPH oxidase activity in dystrophic muscle [99–101]. However, the amino group of taurine can neutralize hypochlorous acid, one of the reactive species generated by myeloperoxidase-halide system in neutrophils [102]. In that reaction, taurine is converted to taurine chloramine, which is less toxic than hypochlorous acid and actually serves as a modulator of the immune system also by interfering with the production of several pro-inflammatory mediators and activation of the transcription factor nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) [102]. In addition, taurine has been proposed to directly activate peroxisome proliferator-activated receptor γ (PPAR γ) in epithelial cells, a mechanism that may account for its protective action against inflammation-related diabetic retinopathy progression [103]. In consideration of the involvement of chronic inflammation and NF- κ B derived mediators in dystrophic muscle [87,

104, 105], the above immunomodulatory actions of taurine are of value. However, whether the anti-inflammatory and anti-oxidant action contributes to the beneficial effect observed in dystrophic animals is not known yet and the evaluation of biomarkers in samples of taurine treated mdx mice will be useful at this regard. Our preliminary results favor a decrease in superoxide anion formation, measured by dihydroethidium staining, in tibialis anterior muscles of exercised mdx mice treated with taurine (De Luca, personal unpublished observations). An attractive hypothesis, currently under study in our laboratory, is that taurine may contrast the impaired SERCA activity in dystrophic muscle either directly or by reducing the damaging effect brought about by oxidation and/or nitrosylation [13, 54, 106]. Interesting recent results of Terrill et al. have shown that a chronic administration of the cysteine precursor 2-oxothiazolidine-4 carboxylate (OTC) markedly decreases the level of thiol oxidation in muscles of mdx mice; in parallel an amelioration of force and muscle morphology has been observed. Importantly the administration was not paralleled by an increase in cysteine or glutathione but rather by an increase in taurine level. The authors underlined that the decrease in taurine content may have a direct causative role in enhanced susceptibility to oxidative stress, disclosing a novel mechanism for beneficial effect of the classical anti-oxidant *N*-acetylcysteine [107].

Considering the mitochondrial sufferance occurring in dystrophic muscle [93], the previously described role of taurine for preserving mitochondrial function has to be taken into account for further studies. Similarly, the potential role of taurine and its chemical chaperone conjugate tauroursodeoxycholic acid in contrasting endoplasmic reticulum stress in various conditions should be considered for the acute and chronic ability of taurine to modulate signaling pathways [108, 109]. In addition, taurine may improve muscle metabolism by contrasting functional ischemia, based on the described vasodilating properties [110]. The clarification of the mechanism of action and the evaluation of long term safety and efficacy also at heart level can add important pre-clinical data to plan clinical trials in DMD patients (Table 1).

Taurine and disuse-related muscle atrophy

Muscle disuse is a general term which describes a condition of inactivity occurring after prolonged bed rest, spaceflight and/or aging. The slow-twitch muscles, devoted to postural maintenance, are the most affected ones, showing a slow-to-fast phenotype transition and severe atrophy, both leading to impaired muscle function. The adaptation of skeletal muscle to different activity includes changes in the expression of structural, metabolic and contractile proteins that fine-tune the

characteristics of this tissue. The hindlimb unloaded (HU) model of disuse in rodents is a widely accepted ground-based model that mimics microgravity condition and is used to study the mechanisms responsible for the disuse-induced modification of skeletal muscle function. The soleus muscle of HU rats and mice becomes atrophic and experiences a slow-to-fast phenotype transition, characterized by an increased expression of the fast myosin heavy chain (MHC) isoform [111, 112]. Along the years, the studies on the HU model have shown that various proteins involved in the control of sarcolemma excitability, calcium ion homeostasis, energy metabolism, and contractile machinery undergo changes in the expression, turnover, and activity in accord with the entering of the slow muscle into a fast program [111, 113–117]. In particular, *ClC-1* chloride and *Nav1.4* sodium channels are differently expressed in fast-twitch and slow-twitch skeletal muscles, the expression of both being higher in the former. Accordingly with the change of phenotype, *ClC-1* channel activity and expression as well as the intracellular resting calcium level in slow-twitch soleus muscle are significantly shifted by HU process toward the values of a fast muscle, even before the modification of MHC expression [111]. Similarly, HU increased sodium current density and sodium channel mRNA level in soleus muscle fibers [113]. All these changes alter the resistance to fatigue of antigravity muscle fibers, an effect that may contribute to the impairment of muscle function, in terms of excitability and contraction. A full understanding of the mechanisms of disuse-induced muscle alterations in humans is still incomplete and few molecules have been proposed for therapy [118, 119]. However, supplementation with essential amino acids and carbohydrates in combination with exercise attenuates muscle protein loss in humans exposed to prolonged inactivity [120, 121]. Based on these considerations and on our previous findings about the action of taurine in the modulation of calcium homeostasis and ion channel function [34, 41, 49], we focused on taurine as a potential candidate to counteract the HU-induced phenotype transition and skeletal muscle function impairment [1, 34].

In agreement with a critical role of taurine in phenotype-specific cellular function, the concentration of the amino acid is twofold higher in soleus compared to EDL muscle. The physiological relevance for this phenotypic difference is still unknown but various hypothesis can be raised based on the essential role of taurine in skeletal muscle and its actions in metabolism and phenotype-dependent properties. Interestingly, our recent findings [59] showed for the first time a marked reduction of taurine content in the soleus muscle of HU rat. This muscle loss would be consistent with an original report of National Aeronautics and Space Administration (NASA)

describing a large excretion of taurine in the urine of the astronauts of the APOLLO mission [122]. In spite of the reduction of taurine in soleus muscle of HU rats, the expression of TauT was unchanged. Indeed, TauT expression was found to be higher in slow-twitch soleus muscle with respect to the fast EDL, and was not reduced during HU, suggesting that the intracellular reduction of taurine is not associated with the change of phenotype. In addition, our data suggest that TauT activity is efficiently maintained during HU, since taurine oral supplementation fully prevents the loss of taurine content in HU-soleus muscle. Thus, we hypothesize that the reduction of intracellular taurine content during HU is likely due to increased taurine efflux. A possible explanation might be that taurine leakage compensates for intracellular osmolarity changes, which likely occurs due to muscle protein degradation and increased catabolism. Accordingly, the production of intracellular osmolytes during muscle disuse atrophy has been described, which may justify taurine escape in this condition [123–125]. Importantly in rats fed with taurine, TauT expression was reduced in soleus muscle, suggesting a negative feedback regulation as a mechanism to control taurine intracellular level. As anticipated the TauT expression is under control of MEF2, a determinant of slow-fiber phenotype [79], thus it is tempting to speculate that TauT expression after taurine supplementation can be reduced by a mechanism involving a complex cross-talk between taurine and CIC-1 modulation during the phenotype transition.

Our findings also highlighted that taurine supplementation in HU rats has preserved resting gCl and resting cytosolic calcium level together with the slow MHC phenotype in the soleus muscle.

However, taurine had little effect on muscle atrophy, which is a severe condition occurring during HU as well as in various muscle diseases [126]. Indeed, it did not prevent the reduction of muscle-to-body weight ratio and of the fiber cross sectional area (CSA), while it partially contrasted the expression of atrogen-1 and mostly of muscle RING-finger protein-1 (MURF-1), two ubiquitin–proteasome pathway enzymes, that are strongly up-regulated as a result of HU-induced atrophy [127]. Such an effect suggests that a longer treatment or a different therapeutic schedule of taurine might have protective effect against muscle atrophy and might be useful to reach a complete muscular recovery. However complex mechanisms control the relative expression of atrogen and MURF-1 in skeletal muscle under various insults [79, 128] and further experiments are needed (Table 1).

Taurine and human skeletal muscle

Taurine has limited use in clinical settings although human use has been considered for specific diseases such

as non-insulin dependent diabetes and related disorders, to treat alcohol withdrawal, congestive heart failure and arrhythmias, rheumatoid arthritis and other chronic inflammatory states, seizure disorders, and liver related disorders [19, 102, 129]. In Table 2 is a brief report of some clinical studies related to taurine supplementation, with relative dosages and outcomes. Most of them focused on diabetes mellitus, insulin resistance and diabetic complications, based on the rationale that plasma taurine concentration is reduced in patients with insulin-dependent diabetes mellitus (IDDM) [129–136]. Taurine was indicated in addition to specific drugs. Other clinical studies tested taurine in congestive heart failure, hypertension, inherited succinic semialdehyde dehydrogenase deficiency, obesity or its supplementation in aged individuals [137–143].

A part for the use in myotonic dystrophy patients [35–37], the potential therapeutic role of taurine for skeletal muscle disorders has yet to be verified in clinical settings. In fact, most of the studies about the role of taurine for skeletal muscle physiology and its potential in pathological conditions have been carried out in animal models. In these conditions taurine depletion or supplementation are directly correlated with changes in the amino acid content in skeletal muscle, which facilitate the drawing of conclusion about amino acid action and potential. However, few studies have been conducted in humans, and some contradictory reports are available, questioning about the actual usefulness of taurine supplementation or on its mechanism of action. Apart for the age-related changes reported in the previous paragraphs, one of the main issue concerns the modulation of taurine concentration in adult skeletal muscle under conditions of exercise and/or metabolic distress. Galloway et al. [144] demonstrated that taurine supplementation to exercised healthy adults leads to a marked increase in the amino acid plasma level that however is not paralleled, after 7 days of supplementation, by an increase in skeletal muscle. They proposed that intramuscular taurine concentration is tightly regulated and that high plasma level may actually work to reduce TauT activity in order to maintain constant the amino acid level. Therefore, even chronic oral taurine supplementation may cause less increase in human muscles than in rodent ones, and the observed muscle effects could be due to extracellular taurine actions. In addition, plasma levels are also tightly regulated via overexpression of TauT in kidney, which may also show specie-specific regulatory pathways [145, 146].

The dose is another important issue. In fact murine pre-clinical studies often require about tenfold higher concentration than in human trials; by the way this has to match the endogenous high level of taurine in target

Table 2 Clinical use of taurine in different pathophysiological conditions

References	Patients	Dose (g/day or mg/kg)	Duration	Result
Franconi et al. [130]	IDDM (Diabetes mellitus type 1)	1.5 g	90 days	No effect
Elizarova and Nedosugova [131]	IDDM	1 g	30 days	Glucose metabolism and trygliceride level improved
Chauncey et al. [133]	NIDDM (DM type 2)	3 g	4 months	Plasma taurine level increased
Brøns et al. [134]	Overweight non-diabetic	1.5 g	8 weeks	No effect
Xiao et al. [136]	Overweight non-diabetic	3 g	2 weeks	Insulin sensitivity improved
Nakamura et al. [132]	NIDDM with microalbuminemia	3 g	12 months	No effect
Moloney et al. [135]	IDDM	1.5 g	2 weeks	Endotelium-dependent reaction improved
Gonzales-Contreras et al. [142]	Cholestasis by parenteral nutrition	~25 mg/kg/day	~50 days	Hepatoprotection with reduction of AST, ALT and GGT
Rosa et al. [143]	Obesity	3 g/day	8 weeks	Increase in plasma levels of taurine and adiponectin; reduction of inflammatory markers
Pearl et al. [141]	Succinic semialdehyde dehydrogenase deficiency (efficacy, safety and tolerability)	50–200 mg/kg/d (age range 12 years)	13 months (mean time from 3 to 50)	No significant effects Tolerability issues at highest doses
Fujita et al. [139]	Hypertension	6 g	7 days	Systolic and diastolic pressure improved
Azuma et al. [138]	Congestive heart failure	6 g	4 weeks	Heart parameters improved
Bergamini et al. [137]	Epilepsy	200 mg–21 g	Various	Seizure frequency reduction
Durelli et al. [36]	Dystrophic myotonia	6–10 g	6 months	Myotonic symptoms improvement
Dunn-Lewis et al. [140]	Elderly	500 mg in multinutrient supplement	4 weeks	Physical function improved

organs. In addition, an accurate muscle exposure to taurine after oral ingestion requires a careful assessment of the pharmacokinetic profile that has not been extensively evaluated in humans. In line with Galloway et al. [144], a single oral dose of 4 g in healthy volunteers allows to get a maximal plasma peak in about 1.5 h and showed an half-life of 1 h with a first-order kinetic clearance; this is in line with kidney being the main organ regulating taurine level [147]. Generally the daily dose of taurine ranges between 3 and 6 g; consequently its fast kinetic can account for some of the puzzling data obtained, suggesting the need of a more careful determination of the optimum dose. It is important to underline that most of the available evidences focus on the usefulness of taurine supplementation in sustaining muscle function in trained individuals. Balshaw et al. have recently evaluated the outcome of 1 g taurine ingestion, evaluated in blind against placebo, on running performance of trained middle-distance runners. They described a modest, although significant, increase in performance in the taurine-treated group, without any change in metabolism parameters [148]. The authors claimed that a similar improvement of

performance after taurine ingestion, without changes in oxygen uptake or plasma lactate, has been found in other studies [144]. Taurine muscle levels were not assessed, thus the correlation between taurine effect and a specific muscle action is rather indirect. Accordingly, they speculated about alternative potential mechanisms, such as the action of taurine at muscle membrane level, in preventing taurine drop during exercise or rather an effect on neuronal function.

In another study, a combination of taurine (2 g) and branched-chain amino acids three times a days for 2 weeks before eccentric exercise, plus 4 days after, has been tested in healthy untreated volunteers. The eccentric exercise protocol consisted of repeated sets elbow flexion at 90° to an extended position, finally leading to uncontrolled damaging stretch. The combination exerted a greater protection against muscle damage and delayed-onset muscle soreness than single administrations, although no detailed investigation has been done to clarify the mechanism of action and/or the amino acid level into the muscle [149]. Similarly, da Silva et al. have recently described the ability of 14 days taurine

administration to increase strength of the elbow flexor subjected to eccentric exercises in young adult males; in parallel, markers of oxidative stress were reduced, without increase in endogenous anti-oxidant expression nor changes in inflammatory markers. Again muscle taurine level were not determined [150]. Therefore the available evidences do not allow to conclude about the ability of supplemented taurine to actually increase its muscle level in adult healthy and trained individuals, suggesting alternative modality of action, i.e. at neuromuscular system. However, it cannot be ruled out that taurine supplementation may effectively enhances muscle taurine levels in conditions characterized by more dramatic fluctuation of its content. This applies to postnatal development and aging, and mostly to pathological conditions such as muscular dystrophy and disuse-related muscle dysfunction (Table 1) [151]. More direct evidences in humans and patients will be helpful, in order to better correlate the effect of exogenous administration of taurine with the ability of residual muscle tissue to uptake the right amount, or rather to disclose taurine actions independent on its intracellular levels [145]. In addition, an inter-individual variation in plasma increase of taurine after supplementation may occur in relation to both nutritional state, age, drug interaction, while gene polymorphism in taurine transporter or modulation of its function and/or expression by cell metabolic state or activation of transcription factors may affect the actual level of taurine being transported into the myofibers [134, 146, 152–154]. Hence caution should be taken when concluding about lack of taurine usefulness for human muscular system without an adequate control of all variables.

Conclusion

We herein summarized the results obtained in about 30 years of research on taurine and skeletal muscle by us and other research groups. Taurine is far from themes of fashion science or from immediate interest in innovative drug development by Pharma Companies. Nevertheless the reason for such a long interest is that taurine acquired over the years a special appeal for its puzzling and multiple effects. We underlined the ability of taurine to control the function of ion channels and consequently membrane excitability as well as calcium homeostasis and excitation–contraction coupling. It has been highlighted that novel evidences are emerging regarding taurine mechanism of action, ranging from modulation of muscle metabolism to control of gene transcription, as well as in the species-specific mechanisms underlying its intracellular levels in both chronic and acute conditions. These make the research on the topic “taurine and skeletal muscle” a continuous source of novel and exciting results allowing to renew the enthusiasm and novel working hypotheses. The

wide and interconnected effects observed support a key role of the amino acid to ensure a proper muscle function and reinforce its interest as therapeutic agent in various inherited and acquired muscular disorders. The available evidences favor a greater effect of taurine in diseased condition accompanied by alterations in taurine concentration in muscle; similar benefit can occur in conditions where fluctuation in taurine level take place such as exercise, protein content in diet or post-natal development. Both acute and chronic effects of taurine supplementation are feasible, and likely occur with different time-scale although similarly interesting and important. Although a careful distinction has not been made, it is predictable that acute effects of taurine are better appreciable in situations of rapid fluctuations such as exercise, or when involving direct modulation of ion channel, or on muscles that are more dependable of external taurine such as fast-twitch ones. In parallel, chronic taurine effects, likely accompanied by changes in intracellular content, could be of value for long term control of neuromuscular function in progressive conditions, such as muscular dystrophy and disuse or aging-related dysfunction. At this regard more evidences are necessary to better understand the interest of taurine for ensuring a proper muscle function in human other than in animals. Consequently, a more clinically-oriented research will help to support the interest of taurine as novel and safer therapeutic approach of rare inherited muscle diseases and other myopathic states.

Authors' contributions

ADL: have made a substantial contribution in designing and writing the review, updating current literature and in interpretation of available data in the field; SP: was significantly involved in writing, in figures and table organizations, literature search and interpretation of available information; DCC: critically revised the manuscript and its organization and gave a substantial support to the finalization of the work. All authors have read and approved the final manuscript.

Acknowledgements

The authors acknowledge the contribution of grants from Telethon-Italy (Conte and De Luca) and Italian Space Agency (Conte) for support of the most recent researches in the taurine field presented herein. The authors wish to thank Dr. Anna Cozzoli for enthusiastic and valuable assistance during the preparation of the present review.

Compliance with ethical guidelines

Competing interests

The authors declare that they have no competing interests.

Received: 26 March 2015 Accepted: 17 July 2015

Published online: 25 July 2015

References

1. Huxtable RJ (1992) Physiological actions of taurine. *Physiol Rev* 72:101–163
2. Barle H, Ahlman B, Nyberg B, Andersson K, Essén P, Wernerman J (1996) The concentrations of free amino acids in human liver tissue obtained during laparoscopic surgery. *Clin Physiol* 16:217–227

3. Huxtable RJ (2000) Expanding the circle 1975-1999: sulfur biochemistry and insights on the biological functions of taurine. *Adv Exp Med Biol* 483:1–25
4. Schaffer SW, Jong CJ, Ramila KC, Azuma J (2010) Physiological roles of taurine in heart and muscle. *J Biomed Sci* 17:S2
5. Faggiano A, Melis D, Alfieri R, De Martino M, Filippella M, Milone F et al (2005) Sulfur amino acids in Cushing's disease: insight in homocysteine and taurine levels in patients with active and cured disease. *J Clin Endocrinol Metab* 90:6616–6622
6. Warskulat U, Flögel U, Jacoby C, Hartwig HG, Thewissen M, Merx MW et al (2004) Taurine transporter knockout depletes muscle taurine levels and results in severe skeletal muscle impairment but leaves cardiac function uncompromised. *FASEB J* 18:577–579
7. Stipanuk MH (2004) Role of the liver in regulation of body cysteine and taurine levels: a brief review. *Neurochem Res* 29:105–110
8. Lambert IH, Kristensen DM, Holm JB, Mortensen OH (2015) Physiological role of taurine—from organism to organelle. *Acta Physiol (Oxf)*. 213:191–212
9. Wu JY, Tang XW, Tsai WH (1992) Taurine receptor: kinetic analysis and pharmacological studies. *Adv Exp Med Biol* 315:263–268
10. Frosini M, Sesti C, Dragoni S, Valoti M, Palmi M, Dixon HB et al (2003) Interactions of taurine and structurally related analogues with the GABAergic system and taurine binding sites of rabbit brain. *Br J Pharmacol* 139:1163–1171
11. Jia F, Yue M, Chandra D, Keramidis A, Goldstein PA, Homanics GE et al (2008) Taurine is a potent activator of extrasynaptic GABA(A) receptors in the thalamus. *J Neurosci* 28:106–115
12. Wu JY, Prentice H (2010) Role of taurine in the central nervous system. *J Biomed Sci* 17:S1
13. Huxtable R, Bressler R (1973) Effect of taurine on a muscle intracellular membrane. *Biochim Biophys Acta* 323:573–583
14. Warskulat U, Heller-Stilb B, Oermann E, Zilles K, Haas H, Lang F et al (2007) Phenotype of the taurine transporter knockout mouse. *Methods Enzymol* 428:439–458
15. Ito T, Kimura Y, Uozumi Y, Takai M, Muraoka S, Matsuda T et al (2008) Taurine depletion caused by knocking out the taurine transporter gene leads to cardiomyopathy with cardiac atrophy. *J Mol Cell Cardiol* 44:927–937
16. Ito T, Oishi S, Takai M, Kimura Y, Uozumi Y, Fujii Y et al (2010) Cardiac and skeletal muscle abnormality in taurine transporter-knockout mice. *J Biomed Sci* 17:S20
17. Lötsch J, Hummel T, Warskulat U, Coste O, Häussinger D, Geisslinger G et al (2014) Congenital taurine deficiency in mice is associated with reduced sensitivity to nociceptive chemical stimulation. *Neuroscience* 259:63–70
18. Schaffer SW, Shimada K, Jong CJ, Ito T, Azuma J, Takahashi K (2014) Effect of taurine and potential interactions with caffeine on cardiovascular function. *Amino Acids* 6:1147–1157
19. Shao A, Hathcock JN (2008) Risk assessment for the amino acids taurine, L-glutamine and L-arginine. *Regul Toxicol Pharmacol* 50:376–399
20. Seifert SM, Schaechter JL, Hershorin ER, Lipshultz SE (2011) Health effects of energy drinks on children, adolescents, and young adults. *Pediatrics* 127:511–528
21. Wolk BJ, Ganetsky M, Babu KM (2012) Toxicity of energy drinks. *Curr Opin Pediatr* 24:243–251
22. Gunja N, Brown JA (2012) Energy drinks: health risks and toxicity. *Med J Aust* 196:46–49
23. Taranukhin AG, Saransaari P, Oja SS (2013) Lethality of taurine and alcohol coadministration in mice. *Adv Exp Med Biol* 776:29–38
24. El Idrissi A, Messing J, Scalia J, Trenkner E (2003) Prevention of epileptic seizures by taurine. *Adv Exp Med Biol* 526:515–525
25. Adrian RH, Bryant SH (1974) On the repetitive discharge in myotonic muscle fibres. *J Physiol* 240:505–515
26. Conte Camerino D, Tricarico D, Desaphy JF (2007) Ion channel pharmacology. *Neurotherapeutics* 4:184–198
27. Jentsch TJ (2008) CLC chloride channels and transporters: from genes to protein structure, pathology and physiology. *Crit Rev Biochem Mol Biol* 43:3–36
28. Conte Camerino D, Franconi F, Mambrini M, Bennardini F, Failli P, Bryant SH et al (1987) The action of taurine on chloride conductance and excitability characteristics of rat striated muscle fibers. *Pharmacol Res Commun* 19:685–701
29. Conte Camerino D, Franconi F, Mambrini M, Mitolo-Chieppa D, Bennardini F, Failli P et al (1987) Effect of taurine on chloride conductance and excitability of rat skeletal muscle fibers. *Adv Exp Med Biol* 217:207–216
30. Conte Camerino D, De Luca A, Mambrini M, Ferrannini E, Franconi F, Giotti A et al (1989) The effects of taurine on pharmacologically induced myotonia. *Muscle Nerve* 12:898–904
31. Pierno S, Tricarico D, De Luca A, Campagna F, Carotti A, Casini G et al (1994) Effects of taurine analogues on chloride channel conductance of rat skeletal muscle fibers: a structure-activity relationship investigation. *Naunyn Schmiedebergs Arch Pharmacol* 349:416–421
32. Pusch M, Accardi A, Liantonio A, Ferrera L, De Luca A, Camerino DC et al (2001) Mechanism of block of single protopores of the Torpedo chloride channel ClC-0 by 2-(*p*-chlorophenoxy)butyric acid (CPB). *J Gen Physiol* 118:45–62
33. Liantonio A, Accardi A, Carbonara G, Fracchiolla G, Loiodice F, Tortorella P et al (2002) Molecular requisites for drug binding to muscle CLC-1 and renal CLC-K channel revealed by the use of phenoxy-alkyl derivatives of 2-(*p*-chlorophenoxy)propionic acid. *Mol Pharmacol* 62:265–271
34. Conte Camerino D, Tricarico D, Pierno S, Desaphy JF, Liantonio A, Pusch M et al (2004) Taurine and skeletal muscle disorders. *Neurochem Res* 29:135–142
35. Durelli L, Mutani R, Fassio F, Satta A, Bartoli E (1982) Taurine and hyperexcitable human muscle: effects of taurine on potassium-induced hyperexcitability of dystrophic myotonic and normal muscles. *Ann Neurol* 11:258–265
36. Durelli L, Mutani R, Fassio F (1983) The treatment of myotonia: evaluation of chronic oral taurine therapy. *Neurology*. 33:599–603
37. Trip J, Drost G, van Engelen BG, Faber CG (2006) Drug treatment for myotonia. *Cochrane Database Syst Rev* (1):CD004762
38. Mankodi A, Takahashi MP, Jiang H, Beck CL, Bowers WJ, Moxley RT et al (2002) Expanded CUG repeats trigger aberrant splicing of CLC-1 chloride channel pre-mRNA and hyperexcitability of skeletal muscle in myotonic dystrophy. *Mol Cell* 10:35–44
39. Lueck JD, Lungu C, Mankodi A, Osborne RJ, Welle SL, Dirksen RT et al (2007) Chloride channelopathy in myotonic dystrophy resulting from loss of posttranscriptional regulation for CLCN1. *Am J Physiol Cell Physiol* 292:C1291–C1297
40. Conte Camerino D, Desaphy JF, Tricarico D, Pierno S, Liantonio A (2008) Therapeutic approaches to ion channel diseases. *Adv Genet* 64:81–145
41. De Luca A, Pierno S, Tricarico D, Desaphy JF, Liantonio A, Barbieri M et al (2000) Taurine and skeletal muscle ion channels. *Adv Exp Med Biol* 483:45–56
42. Schanne OF, Dumaine R (1992) Interaction of taurine with the fast Na⁺-current in isolated rabbit myocytes. *J Pharmacol Exp Ther* 263:1233–1240
43. Satoh H (1998) Inhibition of the fast Na⁺ current by taurine in guinea pig ventricular myocytes. *Gen Pharmacol* 31:155–157
44. De Luca A, Natuzzi F, Desaphy JF, Loni G, Lentini G, Franchini C et al (2000) Molecular determinants of mexiletine structure for potent and use-dependent block of skeletal muscle sodium channels. *Mol Pharmacol* 57:268–277
45. De Luca A, Talon S, De Bellis M, Desaphy JF, Lentini G, Corbo F et al (2003) Optimal requirements for high affinity and use-dependent block of skeletal muscle sodium channel by *N*-benzyl analogs of tocainide-like compounds. *Mol Pharmacol* 64:932–945
46. De Luca A, Pierno S, Liantonio A, Desaphy JF, Natuzzi F, Didonna MP et al (2004) New potent mexiletine and tocainide analogues evaluated in vivo and in vitro as antimyotonic agents on the myotonic ADR mouse. *Neuromuscul Disord* 14:405–416
47. De Luca A, De Bellis M, Corbo F, Franchini C, Muraglia M, Catalano A et al (2012) Searching for novel anti-myotonic agents: pharmacophore requirement for use-dependent block of skeletal muscle sodium channels by *N*-benzylated cyclic derivatives of tocainide. *Neuromuscul Disord* 22:56–65
48. De Luca A, Pierno S, Conte Camerino D (1996) Effect of taurine depletion on excitation–contraction coupling and Cl[−] conductance of rat skeletal muscle. *Eur J Pharmacol* 296:215–222

49. Pierno S, De Luca A, Huxtable RJ, Conte Camerino D (1994) Dual effects of taurine on membrane ionic conductances of rat skeletal muscle fibers. *Adv Exp Med Biol* 359:217–224
50. Hamilton EJ, Berg HM, Easton CJ, Bakker AJ (2006) The effect of taurine depletion on the contractile properties and fatigue in fast-twitch skeletal muscle of the mouse. *Amino Acids* 31:273–278
51. Ito T, Yoshikawa N, Schaffer SW, Azuma J (2014) Tissue taurine depletion alters metabolic response to exercise and reduces running capacity in mice. *J Amino Acids*. 2014:964680
52. Bakker AJ, Berg HM (2002) Effect of taurine on sarcoplasmic reticulum function and force in skinned fast-twitch skeletal muscle fibres of the rat. *J Physiol* 538:185–194
53. Dutka TL, Lamboley CR, Murphy RM, Lamb GD (2014) Acute effects of taurine on sarcoplasmic reticulum Ca^{2+} accumulation and contractility in human type I and type II skeletal muscle fibers. *J Appl Physiol* (1985) 117:797–805
54. Goodman CA, Horvath D, Stathis C, Mori T, Croft K, Murphy RM et al (2009) Taurine supplementation increases skeletal muscle force production and protects muscle function during and after high-frequency in vitro stimulation. *J Appl Physiol* 107:144–154
55. Silva LA, Silveira PC, Ronsani MM, Souza PS, Scheffer D, Vieira LC et al (2011) Taurine supplementation decreases oxidative stress in skeletal muscle after eccentric exercise. *Cell Biochem Funct* 29:43–49
56. Sugiura H, Okita S, Kato T, Naka T, Kawanishi S, Ohnishi S et al (2013) Protection by taurine against iNOS-increase DNA damage in heavily exercised skeletal muscle by inhibition of the NF- κ B signaling pathway. *Adv Exp Med Biol* 775:237–246
57. Tallis J, Higgins MF, Cox VM, Duncan MJ, James RS (2014) Does a physiological concentration of taurine increase acute muscle power output, time to fatigue, and recovery in isolated mouse soleus (slow) muscle with or without the presence of caffeine? *Can J Physiol Pharmacol* 92:42–49
58. Cozzoli A, Rolland JF, Caporosso RF, Sblendorio VT, Longo V, Simonetti S et al (2011) Evaluation of potential synergistic action of a combined treatment with alpha-methyl-prednisolone and taurine on the mdx mouse model of Duchenne muscular dystrophy. *Neuropathol Appl Neurobiol* 37:243–256
59. Pierno S, Liantonio A, Camerino GM, De Bellis M, Cannone M, Gramegna G et al (2012) Potential benefits of taurine in the prevention of skeletal muscle impairment induced by disuse in the hindlimb-unloaded rat. *Amino Acids* 43:431–445
60. Dawson R Jr, Biasseti M, Messina S, Dominy J (2002) The cytoprotective role of taurine in exercise-induced muscle injury. *Amino Acids* 22:309–324
61. Yatabe Y, Miyakawa S, Ohmori H, Mishima H, Adachi T (2009) Effects of taurine administration on exercise. *Adv Exp Med Biol* 643:245–252
62. Nanobashvili J, Neumayer C, Fugl A, Punz A, Blumer R, Prager M et al (2003) Ischemia/reperfusion injury of skeletal muscle: plasma taurine as a measure of tissue damage. *Surgery* 133:91–100
63. Takahashi K, Ohyabu Y, Takahashi K, Solodushko V, Takatani T, Itoh T et al (2003) Taurine renders the cell resistant to ischemia-induced injury in cultured neonatal rat cardiomyocytes. *J Cardiovasc Pharmacol* 41:726–733
64. Tricarico D, Barbieri M, Camerino DC (2000) Taurine blocks ATP-sensitive potassium channels of rat skeletal muscle fibres interfering with the sulphonylurea receptor. *Br J Pharmacol* 130:827–834
65. Tricarico D, Barbieri M, Conte Camerino D (2001) Voltage-dependent antagonist/agonist actions of taurine on Ca^{2+} -activated potassium channels of rat skeletal muscle fibers. *J Pharmacol Exp Ther* 298:1167–1171
66. Suzuki T, Nagao A, Suzuki T (2011) Human mitochondrial diseases caused by lack of taurine modification in mitochondrial tRNAs. *Wiley Interdiscip Rev RNA* 2:376–386
67. Heird WC (2004) Taurine in neonatal nutrition—revisited. *Arch Dis Child Fetal Neonatal Ed* 89:F473–F474
68. Wharton BA, Morley R, Isaacs EB, Cole TJ, Lucas A (2004) Low plasma taurine and later neurodevelopment. *Arch Dis Child Fetal Neonatal Ed* 89:F497–F498
69. Martin-Gronert MS, Ozanne SE (2007) Experimental IUGR and later diabetes. *J Intern Med* 261:437–452
70. Merezak S, Reusens B, Renard A, Goosse K, Kalbe L, Ahn MT et al (2004) Effect of maternal low-protein diet and taurine on the vulnerability of adult Wistar rat islets to cytokines. *Diabetologia* 47:669–675
71. Mortensen OH, Olsen HL, Frandsen L, Nielsen PE, Nielsen FC, Grunnet N et al (2010) Gestational protein restriction in mice has pronounced effects on gene expression in newborn offspring's liver and skeletal muscle; protective effect of taurine. *Pediatr Res* 67:47–53
72. Reusens B, Sparre T, Kalbe L, Bouckennooghe T, Theys N, Kruhøffer M et al (2008) The intrauterine metabolic environment modulates the gene expression pattern in fetal rat islets: prevention by maternal taurine supplementation. *Diabetologia* 51:836–845
73. De Luca A, Conte Camerino D, Failli P, Franconi F, Giotti A (1990) Effects of taurine on mammalian skeletal muscle fiber during development. *Prog Clin Biol Res* 351:163–173
74. Conte Camerino D, De Luca A, Mambrini M, Vrbova G (1989) Membrane ionic conductances in normal and denervated skeletal muscle of the rat during development. *Pflugers Archiv*. 413:568–570
75. Steinmeyer K, Ortland C, Jentsch TJ (1991) Primary structure and functional expression of a developmentally regulated skeletal muscle chloride channel. *Nature* 354:301–304
76. Yoshioka Y, Masuda T, Nakano H, Miura H, Nakaya S, Itazawa S (2002) In vitro ¹H-NMR spectroscopic analysis of metabolites in fast- and slow-twitch muscles of young rats. *Magn Reson Med Sci* 1:7–13
77. Hammarqvist F, Angsten G, Meurling S, Andersson K, Wernerman J (2010) Age-related changes of muscle and plasma amino acids in healthy children. *Amino Acids* 39:359–366
78. de Boo HA, Harding JE (2007) Taurine as a marker for foetal wellbeing? *Neonatology* 91:145–154
79. Uozumi Y, Ito T, Hoshino Y, Mohri T, Maeda M, Takahashi K et al (2006) Myogenic differentiation induces taurine transporter in association with taurine mediated cytoprotection in skeletal muscles. *Biochem J* 394:699–706
80. Miyazaki T, Honda A, Ikegami T, Matsuzaki Y (2013) The role of taurine on skeletal muscle cell differentiation. *Adv Exp Med Biol* 776:321–328
81. Stuerenburg HJ, Stangneth B, Schoser BG (2006) Age related profiles of free amino acids in human skeletal muscle. *Neuro Endocrinol Lett* 27:133–136
82. De Luca A, Conte Camerino D (1992) Effects of aging on the mechanical threshold of rat skeletal muscle fibers. *Pflugers Arch* 420:407–409
83. De Luca A, Tricarico D, Pierno S, Conte Camerino D (1994) Aging and chloride channel regulation in rat fast-twitch muscle fibres. *Pflugers Arch* 427:80–85
84. Pierno S, De Luca A, Camerino C, Huxtable RJ, Conte Camerino D (1998) Chronic administration of taurine to aged rats improves the electrical and contractile properties of skeletal muscle fibers. *J Pharmacol Exp Ther* 286:1183–1190
85. Ito T, Yoshikawa N, Inui T, Miyazaki N, Schaffer SW, Azuma J (2014) Tissue depletion of taurine accelerates skeletal muscle senescence and leads to early death in mice. *PLoS One* 9:e107409
86. Rolland JF, De Luca A, Burdi R, Andreetta F, Confalonieri P, Conte Camerino D (2006) Overactivity of exercise-sensitive cation channels and their impaired modulation by IGF-1 in mdx native muscle fibers: beneficial effect of pentoxifylline. *Neurobiol Dis* 24:466–474
87. Grounds MD, Radley HG, Lynch GS, Nagaraju K, De Luca A (2008) Towards developing standard operating procedures for pre-clinical testing in the mdx mouse model of Duchenne muscular dystrophy. *Neurobiol Dis* 31:1–19
88. Allen DG, Whitehead NP (2011) Duchenne muscular dystrophy—what causes the increased membrane permeability in skeletal muscle? *Int J Biochem Cell Biol* 43:290–294
89. De Luca A, Pierno S, Liantonio A, Cetrone M, Camerino C, Simonetti S et al (2001) Alteration of excitation–contraction coupling mechanism in extensor digitorum longus muscle fibres of dystrophic mdx mouse and potential efficacy of taurine. *Br J Pharmacol* 132:1047–1054
90. De Luca A, Pierno S, Liantonio A, Cetrone M, Camerino C, Frayssé B et al (2003) Enhanced dystrophic progression in mdx mice by exercise and beneficial effects of taurine and insulin-like growth factor-1. *J Pharmacol Exp Ther* 304:453–463
91. McIntosh L, Granberg KE, Brière KM, Anderson JE (1998) Nuclear magnetic resonance spectroscopy study of muscle growth, mdx dystrophy

- and glucocorticoid treatments: correlation with repair. *NMR Biomed* 11:1–10
92. McIntosh LM, Baker RE, Anderson JE (1998) Magnetic resonance imaging of regenerating and dystrophic mouse muscle. *Biochem Cell Biol* 76:532–541
 93. Griffin JL, Des Rosiers C (2009) Applications of metabolomics and proteomics to the mdx mouse model of Duchenne muscular dystrophy: lessons from downstream of the transcriptome. *Genome Med* 1:32
 94. Martins-Bach AB, Bloise AC, Vainzof M, Rahnamaye Rabbani S (2012) Metabolic profile of dystrophic mdx mouse muscles analyzed with in vitro magnetic resonance spectroscopy (MRS). *Magn Reson Imaging* 30:1167–1176
 95. Xu S, Pratt SJ, Spangenburg EE, Lovering RM (2012) Early metabolic changes measured by ¹H MRS in healthy and dystrophic muscle after injury. *J Appl Physiol* 113:808–816
 96. Burdi R, Rolland JF, Frayssse B, Litvinova K, Cozzoli A, Giannuzzi V et al (2009) Multiple pathological events in exercised dystrophic mdx mice are targeted by pentoxifylline: outcome of a large array of in vivo and ex vivo tests. *J Appl Physiol* 106:1311–1324
 97. Horvath DM (2011) The effect of taurine on dystrophic muscle tissue function. PhD thesis. Victoria University
 98. Frayssse B, Liantonio A, Cetrone M, Burdi R, Pierno S, Frigeri A et al (2004) The alteration of calcium homeostasis in adult dystrophic mdx muscle fibers is worsened by a chronic exercise in vivo. *Neurobiol Dis* 17:144–154
 99. Shkryl VM, Martins AS, Ullrich ND, Nowycky MC, Niggli E, Shirokova N (2009) Reciprocal amplification of ROS and Ca(2+) signals in stressed mdx dystrophic skeletal muscle fibers. *Pflugers Arch* 458:915–928
 100. Whitehead NP, Yeung EW, Froehner SC, Allen DG (2010) Skeletal muscle NADPH oxidase is increased and triggers stretch-induced damage in the mdx mouse. *PLoS One* 5:e15354
 101. Khairallah RJ, Shi G, Sbrana F, Prosser BL, Borroto C, Mazaitis MJ et al (2012) Microtubules underlie dysfunction in duchenne muscular dystrophy. *Sci Signal* 5:ra56
 102. Marcinkiewicz J, Kontny E (2014) Taurine and inflammatory diseases. *Amino Acids* 46:7–20
 103. Song MK, Salam NK, Roufogalis BD, Huang TH (2011) Lycium barbarum (Goji Berry) extracts and its taurine component inhibit PPAR- γ -dependent gene transcription in human retinal pigment epithelial cells: possible implications for diabetic retinopathy treatment. *Biochem Pharmacol* 82:1209–1218
 104. Pierno S, Nico B, Burdi R, Liantonio A, Didonna MP, Cipponi V et al (2007) Role of tumour necrosis factor alpha, but not of cyclo-oxygenase-2-derived eicosanoids, on functional and morphological indices of dystrophic progression in mdx mice: a pharmacological approach. *Neuropathol Appl Neurobiol* 33:344–359
 105. De Luca A, Nico B, Rolland JF, Cozzoli A, Burdi R, Mangieri D et al (2008) Gentamicin treatment in exercised mdx mice: identification of dystrophin-sensitive pathways and evaluation of efficacy in work-loaded dystrophic muscle. *Neurobiol Dis* 32:243–253
 106. Gehrig SM, van der Poel C, Sayer TA, Schertzer JD, Henstridge DC, Church JE et al (2012) Hsp72 preserves muscle function and slows progression of severe muscular dystrophy. *Nature* 4(484):394–398
 107. Terrill JR, Boyatzis A, Grounds MD, Arthur PG (2013) Treatment with the cysteine precursor l-2-oxothiazolidine-4-carboxylate (OTC) implicates taurine deficiency in severity of dystropathology in mdx mice. *Int J Biochem Cell Biol* 45:2097–2108
 108. Pan C, Giraldo GS, Prentice H, Wu JY (2010) Taurine protection of PC12 cells against endoplasmic reticulum stress induced by oxidative stress. *J Biomed Sci* 17:S17
 109. Batista TM, da Silva PM, Amaral AG, Ribeiro RA, Boschero AC, Carneiro EM (2013) Taurine supplementation restores insulin secretion and reduces ER stress markers in protein-malnourished mice. *Adv Exp Med Biol* 776:129–139
 110. Abebe W, Mozaffari MS (2011) Role of taurine in the vasculature: an overview of experimental and human studies. *Am J Cardiovasc Dis* 1:293–311
 111. Pierno S, Desaphy JF, Liantonio A, De Bellis M, Bianco G, De Luca A et al (2002) Change of chloride ion channel conductance is an early event of slow-to-fast fibre type transition during unloading-induced muscle disuse. *Brain* 125:1510–1521
 112. Desaphy JF, Pierno S, Liantonio A, Giannuzzi V, Digennaro C, Dinardo MM et al (2010) Antioxidant treatment of hindlimb-unloaded mouse counteracts fiber type transition but not atrophy of disused muscles. *Pharmacol Res* 61:553–563
 113. Desaphy JF, Pierno S, Léoty C, George AL Jr, De Luca A, Camerino DC (2001) Skeletal muscle disuse induces fibre type-dependent enhancement of Na(+)-channel expression. *Brain*. 124:1100–1113
 114. Bastide B, Kischel P, Puterflam J, Stevens L, Pette D, Jin JP et al (2002) Expression and functional implications of troponin T isoforms in soleus muscle fibers of rat after unloading. *Pflugers Arch* 444:345–352
 115. Desaphy JF, Pierno S, Liantonio A, De Luca A, Didonna MP, Frigeri A et al (2005) Recovery of the soleus muscle after short- and long-term disuse induced by hindlimb unloading: effects on the electrical properties and myosin heavy chain profile. *Neurobiol Dis* 18:356–365
 116. Frayssse B, Desaphy JF, Pierno S, De Luca A, Liantonio A, Mitolo CI et al (2003) Decrease in resting calcium and calcium entry associated with slow-to-fast transition in unloaded rat soleus muscle. *FASEB J*. 17:1916–1918
 117. Schulte LM, Navarro J, Kandarian SC (1993) Regulation of sarcoplasmic reticulum calcium pump gene expression by hindlimb unweighting. *Am J Physiol* 264:C1308–C1315
 118. Fitts RH, Riley DR, Widrick JJ (2001) Functional and structural adaptations of skeletal muscle to microgravity. *J Exp Biol* 204:3201–3208
 119. Adams GR, Caiozzo VJ, Baldwin KM (2003) Skeletal muscle unweighting: spaceflight and ground-based models. *J Appl Physiol* 95:2185–2201
 120. Paddon-Jones D, Sheffield-Moore M, Urban RJ, Sanford AP, Aarsland A, Wolfe RR et al (2004) Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bedrest. *J Clin Endocrinol Metab* 89:4351–4358
 121. Fitts RH, Romatowski JG, Peters JR, Paddon-Jones D, Wolfe RR, Ferrando AA (2007) The deleterious effects of bed rest on human skeletal muscle fibers are exacerbated by hypercortisolemia and ameliorated by dietary supplementation. *Am J Physiol Cell Physiol* 293:C313–C320
 122. Leach CS, Rambaut PC, Fischer CL (1975) A comparative study of two methods of urine preservation. *Clin Biochem* 8:108–117
 123. Grichko VP, Heywood-Cooksey A, Kidd KR, Fitts RH (2000) Substrate profile in rat soleus muscle fibers after hindlimb unloading and fatigue. *J Appl Physiol* 88:473–478
 124. Ojala BE, Page LA, Moore MA, Thompson LV (2001) Effects of inactivity on glycolytic capacity of single skeletal muscle fibers in adult and aged rats. *Biol Res Nurs* 3:88–95
 125. Stein TP, Wade CE (2005) Metabolic consequences of muscle disuse atrophy. *J Nutr* 135:1824S–1828S
 126. Murton AJ, Constantin D, Greenhaff PL (2008) The involvement of the ubiquitin proteasome system in human skeletal muscle remodelling and atrophy. *Biochim Biophys Acta* 1782:730–743
 127. Bodine SC, Stitt TN, Gonzalez M, Kline WO, Stover GL, Bauerlein R et al (2001) Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol* 3:1014–1019
 128. Yamamoto D, Ikeshita N, Matsubara T, Tasaki H, Heringtyas EH, Toda K et al (2008) GHRP-2, a GHS-R agonist, directly acts on myocytes to attenuate the dexamethasone-induced expressions of muscle-specific ubiquitin ligases, Atrogin-1 and MuRF1. *Life Sci* 82:460–466
 129. Ito T, Schaffer SW, Azuma J (2012) The potential usefulness of taurine on diabetes mellitus and its complications. *Amino Acids* 42:1529–1539
 130. Franconi F, Bennardini F, Mattana A, Miceli M, Ciuti M, Mian M et al (1995) Plasma and platelet taurine are reduced in subjects with insulin-dependent diabetes mellitus: effects of taurine supplementation. *Am J Clin Nutr* 61:1115–1119
 131. Elizavova EP, Nedosugova LV (1996) First experiments in taurine administration for diabetes mellitus. The effect on erythrocyte membranes. *Adv Exp Med Biol* 403:583–588
 132. Nakamura T, Ushiyama C, Suzuki S, Shimada N, Ohmuro H, Ebihara I et al (1999) Effects of taurine and vitamin E on microalbuminuria, plasma metalloproteinase-9, and serum type IV collagen concentrations in patients with diabetic nephropathy. *Nephron*. 83:361–362

133. Chauncey KB, Tenner TE Jr, Lombardini JB, Jones BG, Brooks ML, Warner RD et al (2003) The effect of taurine supplementation on patients with type 2 diabetes mellitus. *Adv Exp Med Biol* 526:91–96
134. Brøns C, Spohr C, Storgaard H, Dyerberg J, Vaag A (2004) Effect of taurine treatment on insulin secretion and action, and on serum lipid levels in overweight men with a genetic predisposition for type II diabetes mellitus. *Eur J Clin Nutr* 58:1239–1247
135. Moloney MA, Casey RG, O'Donnell DH, Fitzgerald P, Thompson C, Bouchier-Hayes DJ (2010) Two weeks taurine supplementation reverses endothelial dysfunction in young male type 1 diabetics. *Diab Vasc Dis Res* 7:300–310
136. Xiao C, Giacca A, Lewis GF (2008) Oral taurine but not *N*-acetylcysteine ameliorates NEFA-induced impairment in insulin sensitivity and beta cell function in obese and overweight, non-diabetic men. *Diabetologia* 51:139–146
137. Bergamini L, Mutani R, Delsedime M, Durelli L (1974) First clinical experience on the antiepileptic action of taurine. *Eur Neurol* 11:261–269
138. Azuma J, Sawamura A, Awata N, Ohta H, Hamaguchi T, Harada H et al (1985) Therapeutic effect of taurine in congestive heart failure: a double-blind crossover trial. *Clin Cardiol* 8:276–282
139. Fujita T, Ando K, Noda H, Ito Y, Sato Y (1987) Effects of increased adrenomedullary activity and taurine in young patients with borderline hypertension. *Circulation* 75:525–532
140. Dunn-Lewis C, Kraemer WJ, Kupchak BR, Kelly NA, Creighton BA, Luk HY et al (2011) A multi-nutrient supplement reduced markers of inflammation and improved physical performance in active individuals of middle to older age: a randomized, double-blind, placebo-controlled study. *Nutr J* 10:90
141. Pearl PL, Schreiber J, Theodore WH, McCarter R, Barrios ES, Yu J et al (2014) Taurine trial in succinic semialdehyde dehydrogenase deficiency and elevated CNS GABA. *Neurology* 18(82):940–944
142. González-Contreras J, Villalobos Gámez JL, Gómez-Sánchez AI, García-Almolda JM, Enguix Armanda A, Rius Díaz F et al (2012) Cholestasis induced by total parenteral nutrition: effects of the addition of Taurine (Tauramin®) on hepatic function parameters; possible synergistic action of structured lipids (SMOFlipid®). *Nutr Hosp* 27:1900–1907
143. Rosa FT, Freitas EC, Deminice R, Jordão AA, Marchini JS (2014) Oxidative stress and inflammation in obesity after taurine supplementation: a double-blind, placebo-controlled study. *Eur J Nutr* 53:823–830
144. Galloway SD, Talanian JL, Shoveller AK, Helgenhauser GJ, Spriet LL (2008) Seven days of oral taurine supplementation does not increase muscle taurine content or alter substrate metabolism during prolonged exercise in humans. *J Appl Physiol* 105:643–651
145. Spriet LL, Whitfield J (2015) Taurine and skeletal muscle function. *Curr Opin Clin Nutr Metab Care*. 18:96–101
146. Tappaz ML (2004) Taurine biosynthetic enzymes and taurine transporter: molecular identification and regulations. *Neurochem Res* 29:83–96
147. Ghandforoush-Sattari M, Mashayekhi S, Krishna CV, Thompson JP, Routledge PA (2010) Pharmacokinetics of oral taurine in healthy volunteers. *J AmIno Acids* 346237
148. Balshaw TG, Bampouras TM, Barry TJ, Sparks SA (2013) The effect of acute taurine ingestion on 3-km running performance in trained middle-distance runners. *Amino Acids* 44:555–561
149. Ra SG, Miyazaki T, Ishikura K, Nagayama H, Suzuki T, Maeda S et al (2013) Additional effects of taurine on the benefits of BCAA intake for the delayed-onset muscle soreness and muscle damage induced by high-intensity eccentric exercise. *Adv Exp Med Biol* 776:179–187
150. da Silva LA, Tromm CB, Bom KF, Mariano I, Pozzi B, da Rosa GL et al (2014) Effects of taurine supplementation following eccentric exercise in young adults. *Appl Physiol Nutr Metab* 39:101–104
151. Pechlivanis A, Kostidis S, Sarasilandis P, Petridou A, Tsalis G, Veselkov K et al (2013) 1H NMR study on the short- and long-term impact of two training programs of sprint running on the metabolic fingerprint of human serum. *J Proteome Res* 4(12):470–480
152. Gregor P, Hoff M, Holik J, Hadley D, Fang N, Coon H et al (1994) Dinucleotide repeat polymorphism in the human taurine transporter gene (TAUT). *Hum Mol Genet* 3:2263
153. Han X, Patters AB, Jones DP, Zellkovic I, Chesney RW (2006) The taurine transporter: mechanisms of regulation. *Acta Physiol (Oxf)* 187:61–73
154. Finlay EK, Berry DP, Wickham B, Gormley EP, Bradley DG (2012) A genome wide association scan of bovine tuberculosis susceptibility in Holstein-Friesian dairy cattle. *PLoS One* 7:e30545

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



From: [Rotstein, David](#)
To: [Jones, Jennifer L](#)
Subject: EON-266814; EON-266821 and EON-266827-Merrick-FW: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action
Date: Wednesday, August 24, 2016 6:35:16 AM

Here you go Jen.

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/ICERT
7519 Standish Place, RM 120

240-402-5613 (Office) (NEW NUMBER)

(b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: Burkholder, William
Sent: Wednesday, July 27, 2016 4:30 PM
To: Rotstein, David; Benjamin, Linda
Cc: Krieger, Darlene; Queen, Jackie L; Hodges, April; Conway, Charlotte
Subject: RE: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

OK Everyone. The product appears to be a dry extruded product, for which the AAFCO Cat Food Nutrient Profiles content for taurine is 0.10% on a dry matter basis. Clearly all three samples were analyzed to contain more than that amount of taurine. **On a dry matter basis the concentration of taurine in the samples was analyzed to be:**

FACTS #	Amount Taurine Found	%Moisture	%Dry Matter	Amount
Taurine on a Dry Matter Basis				
958500	0.183g/100g \approx 0.18%	2.20%	100 - 2.20 = 97.80%	
	0.183/0.9780 = 0.187%			
958501	0.153g/100g \approx 0.15%	1.99%	100 - 1.99 = 98.01%	
	0.153/0.9801 = 0.156%			
958504	0.171g/100g \approx 0.17%	2.79%	100 - 2.79 = 97.21%	
	0.171/0.9721 = 0.176%			

All of the Dry Matter Taurine percentages are above 0.10%. IF any of the samples were canned cat food, they would not be in compliance with the AAFCO Cat Food Nutrient Profiles for the recommended minimum taurine content and IF the label indicated the product was formulated to meet the AAFCO Cat Food Nutrient Profiles the product would be misbranded.

The answer to the question of consequence/causation of the taurine content in the product from which these three samples originated to the cats in the consumer complaint is that this(ese) lot(s) of product are not indicated to be causative. However, dilated cardiomyopathy from taurine deficiency

occurs over a long period of exposure to a deficient diet (months to a year or more), so, if these cats were eating the Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry for the 3 years indicated in the complaint, it is possible that the product was deficient for some long interval of time during that three year period and that a return to “normal” taurine levels in the diet were insufficient to correct the problem in the three cats that developed low blood taurine and the two with dilated cardiomyopathy. Treatment for dilated cardiomyopathy caused by taurine deficiency takes higher daily doses of taurine for several months than normal dietary amounts and is not completely curative.

Recommendations for regulatory steps to consider: [REDACTED] (b) (5)

[REDACTED]

Consider recommending the owner have an ophthalmic exam performed on the cat being treated for low blood taurine to see if there are signs of retinal degeneration due to taurine deficiency.

William J. Burkholder, DVM, PhD, DACVN
Leader, Nutrition and Labeling Team I, HFV-228
Division of Animal Feeds
Center for Veterinary Medicine
United States Food and Drug Administration
7519 Standish Place
Rockville, Maryland 20855
Phone: 240-402-5900
Fax: 240-453-6882
E-mail: william.burkholder@fda.hhs.gov

The opinions and information in this message are those of the author and do not necessarily reflect the views and policies of the U.S. Food and Drug Administration. Because of the nature of electronically transferred information, the integrity or security of this message cannot be guaranteed. This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution or copying is strictly prohibited. If you think you have received this e-mail message in error, please e-mail the sender immediately at William.Burkholder@fda.hhs.gov.

From: Rotstein, David
Sent: Wednesday, July 27, 2016 2:23 PM
To: Benjamin, Linda; Burkholder, William
Cc: Krieger, Darlene; Queen, Jackie L; Hodges, April
Subject: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Please see the moisture content below:

The moisture content for the samples are as follows:

FACTS #	Amount Taurine Found	%Moisture
958500	0.183g/100g ≈ 0.18%	2 20%
958501	0.153g/100g ≈ 0.15%	1 99%
958504	0.171g/100g ≈ 0.17%	2 79%

David Rotstein, DVM, MPVM, Dipl. ACVP
 CVM Vet-LIRN Liaison
 CVM OSC/DC/ICERT
 7519 Standish Place, RM 120

240-402-5613 (Office) (NEW NUMBER)

(b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: Benjamin, Linda
Sent: Wednesday, July 27, 2016 7:54 AM
To: Burkholder, William
Cc: Rotstein, David; Krieger, Darlene; Queen, Jackie L; Hodges, April
Subject: FW: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Hi Bill - Could you please respond to Dave Rotstein.

Dave - As ORS's numbers are very close to the 0.2% guarantee, it might be helpful to know the AV, CV, and/or 95% confidence limit for the analytical method. Additionally, do you know if the numbers below are being reported on a dry matter basis? FYI, the sample description on the collection reports (first 3 attachments) has either "One unopened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe weighing 5.4kg" or "Opened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe that only had 0.15kg of product. This sample was used by the consumer" but below **my green highlight** you referenced taurine # for canned products.

Sorry Bill - I just want to make sure you have everything you need.

Thanks for the opportunity to comment,
 Linda

From: Rotstein, David
Sent: Wednesday, July 27, 2016 7:22 AM
To: Benjamin, Linda
Cc: Krieger, Darlene; Queen, Jackie L; Hodges, April
Subject: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Linda,

We received an email from ORS with results for taurine for a cat food. Testing was based on a consumer complaint for 3 cats with cardiac disease and low taurine.

ORS has not finalized the results, but sent on the findings for the DRY cat food and asked whether CVM considers the results to be low based on the AAFCO requirements for wet cat food.

REQUEST: To answer the following questions:

- 1) Is the taurine low for a dry cat food based on AAFCO nutrient profiles?
- 2) If the taurine is low, would it be biologically significant for cats that ate this as their sole/primary diet?

The responses will [REDACTED] (b) (5).

REVIEWERS: Bill Burkholder, Krisztina Atkinson, Randall Lovell.

Date Needed: [REDACTED] (b) (5)

Email from the ORS Lab:

David I hope you can help us with these findings.

We received three consumer complaint Dry Cat Food products for Amino Acid analysis. We assayed the products for the Amino Acid profile and found only Taurine low.

FACTS #	Amount Taurine Found
958500	0.183g/100g ≈ 0.18%
958501	0.153g/100g ≈ 0.15%
958504	0.171g/100g ≈ 0.17%

The label for all of the samples are the same and Taurine is declared 0.20% minimum. The AAFCO Nutrient Profile from August 2015 states that the minimum limits for Taurine is 0.20% in canned products. Do you consider these product violated?

Attachments:

**Collection Reports
Pet Food Report
Vet-LIRN Summary**

Medical records were collected and evaluated by Vet-LIRN. These can be provided by request.

Thank you

David Rotstein, DVM, MPVM, Dipl. ACVP

CVM Vet-LIRN Liaison

CVM OSC/DC/ICERT

7519 Standish Place, RM 120

240-402-5613 (Office) (**NEW NUMBER**)

(b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david_rotstein@fda.hhs.gov.

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958500

This is an accurate reproduction of the original electronic record as of 07/27/2016

Flag	Flag Remarks				
Episode Number	Origin	Basis	Sample Type	FIS Smpl Num	Status
	Domestic	Surveillance	Official	16260261	Completed
FEI	Date Collected	Product Code	Responsible Firm	PAC	Hours
3004211953	06/23/2016	72AYT02	Manufacturer	71R801	11
Compliance Num	Country of Origin				
	United States				
Related Smpl Num	Position Class	Sampling District	NDC Number	Permit Number	Storage Rqrmnt.
	INV	NWJ-DO			Ambient
Dealer is Consumer	Crx/DEA Schedule	Recall Num	Consumer Compl. Num	Brand Name	
No			146048	Merrick	

Product Description
See Remarks Section.

Product Label
See continuation.

Reason for Collection	MFG Codes	Expiration Date
Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.	"16025DL1 38310 14131"	07/26/17

Firm Legal Name	Address	Type of Firm	Firm FEI	FCE
Merrick Pet Care, Inc.	3275 Tierra Blanca Rd Hereford, TX 79045-7823 US	Manufacturer	3004211953	02944
(b) (6)	(b) (6) US	Dealer	(b) (6)	

Size of Lot	Est. Value	Rcpt Type	Carrier Name	Date Shipped
One paper bag weighing 5.4kg	\$.00	FDA484		

Description of Sample
One unopened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe weighing 5.4kg

Method of Collection
See continuation.

How Prepared
See continuation.

Collector's Identification on Package and/or Label	Collector's Identification on Seal
"958500 06/23/2016 EB"	"958500 06/23/2016 Esteban Beltran"

Sample Delivered To	Date Delivered	Orig C/R & Records To
SRL-ACNA	06/28/2016	NWJ-DO
	Lab w/Split Sample	Lab
	0	SRL

Document Number	Document Date	Document Type	Document Remarks
1	06/23/2016	Other	FDA 484, Receipt for Sample, 1 page
2	06/23/2016	Other	FDA 484, Receipt for Sample Amend, 1 page
3	06/23/2016	Other	Photos of Product Labeling, 3 pages

Date: 07/27/2016

Page: 1 of 3

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958500

This is an accurate reproduction of the original electronic record as of 07/27/2016

Remarks

See continuation.

Payment Amount	Payment Method	704(d) Sample	702(b) Portion	Collector's Name	Date & Time of Signature	Meaning
\$27.99	Cash	No	No	Esteban Beltran	07/07/2016 08:54 AM ET	Collector

Name of Signer
Esteban Beltran

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958500

This is an accurate reproduction of the original electronic record as of 07/27/2016

Continuation:

Product Label

Finished Product: Label on bag read in parts: "****Lot #: 16025DL1 38310 14131 *** Merrick Whole Health Made Right Purrfect Bistro Grain Free *** REAL CHICKEN RECIPE FISH-FREE *** Net Wt 12Lb (5.4kg) *** WHOLE HEALTH MADE RIGHT! *** SQF INSTITUE CERTIFIED *** MERRICK PET CARE, INC. P.O. BOX 9800 AMARILLO, TX 79105 USA WWW.MERRICKCARE.COM *** Best By: July 26, 2017****"

Method of Collection

On 06/23/2016, I collected a sample from the storage area of a retail store. The sample was placed in a clear plastic bag. I officially sealed the clear plastic bag containing the sample with a FDA415a on site. The sample was transported via GOV to the NBRP sample prep room.

How Prepared

On 06/28/16, I attached an FDA 525 envelope to the sample and I placed the sample into a shipping box with bubble wrap. I secured the shipping box for shipment to SRL-ACNA via UPS (Tracking #: 1Z A47 51E 01 9888 4498). I delivered the sample inside to the NBRP reception area for UPS pickup.

Remarks

Product Description: Poultry based dry cat food packed in a dark orange paper bag with brown letters and an image of a cat on the front.

An amendment to the original FDA 484 was done in order to further describe what each sample number consists of and to identify what lot number of the product pertains to the sample number. CSO Gobiga Vanniyasingam assisted during the assignment. Related samples include 958501, 958502, 958503, 958504.

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958504

This is an accurate reproduction of the original electronic record as of 07/27/2016

Flag	Flag Remarks				
Episode Number	Origin	Basis	Sample Type	FIS Smpl Num	Status
	Domestic	Surveillance	Official	16260366	Completed
FEI	Date Collected	Product Code	Responsible Firm	PAC	Hours
3004211953	06/28/2016	72AYT02	Manufacturer	71R801	11
Compliance Num	Country of Origin				
	United States				
Related Smpl Num	Position Class	Sampling District	NDC Number	Permit Number	Storage Rqrmnt.
958500	INV	NWJ-DO			Ambient
Dealer is Consumer	Crx/DEA Schedule	Recall Num	Consumer Compl. Num	Brand Name	
No			146048	Merrick	

Product Description
See Remarks Section.

Product Label
See continuation.

Reason for Collection	MFG Codes	Expiration Date
Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.	"16025DL1 38310 14131"	07/26/17

Firm Legal Name	Address	Type of Firm	Firm FEI	FCE
Merrick Pet Care, Inc.	3275 Tierra Blanca Rd Hereford, TX 79045-7823 US	Manufacturer	3004211953	02944
(b) (6)	(b) (6)	Dealer	(b) (6)	

Size of Lot	Est. Value	Rcpt Type	Carrier Name	Date Shipped
One paper bag weighing 0.15kg	\$.00	FDA484		

Description of Sample
Opened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe that only had 0.15kg of product. This sample was used by the consumer.

Method of Collection
See continuation.

How Prepared
See continuation.

Collector's Identification on Package and/or Label	Collector's Identification on Seal
"958504 06/28/2016 EB"	"958504 06/28/2016 Esteban Beltran"

Sample Delivered To	Date Delivered	Orig C/R & Records To
SRL-ACNA	06/28/2016	NWJ-DO
	Lab w/Split Sample	Lab
	0	SRL

Document Number	Document Date	Document Type	Document Remarks
1	06/28/2016	Other	FDA 484, Receipt for Sample, 1pg.
2	06/28/2016	Other	Photos of Product Labeling, 2pgs.

Date: 07/27/2016

Page: 1 of 3

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958504

This is an accurate reproduction of the original electronic record as of 07/27/2016

Remarks

See continuation.

Payment Amount	Payment Method	704(d) Sample	702(b) Portion	Collector's Name
\$0.00	No Charge	No	No	Esteban Beltran
Name of Signer	Date & Time of Signature			Meaning
Esteban Beltran	07/07/2016 08:57 AM ET			Collector

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958504

This is an accurate reproduction of the original electronic record as of 07/27/2016

Continuation:

Product Label

Finished Product: Label on bag read in parts: "****Lot #:16025DL3 38310 14131****Merrick Whole Health Made Right Purrfect Bistro Grain Free *** REAL CHICKEN RECIPE FISH-FREE *** Net Wt 12Lb (5.4kg) *** WHOLE HEALTH MADE RIGHT! *** SQF INSTITUTE CERTIFIED *** MERRICK PET CARE, INC. P.O. BOX 9800 AMARILLO, TX 79105 USA WWW.MERRICKCARE.COM *** Best By: July 26, 2017****"

Method of Collection

On 06/28/16, I collected the used, opened bag which had been provided by the consumer to the (b) (6) (b) (6) The sample was placed in a clear plastic bag. I officially sealed the clear plastic bag containing the sample with a FDA415a on site. The sample was transported via GOV to the NBRP sample prep room.

How Prepared

On 06/28/16, I attached an FDA 525 envelope to the sample and I placed the sample into a shipping box with bubble wrap. I secured the shipping box for shipment to SRL-ACNA via UPS (Tracking #: 1Z A47 51E 01 9888 4498). I delivered the sample inside to the NBRP reception area for UPS pickup.

Remarks

Product Description: Poultry based dry cat food packed in a dark orange paper bag with brown letters and an image of a cat on the front.

CSO Gobiga Vanniyasingam assisted during the assignment. Related samples include 958500, 958501, 958502, 958503.

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
6/6/2016	C	(b) (6)	RECEPTION ACTIONS NOTE Sympathy card sent- (b) (6)
6/6/2016	C	(b) (6)	MEDICAL COMMENTS 6/6/2016 11:15 FDA complaint submitted: Pet Food Safety Report, ID 53897, was successfully submitted on 6/6/2016 11:15:17 AM EST to the FDA, and it was issued an Individual Case Safety Report Number (ICSR) of 1053335.
(b) (6)	C	(b) (6)	MEDICAL COMMENTS (b) (6) 16:26 (b) (6) AllForwardActions To: (b) (6) Sent Items Thursday, (b) (6) 4:24 PM Hi (b) (6), Sorry for the delay in getting back with you, I needed to get permission from the owner's before providing you with their contact information. Below is the their information as well as the names of the individual cats. The cat, (b) (6), with dilated cardiomyopathy was euthanized yesterday. Owners: (b) (6) Cats: (b) (6): 5/9/2016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016, euthanized on (b) (6) 5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016 - (b) (6): 9yr male neutered domestic long hair: 196 nmol/ml - (b) (6): 8y female spayed domestic short hair: 368 nmol/ml (b) (6): 9yr male neutered domestic long hair: 124 nmol/ml - (b) (6): 9yr male neutered domestic long hair: 536 nmol/ml Please let me know if you have any other questions.

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Sincerely,

(b) (6)
(b) (6)

(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)

(b) (6) Reply All

Wednesday, (b) (6) 12:56 PM

Thank You for providing me this information (b) (6). Could you provide us the pet parents information as well. We would like to reach out to the pet parent as well and speak with her. Thanks.

(b) (6) C (b) (6)

COMMUNICATIONS WITH CLIENT

(b) (6) 15:55

(b) (6) - expressed my condolences. asked for permission to provide contact info to company and the FDA - owner consented. Discussed what to expect when talking to company. Owner thankful for call.

(b) (6) R (b) (6)

Euthanasia Notice - FINAL (b) (6) - Euthanasia Notice

(b) (6)



B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

TO: (b) (6)
(b) (6)
(b) (6)
(b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: (b) (6)

RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 5 Mos. **Sex:** Spayed Female

~ EUTHANASIA NOTIFICATION ~

Dear Dr. (b) (6):

This letter is to inform you that your patient, (b) (6) was visited by (b) (6) At Home house call service today for end-of-life care.

If you have any questions, please feel free to contact me at the location noted above.

Thank you,

(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	C	(b) (6)	<p>MEDICAL COMMENTS - Closed Jun 04/2016 (b) (6) 18:03 Seen on emergency today, embolic event secondary to DCM. Discussed necropsy and advised that nutritionist and cardiologist agreed that prior test results were sufficient; necropsy would not reveal anything not already documented. Owner had already admin 0.2ml buprenex sublingually, requested I admin the remaining 0.3ml dispensed today which I did. They then spent time privately with the patient prior to euthanasia.</p> <p>Flushed cephalic catheter in right front leg; patent. Admin 20mg (2ml) expired propofol IV, apneic and unresponsive Admin 975mg (2.5ml) beuthanasia IV, 3 exhalation spasms followed Confirmed deceased by prolonged thoracic auscultation Removed IVC, placed (b) (6) in coffin, nested in owner's blanket</p>
(b) (6)	D	(b) (6)	Pleural Effusion Final
(b) (6)	D	(b) (6)	Feline Arterial Thromboembolic Disease Final
(b) (6)	R	(b) (6)	Referral Letter - Cardio Resident Eval and labs - FINAL (b) (6) - REF fxd

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

TO: (b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: Wednesday, (b) (6)
RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 5 Mos. Sex: Spayed Female
Current Weight: 5.3 kilograms as of 5/25/2016

Thank you for referring (b) (6). The following is a case summary.

Date of evaluation: Wednesday, (b) (6)

Date of previous cardiac evaluation: Wednesday, (b) (6)

CHIEF COMPLAINT: heavy breathing, dragging RH limb

HISTORY: (b) (6) previously doing well. Appetite had improved, eating 2/3 can max cal per day. Normal breathing. Owner noted acute onset of dragging RH limb this morning and heavy breathing. No interest in food this morning. Brought in to ER immediately. Received 9mg lasix IV total and 0.075mg buprenorphine IV on presentation. Previous hx: Diagnosed with DCM (b) (6). Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4. Taurine level confirmed taurine deficiency

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was tachypnic with moderate increased effort, RR 48. Normal BV sounds, no crackles on auscultation. Unable to bear weight on RH limb, dragging. Femoral pulses were fair in left hind, absent in left hind limb. Right paw pads cold to the touch. Heart rate was 160 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) 5/8/16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

k, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, age cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Cursory Ultrasound: small volume pleural effusion. No pericardial effusion. Large thrombus in LV.

Brief Echo 5/25/16: moderate volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle unchanged from previous.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM (b) (6) 2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm

IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %

Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), taurine deficiency, moderate left atrial enlargement, pleural effusion, hx azotemia, LV thrombus, FATE (partial)

THORACOCENTESIS 4/10/16: a total of 180ml of yellow fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed scant effusion remained in the patient.

THORACOCENTESIS (b) (6): a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: (b) (6) clinical signs are due to a partial aortic thromboembolus and recurrent left sided congestive heart failure. Thrombus previously noted in LV is still present. There is a very guarded prognosis of recovery with ATE. Discussed with owners, if they wants to go forward with treatment recommend hospitalization to manage CHF and pain. Discussed risk of reperfusion injury, main concern hyperkalemia which can be fatal. At home care will consist primarily of pain medication and nursing care. She may or may not gain function back to hind limb, and this could take up to 1-2 months to see improvement if any. Owner elected to take home to spend a few hours to spend time and have (b) (6) come for euthanasia this evening. IVC left in place (right cephalic). Sent home with additional dose of buprenorphine to keep her comfortable until euthanasia. Owners understand she may pass on her own. Gave owners my condolences.

MEDICATIONS:

Buprenorphine 0.3mg/ml- give entire contents of syringe (0.5ml) at 3pm today for pain control

Thank you for the courtesy of this interesting referral. Please feel free to contact me

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

with any questions or comments.

Sincerely,

(b) (6)

(b) (6)

Sent electronically - no signature required

(b) (6)

(b) (6)

Client ID: (b) (6) Patient ID: (b) (6) Patient Name: (b) (6)

Pain Condition Scores

Have you noticed the "PCS" aka Pain Condition Score in our medical records? As part of our AAHA specialty accreditation, we evaluate the level of pain or discomfort for each and every patient we see. You will see this reflected in the physical exam recordings as a "PCS". We follow the guidelines created and published by Colorado State University. If you have questions about interpreting pain or treatment options, or you need to schedule a pain management consult, please speak with our Anesthesiology & Pain Management Department by calling (b) (6).

Clinical Studies

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

We are recruiting for a variety of clinical studies across multiple specialties and are looking for eligible patients. For a complete list, please visit our website at [and click on Veterinary Professionals](#) and then Current Clinical Studies.

Avian & Exotics

(b) (6) is happy to re-introduce our Avian & Exotics Department and welcome (b) (6) DVM, ABVP (Avian Practice) to our team! We offer advanced diagnostics and treatment methods to complement the care provided by referring veterinarians. In addition to referral services, we provide primary care and medical boarding. Our facility maintains a dedicated exotics ward and a 24-hour monitored critical care unit. The Avian & Exotics Department accepts appointments 5 days a week (Tuesday-Saturday) and our emergency service receives patients, including exotic species, 24 hours a day. You can contact (b) (6)

(b) (6) C

(b) (6)

EMERGENCY PHYSICAL EXAM - Closed (b) (6)
(b) (6)

Chief Complaint: Respiratory distress

History: (b) (6) presented for STAT evaluation of respiratory distress. Owner noticed progressive tachypnea this AM and difficulty using right hindlimb. She did not want to eat this AM so she did not receive her AM medication. She is currently under the care of our Cardiology Service for Dilated Cardiomyopathy (DCM) (suspect secondary to taurine deficiency), moderate left atrial enlargement, pleural effusion, azotemia, and LV thrombus.

Other Medical Problems: None

Medications/Supplements: Pimobendan, Lasix, taurine supplementation, appetite stimulant

Environment: indoors only, several other cats

Vaccination Status: UTD

Current Diet (Type): Tempting to eat
- Frequency:

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

(b) (6)

Page 8 of 47

Date: 6/7/2016 2:33 PM

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

- Amount:

Physical Examination:

S(ubjective): BAR/distressed, hydration WNL, BCS 7/9, pain score: 1/4

O(b)jective):

Weight: 5.3 kilograms

TPR: T: 94.8 HR: 188, RR/RE: 60/rapid/shallow

EENT: clear AU/OU, no nasal discharge, normal cervical palpation, mm pink, moist/CRT<2 sec

INTEG: Hair coat ok

PLN: WNL

CV: NSR, no murmur ausculted, left femoral pulse moderate/synchronous, right very difficult to feel to absent

RESP: tachypneic, sl. dull ventrally, no crackles/wheezes

GI: soft, nonpainful, no masses

UG: FS, NSF

M/S: laterally recumbent, a

Neuro: alert/appropriate, cranial nerves intact, no placing deficits or spinal/neck pain

Problems/Differential Diagnoses:

Respiratory distress, decreased motor/absent femoral pulse RHL

Diagnostics:

None performed

Assessment:

12yo FS DSH

- absent to faint femoral pulse RHL, decreased motor, hypothermia, hx: DCM with LV thrombus- r/o saddle thrombus vs. other

- respiratory distress, mild amount pleural effusion on TFAST- r/o secondary to CHF secondary to DCM (suspect taurine deficiency)

- hx: Dilated Cardiomyopathy (DCM) (suspect secondary to taurine deficiency), moderate left atrial enlargement, pleural effusion, azotemia, and LV thrombus.

Treatment:

Placed in oxygen. IVC placed. 4mg Lasix IV, followed by additional 5mg IV. 0.015mg/kg Buprenorphine IV. Improved rr/re with above.

Plan/Recommendations:

Discussed PE at length with owner. Concerned for partial vs. full saddle thrombus RHL secondary to LV thrombus we know she has. Discussed options- point to

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Page 9 of 47

Date: 6/7/2016 2:33 PM

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

severity of underlying disease- ATH, repeat Echo, supportive care, vs. euthanasia. Owner elected to continue supportive care until they could speak with (b) (6), considering euthanasia. Elected RED code, transferred to cardiology.

(b) (6)	P	(b) (6)	0.50 mg of Buprenex (Buprenorphine) 0.3mg/mL MG C3 (MOBHL0) Rx #: 2579780 0 Of 0 Refills Give the entire contents of the syringe (0.5ml) under the tongue at 3pm.
---------	---	---------	---

(b) (6)	C	(b) (6)	CARDIAC EVALUTION - CLOSED 06/04/2016 - Cardiac Evaluation
---------	---	---------	--

Date of evaluation: Wednesday, (b) (6)

Date of previous cardiac evaluation: Wednesday, May 25, 2016

CHIEF COMPLAINT: heavy breathing, dragging RH limb

HISTORY: (b) (6) previously doing well. Appetite had improved, eating 2/3 can max cal per day. Normal breathing. Owner noted acute onset of dragging RH limb this morning and heavy breathing. No interest in food this morning. Brought in to ER immediately. Received 9mg lasix IV total and 0.075mg buprenohpine IV on presentation.

Previous hx: Diagnosed with DCM 5/9/16. Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4. Taurine level confirmed taurine deficiency

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was tachypnic with moderate increased effort, RR 48. Normal BV sounds, no crackles on auscultation. Unable to bear weight on RH limb, dragging. Femoral pulses were fair in left hind, absent in left hind limb. Right paw pads cold to the touch. Heart rate was 160 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) 5/8/16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Cursory Ultrasound: small volume pleural effusion. No pericardial effusion. Large thrombus in LV.

Brief Echo 5/25/16: moderate volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle unchanged from previous.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM 5/9/2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), taurine deficiency, moderate left atrial enlargement, pleural effusion, hx azotemia, LV thrombus, FATE (partial)

THORACOCENTESIS 4/10/16: a total of 180ml of yellow fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed scant effusion remained in the patient.

THORACOCENTESIS 5/9/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: (b) (6) clinical signs are due to a partial aortic thromboembolus and recurrent left sided congestive heart failure. Thrombus previously noted in LV is still present. There is a very guarded prognosis of recovery with ATE. Discussed with owners, if they want to go forward with treatment recommend hospitalization to manage CHF and pain. Discussed risk of reperfusion injury, main concern hyperkalemia which can be fatal. At home care will consist primarily of pain medication and nursing care. She may or may not gain function back to hind limb, and this could take up to 1-2 months to see improvement if any. Owner elected to take home to spend a few hours to spend time and have (b) (6) come for euthanasia this evening. IVC left in place (right cephalic). Sent home with additional dose of buprenorphine to keep her comfortable until euthanasia. Owners understand she may pass on her own. Gave owners my condolences.

MEDICATIONS:

Buprenorphine 0.3mg/ml- give entire contents of syringe (0.5ml) at 3pm today for pain control

(b) (6) CK (b) STAT
Reason for Visit: Emergency
Date Patient Checked Out: (b) (6) Practice TF

(b) (6) TC (b) (6) MEDICAL COMMENTS - TENTATIVE
(b) (6) 10:40
SW (b) (6) at Merrick - updated company that we have documented taurine deficiency in 2 other cats in house hold. The quality assurance team indicated that the level of taurine in the lot # I gave them was sufficient - discussed that this likely takes 3-6 month to develop and likely to be related to earlier lot and they need to

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

investigate further. Asked if the level they gave me was from them retesting the food after I called or from an earlier test - not sure. Asked for information from the taurine tests - told I can send summary of lab results. Also indicated that I am reporting this to the FDA. She give new info to her manager and quality assurance. Told them I expect them to follow up with me. Below email sent to Merrick:

Taurine Levels

(b) (6)
To:
(b) (6)@merrickpetcare.com
Hi (b) (6)

Thank you for your help with these cases. Here is the summary of the lab results:

12yr female spayed domestic short hair diagnosed and clinical for dilated cardiomyopathy
-5/9/2016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016

5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016
-9yr male neutered domestic long hair: 196 nmol/ml
-8y female spayed domestic short hair: 368 nmol/ml
-9yr male neutered domestic long hair: 124 nmol/ml
-9yr male neutered domestic long hair: 536 nmol/ml

Please let me know if you have any other questions.

Sincerely,

(b) (6)
(b) (6)
Clinical Nutrition Department

(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE (b) (6) 08:42 O Imom- was previously doing well. Eating ~2/3 can of max cal per day, sRR 6-7breaths/15sec. Now, this morning, dragging RH limb and breathing heavier.</p> <p>SWO- recommended to come in as soon as possible since (b) (6) breathing heavy. Unfortunately, cardiology will be in surgery this morning. Should go through emergency and I will consult.</p>
(b) (6)	B	(b) (6)	.08 mg of Buprenex (Buprenorphine) 0.3mg/mL MG C3 (MOBHL0) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Specialty/Referral Exam Level 3 (REF03) by (b) (6)
(b) (6)	B	(b) (6)	Echo Guided Thoracocentesis Group (EGT) by (b) (6)
(b) (6)	B	(b) (6)	1.00 EGT Procedure (USSC50) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Cared for by (b) (6)
(b) (6)	B	(b) (6)	.50 mg of Buprenex (Buprenorphine) 0.3mg/mL MG C3 (MOBHL0) by (b) (6)
(b) (6)	B	(b) (6)	At Home Euthanasia Group (HCEUTH) by (b) (6)
(b) (6)	B	(b) (6)	1.00 At Home Euthanasia Service (HC08) by (b) (6)
(b) (6)	B	(b) (6)	1.00 At Home Burial (HC10) by (b) (6)
(b) (6)	B	(b) (6)	100.00 mg of Telazol 100mg/mL inj per mg (C3-N) 2103 (MLTZL1) by (b) (6)
(b) (6)	B	(b) (6)	10.00 mg of Acepromazine 10mg/mL Inj per mg (MLA2L1) by (b) (6)
(b) (6)	B	(b) (6)	1.00 mg of Butorphanol 10mg/mL Inj per mg (C4) (MOB2L10) by (b) (6)
(b) (6)	B	(b) (6)	100.00 mg of Beuthanasia Soln 390 mg/mL (C-3N) / MG (MOB2L1) by (b) (6)
(b) (6)	B	(b) (6)	1.00 IV Catheter Placement (CATH) by (b) (6)
(b) (6)	B	(b) (6)	1.00 each of Tx Catheter IV 22g x 1" Surflo (BLUE) (H113) by (b) (6)
(b) (6)	B	(b) (6)	1.00 each of Tx IV Ext T Set Hospira 1265028 (H027) by (b) (6)
(b) (6)	B	(b) (6)	At Home Euthanasia Group (HCEUTH) by (b) (6)
(b) (6)	B	(b) (6)	1.00 At Home Euthanasia Service (HC08) by (b) (6)
(b) (6)	B	(b) (6)	1.00 At Home Burial (HC10) by (b) (6)
(b) (6)	B	(b) (6)	-100.00 mg of Telazol 100mg/mL inj per mg (C3-N) 2103 (MLTZL1) by (b) (6)
(b) (6)	B	(b) (6)	-10.00 mg of Acepromazine 10mg/mL Inj per mg (MLA2L1) by (b) (6)
(b) (6)	B	(b) (6)	-1.00 mg of Butorphanol 10mg/mL Inj per mg (C4) (MOB2L10) by (b) (6)
(b) (6)	B	(b) (6)	875.00 mg of Beuthanasia Soln 390 mg/mL (C-3N) / MG (MOB2L1) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)
(b) (6)	B	(b) (6)	1.00 each of Tx Injection Cap/Plug Termo 007110 (H118) by (b) (6)
(b) (6)	B	(b) (6)	Oxygen Therapy (Caged)/Hour (Group) (O2CAGE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 O2 Therapy Per Hour (T044) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Oxygen-related Patient Care / Hour (O2CARE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
(b) (6)	B	(b) (6)	4.00 mg of Furosemide (Lasix) 5% Injection per mg (MMF2L5) by (b) (6)
(b) (6)	B	(b) (6)	5.00 mg of Furosemide (Lasix) 5% Injection per mg (MMF2L5) by (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	B	(b) (6)	1.00 Cared for by (b) (6) (EGB) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Emergency Exam Level 4 (EE04) by (b) (6)
(b) (6)	P	(b) (6)	7.00 can of lams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) Rx #: 2578487 0 Of 6 Refills Feed up to 1 can daily.
(b) (6)	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE</p> <p>(b) (6) 13:52</p> <p>(b) (6) - (b) (6) doing ok. sRR7breaths/15 sec. Ate ~3/4 can of the lams max cal last night. Had normal BM yesterday. Hind limbs are very weak, one is worse than the other, but able to take a few steps on it before needing a rest. Does not see painful or distressed. Rec continue lasix 1/4 tab SID for now until appetite is consistent, then may consider increasasing. Continue pimo and taurine. Will put refill through for max cal.</p> <p>Emailed Client: I put through a prescription for 7 cans of food for (b) (6). She needs just under 1 can per day (although if she eats a whole can per day that is fine). There are also refills on the prescription if you need more. I will call to check in on her in a few days. Please call me with any concerns.</p>
(b) (6)	B	(b) (6)	7.00 can of lams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) by (b) (6)
(b) (6)	C	(b) (6)	<p>PHARMACY NOTE</p> <p>Returned O call, left voice message that medication is ready for pick up</p>
(b) (6)	P	(b) (6)	21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472) Rx #: 2569385 1 Of 12 Refills Filled by: (b) (6) Give 1 tablet by mouth twice daily with food.
(b) (6)	B	(b) (6)	21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472) by (b) (6)
(b) (6)	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE</p> <p>(b) (6) 10:12</p> <p>(b) (6) - (b) (6) back to licking gravy, not eating a lot of solid food. Hind legs are weak. Owner not able to get sRR yet but seems comfortable. A/o to continue with current meds. If stops eating, then stop lasix. Otherwise will touch base in a few days. Gave owner my cell phone number if they need anything.</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/25/2016	TC	(b)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/25/2016 18:38 (b) (6) - BW wnl, kidney values have decreased to normal. A/o to give lasix 12.5mg tabs- 1/4 tab SID. Will adjust based on appetite and breathing. Continue with other meds (pimo, taurine and app stimulant). Owner thankful.
5/25/2016	D	(b)	Pleural Effusion Final
5/25/2016	C	(b)	CARDIAC EVALUTION - CLOSED 05/28/2016 - Cardiac Evaluation

Date of evaluation: Wednesday, May 25, 2016

Date of previous evaluation: Sunday, May 15, 2016

CHIEF COMPLAINT: heavy breathing

HISTORY: Owners noted heavy breathing yesterday. Decreased appetite yesterday and today. Prior to that her appetite was improving. Owners transitioned her to royal canin and she started eating small amounts of solid food, previously only licking gravy.

Previous hx: Diagnosed with DCM 5/9/16. Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4. Taurine level confirmed taurine deficiency

PHYSICAL EXAM: The patient was bright, alert and responsive. No murmur on auscultation. The patient was tachypnic with mild increased effort, RR 48. Normal BV sounds, no crackles on auscultation. Faint referred upper airway noise. Femoral pulses were fair and synchronous. Heart rate was 180 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) 5/8/16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Brief Echo 5/25/16: moderate volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle unchanged from previous.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM 5/9/2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).
Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia, LV thrombus

THORACOCENTESIS 4/10/16: a total of 180ml of yellow fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed scant effusion remained in the patient.

THORACOCENTESIS 5/9/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.
ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: (b) (6) had 180ml fluid removed from her chest today. A renal panel showed normal renal values (BUN 31, creatinine 1.3)- previous azotemia. Will start with very low dose of lasix since decreased appetite right now (decreased appetite seemed to correlate with onset of heavy breathing). If appetite improves, can consider increasing lasix dose. Continue other medications as below. Recheck in 2 weeks, sooner if concerns.

MEDICATIONS:

START:

Lasix 12.5mg tablets- give 1/4 tablet by mouth once daily

CONTINUE:

Taurine 250mg by mouth twice daily

Mirtazepine 15mg tablets: Give 1/4 tablet by mouth every 3 days as needed.

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

5/25/2016 I

(b) (6)

Cardiology Discharge Instructions

(b) (6)

(b) (6)

(b) (6)

(b) (6)

(b) (6) had 160ml of fluid removed from her chest tonight. The clot in her heart appears similar to previous. I will call you with the bloodwork results and we can determine what to do with her lasix dose.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Recheck in 2 weeks, sooner if concerns.

MEDICATIONS:
CONTINUE:

Taurine 250mg by mouth twice daily
 Mirtazepine 15mg tablets: Give ¼ tablet by mouth every 3 days as needed.
 Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

WAIT TO BE INSTRUCTED FURTHER ON LASIX DOSE

Watch (b) (6) for the following clinical signs and call a veterinarian if you see any of these:

- Initiation of or increase in cough
- Excessive panting or wheezing
- Restlessness, unable to get comfortable
- Decreased appetite
- Lethargy/weakness
- Collapse or fainting

It has been a pleasure caring for (b) (6). Thank you for entrusting us with his care. If you have any further questions or problems, please don't hesitate to call.

5/25/2016	L	(b) (6)	(b) (6), (b) (6)	Cardiac Panel #10 results from (b) (6) Requisition ID: 0 <table border="0" style="width: 100%;"> <thead> <tr> <th style="text-align: left;">Test</th> <th style="text-align: left;">Result</th> <th style="text-align: left;">Reference Range</th> </tr> </thead> <tbody> <tr> <td>HCT =</td> <td>34 %</td> <td></td> </tr> <tr> <td>NA+ =</td> <td>147.3 mmol/L</td> <td>146.2 - 156.2</td> </tr> <tr> <td>K+ =</td> <td>6.65 mmol/L H</td> <td>3.41 - 4.71</td> </tr> <tr> <td>CL- =</td> <td>115.1 mmol/L L</td> <td>117.0 - 125.3</td> </tr> <tr> <td>BUN =</td> <td>31 mg/dL</td> <td>22 - 33</td> </tr> <tr> <td>CREA =</td> <td>1.3 mg/dL</td> <td>0.07 - 1.9</td> </tr> </tbody> </table> Manually entered. PCV = 32% TS = 6.0g/dL	Test	Result	Reference Range	HCT =	34 %		NA+ =	147.3 mmol/L	146.2 - 156.2	K+ =	6.65 mmol/L H	3.41 - 4.71	CL- =	115.1 mmol/L L	117.0 - 125.3	BUN =	31 mg/dL	22 - 33	CREA =	1.3 mg/dL	0.07 - 1.9
Test	Result	Reference Range																							
HCT =	34 %																								
NA+ =	147.3 mmol/L	146.2 - 156.2																							
K+ =	6.65 mmol/L H	3.41 - 4.71																							
CL- =	115.1 mmol/L L	117.0 - 125.3																							
BUN =	31 mg/dL	22 - 33																							
CREA =	1.3 mg/dL	0.07 - 1.9																							

5/25/2016	P	(b) (6)	30.00 ml of DNULsix 10mg/ml/ML (M0568) Rx #: 2576809 0 Of 12 Refills Give 0.5ml by mouth once daily or as directed by your veterinarian.
-----------	---	---------	--

5/25/2016	V	(b)	May 25, 2016 04:26 PM Staff: (b) ----- Weight : 5.30 kilograms Rm. 14
-----------	---	-----	--

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/25/2016	CK	(b) (6)	breathing heavy Reason for Visit: Recheck Date Patient Checked Out: 05/25/16 Practice (b) (6)
5/25/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/25/2016 14:40 (b) (6) - (b) (6) breathing heavy today. No interest in food yesterday or today. Owner to bring in this afternoon.
5/25/2016	B	(b) (6)	1.00 Specialty/Referral Exam Level 2 (REF02) by (b) (6)
5/25/2016	B	(b) (6)	Echo Guided Thoracocentesis Group (EGT) by (b) (6)
5/25/2016	B	(b) (6)	1.00 EGT Procedure (USSC50) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Thoracocentesis Therapeutic (R33) by (b) (6)
5/25/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/25/2016	B	(b) (6)	1.00 In-house lab (XNBALIX) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Cardiac (b) (6) Panel #10 ((b) (6)) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)
5/25/2016	B	(b) (6)	30.00 ml of DNULsix 10mg/ml/ML (M0568) by (b) (6)
5/25/2016	B	(b) (6)	-30.00 ml of DNULsix 10mg/ml/ML (M0568) by (b) (6)
5/22/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/22/2016 16:23 (b) (6) - Appetite is still the same, but now (b) (6) will go to the food on her own instead of owners bringing it to her. Still only eating gravy, no solid food yet. Owner has not tried Max Cal, recommended trying that. Cats who go prolonged period without eating at risk for hepatic lipidosis. Personatlity wise, she is much improved, almost back to normal self. Ambulating around the house as before. Very social. Owner bought Royal Canin as new diet. Gets taurine and pimo BID now. Told owner to continue appetite stimulant for now (had stopped this). Urinating and defecating outside of the litter box, not a SE of meds, likley behavioral. Soft stools. A/o let me know if soft stools continue.
5/19/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT 5/19/2016 16:18

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
<p>(b) (6) - updating that I did talk with Merrick customer service today to take my complaint and are filing it with their quality assurance. I am not sure when they will get back with me, but I will let them know as soon as I hear anything. Owner thankful for call.</p>			
5/19/2016	C	(b) (6)	<p>MEDICAL COMMENTS ***ADDENDUM 5/19/2016 5/19/2016 11:49 Called Merrick at 1(800)664-7387 to report taurine deficiency possibly related to consumption of their product, Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry (best by 7/26/2017, lot #16025 DL1 38310 14131 - lost # difficult to read), USB# 22808 38310). Owner has been feeding this food for approximately 3 years, 5 cats total in household, product has been purchased from the (b) (6). Requesting that the company investigate this possible deficiency, also discussed that I would like for the other cats in the household to be tested. (b) (6) @ Merrick - said I could expect call back in 2 weeks, let her know I would like to know when to expect a call. She will submit complaint and let me know. ADDENDUM on 5/19/2016 at 15:28:19 from (b) (6) DVM, DACVN Merrick called back - additional questions of how long the cat has been sick - presented to ER on 5/8 and sick day before; also wanted to know if bag was new - yes bag was purchased about 2 weeks prior per owner. My concern however is that it takes several months for this to develop and I do not believe this is a single bag/lot issue.</p>
5/19/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT 5/19/2016 10:10 (b) (6) - introduce3d myself, asked owner about diet history, has been feeding Merrick Purrfect Bistro Grain Free Real Chicken Recipe for approximately 3 years and purchasing from the (b) (6) Prior to this feeding Dick Van Pattons Indoor Formula Dry, chicken and salmon flavor. Discussed with owner that I will contact the company and also report to the FDA. Will let owner know of communication. In my experience sometimes the company will also want to reach out to the client. Owner thankful for call.</p>
5/18/2016	TC	(b)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE 5/18/2016 18:16 (b) (6)- (b) (6) still not eating, had a little gravy this morning. Drank a lot of water today. Breathing is normal. Owner dropping food off tonight.</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/18/2016	TC	(b) (6)	COMMUNICATIONS WITH DOCTOR - TENTATIVE 5/18/2016 18:15 Imom for (b) (6) regarding appt on Saturday with other cats in house. Would like whole blood taurine levels sent to the UC Davis amino acid lab. Call me or speak with nutrition regarding any questions.
5/17/2016	P	(b) (6)	3.00 can of Iams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) Rx #: 2573240 0 Of 3 Refills Feed as directed
5/17/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/17/2016 15:58 (b) (6) - did not eat much this morning. O gave appetite stimulant this morning and then left her alone with some food. Has not checked on her yet. Normal BM last night. sRR6brs/15sec last night. Vet coming for house call Saturday morning to take taurine sample for other cats. A/o to transition after that- recommended Hills Science Diet, Purina, Royal Canin. Will also rx Iams max cal for her to pick up here and offer (b) (6).
5/17/2016	B	(b) (6)	3.00 can of Iams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) by (b) (6)
5/16/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/16/2016 18:04 (b) (6) - (b) (6) seemed to be doing better last night. A little brighter when they got home. Started taurine supplementation last night. Not eating solid foods yet, but licking gravy- had the gravy from almost 3 cans last night. Owner put solid food in blender, but (b) (6) not interested (may have been too thick still). Drinking water. No BM, not a concern because she is not eating. Asked owner to bring in the food in original package as soon as possible, owner was planning on dropping off tomorrow. Also discussed to get other 4 cats tested for taurine levels this week, as since we are changing their diet we would like to know levels on current diet. Owner will call to either have mobile vet come to house or schedule here with GP this week. Told owner I will talk to nutrition about recommended diets.
5/15/2016	R	(b) (6)	Referral Letter - Cardio Resident Eval and labs - FINAL 05/15/2016 - (b) (6)

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

TO: (b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: Sunday, May 15, 2016
RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 4 Mos. Sex: Spayed Female
Current Weight: 5.2 kilograms as of 5/15/2016

Thank you for referring (b) (6). The following is a case summary.

Date of evaluation: Sunday, May 15, 2016

Date of previous cardiac evaluation: Monday, (b) (6)

CHIEF COMPLAINT: Recheck, not eating

HISTORY: (b) (6) has not eaten since discharge on (b) (6). Owner gave mirtazapine yesterday. Today licking some of the liquid off the food and very polydyspic, but no interest in eating any solid food. Will take a few steps and then lay down, very weak. Owner not able to get sRR, awake RR 6breaths/15sec. No heavy breathing noted.

Previous hx: Diagnosed with DCM (b) (6). Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4.

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was eupnic, RR 32. Normal BV sounds, no crackles on auscultation. Femoral pulses were fair and synchronous. Heart rate was 180 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) 5/8/16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted

CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, age cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, tentative med note, V:Vital signs

(b) (6)

Page 21 of 47

Date: 6/7/2016 2:33 PM

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

in left ventricle.

ECHOCARDIOGRAM (b) (6):

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).
Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia, LV thrombus

THORACOCENTESIS 5/9/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.
ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: The small amount of pleural effusion was not enough to warrant thoracocentesis, especially since patient eupnic. (b) (6) taurine level came back low (24nmol/ml). Recommended supplementing taurine at below dose. Discussed with owner it can take up to 2-3 weeks to see an effect of this and even longer (3 months) to see changes on echocardiogram. There is a chance (b) (6) will succumb to congestive heart failure before we see the positive effects of the taurine. A renal panel revealed azotemia. Discussed with owner the challenge going forward of managing the heart failure with azotemia. If (b) (6) develops trouble breathing, may not have options to treat. Euthanasia may be the most humane option for (b) (6) at that time.

The echocardiogram showed a large mass in the left ventricle, consistent with a thrombus. There is a risk that this clot, or a piece of it, leaves the heart and causes a thromboembolic event. The most common place for thromboembolic disease in cats is the aortic bifurcation, but can occur anywhere including the lungs which can be fatal. Advised owners, if FATE suspected at home, bring (b) (6) to closest emergency facility. At that time, euthanasia is recommended. We discussed holding off on an aspirin or Plavix for now, as it will not affect the current clot, and (b) (6) is not yet eating.

Owners willing to start taurine supplementation and take it day by day. Owners understand the possible outcomes we discussed today. I will call to check in tomorrow.

MEDICATIONS:

START:

Taurine 250mg by mouth twice daily

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

CONTINUE:

Mirtazepine 15mg tablets: Give 1/4 tablet by mouth every 3 days as needed.

HOLD FOR NOW:

Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

Thank you for the courtesy of this interesting referral. Please feel free to contact me with any questions or comments.

Sincerely,

(b) (6) (Cardiology Resident)

(b) (6), BVSc, MRCVS, ACVIM (Cardiology)
Sent electronically - no signature required

(b) (6)

Client ID: (b) (6) Patient ID: (b) (6) Patient Name (b) (6)

DATE/TIME	TEST	RESULT	REFERENCE RANGE
5/15/2016	BUN	= 61 mg/dL (H)	22 - 33
5/15/2016	CL-	= 104.5 mmol/L (L)	117.0 - 125.3
5/15/2016	CREA	= 3.1 mg/dL (H)	0.07 - 1.9
5/15/2016	HCT	= 43 %	
5/15/2016	K+	= 3.33 mmol/L (L)	3.41 - 4.71
5/15/2016	NA+	= 145.3 mmol/L (L)	146.2 - 156.2

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Page 23 of 47

Date: 6/7/2016 2:33 PM

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Lab Comments: Manually entered.

Additional Comments: BUN/Cre=19.7mg/mg PCV=39% TS=7.8g/dL Normal

Pain Condition Scores

Have you noticed the "PCS" aka Pain Condition Score in our medical records? As part of our AAHA specialty accreditation, we evaluate the level of pain or discomfort for each and every patient we see. You will see this reflected in the physical exam recordings as a "PCS". We follow the guidelines created and published by Colorado State University. If you have questions about interpreting pain or treatment options, or you need to schedule a pain management consult, please speak with our Anesthesiology & Pain Management Department by calling (b) (6).

Clinical Studies

We are recruiting for a variety of clinical studies across multiple specialties and are looking for eligible patients. For a complete list, please visit our website at [and click on Veterinary Professionals and then Current Clinical Studies.](#)

Avian & Exotics

(b) (6) is happy to re-introduce our Avian & Exotics Department and welcome (b) (6), ABVP (Avian Practice) to our team! We offer advanced diagnostics and treatment methods to complement the care provided by referring veterinarians. In addition to referral services, we provide primary care and medical boarding. Our facility maintains a dedicated exotics ward and a 24-hour monitored critical care unit. The Avian & Exotics Department accepts appointments 5 days a week (Tuesday-Saturday) and our emergency service receives patients, including exotic species, 24 hours a day. You can contact (b) (6)

5/15/2016 TC

(b) (6)

COMMUNICATIONS WITH CLIENT - TENTATIVE
5/15/2016 16:34

(b) (6) - discussed azotemia. Since (b) (6) is eupnic, would hold off on lasix for now. Hope would be that she may be able to breathe comfortably without lasix for enough time that taurine may start to help. Otherwise may give low dose of lasix, but going to be a big challenge with azotemia. Owners are to start taurine tonight. Discussed case with nutrition. Will file a complaint about the food. Will have more

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			information on this tomorrow. Will call to check in tomorrow.
5/15/2016	D	(b) (6)	Taurine Deficiency Final
5/15/2016	D	(b) (6)	Azotemia Tentative
5/15/2016	D	(b) (6)	Pleural Effusion Final

5/15/2016 C (b) (6) CARDIAC EVALUTION - CLOSED 05/18/2016 - Cardiac Evaluation

Date of evaluation: Sunday, May 15, 2016

Date of previous cardiac evaluation: Monday, (b) (6)

CHIEF COMPLAINT: Recheck, not eating

HISTORY: (b) (6) has not eaten since discharge on (b) (6). Owner gave mirtazapine yesterday. Today licking some of the liquid off the food and very polydyspic, but no interest in eating any solid food. Will take a few steps and then lay down, very weak. Owner not able to get sRR, awake RR 6breaths/15sec. No heavy breathing noted.

Previous hx: Diagnosed with DCM (b) (6). Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4.

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was eupnic, RR 32. Normal BV sounds, no crackles on auscultation. Femoral pulses were fair and synchronous. Heart rate was 180 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) 5/8/16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM 5/9/2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (b) (6).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia, LV thrombus

THORACOCENTESIS 5/9/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.
ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: The small amount of pleural effusion was not enough to warrant thoracocentesis, especially since patient eupnic. (b) (6) taurine level came back low (24nmol/ml). Recommended supplementing taurine at below dose. Discussed with owner it can take up to 2-3 weeks to see an effect of this and even longer (3 months) to see changes on echocardiogram. There is a chance (b) (6) will succumb to congestive heart failure before we see the positive effects of the taurine. A renal panel revealed azotemia. Discussed with owner the challenge going forward of managing the heart failure with azotemia. If (b) (6) develops trouble breathing, may not have options to treat. Euthanasia may be the most humane option for (b) (6) at that time. The echocardiogram showed a large mass in the left ventricle, consistent with a thrombus. There is a risk that this clot, or a piece of it, leaves the heart and causes a thromboembolic event. The most common place for thromboembolic disease in cats is the aortic bifurcation, but can occur anywhere including the lungs which can be fatal. Advised owners, if FATE suspected at home, bring (b) (6) to closest emergency facility. At that time, euthanasia is recommended. We discussed holding off on an aspirin or Plavix for now, as it will not affect the current clot, and (b) (6) is not yet eating. Owners willing to start taurine supplementation and take it day by day. Owners understand the possible outcomes we discussed today. I will call to check in tomorrow.

MEDICATIONS:

START:

Taurine 250mg by mouth twice daily

CONTINUE:

Mirtazepine 15mg tablets: Give 1/4 tablet by mouth every 3 days as needed.

HOLD FOR NOW:

Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

5/15/2016	I	(b) (6)	Cardiology Discharge Instructions
		(b) (6)	
		(b) (6)	
		(b) (6)	
		(b) (6)	

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6) has a small amount of fluid in her chest today. It was not enough to warrant draining today.

Her taurine levels came back low. Please start supplementing taurine as below. It can take up to 2-3 weeks to see an effect of this.

The echocardiogram showed a large mass in one of the chambers of her heart (the left ventricle). There is a risk that this clot, or a piece of it, leaves the heart. If that happens, it can travel to any part of the body (lungs, hind legs, etc) and this can be fatal. We discussed holding off on an aspirin or Plavix medication for now, as it will not do anything for the current clot, and (b) (6) is not yet eating. I will call you with her bloodwork results this afternoon.

MEDICATIONS:

START:

Taurine 250mg by mouth twice daily

Watch (b) (6) for the following clinical signs and call a veterinarian if you see any of these:

- Initiation of or increase in cough
- Excessive panting or wheezing
- Restlessness, unable to get comfortable
- Decreased appetite
- Lethargy/weakness
- Collapse or fainting

It has been a pleasure caring for (b) (6). Thank you for entrusting us with his care. If you have any further questions or problems, please don't hesitate to call.

5/15/2016	L	(b) (6)	(b) (6), (b) (6)	Cardiac Panel #10 results from (b) (6) In-Clinic Requisition ID: 0 <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Test</th> <th>Posted</th> <th>Result</th> <th>Final</th> <th>Reference Range</th> </tr> </thead> <tbody> <tr> <td>HCT =</td> <td></td> <td>43 %</td> <td></td> <td></td> </tr> <tr> <td>NA+ =</td> <td></td> <td>145.3 mmol/L</td> <td>L</td> <td>146.2 - 156.2</td> </tr> <tr> <td>K+ =</td> <td></td> <td>3.33 mmol/L</td> <td>L</td> <td>3.41 - 4.71</td> </tr> <tr> <td>CL- =</td> <td></td> <td>104.5 mmol/L</td> <td>L</td> <td>117.0 - 125.3</td> </tr> <tr> <td>BUN =</td> <td></td> <td>61 mg/dL</td> <td>H</td> <td>22 - 33</td> </tr> <tr> <td>CREA =</td> <td></td> <td>3.1 mg/dL</td> <td>H</td> <td>0.07 - 1.9</td> </tr> </tbody> </table> <p>Manually entered. BUN/Cre=19.7mg/mg PCV=39% TS=7.8g/dL Normal</p>	Test	Posted	Result	Final	Reference Range	HCT =		43 %			NA+ =		145.3 mmol/L	L	146.2 - 156.2	K+ =		3.33 mmol/L	L	3.41 - 4.71	CL- =		104.5 mmol/L	L	117.0 - 125.3	BUN =		61 mg/dL	H	22 - 33	CREA =		3.1 mg/dL	H	0.07 - 1.9
Test	Posted	Result	Final	Reference Range																																			
HCT =		43 %																																					
NA+ =		145.3 mmol/L	L	146.2 - 156.2																																			
K+ =		3.33 mmol/L	L	3.41 - 4.71																																			
CL- =		104.5 mmol/L	L	117.0 - 125.3																																			
BUN =		61 mg/dL	H	22 - 33																																			
CREA =		3.1 mg/dL	H	0.07 - 1.9																																			

5/15/2016 V (b) May 15, 2016 03:24 PM Staff: (b)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History

			Weight : 5.20 kilograms cardio baby scale
5/15/2016	CK	(b) (6)	Reason for Visit: Recheck Date Patient Checked Out: (b) (6) Practice (b) (6)
5/15/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE (b) (6) 13:29 (b) (6) (mrs)- owner gave mirtazapine, no improvement in appetite. Drinking excessively. Having a hard time walking, very weak. Owner not able to get sRR, awake breathing 6breaths/15sec. Offered to see (b) (6) today. Made appt for 3pm
5/15/2016	B	(b) (6)	1.00 Specialty/Referral Exam Level 2 (REF02) by (b) (6)
5/15/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/15/2016	B	(b) (6)	1.00 In-house lab (XNBALIX) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Cardiac (b) (6) Panel #10 ((b) (6)) by (b) (6)
5/15/2016	B	(b) (6)	Echo Guided Thoracocentesis Group (EGT) by (b) (6)
5/15/2016	B	(b) (6)	1.00 EGT Procedure (USSC50) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Cared for by (b) (6) by (b) (6)
5/14/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/14/2016 17:42 Mrs called and Imovm that (b) (6) hasn't been eating well. I called back and sw Mr. He said she is eating only very tiny amounts and not improving, wanted to know if I had suggestions. Owners feel breathing is still ok, 6 breaths/15 seconds but coughed a little today. I told Mr she could have poor app due to fluid reforming or azotemia or her heart disease in general. She may need to be rechecked sooner than later to evaluate this and r/o fluid and worsening azo. Owners plan to discuss w/ (b) (6) but wanted to know if there is something they could give her before morning. I offered to prescribe appetite stimulant, explained that this may not work b/c it doesn't override what is causing the inappetance in the 1st place but it's fine to try. Mr was thankful, said he may or may not pick it up tonight but is glad to have the option.
5/14/2016	P	(b) (6)	2.00 tablet of Mirtazapine 15mg Tablet (M1052)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/14/2016	B	(b) (6)	Rx #: 2571986 0 Of 3 Refills Give 1/4 tablet once every 3 days as needed to stimulate appetite. 2.00 tablet of Mirtazapine 15mg Tablet (M1052) by (b) (6)
5/12/2016	C	(b) (6)	TRIAGE CALL 5/12/2016 21:23 Per owner, (b) (6) appetite has been decreasing over the last couple days. Yesterday only ate about 2 tablespoons, today less. Let owner know that if the appetite has been decreasing recommend a recheck. Owner wants to talk to cardio first to see about an appetite stimulant. She will call tomorrow to speak with cardio department.
5/11/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/11/2016 13:16 (b) (6) (mrs)- still weak and unstable, but up and walking around short distances. sRR was 5breaths/15sec this morning. Ate 2 teaspoons canned food last night, so owner gave 1/4 tab lasix. Has not eaten yet this morning. Advised owner since sRR wnl, hold off on lasix for now. Will restart when either (b) (6) has a good appetite or if sRR >8br/15esc. Owner understands. Also hold off on pimo and taurine supplement. Should have taurine level back by recheck in 2 weeks. Owner asked about starting aspirin. Can consider aspirin/plavix at recheck if appetite is good. Discussed that they may lower risk, but do not prevent risk of clot formation. Will call to check on appetite in a few days. Owner to call sooner with concerns.
5/10/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/10/2016 10:13 (b) (6) (mrs)- (b) (6) was drinking a lot last night. Has not eaten anything yet. sRR 7breaths/15sec. Has gotten up and walked around, otherwise sleeping. Advised owner to hold off on meds today, would like her eating before restarting them. Will call tomorrow to check on appetite and advised what to do with lasix.
5/10/2016	L	(b) (6)	Miscellaneous results from (b) (6) (East) Requisition ID: 189206 Posted Final Asc: (b) (6) Profile: Taurine RE: 16758 Sample: PLASMA, HEPARIN RE: 16759 Taurine 24 NMOL/ML nmol/ml Feline taurine ranges: normal plasma 60-120 nmol/mL critical level <40 nmol/mL; whole blood normal 300-600 nmol/mL

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			<p>critical level <200 nmol/mL. TEST PERFORMED AT THE UNIVERSITY OF WISCONSIN</p> <p>FELINE PLASMA NORMALS: 60-120 nmol/ml WHOLE BLOOD NORMALS: 300-600 nmol/ml PLASMA CRITICAL: LESS THAN 40 nmol/ml WHOLE BLOOD CRITICAL: LESS THAN 200 nmol/ml</p> <p>K9 PLASMA NORMALS: 60-120 nmol/ml WHOLE BLOOD NORMALS: 200-350 nmol/ml PLASMA CRITICAL: LESS THAN 40 nmol/ml WHOLE BLOOD CRITICAL: LESS THAN 150 nmol/ml</p>

(b) (6) R (b) Referral Letter - Cardio Resident Eval and labs - FINAL (b) (6) - (b)

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

TO: (b) (6)
(b) (6)
(b) (6)
(b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: Monday, May 09, 2016
RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 4 Mos. **Sex:** Spayed Female
Current Weight: 15.6 pounds as of 12/28/2009

Thank you for referring (b) (6). The following is a case summary.

Date of evaluation: Monday, (b) (6)

CHIEF COMPLAINT: pleural effusion

HISTORY: Presented to ER last night for lethargy and ADR. Cursorsy ultrasound revealed pleural effusion. Thoracocentesis yielded 25ml yellow tinged fluid from the right side. Patient received 12mg lasix IV and was placed in oxygen overnight. Her RR was wnl, with slight effort noted overnight. She ate a small amount. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3).

PHYSICAL EXAM: The patient was bright, alert and responsive. No murmur on auscultation, but heart sound slightly muffled. The patient was eupnic, RR 30 with slight effort. Normal BV sounds, no crackles on auscultation. Pulses were fair and synchronous. Heart rate was 170 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) 5/8/16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM). Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial

(b) (6)

ack, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, :Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, C:Tentative medl note, V:Vital signs

(b) (6)

Page 31 of 47

Date: 6/7/2016 2:33 PM

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia

THORACOCENTESIS 5/9/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

SUMMARY: (b) (6) has dilated cardiomyopathy (DCM) and has developed pleural effusion due to congestive heart failure. The pleural effusion was removed today and we are starting the medications as described below. Although taurine deficiency is a rare cause for dilated cardiomyopathy in cats, we submitted a taurine level to the lab today to look for this. The test results can take up to 2 weeks; we will fax results when they are available. Unfortunately the prognosis for DCM (with no taurine deficiency) in cats is poor, with survival time 3-6 weeks or in rare cases 4-6 months. A renal panel was recheck today and showed elevated BUN (74) and normal creatinine (1.4). Discussed with owner the azotemia may make treating the congestive heart failure challenging.

We discussed that with heart enlargement there is a risk for developing a blood clot, or stroke. Although aspirin +/- plavix can be given in hopes of reducing clot formation, they have not been definitively proven to prevent blood clot formation in cats. If the owners elect to start this medication I would recommend waiting until we know she is eating and feeling well on the below medications as aspirin/plavix can cause GI side effects (vomiting, inappetance) in some cats. A recheck with cardiology is recommended in 2 weeks, or sooner if clinical signs.

Advised owner congestive heart failure likely contributing to lethargy, but concern that weakness may be secondary to the DCM and low cardiac output. If that is the case, may not improve once we get her out of failure. Owner comfortable taking (b) (6) home to see how she does. (b) (6) was discharged on the below medications. If she does not improve at home over the next few days, owner to consider euthanasia.

MEDICATIONS:

START: Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

START in 3 DAYS IF EATING

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.
If eating and easy to medicate, start: Taurine 250 mg by mouth twice a day with food.

Thank you for the courtesy of this interesting referral. Please feel free to contact me

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

with any questions or comments.

Sincerely,

(b) (6)

(b) (6)

(b) (6)

(b) (6)

(b) (6)

Client ID: (b) (6) Patient ID: (b) (6) Patient Name: (b) (6)

DATE/TIME	TEST	RESULT	REFERENCE RANGE
(b) (6)	CREA	= 1.4 mg/dL	0.8 - 2.4

Lab Comments: CREA: Test results for the latest analyzer run have been multiplied by the dilution factor for a dilution of 1 in 4 total.

DATE/TIME	TEST	RESULT	REFERENCE RANGE
(b) (6)	ALB	= 2.8 g/dL	2.3 - 3.9
(b) (6)	ALB/GLOB	= 0.8	
(b) (6)	ALKP	= 11 U/L (L)	14 - 111
(b) (6)	ALT	= 140 U/L (H)	12 - 130
(b) (6)	BUN/UREA	= 74 mg/dL (H)	16 - 36
(b) (6)	Chloride	= 100 mmol/L (L)	112 - 129
(b) (6)	CREA	= --- mg/dL	0.8 - 2.4
(b) (6)	GLOB	= 3.3 g/dL	2.8 - 5.1
(b) (6)	GLU	= 105 mg/dL	71 - 159
(b) (6)	Na/K	= 29	
(b) (6)	OSM calc	= 298 mmol/kg	
(b) (6)	PHOS	= 7.0 mg/dL	3.1 - 7.5
(b) (6)	Potassium	= 4.7 mmol/L	3.5 - 5.8
(b) (6)	Sodium	= 138 mmol/L (L)	150 - 165

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	TP		= 6.1 g/dL 5.7 - 8.9

Pain Condition Scores

Have you noticed the "PCS" aka Pain Condition Score in our medical records? As part of our AAHA specialty accreditation, we evaluate the level of pain or discomfort for each and every patient we see. You will see this reflected in the physical exam recordings as a "PCS". We follow the guidelines created and published by Colorado State University. If you have questions about interpreting pain or treatment options, or you need to schedule a pain management consult, please speak with our Anesthesiology & Pain Management Department by calling (b) (6)

Clinical Studies

We are recruiting for a variety of clinical studies across multiple specialties and are looking for eligible patients. For a complete list, please visit our website at [and click on Veterinary Professionals and then Current Clinical Studies.](#)

Avian & Exotics

(b) (6) is happy to re-introduce our Avian & Exotics Department and welcome (b) (6), DVM, ABVP (Avian Practice) to our team! We offer advanced diagnostics and treatment methods to complement the care provided by referring veterinarians. In addition to referral services, we provide primary care and medical boarding. Our facility maintains a dedicated exotics ward and a 24-hour monitored critical care unit. The Avian & Exotics Department accepts appointments 5 days a week (Tuesday-Saturday) and our emergency service receives patients, including exotic species, 24 hours a day. You can contact (b) (6)

(b) (6) | (b) Cardiology Discharge Instructions
(b) (6)
(b) (6)
(b) (6)
(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

A cardiologist has evaluated (b) (6) and has diagnosed her with Dilated Cardiomyopathy (DCM). DCM means she has poor muscle contraction of the heart and she has developed significant heart enlargement over time. Her clinical signs were due to congestive heart failure (fluid buildup around the lungs called pleural effusion), which developed secondary to the enlarged heart. We removed all the pleural effusion today. The fluid will reform but how fast this occurs is unpredictable. Please start the medications as below to help clear fluid and slow the fluid formation.

Although taurine deficiency is a rare cause for cardiomyopathy in cats, we submitted a taurine level to the lab today to look for this. The test results can take up to 2 weeks; we will call you when the results are available.

As we discussed, (b) (6) has elevations in her kidney values. This can make treating her heart disease challenging because she may not tolerate the lasix. If her kidney values become elevated to a certain degree, it will make her feel sick and she will likely have a decreased appetite or stop eating. We will monitor her kidney values with bloodwork over time.

Cats with heart enlargement are at risk for developing a blood clot, or stroke. Although aspirin and/or plavix can be given in hopes of reducing clot formation, they have not been proven to prevent blood clot formation in cats. If you elect to start this medication I would recommend waiting until she is eating and feeling well at home as these medications can cause GI side effects (vomiting, inappetance) in some cats.

Please periodically take a sleeping respiratory rate (sRR) at home. WHILE (b) (6) IS SLEEPING, count the number of times she breathes in over 15 seconds. She should breathe 8 or fewer breaths in 15 seconds.

A recheck with cardiology is recommended in 2 weeks, or sooner if you see any of the below signs.

MEDICATIONS:

START TODAY: Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet by mouth once a day

Furosemide: Also called Salix or Lasix. This is a diuretic and will help clear the fluid from your pet's lungs. Side effects include electrolyte abnormalities (if they stop eating), dehydration and kidney enzyme elevations. Blood work can be done to monitor these. This medication will be probably given for the life of your pet.

START IN 3 DAYS IF EATING:

Pimobendan 1.5 mg tiny tabs: Give 1 tablet by mouth two times a day WITH

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

FOOD.

Pimobendan is a phosphodiesterase inhibitor that gives increased contractility and arterial vasodilation. This will help the heart function better, allow your cat to feel better and live longer. Any medication can upset the stomach. This drug does not typically cause this, but if you see any changes, please stop the drug till you talk to a doctor here at (b) (6). Please give this with (b) (6) meals. Giving on empty stomach is more likely to make her nauseous.

We have called this medication into (b) (6). Please call them to order it and they will mail it to you.

If eating, start: Taurine 250 mg by mouth twice a day with food. I have submitted blood for a taurine level. The result may not return for 2 weeks. In the meantime, please start Taurine at home, 250 mg two times a day with food. This can be purchased at any health food store. If she is not eating well or if it is difficult to give her this medication, you can skip this until we get the taurine result from the blood work.

Watch (b) (6) for the following clinical signs and call a veterinarian if you see any of these:

- Initiation of or increase in cough
- Excessive panting or wheezing
- Restlessness, unable to get comfortable
- Decreased appetite
- Lethargy/weakness
- Collapse or fainting

It has been a pleasure caring for (b) (6). Thank you for entrusting us with his care. If you have any further questions or problems, please don't hesitate to call.

(b) (6) L

(b) (6)

Chemistry results from (b) (6) In-clinic

Laboratory Test	Requisition ID: 197	Posted	Final
Test	Result	Reference Range	
CREA =	1.4 mg/dL	0.8 - 2.4	

CREA: Test results for the latest analyzer run have been multiplied by the dilution factor for a dilution of 1 in 4 total.

(b) (6) C

(b) (6)

PHARMACY NOTE

Called (b) (6) and spoke to (b) (6). Ordered Pimobendan

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			1.5mg tiny tabs. Give 1 tablet by mouth twice daily with food. #100, 12 refills

(b) (6) L	Chemistry results from Laboratory Requisition	(b) (6)	In-clinic Posted	Final Range
Test	Result		Reference	
ALB =	2.8 g/dL		2.3 - 3.9	
ALKP =	11 U/L L		14 - 111	
ALT =	140 U/L H		12 - 130	
BUN/UREA =	74 mg/dL H		16 - 36	
Chloride =	100 mmol/L L		112 - 129	
CREA -	--.-- mg/dL		0.8 - 2.4	
GLU =	105 mg/dL		71 - 159	
PHOS =	7.0 mg/dL		3.1 - 7.5	
Potassium =	4.7 mmol/L		3.5 - 5.8	
Sodium =	138 mmol/L L		150 - 165	
TP =	6.1 g/dL		5.7 - 8.9	
GLOB =	3.3 g/dL		2.8 - 5.1	
ALB/GLOB =	0.8			
Na/K =	29			
OSM calc =	298 mmol/kg			

(b) (6) TC (b) COMMUNICATIONS WITH CLIENT - TENTATIVE
 (b) (6) 12:35
 10am- (b) (6) (mrs)- Discussed echo confirmed heart disease, DCM. Reviewed causes of DCM (unlikely taurine def, but will submit for levels) and prognosis with owner. Risk of future episodes of CHF, when is unpredictable. Owner consented to thoracocentesis. If continues to breath comfortably out of oxygen can go home this afternoon.

 12:30pm- (b) (6)- (b) (6) breathing is stable out of oxygen. Very weak and lethargic. Ate a small amount of food this morning. Discussed since breathing is comfortable, can try at home. If energy level does not improve at home over the next few days, may consider euthanasia. Discussed elevated kidney values and how that is going to make treating CHF with lasix challenging. Owner is comfortable with trying (b) (6) at home to see how she does. Will have husband call back to set up a time.

(b) (6) P (b) (6) 21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472)
 Rx #: 2569385 0 Of 12 Refills
 Give 1 tablet by mouth twice daily with food.
 (b) (6) P (b) (6) 60.00 tablet of Lasix (Salix / Furosemide) 12.5mg Tablet (M568)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	D	(b) (6)	Rx #: 2569382 0 Of 0 Refills Give 1/2 tablet by mouth twice daily. Pleural Effusion Final
(b) (6)	D		Left Atrial Enlargement Final
(b) (6)	D		Dilated Cardiomyopathy Final

(b) (6) C (b) RADIOLOGY REVIEW - FINAL (b) (6)
The DV view of the thorax obtained on (b) (6) has been reviewed and there is a moderate to large amount of pleural effusion that obscures visualization of the cardiac silhouette. There is also an area of increased opacity in the left hemithorax in the region of the caudal segment of left cranial lung lobe. The remaining lung parenchyma appears to be within normal limits and the pulmonary vessels appear normal. This combination of findings may be the result of heart failure or neoplasia and a cardiac consult is recommended for further evaluation.

This review was written by: (b) (6), DVM, DACVR, DACVS

(b) (6) C (b) CARDIAC EVALUTION - CLOSED (b) (6) - Cardiac Evaluation

Date of evaluation: Monday, (b) (6)

CHIEF COMPLAINT: pleural effusion

HISTORY: Presented to ER last night for lethargy and ADR. cursory ultrasound revealed pleural effusion. Thoracocentesis yielded 25ml yellow tinged fluid from the right side. Patient received 12mg lasix IV and was placed in oxygen overnight. Her RR was wnl, with slight effort noted overnight. She ate a small amount. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3).

PHYSICAL EXAM: The patient was bright, alert and responsive. No murmur on auscultation, but heart sound slightly muffled. The patient was eupnic, RR 30 with slight effort. Normal BV sounds, no crackles on auscultation. Pulses were fair and synchronous. Heart rate was 170 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) (b) (6): Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

ECHOCARDIOGRAM (b) (6):
IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM). Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia

THORACOCENTESIS (b) (6): a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

SUMMARY: (b) (6) has dilated cardiomyopathy (DCM) and has developed pleural effusion due to congestive heart failure. The pleural effusion was removed today and we are starting the medications as described below. Although taurine deficiency is a rare cause for dilated cardiomyopathy in cats, we submitted a taurine level to the lab today to look for this. The test results can take up to 2 weeks; we will fax results when they are available. Unfortunately the prognosis for DCM (with no taurine deficiency) in cats is poor, with survival time 3-6 weeks or in rare cases 4-6 months. A renal panel was recheck today and showed elevated BUN (74) and normal creatinine (1.4). Discussed with owner the azotemia may make treating the congestive heart failure challenging.

We discussed that with heart enlargement there is a risk for developing a blood clot, or stroke. Although aspirin +/- plavix can be given in hopes of reducing clot formation, they have not been definitively proven to prevent blood clot formation in cats. If the owners elect to start this medication I would recommend waiting until we know she is eating and feeling well on the below medications as aspirin/plavix can cause GI side effects (vomiting, inappetance) in some cats. A recheck with cardiology is recommended in 2 weeks, or sooner if clinical signs.

Advised owner congestive heart failure likely contributing to lethargy, but concern that weakness may be secondary to the DCM and low cardiac output. If that is the case, may not improve once we get her out of failure. Owner comfortable taking (b) (6) home to see how she does. (b) (6) was discharged on the below medications. If she does not improve at home over the next few days, owner to consider euthanasia.

MEDICATIONS:

START: Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

START in 3 DAYS IF EATING

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
If eating and easy to medicate, start: Taurine 250 mg by mouth twice a day with food.			

(b) (6)	L	(b) (6)	Basic Metabolic Profile, (b) (6) results from (b) (6) In-Clinic Requisition ID: 0 Posted Final <table border="0" style="width: 100%;"> <thead> <tr> <th style="width: 30%;">Test</th> <th style="width: 30%;">Result</th> <th style="width: 40%;">Reference Range</th> </tr> </thead> <tbody> <tr> <td>HCT =</td> <td>40 %</td> <td>12 - 70</td> </tr> <tr> <td>HB =</td> <td>13.1 g/dL</td> <td>9.9 - 14.9</td> </tr> <tr> <td>NA+ =</td> <td>146.3 mmol/L</td> <td>146.2 - 156.2</td> </tr> <tr> <td>K+ =</td> <td>4.99 mmol/L H</td> <td>3.41 - 4.71</td> </tr> <tr> <td>CL+ =</td> <td>107.8 mmol/L L</td> <td>117.0 - 125.3</td> </tr> <tr> <td>CA++ =</td> <td>1.17 mmol/L</td> <td>1.16 - 1.35</td> </tr> <tr> <td>MG++ =</td> <td>1.08 mmol/L H</td> <td>0.33 - 0.49</td> </tr> <tr> <td>GLU =</td> <td>156 mg/dL H</td> <td>72 - 132</td> </tr> <tr> <td>LAC =</td> <td>9.7 mg/dL H</td> <td>0.7 - 1.9</td> </tr> <tr> <td>BUN =</td> <td>67 mg/dL H</td> <td>22 - 33</td> </tr> <tr> <td>CREAT =</td> <td>5.3 mmo/L H</td> <td>1.1 - 3.5</td> </tr> <tr> <td>O2CAP =</td> <td>18.2 mL/dL</td> <td></td> </tr> <tr> <td>TCO2 =</td> <td>19.9 mmol/L</td> <td></td> </tr> <tr> <td>GAP =</td> <td>20.3 mmol/L</td> <td></td> </tr> <tr> <td>CA/MG =</td> <td>1.1 mol/mol</td> <td></td> </tr> <tr> <td>OSM =</td> <td>313.5 mOsm/kg</td> <td></td> </tr> <tr> <td>BUN/CREA =</td> <td>12.7 mg/mg</td> <td></td> </tr> </tbody> </table> Manually entered. PCV: 43% T.S: 6.6mg/dl	Test	Result	Reference Range	HCT =	40 %	12 - 70	HB =	13.1 g/dL	9.9 - 14.9	NA+ =	146.3 mmol/L	146.2 - 156.2	K+ =	4.99 mmol/L H	3.41 - 4.71	CL+ =	107.8 mmol/L L	117.0 - 125.3	CA++ =	1.17 mmol/L	1.16 - 1.35	MG++ =	1.08 mmol/L H	0.33 - 0.49	GLU =	156 mg/dL H	72 - 132	LAC =	9.7 mg/dL H	0.7 - 1.9	BUN =	67 mg/dL H	22 - 33	CREAT =	5.3 mmo/L H	1.1 - 3.5	O2CAP =	18.2 mL/dL		TCO2 =	19.9 mmol/L		GAP =	20.3 mmol/L		CA/MG =	1.1 mol/mol		OSM =	313.5 mOsm/kg		BUN/CREA =	12.7 mg/mg	
Test	Result	Reference Range																																																							
HCT =	40 %	12 - 70																																																							
HB =	13.1 g/dL	9.9 - 14.9																																																							
NA+ =	146.3 mmol/L	146.2 - 156.2																																																							
K+ =	4.99 mmol/L H	3.41 - 4.71																																																							
CL+ =	107.8 mmol/L L	117.0 - 125.3																																																							
CA++ =	1.17 mmol/L	1.16 - 1.35																																																							
MG++ =	1.08 mmol/L H	0.33 - 0.49																																																							
GLU =	156 mg/dL H	72 - 132																																																							
LAC =	9.7 mg/dL H	0.7 - 1.9																																																							
BUN =	67 mg/dL H	22 - 33																																																							
CREAT =	5.3 mmo/L H	1.1 - 3.5																																																							
O2CAP =	18.2 mL/dL																																																								
TCO2 =	19.9 mmol/L																																																								
GAP =	20.3 mmol/L																																																								
CA/MG =	1.1 mol/mol																																																								
OSM =	313.5 mOsm/kg																																																								
BUN/CREA =	12.7 mg/mg																																																								

(b) (6)	TC	(b) (6)	LAB RESULTS - NOTES - TENTATIVE (b) (6) 00:00 Lab Results: PCV: 42% TS g/dl: 6.8 Serum: Normal Original Lab Date:
---------	----	---------	---

(b) (6)	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Basic Metabolic (b) (6) Panel # 2 (b) (6) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Specialty/Referral Exam Level 3 (REF03) by (b) (6)
(b) (6)	B	(b) (6)	Echocardiogram Level 3 Group (USSC19) by (b) (6)
(b) (6)	B	(b) (6)	1.00 VRC04 Procedure (VRC04) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Thoracocentesis Therapeutic (R33) by (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	B	(b) (6)	60.00 tablet of Lasix (Salix / Furosemide) 12.5mg Tablet (M568) by (b) (6)
(b) (6)	B	(b) (6)	21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472) by (b) (6)
(b) (6)	B	(b) (6)	Oxygen Therapy (Caged)/Hour (Group) (O2CAGE) by (b) (6)
(b) (6)	B	(b) (6)	7.00 O2 Therapy Per Hour (T044) by (b) (6)
(b) (6)	B	(b) (6)	7.00 Oxygen-related Patient Care / Hour (O2CARE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
(b) (6)	B	(b) (6)	Hospitalization Hours Smart Group (HOSPIT) by (b) (6)
(b) (6)	B	(b) (6)	17.00 Hospitalization Hours- Feline (H01) by (b) (6)
(b) (6)	B	(b) (6)	1.00 In-House Nutrition Assessment Level 1 (NTR012) by (b) (6)
(b) (6)	B	(b) (6)	17.00 Critical Care Level 2- Hours (CCU2) by (b) (6)
(b) (6)	B	(b) (6)	.20 ml of DNULsix 50mg/ml/ML (T106) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Cared by (b) (6) - Cardiology (b) (6) by (b) (6)
(b) (6)	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Taurine U of Wisc (via (b) (6) S16755) (L0245) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Taurine U of Wisc (via (b) (6) S16755) (L0245) by (b) (6)
(b) (6)	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Chemistry IV Renal Panel (b) (6) (CH25) by (b) (6)
(b) (6)	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Dilution Verification Catalyst CREA (CH11DV) by (b) (6)

(b) (6) C (b) (6) EMERGENCY PHYSICAL EXAM - Closed May (b) (6)
 (b) (6)

Chief Complaint: Lethargic

History: Starting yesterday patient was noted to be lethargic and not herself. 4 other cats so difficult to say if she was eating but they think she was. Not sure about U/BM. Indoor only. Did not notice she was having issues breathing.

Other Medical Problems: None

Medications/Supplements: None

Environment: Indoor only

Vaccination Status:

Current Diet (Type):

- Frequency:

- Amount:

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Physical Examination:

S(ubjective): QAR, hydration WNL, BCS 7/9, pain score: 0/4

O(b)jective:

Weight: 15.6 pounds

TPR: [temp - 93]F, [HR - 150] bpm, [RR - 60] bpm

EENT: clear AU/OU, no nasal discharge, normal cervical palpation, mm pink, moist/CRT < 2 sec

INTEG: Hair coat ok

PLN: WNL

CV: Heart sounds muffled

RESP: Increased RE, dull lung sounds

GI: soft, nonpainful, no masses

UG: SF, WNL

M/S: amb x 4

Neuro: alert/appropriate, cranial nerves intact

Problems/Differential Diagnoses:

Dyspnea

Lethargy

Diagnostics:

Cursory ultrasound - mild to moderate amount of pleural effusion R>L

DV thoracic radiograph - cardiac silhouette difficult to visualize, pleural effusion

(b) (6) 2

Assessment:

12 yr SF DSH

1. Pleural effusion, dyspnea - r/o cardiac (HCM) vs neoplasia (lymphoma vs other)

Treatment:

12 mg Lasix IM at 10 PM

Place in O2 cage

Thoracocentesis - 25 mL clear to yellow fluid removed from the right side

Place IVC, 12 mg Lasix IV at 2 AM

Plan/Recommendations:

Discussed differentials for pleural effusion - cardiac vs neoplasia. Due to pleural effusion cannot tell on radiographs if this is cardiac over neoplastic. Rec thoracocentesis to make (b) (6) breathe more comfortably - o consents. Rec echocardiogram in the morning to see if this is heart disease. If this is CHF spoke about disease process and prognosis. If this is neoplasia owner's may decide to

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6) Phone: (b) (6) Address: (b) (6) (b) (6)	Patient: (b) (6) Species: Feline Age: 12 Yrs. 5 Mos. Color: Black	Breed: Shorthair, Domestic Sex: Spayed Female
---	--	--

Date	Type	Staff	History
------	------	-------	---------

stop. Discussed possibility of taking repeat radiographs tomorrow if needed.

(b) (6) R (b) (6) Tx Template (blank)- Old WTS - TENTATIVE
WARD TREATMENT SHEET

DATE: (b) (6)		WARD: ccu	
LAST NAME: (b) (6)		CLINICIAN: (b) (6)	(b) (6)
PATIENT NAME: (b) (6)		TRANSFER DOCTOR:	
BREED: Shorthair, Domestic	COLOR: Black	LEGEND: O = scheduled X = performed D/C = discontinued inc = increase dec = decrease ⊖ = not given	
AGE: 12 Yrs. 4 Mos.	SEX: Spayed Female		
WEIGHT: 15.6 pounds as of: 12/28/2009			
PROBLEM/WORKING DIAGNOSIS:		SURGERY:	
Pleural effusion		IV CATHETER:	
ALERT: increased RR		CODE: RED	

Technician STAFF ID ==>			8	9	10	11	N	1	2	3	4	5	6	7	8	9	10	11	M	1	2	3	4	5	6	7
DR Staff ID	TREATMENTS	Call Parameters																								
	Temperature	<99, >102.5																								
	HR/MM/CRT	<140, >240																								
	RR/RE	>40, increased RE	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O
	Pulse Quality	wk																								
	Assess Pain & Note Score	2																								
	WEIGH																									
	RER: 262 kcal/day Diet: 115 g EN BID - any -				O											O										
	Quantify Food Intake							O									O									
	Lasix 12 mg IV - ask		O																			O				
	O2 cage																									

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date Type Staff History

	AM SHIFT
	MID SHIFT
	OVER

TIME	Init. TECH	Temp	HR	Pulse qual.	RR/effort	MM color	CRT	PCV/ts	BG	Turn	Blood Pressure	Pain Score	BM	Urine	Vomit	Grams Consumed	Init. AA
8 a																	
9 a																	
10 a																	
11a																	
N																	
1 p																	
2 p																	
3 p																	
4 p																	
5 p																	
6 p																	
7 p																	
8 p																	
9 p																	
10 p																	
11 p																	
M																	
1 a																	
2 a																	
3 a																	
4 a																	
5 a																	
6 a																	
7 a																	

TIME	ADDITIONAL COMMENTS

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	B	(b) (6)	3.00 O2 Therapy Per Hour (T044) by (b) (6)
(b) (6)	B	(b) (6)	3.00 Oxygen-related Patient Care / Hour (O2CARE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
(b) (6)	B	(b) (6)	IV Catheter with Injection Cap (IVCATCP) by (b) (6)
(b) (6)	B	(b) (6)	1.00 IV Catheter Placement (CATH) by (b) (6)
(b) (6)	B	(b) (6)	1.00 each of Tx Catheter IV 20g x 2" Surflo (PINK) (H0112) by (b) (6)
(b) (6)	B	(b) (6)	1.00 each of Tx IV Ext T Set Hospira 1265028 (H027) by (b) (6)
(b) (6)	B	(b) (6)	1.00 each of Tx Injection Cap/Plug Termo 007110 (H118) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)
(b) (6)	B	(b) (6)	Thorax Radiographic Study Group (RADTH) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Radiograph Preparation (XFEE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 One view rdgh stdy (RAD1V) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Radiologist Review Fee (RADGN) by (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates,
 I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended,
 R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
6/7/2016	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE 6/7/2016 12:33 On the phone with client discussing (b) (6) also discussed (b) (6) T4 and liver values need to be rechecked 6 weeks after his meds began to test for any needed dose adjustments. We can come to the home or he can schedule with a (b) (6) in the hospital. We discussed that he's enjoyed working with (b) (6) before.</p>
6/6/2016	C	(b) (6)	<p>MEDICAL COMMENTS 6/6/2016 11:47 FDA complaint submitted: Pet Food Safety Report, ID 54405, was successfully submitted on 6/6/2016 11:44:41 AM EST to the FDA, and it was issued an Individual Case Safety Report Number (ICSR) of 1053339</p>
6/1/2016	TC	(b) (6)	<p>MEDICAL COMMENTS - TENTATIVE 6/1/2016 10:46 SW (b) (6) at Merrick - updated company that we have documented taurine deficiency in 2 other cats in house hold. The quality assurance team indicated that the level of taurine in the lot # I gave them was sufficient - discussed that this likely takes 3-6 month to develop and likely to be related to earlier lot and they need to investigate further. Asked if the level they gave me was from them retesting the food after I called or from an earlier test - not sure. Asked for information from the taurine tests - told I can send summary of lab results. Also indicated that I am reporting this to the FDA. She give new info to her manager and quality assurance. Told them I expect them to follow up with me. Below email sent to Merrick:</p> <p>Taurine Levels (b) (6) To: (b) (6)@merrickpetcare.com Hi (b) (6)</p> <p>Thank you for your help with these cases. Here is the summary of the lab results:</p> <p>12yr female spayed domestic short hair diagnosed and clinical for dilated cardiomyopathy -5/9/2016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016</p> <p>5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016

-9yr male neutered domestic long hair: 196 nmol/ml

-8y female spayed domestic short hair: 368 nmol/ml

-9yr male neutered domestic long hair: 124 nmol/ml

-9yr male neutered domestic long hair: 536 nmol/ml

Please let me know if you have any other questions.

Sincerely,

(b) (6)

(b) (6), DVM Diplomate ACVN
Clinical Nutrition Department

(b) (6)

5/31/2016 C

(b) (6)

COMMUNICATIONS WITH CLIENT - Closed Jun 02/2016

5/31/2016 16:14

Spoke with husband, he confirmed email was received and (b) (6) and (b) (6) have begun taurine supplementation. Discussed results are a different normal range for the whole blood testing I did vs the plasma testing the cardiologist did on (b) (6). He notes (b) (6) had been getting some Fancy Feast so that likely explains why his values are top of the normal range. Discussed (b) (6) elevated globulins, need protein electrophoresis to better define the issue, can be chronic inflammatory response or a few types of cancer. If cancer, because they are (b) (6) clients they can consult with oncology at no charge to hear what the treatment options would be. They can bring (b) (6) into the office with a (b) (6) doctor for the bloodwork or I can do the bloodwork at the house. If I am doing it, I would not charge any recheck exam, just the travel and diagnostic test costs. If a (b) (6) in the hospital, they shouldn't have to charge for an exam either because she was just checked in late May. Owner likes (b) (6); advised he could schedule that with her. Discussed rechecking (b) (6) T4 and liver values 6 weeks after starting meds to check if dose needs adjusting; can be done at the house or in the office as well. Advised Nutrition is contacting Merrick about the taurine results and she feels the owner shouldn't have to pay for the taurine testing; she wants Merrick to have to pay for it directly. So I

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
5/31/2016	C	(b) (6)	<p>told the owner that first and foremost, they are responsible for payment of the testing to (b) (6) and once we advise them of a charge, they would be required to pay it. If the Nutritionist is able to circumvent that by having Merrick pay us directly, that would be a nice advantage for the client. He understands.</p>
5/31/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed Jun 02/2016 5/31/2016 15:39 LMOM on husband cell making sure they received my treatment advice in the email from over the weekend. Please call back or reply to email so I can be certain the treatment guidelines were received.</p>
5/28/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 30/2016 5/28/2016 15:46 Email to client: Hi Mr and Mrs (b) (6) –</p> <p>I left a (long-winded) message on (b) (6) voicemail earlier today. The nutritionist has since been in contact with me and advised that both (b) (6) and (b) (6) should be started on taurine supplementation. She recommends 250mg taurine twice daily for 2-3 weeks. Because you've already switched them to another diet, after 2-3 weeks, supplementation can be discontinued. (b) (6) and (b) (6) tested safely within the normal range for taurine, so they do not require any supplementation.</p> <p>I presume since you have already been treating (b) (6), you likely have a supply of taurine supplement. If not, feel free to contact me (b) (6) or the nutritionist or cardiologist to get a larger supply in order to treat the brothers.</p> <p>The nutritionist also advised she'll be contacting Merrick again now that the data has been received. Once she has heard more from them, she'll be in contact with you, as well.</p> <p>I also mentioned in the voicemail that (b) (6) blood test was repeated and verified that she does have elevated globulins. The most harmless reason would be chronic inflammation, but since she's been otherwise healthy, it is valuable to pursue further diagnostic inquiry. Unfortunately, elevated globulins can also indicate cancer, so we want to determine precisely what is happening with her. We can collect another blood sample from her at any time in order to perform a test called protein electrophoresis which further defines which specific immunoglobulins are elevated. You may choose to bring her into the office or have us out to the home again. (b) (6) will need repeat bloodwork after he's been on his thyroid supplement for 6 weeks, we could collect her second sample at that time as well, if</p>

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
5/28/2016	C	(b) (6)	<p>you choose.</p> <p>Feel free to contact me with any questions. I will next be in the office on Tuesday May 31st. Happy Memorial Day weekend – (b) (6)</p> <hr/> <p>COMMUNICATIONS WITH DOCTOR - Closed May 30/2016 (b) (6) 15:45 Email response from (b) (6) to (b) (6): I'm also going to talk with (b) (6) this week about the cost of the taurine test. In my opinion this should be paid for by the company. I don't want the owner to pay the cost yet until I talk with (b) (6) and the company again.</p> <p>(b) (6)</p> <hr/> <p>From: (b) (6) Sent: Saturday, May 28, 2016 2:16 PM To: (b) (6) Subject: RE: price for taurine test There is a risk of deficiency with anything <200, so that's why I would go ahead and supplement both cats...and it's harmless:-)</p> <hr/> <p>From: (b) (6) Sent: Saturday, May 28, 2016 2:13 PM To: (b) (6) Subject: RE: price for taurine test It is really interesting...probably the same reason some puppies raised on an unbalanced home cooked diet never have issues and other do.</p> <p>Great the diet has been changed. We should get the cats that tested low on some supplementation for 2-3 weeks just to cover our bases. 250mg taurine PO BID...if she needs to use a powder form and mix with the cats food that's fine</p> <p>Let the owner know I will touch base with the company after Memorial day...I have not heard back from them yet. This will also give me much more to go on when reporting to the FDA...who know this might turn into a pet food recall (it should turn into a recall)!</p> <p>Thank you so much for the update!</p> <p>(b) (6)</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

From: (b) (6)
 Sent: Saturday, May 28, 2016 1:55 PM
 To: (b) (6)
 Subject: RE: price for taurine test
 Hi (b) (6) –
 Thanks for providing this justification for the lab decision; I really appreciate it! I left you a voicemail earlier today – the results are in. 2 cats tested within the normal range [(b) (6) 368, (b) (6) 536 (300-600)]. (b) (6) was 196 and (b) (6) was way down at 124. All 4 cats were switched to Royal Canin food about 7 days ago. I left a voicemail for the client advising of the results, but told him I wanted your input before devising a treatment strategy. I would think of these 4, only (b) (6) would benefit substantially from taurine supplementation. I presume (b) (6) levels are sufficient now that he's been put on a properly formulated diet. Do you agree? This case is so interesting... how the cats fall all along the clinical spectrum, including some that have sufficient taurine, despite all eating the same presumably flawed diet.
 Thanks for your input,
 (b) (6)

5/28/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 30/2016 5/28/2016 13:44 LMOM - advised client taurine results have been received, I have notified nutrition dept who will weigh in on treatment decision-making. Normal range is 300-600 and (b) (6) and (b) (6) tested within that range. Clinical signs are unlikely above 200, (b) (6) was 196, so it is likely he wouldn't show any issues. (b) (6) tested at 124 so he might be the one to benefit from additional supplementation, aside from just the diet change to the Royal Canin food. We will wait to initiate any therapy until the nutritionist has a chance to comment; we are working as a team on this. Since we have results, we likely have an invoice from the lab as well, so we should be able to advise of the cost of this testing in the short-term. I had spoken with his wife about (b) (6) having elevated globulins and on the re-test that status persists, was verified. Recommend additional blood testing for further work-up, could be collected when we visit (b) (6) for bloodwork 6 weeks after starting his thyroid meds. Please call back to discuss these results.</p>
-----------	---	---------	---

5/27/2016	TC	(b) (6)	<p>LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:32 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)</p>
-----------	----	---------	--

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)	Patient: (b) (6)	
Phone: (b) (6)	Species: Feline	Breed: Longhair, Domestic
Address: (b) (6)	Age: 9 Yrs. 10 Mos.	Sex: Neutered Male
(b) (6)	Color: Calico	

Date	Type	Staff	History
5/24/2016	P	(b) (6)	100.00 tablet of Tapazole (Methimazole) 5mg Tablet (M066) Rx #: 2576298 0 Of 0 Refills Give 1 tablet by mouth twice daily. Check bloodwork for dose adjustment 6 weeks after starting medication.
5/24/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:41 Spoke with Mrs; advised hyperthyroid with some liver elevations. Reviewed life-long treatment, bloodwork 6 weeks after med started and then twice yearly if stable. If dose is changed after first bloodwork, we repeat bloodwork again 6 weeks later until properly regulated. Med can be tablet, liquid or transdermal. Owner wants to crush tablet into canned food; advised this is fine as long as we're certain he's the only one who might consume the medicated food within their group-housing situation. Owner feels she can guarantee that. Meds will be at 197 pharmacy. Taurine pending, will call. Advised final pricing on taurine at CA lab not yet determined, will be in touch with that info as soon as finalized. Owner asked why use a diff lab; advised nutritionists recommended this lab, specialized testing at university, two labs finding low levels strengthens case against food company.
5/24/2016	B	(b) (6)	100.00 tablet of Tapazole (Methimazole) 5mg Tablet (M066) by (b) (6)
5/24/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)
5/21/2016	C	(b) (6)	GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Weight loss/Taurine check History: Owner notes chronic weight loss across the recent months. Was losing hair for over a year, but was told it was related to anxiety. Eats with voracious appetite. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; he is one of 4 cats that live together in a room above the garage. Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household. Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food Meds: none S: BAR, PCS 0/4 - slightly feisty O: MM / ORPH: Pink, moist, crt <2 sec, mild tartar E/E: mild black debris in outer cartilages of left ear, deep canal WNL, right ear WNL. ophtho WNL.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

INT: alopecia caudal dorsum, ventrum, lateral thighs. no ectoparasites observed.
 PLN: WNL
 CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic
 GI/UG: Compliant, no masses
 MS/NS: Normal amb x4
 BCS: 1.5-2/5 4.4kg
 A: 9yr9mo MN DLH
 1) weight loss - r/o hyperthyroid, diabetes mellitus, organ dz (kidney, liver), other endocrinopathy, neoplasia, nutritional problem
 2) alopecia - r/o FAD, other derm issue, psychogenic (stress, pain-related)
 3) dental disease
 4) otic debris - r/o infection vs inadequate grooming
 P: PE
 Taurine level
 CBC/Superchem/T4

Advised client of marked weight loss from last documented weight. Systemic bloodwork may illuminate the reason; will call with results next week. Taurine level will take 7-10 days.

Advised client we are sending taurine test to a different lab than the one that tested (b) (6) sample, at the advice of the nutrition service. We do not have a price in our computer system for this test through this lab, so the client will be invoiced for the taurine level (for all 4 cats) once that is established. Client paid today's services during the visit and is aware of the pending charge; advised the (b) (6) charge was \$214 and the charge at the other lab will likely be within \$50 under/over that fee. He commits to paying taurine test fees once advised of final fee. Stated we want to submit samples for testing ASAP and he understands fee structure will not be set until after tests are underway.

5/21/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 23/2016 5/21/2016 16:25 mailed welcome card, magnet, Rabies certificates (b) (6), (b) (6) and feedback postcard
-----------	---	---------	---

5/21/2016	V	(b) (6)	May 21, 2016 11:21 AM Staff: (b) (6) ----- Weight : 4.40 kilograms HC-RS scale
-----------	---	---------	---

5/21/2016	L	(b) (6)	Hematology results from (b) (6) (East) Requisition ID: 209396 Posted Final
-----------	---	---------	---

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

Test	Result	Reference Range
HCT	42 %	29 - 48
HGB	15.0 g/dL	9.3 - 15.9
MCHC	35.7 g/dL	30 - 38
WBC	14.2 10 ³ /uL	3.5 - 16.0
Bands	0 %	0 - 3
RBC	9.5 10 ⁶ /uL	5.92 - 9.93
MCV	44 fL	37 - 61
MCH	15.8 pg	11 - 21
ABS BASO	0 /uL	0 - 150
ABS NEUTB	0 /uL	0 - 150
Platelet C	254 10 ³ /uL	200 - 500
Platelet E	ADEQUATE	ADEQUATE -
Neutrophil	53 %	35 - 75
Lymphocyte	41 %	20 - 45
Monocytes	2 %	1 - 4
Eosinophil	4 %	2 - 12
Basophils	0 %	0 - 1
Absolute N	7526 /uL	2500 - 8500
Absolute L	5822 /uL	1200 - 8000
Absolute M	284 /uL	0 - 600
Absolute E	568 /uL	0 - 1000

Ascn: (b) (6) Profile: Complete Blood Count

5/21/2016 L

Chemistry results from (b) (6) (East) Requisition

ID: 209396	Posted	Final	Reference Range
Test	Result		Reference Range
ALB	3.4 g/dL		2.5 - 3.9
ALKP	174 U/L H		6 - 102
ALT	243 U/L H		10 - 100
AMYL	882 U/L		100 - 1200
AST	46 U/L		10 - 100
BUN/UREA	17 mg/dL		14 - 36
Ca	9.2 mg/dL		8.2 - 10.8
Chloride	111 mEq/L		104 - 128
CHOL	181 mg/dL		75 - 220
CK	124 U/L		56 - 529
CREA	0.6 mg/dL		0.6 - 2.4
GGT	3 U/L		1 - 10
GLU	80 mg/dL		64 - 170
Mg	1.7 mEq/L		1.5 - 2.5
PHOS	5.9 mg/dL		2.4 - 8.2
Potassium	4.5 mEq/L		3.4 - 5.6
Sodium	150 mEq/L		145 - 158
TBIL	0.1 mg/dL		0.1 - 0.4
TP	5.9 g/dL		5.2 - 8.8
TRIG	58 mg/dL		25 - 160
GLOB	2.5 g/dL		2.3 - 5.3

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates,
 I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended,
 R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

A/G Ratio 1.4 Ratio 0.35 - 1.5
B/C Ratio 28 Ratio 4 - 33
Na/K Ratio 33

5/21/2016 L

Endocrinology results from (b) (6)
 (East) Requisition ID: 209396 Posted Final

Test	Result	Reference Range
T4	20.2 ug/dL H	0.8 - 4.0

 Ascn: (b) (6) Profile: Total T4

 Result verified.

5/21/2016 L

Miscellaneous results from (b) (6) ics
 (East) Requisition ID: 209396 Posted Final
 Ascn: (b) (6) Profile: Superchem
 RE: 1045 PrecisionP 28 U/L 8 - 26
 PrecisionPSL elevations correlate closely with abnormal PLI concentrations. In cats with appropriate clinical signs, this PrecisionPSL is supportive of, but not definitive, for a diagnosis of pancreatitis. In cats without clinical signs of pancreatitis, a mild elevation is an insignificant finding.
 RE: 11067 Comment
 Hemolysis 1+ No significant interference.

5/21/2016 B
 5/21/2016 B
 5/21/2016 B
 5/21/2016 B
 5/21/2016 B
 5/21/2016 B
 5/21/2016 B
 5/21/2016 B

(b) (6) 1.00 Superchem Cbc T4 (b) (6) Sa120 (L85) by (b) (6)
 (b) (6) 1.00 House Call Travel Level 2 (HC06) by (b) (6)
 (b) (6) 1.00 At Home Appointment (HC04) by (b) (6)
 (b) (6) Laboratory Request / Sample Handling (LABS) by (b) (6)
 (b) (6) 1.00 Outside Lab (XTBALUO) by (b) (6)
 (b) (6) 1.00 Sample Handling & Disposal (LFEE) by (b) (6)
 (b) (6) 1.00 Lab Sample Label (TL) by (b) (6)
 (b) (6) 1.00 Cared for by (b) (6) ((b) (6)) by (b) (6)

5/20/2016 C

(b) (6) COMMUNICATIONS WITH CLIENT - Closed May 21/2016
 5/20/2016 15:00
 Called to confirm tomorrow's appointment from (b) (6), (b) (6) (b) (6) and (b) (6) at 9 am. I also mentioned in my message that we should use the address (b) (6) (b) (6) in the GPS. If any questions please call (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

5/17/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 19/2016 5/17/2016 12:29 Responding to owner's message, booked (b) (6), (b) (6), (b) (6), (b) (6) for House Call on Saturday (b) (6). (b) (6) was seen on emergency and diagnosed with low taurine, so all cats need to be screened. Had been eating Merrick dry food. Cats are kept in a finished room above the garage; he thinks they won't need to be confined/isolated more than that in order to work on them. Discussed senior bloodwork as well. He notes this emergency with (b) (6) was a wake-up call and he'd like to thoroughly have everyone checked out. (b) (6) is losing weight. (b) (6) and (b) (6) have been to another vet and have a current Rabies; (b) (6) and (b) (6) haven't been to the vet in a long time. Discussed PureVax 1yr vs 3yr vs standard RabVac, vaccine-associated sarcoma issue - owner wants the purified vaccine, prefers the one year since they should be examined annually anyway. Advised if (b) (6) status progresses and we need to be checking her as well, please call to inform us in case we need special items/supplies for her care. Owner notes we should use (b) (6) with the GPS; his home address was renamed/renumbered a few years ago, but GPS cannot often find (b) (6)</p>
-----------	---	---------	---

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE</p> <p>(b) (6) 12:25</p> <p>Spoke with owner, advised of specialist's comments about protein electrophoresis. Can conduct further testing through our Internal Medicine service of ultrasounds of chest and abdomen, essentially searching for origin of chronic inflammation. It is possible that testing will come back normal, despite the bloodwork indicating the elevated globulins. Another option is to track her body weight at home once monthly and report any weight loss promptly; if none noted, recheck bloodwork in 3 months to see if globulins are resolved. Specific cancers cause specific spikes and her results do not show those elevations, so that is good news. But we don't have a definite reason for her chronic inflammation. Owner understands and will relate to wife.</p>
---------	----	---------	---

(b) (6)	6 TC	(b) (6)	<p>COMMUNICATIONS WITH DOCTOR - TENTATIVE</p> <p>(b) (6) 12:11</p> <p>Spoke with (b) (6) at (b) (6); she notes monoclonal globulin spikes are most concerning because they define lymphoma, myeloma. Polyclonal spikes can be FIP, but also any cause of chronic inflammation. There is a chance chronic inflammation can be associated with neoplastic process though. She notes the degree of elevation is mild. If the owners want to work this up aggressively, full body imaging, best with ultrasound, to search for cancer. If they would like to monitor, recheck chemistry panel in 3 months and assess globulin count at that time. Only repeat electrophoresis if significantly higher elevation. If pet is losing weight, neoplasia moves up the list of differentials. Advised chronic otitis externa in this otherwise healthy patient; doctor concurs that wouldn't be sufficient to cause systemic response.</p>
---------	------	---------	---

(b) (6)	L	<p>Chemistry results from (b) (6) (East) Requisition</p> <p>ID: 209396</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Test</th> <th>Result</th> <th>Final</th> <th>Reference Range</th> </tr> </thead> <tbody> <tr> <td>ALB</td> <td>2.9 g/dL</td> <td></td> <td>2.5 - 3.9</td> </tr> <tr> <td>TP</td> <td>8.2 g/dL</td> <td></td> <td>5.2 - 8.8</td> </tr> <tr> <td>GLOB</td> <td>5.3 g/dL</td> <td></td> <td>2.3 - 5.3</td> </tr> <tr> <td>ALPHA 1</td> <td>0.3 g/dL</td> <td></td> <td>0.2 - 1.1</td> </tr> <tr> <td>ALPHA 2</td> <td>0.7 g/dL</td> <td></td> <td>0.4 - 0.9</td> </tr> <tr> <td>BETA</td> <td>0.6 g/dL</td> <td></td> <td>0.3 - 0.9</td> </tr> <tr> <td>GAMMA</td> <td>3.6 g/dL H</td> <td></td> <td>0.3 - 2.5</td> </tr> </tbody> </table>		Test	Result	Final	Reference Range	ALB	2.9 g/dL		2.5 - 3.9	TP	8.2 g/dL		5.2 - 8.8	GLOB	5.3 g/dL		2.3 - 5.3	ALPHA 1	0.3 g/dL		0.2 - 1.1	ALPHA 2	0.7 g/dL		0.4 - 0.9	BETA	0.6 g/dL		0.3 - 0.9	GAMMA	3.6 g/dL H		0.3 - 2.5
Test	Result	Final	Reference Range																																
ALB	2.9 g/dL		2.5 - 3.9																																
TP	8.2 g/dL		5.2 - 8.8																																
GLOB	5.3 g/dL		2.3 - 5.3																																
ALPHA 1	0.3 g/dL		0.2 - 1.1																																
ALPHA 2	0.7 g/dL		0.4 - 0.9																																
BETA	0.6 g/dL		0.3 - 0.9																																
GAMMA	3.6 g/dL H		0.3 - 2.5																																

6/2/2016	L	<p>Miscellaneous results from (b) (6)</p>
----------	---	--

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(East) Requisition ID: (b) (6) **Posted** **Final**
Ascn: (b) (6) **Profile:** Protein Electrophoresis, Serum
RE: 1140 Interpreta
 The gamma globulin fraction is elevated, characterized by a broad polyclonal band, resulting from a mixture of increased immunoglobulins associated with an immune response. Potential causes include suppurative disease, chronic infectious disease (bacterial; protozoal; viral; rickettsial; fungal), connective tissue disease, chronic granulomatous disease, etc. Correlate with clinical findings.
PATHOLOGIST:

(b) (6), BVSc (Hons 1), DACVP
 (b) (6)
 (b) (6)

Due to difference in method of analysis, there may be slight differences in the quantitative albumin and calculated globulin results between serum electrophoresis results compared to a generalchemistry panel.

(b) (6)	C	(b) (6)	MEDICAL COMMENTS - Closed (b) (6) (b) (6) 18:35 Drew sample for protein electrophoresis while at the home for EOL care for (b) (6).
---------	---	---------	---

(b) (6)	B	(b) (6)	1.00 House Call Travel Level 2 (HC06) by (b) (6)
(b) (6)	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Protein Electrophor. Serum (b) (6) T240 (L018) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
(b) (6)	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)

(b) (6)	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed (b) (6) (b) (6) 16:21 (See full phone call under (b) (6) record)
---------	---	---------	--

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
<p>Discussed (b) (6) elevated globulins, need protein electrophoresis to better define the issue, can be chronic inflammatory response or a few types of cancer. If cancer, because they are (b) (6) clients they can consult with oncology at no charge to hear what the treatment options would be. They can bring (b) (6) into the office with a (b) (6) doctor for the bloodwork or I can do the bloodwork at the house. If I am doing it, I would not charge any recheck exam, just the travel and diagnostic test costs. If a (b) (6) in the hospital, they shouldn't have to charge for an exam either because she was just checked in late May. Owner likes (b) (6); advised he could schedule that with her.</p>			
(b) (6)	TC	(b) (6)	<p>LAB RESULTS - NOTES - TENTATIVE (b) (6) 15:36 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)</p>
(b) (6)	C	(b) (6)	<p>COMMUNICATIONS WITH DOCTOR - Closed (b) (6) (b) (6) 16:48 Spoke with doctor at (b) (6) consult line - she opted to rerun the full chemistry profile to validate the results since (b) (6) remaining profile is so normal. If globulins are truly elevated, protein electrophoresis is the next step. Ddx: myeloma, lymphoma, FIP, other neoplasia, chronic inflammatory condition. Asked specifically about taurine based on (b) (6) and current investigation into whole household's taurine status; not aware of any relationship between globulins and taurine.</p>
(b) (6)	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed (b) (6) (b) (6) 16:46 Spoke with Mrs; (b) (6) has elevated globulins which can indicate cancer or a chronic inflammatory condition. Spoke with specialist and no correlation with taurine deficiency. Lab is going to re-run her full profile to validate the results. Expect an update in 1-2 days. If verified, we may need to collect additional blood for the next level of testing which tells us which specific pattern of globulins is elevated. Taurine pending, will call.</p>
(b) (6)	C	(b) (6)	<p>GP PHYSICAL EXAM - Closed (b) (6) Date Presented: (b) (6) Chief Complaint: Wellness/Taurine check History: Doing well. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; she is one of 4 cats that live together in a room above the</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

garage. Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household.
 Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food
 Meds: none
 S: BAR, PCS 0/4
 O:
 MM / ORPH: Pink, moist, crt <2 sec, small suspect FORL right upper PM3, mild tartar overall.
 E/E: copious black debris AU, mildly pruritic while cleaning. ophtho WNL.
 INT: WNL; no evidence of ectoparasites observed
 PLN: WNL
 CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic
 GI/UG: Compliant, no masses
 MS/NS: Normal amb x4
 BCS: 3/5 4.15kg
 A: 8yr FS DSH
 1) otitis externa - r/o bacterial/fungal vs ear mites
 2) dental disease
 P: PE
 Taurine level
 CBC/Vetscreen
 Disp Tresaderm 7.5ml - apply 2-3 drops in each ear twice daily for 7-10 days, keep in fridge
 ear cleaning
 PureVax Rabies 1yr SQ right hind (lot#17390B, exp 12/11/2016)

Discussed ear infection and treatment. Will call with lab results; systemic early next week, taurine in 7-10 days.

5/21/2016	I	(b) (6)	An animal is not considered immunized for at least 28 days after the initial or primary vaccination is administered. For this reason, pets receiving their first rabies vaccine should not be left outdoors unattended.								
5/21/2016	P	(b) (6)	1.00 bottle of Tresaderm 7.5ml (Merial] (M225) Rx #: 2574865 0 Of 0 Refills Apply 2-3 drops in each ear twice daily for 7-10 days.								
5/21/2016	V	(b)	May 21, 2016 11:15 AM Staff: (b) ----- Weight : 4.15 kilograms HC-RS scale								
5/21/2016	L		Hematology results from (b) (6) (East) Requisition <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">ID: 209396</th> <th style="text-align: left;">Posted</th> <th style="text-align: left;">Final</th> <th style="text-align: left;">Reference Range</th> </tr> <tr> <th style="text-align: left;">Test</th> <th style="text-align: left;">Result</th> <th></th> <th></th> </tr> </thead> </table>	ID: 209396	Posted	Final	Reference Range	Test	Result		
ID: 209396	Posted	Final	Reference Range								
Test	Result										

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			HCT 31 % 29 - 48 HGB 11.0 g/dL 9.3 - 15.9 MCHC 35.5 g/dL 30 - 38 WBC 13.4 10 ³ /uL 3.5 - 16.0 Bands 0 % 0 - 3 RBC 6.7 10 ⁶ /uL 5.92 - 9.93 MCV 46 fL 37 - 61 MCH 16.4 pg 11 - 21 ABS BASO 0 /uL 0 - 150 ABS NEUTB 0 /uL 0 - 150 Platelet C 375 10 ³ /uL 200 - 500 Platelet E ADEQUATE ADEQUATE - Neutrophil 55 % 35 - 75 Lymphocyte 37 % 20 - 45 Monocytes 2 % 1 - 4 Eosinophil 6 % 2 - 12 Basophils 0 % 0 - 1 Absolute N 7370 /uL 2500 - 8500 Absolute L 4958 /uL 1200 - 8000 Absolute M 268 /uL 0 - 600 Absolute E 804 /uL 0 - 1000 Ascn: (b) (6) Profile: Complete Blood Count

5/21/2016 L

Chemistry results from (b) (6) (East) Requisition

ID: 209396	Posted	Final	Reference Range
Test	Result		
ALB	2.6 g/dL		2.5 - 3.9
ALKP	18 U/L		6 - 102
ALT	14 U/L		10 - 100
AST	11 U/L		10 - 100
BUN/UREA	29 mg/dL		14 - 36
Ca	8.8 mg/dL		8.2 - 10.8
Chloride	113 mEq/L		104 - 128
CHOL	94 mg/dL		75 - 220
CK	62 U/L		56 - 529
CREA	1.2 mg/dL		0.6 - 2.4
GLU	79 mg/dL		64 - 170
PHOS	5.7 mg/dL		2.4 - 8.2
Potassium	4.8 mEq/L		3.4 - 5.6
Sodium	148 mEq/L		145 - 158
TBIL	0.1 mg/dL		0.1 - 0.4
TP	9.1 g/dL H		5.2 - 8.8
GLOB	6.5 g/dL H		2.3 - 5.3
A/G Ratio	0.4 Ratio		0.35 - 1.5
B/C Ratio	24 Ratio		4 - 33
Na/K Ratio	31		

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/21/2016	L		Miscellaneous results from (b) (4) (East) Requisition ID: 209396 Posted Final Ascn: (b) (6) Profile: Vet Screen RE: 11067 Comment Hemolysis 1+ No significant interference.
5/21/2016	B	(b) (6)	1.00 At Home Additional Pet Appointment (HC03) by (b) (6)
5/21/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Vetscreen Cbc Antec SA030 (L00030) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Fel Vax Rabies 1 Year Purevax (Merial) (V21) by (b) (6)
5/21/2016	B	(b) (6)	1.00 bottle of Tresaderm 7.5ml (Merial] (M225) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Cared for by (b) (6) by (b) (6)

(b) (6) C (b) (6) **COMMUNICATIONS WITH CLIENT - Closed** (b) (6)
 (b) (6) 15:04
 Called to confirm tomorrow's appointment from (b) (6), (b) (6), (b) (6) and (b) (6) at 9 am. I also mentioned in my message that we should use the address (b) (6) (b) (6) in the GPS. If any questions please call (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
6/1/2016	TC	(b) (6)	<p>MEDICAL COMMENTS - TENTATIVE</p> <p>(b) (6) 10:46 SW (b) (6) at Merrick - updated company that we have documented taurine deficiency in 2 other cats in house hold. The quality assurance team indicated that the level of taurine in the lot # I gave them was sufficient - discussed that this likely takes 3-6 month to develop and likely to be related to earlier lot and they need to investigate further. Asked if the level they gave me was from them retesting the food after I called or from an earlier test - not sure. Asked for information from the taurine tests - told I can send summary of lab results. Also indicated that I am reporting this to the FDA. She give new info to her manager and quality assurance. Told them I expect them to follow up with me. Below email sent to Merrick:</p> <p>Taurine Levels (b) (6) To: (b) (6)@merrickpetcare.com Hi (b) (6),</p> <p>Thank you for your help with these cases. Here is the summary of the lab results:</p> <p>12yr female spayed domestic short hair diagnosed and clinical for dilated cardiomyopathy -5/9/2016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016</p> <p>5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016 -9yr male neutered domestic long hair: 196 nmol/ml -8y female spayed domestic short hair: 368 nmol/ml -9yr male neutered domestic long hair: 124 nmol/ml -9yr male neutered domestic long hair: 536 nmol/ml</p> <p>Please let me know if you have any other questions.</p> <p>Sincerely, (b) (6) (b) (6), DVM Diplomate ACVN Clinical Nutrition Department</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
			(b) (6)
5/27/2016	TC	(b) (6)	LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:34 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)
5/24/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:50 Spoke with Mrs; systemic blood results WNL for (b) (6). Taurine pending.
5/21/2016	C	(b) (6)	GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Wellness/Taurine check History: Doing well. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; he is one of 4 cats that live together in a room above the garage. Indoor only (b) (6) has DCM from taurine deficiency, checking status of other cats in household. Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food Meds: none S: BAR, PCS 0/4 O: MM / ORPH: Pink, moist, crt <2 sec, moderate tartar overall E/E: ophtho/otoscopic exams WNL INT: no evidence of ectoparasites observed. matted hair present. PLN: WNL CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic GI/UG: Compliant, no masses MS/NS: Normal amb x4 BCS: 4/5 9.5kg A: 9yr9mo MN DLH 1) overweight 2) dental disease

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

3) matted hair
 P: PE
 Taurine level
 CBC/Vetscreen

Recommend weight loss, frequent brushing +/- clippers to remove mats or using a groomer. Dental condition warrants treatment. Will call with systemic blood results early next week, taurine level in 7-10 days.

5/21/2016 V (b) (6) May 21, 2016 11:21 AM Staff: (b) (6)

 Weight : 9.50 kilograms
 HC-RS scale

5/21/2016 L (b) (6) **Hematology results from (b) (6) (East) Requisition**
ID: 209396 Posted Final

Test	Result	Reference Range
HCT	40 %	29 - 48
HGB	12.3 g/dL	9.3 - 15.9
MCHC	30.8 g/dL	30 - 38
WBC	11.6 10 ³ /uL	3.5 - 16.0
Bands	0 %	0 - 3
RBC	7.9 10 ⁶ /uL	5.92 - 9.93
MCV	51 fL	37 - 61
MCH	15.6 pg	11 - 21
ABS BASO	0 /uL	0 - 150
ABS NEUTB	0 /uL	0 - 150
Platelet C	188 10 ³ /uL L	200 - 500
Platelet E	ADEQUATE	ADEQUATE -
Neutrophil	72 %	35 - 75
Lymphocyte	21 %	20 - 45
Monocytes	3 %	1 - 4
Eosinophil	4 %	2 - 12
Basophils	0 %	0 - 1
Absolute N	8352 /uL	2500 - 8500
Absolute L	2436 /uL	1200 - 8000
Absolute M	348 /uL	0 - 600
Absolute E	464 /uL	0 - 1000

Ascn: (b) (6) Profile: Complete Blood Count Ascn:
 (b) (6) Profile: Complete Blood Count

Platelet count reflects the minimum number due to platelet clumping.

5/21/2016 L (b) (6) **Chemistry results from (b) (6) (East) Requisition**

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

ID: (b) (6)	Posted	Final	
Test	Result	Reference	Range
ALB	3.1 g/dL	2.5 - 3.9	
ALKP	27 U/L	6 - 102	
ALT	64 U/L	10 - 100	
AST	44 U/L	10 - 100	
BUN/UREA	26 mg/dL	14 - 36	
Ca	9.3 mg/dL	8.2 - 10.8	
Chloride	112 mEq/L	104 - 128	
CHOL	98 mg/dL	75 - 220	
CK	157 U/L	56 - 529	
CREA	1.2 mg/dL	0.6 - 2.4	
GLU	90 mg/dL	64 - 170	
PHOS	5.9 mg/dL	2.4 - 8.2	
Potassium	5.1 mEq/L	3.4 - 5.6	
Sodium	150 mEq/L	145 - 158	
TBIL	0.1 mg/dL	0.1 - 0.4	
TP	8.4 g/dL	5.2 - 8.8	
GLOB	5.3 g/dL	2.3 - 5.3	
A/G Ratio	0.6 Ratio	0.35 - 1.5	
B/C Ratio	22 Ratio	4 - 33	
Na/K Ratio	29		

5/21/2016	L	(b) (6)	Miscellaneous results from (b) (6) (East) Requisition ID: 209396 Posted Final Asc: (b) (6) Profile: Vet Screen RE: 11067 Comment Hemolysis 1+ No significant interference. Asc: (b) (6) Profile: Vet Screen RE: 11067 Comment Hemolysis 1+ No significant interference.
-----------	---	---------	--

5/21/2016	B	(b) (6)	1.00 At Home Additional Pet Appointment (HC03) by (b) (6)
5/21/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Vetscreen Cbc Antec SA030 (L00030) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)

5/20/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 21/2016 5/20/2016 15:04 Called to confirm tomorrow's appointment fro (b) (6), (b) (6), (b) (6) and (b) (6) at 9
-----------	---	---------	---

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates,
 I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended,
 R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

am. I also mentioned in my message that we should use the address (b) (6) (b) (6) in the GPS. If any questions please call (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 7 Mos.
Color:

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
6/7/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 6/7/2016 12:32 While speaking with owner about (b) (6), discussed (b) (6) dental. Spends the day at 197, but most often home that same night after procedure. Bloodwork is good for 2 months. Can schedule with GP or dentistry according to owner's preference.
5/27/2016	TC	(b) (6)	LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:38 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)
5/24/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:51 Spoke with Mrs; bloodwork WNL, excellent news for planning anesthesia and dental work. Important that taurine status is addressed prior to anesthesia, but dental work should be planned for the next 4-8 weeks. Taurine pending, will call.
5/21/2016	C	(b) (6)	GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Wellness/Taurine check History: Doing well. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; he is the one cat who lives in the house (b) (6) is aggressive toward (b) (6), so he lives away from other cats). Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household. Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food Meds: none S: BAR, PCS 0/4 O: MM / ORPH: Pink, moist, crt <2 sec; right upper canine tooth loose, significant gingivitis locally. heavy tartar on PM3s bilaterally. missing incisors. E/E: brown debris in outer ear cartilages bilaterally, but canals clean/free of debris. ophtho exam WNL. INT: matted hair. no evidence of ectoparasites observed. PLN: WNL CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic GI/UG: Compliant, no masses

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 7 Mos.
Color:

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

MS/NS: Normal amb x4
 BCS: 3-3.5/5 6.7kg
 A: 9yr7mo MN DLH
 1) dental disease
 2) matted hair
 P: PE
 Taurine level
 CBC/Superchem
 PureVax Rabies 1yr SQ right hind (lot# 17390B, exp 12/11/2016)

Advised dental status is poor and likely painful; recommend prompt dental cleaning under general anesthesia with extraction of canine +/- other teeth. Can use dental specialists or general practitioner depending on owner's preference. Recommend waiting for taurine level and any management pertaining to that issue before scheduling anesthetic procedure. Can be shaved down during anesthesia; frequent brushing +/- intermittent clipping or taking to a groomer is needed. Will call with systemic blood results early next week; taurine level will take 7-10 days.

5/21/2016	I	(b) (6)	An animal is not considered immunized for at least 28 days after the initial or primary vaccination is administered. For this reason, pets receiving their first rabies vaccine should not be left outdoors unattended.
-----------	---	---------	---

5/21/2016	V	(b)	May 21, 2016 11:24 AM Staff: (b) ----- Weight : 6.70 kilograms HC-RS scale
-----------	---	-----	---

5/21/2016	L	(b) (6)	Hematology results from (b) (6) (East) Requisition ID: 209396																																																																				
			<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Test</th> <th>Posted</th> <th>Final</th> <th>Reference Range</th> </tr> </thead> <tbody> <tr> <td>HCT</td> <td>36 %</td> <td></td> <td>29 - 48</td> </tr> <tr> <td>HGB</td> <td>11.8 g/dL</td> <td></td> <td>9.3 - 15.9</td> </tr> <tr> <td>MCHC</td> <td>32.8 g/dL</td> <td></td> <td>30 - 38</td> </tr> <tr> <td>WBC</td> <td>9.8 10³/uL</td> <td></td> <td>3.5 - 16.0</td> </tr> <tr> <td>Bands</td> <td>0 %</td> <td></td> <td>0 - 3</td> </tr> <tr> <td>RBC</td> <td>7.9 10⁶/uL</td> <td></td> <td>5.92 - 9.93</td> </tr> <tr> <td>MCV</td> <td>46 fL</td> <td></td> <td>37 - 61</td> </tr> <tr> <td>MCH</td> <td>14.9 pg</td> <td></td> <td>11 - 21</td> </tr> <tr> <td>ABS BASO</td> <td>0 /uL</td> <td></td> <td>0 - 150</td> </tr> <tr> <td>ABS NEUTB</td> <td>0 /uL</td> <td></td> <td>0 - 150</td> </tr> <tr> <td>Platelet C</td> <td>490 10³/uL</td> <td></td> <td>200 - 500</td> </tr> <tr> <td>Platelet E</td> <td>ADEQUATE</td> <td></td> <td>ADEQUATE -</td> </tr> <tr> <td>Neutrophil</td> <td>59 %</td> <td></td> <td>35 - 75</td> </tr> <tr> <td>Lymphocyte</td> <td>33 %</td> <td></td> <td>20 - 45</td> </tr> <tr> <td>Monocytes</td> <td>2 %</td> <td></td> <td>1 - 4</td> </tr> <tr> <td>Eosinophil</td> <td>6 %</td> <td></td> <td>2 - 12</td> </tr> </tbody> </table>	Test	Posted	Final	Reference Range	HCT	36 %		29 - 48	HGB	11.8 g/dL		9.3 - 15.9	MCHC	32.8 g/dL		30 - 38	WBC	9.8 10 ³ /uL		3.5 - 16.0	Bands	0 %		0 - 3	RBC	7.9 10 ⁶ /uL		5.92 - 9.93	MCV	46 fL		37 - 61	MCH	14.9 pg		11 - 21	ABS BASO	0 /uL		0 - 150	ABS NEUTB	0 /uL		0 - 150	Platelet C	490 10 ³ /uL		200 - 500	Platelet E	ADEQUATE		ADEQUATE -	Neutrophil	59 %		35 - 75	Lymphocyte	33 %		20 - 45	Monocytes	2 %		1 - 4	Eosinophil	6 %		2 - 12
Test	Posted	Final	Reference Range																																																																				
HCT	36 %		29 - 48																																																																				
HGB	11.8 g/dL		9.3 - 15.9																																																																				
MCHC	32.8 g/dL		30 - 38																																																																				
WBC	9.8 10 ³ /uL		3.5 - 16.0																																																																				
Bands	0 %		0 - 3																																																																				
RBC	7.9 10 ⁶ /uL		5.92 - 9.93																																																																				
MCV	46 fL		37 - 61																																																																				
MCH	14.9 pg		11 - 21																																																																				
ABS BASO	0 /uL		0 - 150																																																																				
ABS NEUTB	0 /uL		0 - 150																																																																				
Platelet C	490 10 ³ /uL		200 - 500																																																																				
Platelet E	ADEQUATE		ADEQUATE -																																																																				
Neutrophil	59 %		35 - 75																																																																				
Lymphocyte	33 %		20 - 45																																																																				
Monocytes	2 %		1 - 4																																																																				
Eosinophil	6 %		2 - 12																																																																				

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)	Patient: (b) (6)	
Phone: (b) (6)	Species: Feline	Breed: Longhair, Domestic
Address: (b) (6)	Age: 9 Yrs. 7 Mos.	Sex: Neutered Male
(b) (6)	Color:	

Date	Type	Staff	History
------	------	-------	---------

Basophils 0 % 0 - 1
Absolute N 5782 /uL 2500 - 8500
Absolute L 3234 /uL 1200 - 8000
Absolute M 196 /uL 0 - 600
Absolute E 588 /uL 0 - 1000
Ascن: (b) (6) **Profile:** Complete Blood Count

5/21/2016 L

Chemistry results from (b) (6) **(East) Requisition**
ID: 209396 **Posted** **Final**

Test	Result	Reference Range
ALB	3.8 g/dL	2.5 - 3.9
ALKP	25 U/L	6 - 102
ALT	32 U/L	10 - 100
AMYL	1067 U/L	100 - 1200
AST	14 U/L	10 - 100
BUN/UREA	32 mg/dL	14 - 36
Ca	9.9 mg/dL	8.2 - 10.8
Chloride	112 mEq/L	104 - 128
CHOL	125 mg/dL	75 - 220
CK	76 U/L	56 - 529
CREA	1.3 mg/dL	0.6 - 2.4
GGT	1 U/L	1 - 10
GLU	99 mg/dL	64 - 170
Mg	2.2 mEq/L	1.5 - 2.5
PHOS	5.5 mg/dL	2.4 - 8.2
Potassium	5.1 mEq/L	3.4 - 5.6
Sodium	150 mEq/L	145 - 158
TBIL	0.1 mg/dL	0.1 - 0.4
TP	7.8 g/dL	5.2 - 8.8
TRIG	97 mg/dL	25 - 160
GLOB	4.0 g/dL	2.3 - 5.3
A/G Ratio	1.0 Ratio	0.35 - 1.5
B/C Ratio	25 Ratio	4 - 33
Na/K Ratio	29	

5/21/2016 L

Miscellaneous results from (b) (6)
(East) Requisition ID: 209396 **Posted** **Final**
Ascن: (b) (6) **Profile:** Superchem
RE: 1045 PrecisionP 17 U/L 8 - 26
Acute pancreatitis is unlikely. Chronic pancreatitis is not excluded by a normal PrecisionPSL.
RE: 11067 **Comment**
Hemolysis 1+ No significant interference.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 7 Mos.
Color:

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
5/21/2016	B	(b) (6)	1.00 At Home Additional Pet Appointment (HC03) by (b) (6)
5/21/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Superchem Cbc (b) (6) Sa020 (L07) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Fel Vax Rabies 1 Year Purevax (Merial) (V21) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Cared for by (b) (6) by (b) (6)
5/20/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 21/2016 5/20/2016 15:05 Called to confirm tomorrow's appointment from (b) (6), (b) (6), (b) (6) and (b) (6) at 9 am. I also mentioned in my message that we should use the address (b) (6) (b) (6) in the GPS. If any questions please call (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Report Details - EON-266821					
ICSR:	1053339				
Type Of Submission:	Initial				
Report Version:	FPSR.FDA.PETF.V.V1				
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)				
Reporting Type:	Voluntary				
Report Submission Date:	2016-06-06 11:44:41 EDT				
Reported Problem:	Problem Description: Another household cat diagnosed with dilated cardiomyopathy and taurine deficiency - separate report filed (FDA ICSR ID 1053335). Euthanized on (b) (6) due to aortic thromboembolism. Review of the patient's diet history revealed that all 5 cats in household had been fed Merrick Purrrfect Bistro Grain Free Real Chicken Recipe Feline dry for approximately 3 years. Remaining 4 cats in household tested for taurine deficiency - whole blood samples submitted to University of California Davis (normal 300-600 nmol/ml, no known risk for deficiency >200), results received on 5/27/16 - (b) (6) 196nmol/ml - started on taurine supplementation 250mg PO BID for 2-3 weeks. Diet was changed at the time of other cat's diagnosis (5/15/15). Patient also diagnosed with hyperthyroidism on same day as blood submitted for taurine testing - history of weight loss. An echo was not performed on this patient therefore it is unknown if he had evidence of DCM.				
	Date Problem Started: 05/27/2016				
	Concurrent Medical Problem: No				
	Outcome to Date: Not Applicable				
Product Information:	Product Name: Merrick Purrrfect Bistro Grain Free Real Chicken Recipe				
	Product Type: Pet Food				
	Lot Number: <table border="1" style="display: inline-table; vertical-align: top;"> <tr> <td>Lot Number:</td> <td>16025 DL1 38310 14131</td> </tr> <tr> <td>Expiration Date:</td> <td>07/26/2017</td> </tr> </table>	Lot Number:	16025 DL1 38310 14131	Expiration Date:	07/26/2017
	Lot Number:	16025 DL1 38310 14131			
	Expiration Date:	07/26/2017			
	UPC: 2280838310				
	Package Type: BAG				
	Package Size: 5.4 kilogram				
	Number Purchased: 1				
	Possess Opened Product: Yes				
	Storage Conditions: stored in bag indoors				
	Product Use Information:	Description: fed to cats in bowl			
		Last Exposure Date: 05/15/2016			
		Product Use Stopped After the Onset of the Adverse Event: Yes			
Perceived Relatedness to Adverse Event: Definitely related					
Other Foods or Products Given to the Animal During This Time Period: No					
Manufacturer/Distributor Information:	Name: Merrick Pet Care, Inc				
	Type(s): Manufacturer				
	Address: P.O. Box 9800 Amarillo Texas 79105 United States				
FDA-CVM-FOIA-2019-1704-000253					

		Contact:	Phone: 18006647387	
			Web Address: www.merrickpetcare.com	
		Possess One or More Labels from This Product:	Yes	
	Purchase Location Information:	Name:	(b) (6)	
		Address:	(b) (6) United States	
Animal Information:	Name:	(b) (6)		
	Type Of Species:	Cat		
	Type Of Breed:	Mixed (Cat)		
	Gender:	Male		
	Reproductive Status:	Neutered		
	Weight:	4.4 Kilogram		
	Age:	9 Years		
	Assessment of Prior Health:	Fair		
	Number of Animals Given the Product:	5		
	Number of Animals Reacted:	3		
	Owner Information:	Owner Information provided:	Yes	
		Contact:	Name:	(b) (6)
			Phone:	(b) (6)
	Address:	(b) (6) United States		
Healthcare Professional Information:	Practice Name:	(b) (6)		
	Contact:	Name:	(b) (6)	
		Phone:	(b) (6)	
		Email:	(b) (6)	
Address:	(b) (6) United States			
Sender Information:	Name:	(b) (6)		
	Address:	(b) (6)		
		United States		
	Contact:	Phone:	(b) (6)	
		Email:	(b) (6)	
Permission To Contact Sender:	Yes			

FDA-CVM-FOIA-2019-1704-000254

	Preferred Method Of Contact:	Email
	Reported to Other Parties:	Manufacturer

Additional Documents:

Report Details - EON-345822		
ICSR:	2040525	
Type Of Submission:	Initial	
Report Version:	FPSR.FDA.PETF.V.V1	
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)	
Reporting Type:	Voluntary	
Report Submission Date:	2018-01-22 17:19:17 EST	
Reported Problem:	Problem Description: (b) (6) was presented for evaluation of cough, labored breathing, multiple episodes of collapse, cardiomegaly, and suspected congestive heart failure. Congestive heart failure was confirmed with thoracic radiographs and echocardiogram revealed dilated cardiomyopathy.	
	Date Problem Started: 12/30/2017	
	Concurrent Medical Problem: Yes	
	Pre Existing Conditions: Canine atopy controlled with current treatment of sublingual immunotherapy and Apoquel.	
	Outcome to Date: Better/Improved/Recovering	
Product Information:	Product Name: California Natural Grain-Free Kangaroo and Red Lentils Recipe	
	Product Type: Pet Food	
	Lot Number:	
	Package Type: BAG	
	Purchase Date: 01/02/2018	
	Possess Unopened Product: No	
	Possess Opened Product: Yes	
	Product Use Information:	Description: (b) (6) had been eating this dog food since she first displaced signs of pruritis as a puppy and food allergy was considered as a potential contributor.
		Last Exposure Date: 01/10/2018
		Time Interval between Product Use and Adverse Event: 6 Years
		Product Use Stopped After the Onset of the Adverse Event: Yes
		Adverse Event Abate After Product Stop: Unknown
		Product Use Started Again: No
		Perceived Relatedness to Adverse Event: Probably related
Other Foods or Products Given to the Animal During This Time Period: Yes		
Manufacturer /Distributor Information:		
Purchase Location Information:		

Animal Information:	Name:	(b) (6)		
	Type Of Species:	Dog		
	Type Of Breed:	Mixed (Dog)		
	Gender:	Female		
	Reproductive Status:	Neutered		
	Weight:	25.1 Kilogram		
	Age:	6 Years		
	Assessment of Prior Health:	Good		
	Number of Animals Given the Product:	4		
	Number of Animals Reacted:	4		
	Owner Information:	Owner Information provided:	Yes	
		Contact:	Name:	(b) (6)
			Phone:	(b) (6)
			Email:	(b) (6)
	Address:	(b) (6) United States		
Healthcare Professional Information:	Practice Name:	(b) (6)		
	Contact:	Name:	(b) (6)	
		Phone:	(b) (6)	
		Other Phone:	(b) (6)	
		Email:	(b) (6)	
	Address:	(b) (6) United States		
Sender Information:	Name:	(b) (6)		
	Address:	(b) (6) United States		
	Contact:	Phone:	(b) (6)	
		Other Phone:	(b) (6)	
		Email:	(b) (6)	
	Permission To Contact Sender:	Yes		
	Preferred Method Of Contact:	Phone		
	Reported to Other Parties:	Other		
Additional Documents:	Attachment:	Medical record.pdf		
	Description:	Medical record and echo report		
	Type:	Medical Records		

FDA-CVM-FOIA-2019-1704-000257

Attachment:	Taurine level.pdf
Description:	Taurine level
Type:	Laboratory Report
Attachment:	Listserve on kangaroo and lentil diets.pdf
Description:	Discussion amongst veterinary cardiologists of dilated cardiomyopathy in patients eating either kangaroo and lentil or vegan diets with lentils
Type:	Other

17962

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/aal/aal.html>

Vet/Tech Contact: Account # _____ (b) (6) / Contact: _____ (b) (6) Date: 1-10-18

Company Name: _____ (b) (6)

Address: _____ (b) (6)
_____ (b) (6)

Email: _____ (b) (6)

Tel: _____ (b) (6) Fax: _____ (b) (6)

Billing Contact: _____ (b) (6) TAX ID: _____

Email: _____ (b) (6) Tel: _____ (b) (6)

Patient Name: _____ (b) (6)

Species: kg

Owner's Name: _____ (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____

Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _ _ Whole Blood: 292 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

From: [Norris, Anne](#)
To: [Rotstein, David](#); [Hartogenesis, Martine](#); [DeLancey, Siobhan](#)
Subject: RE: Calls Complete
Date: Tuesday, June 25, 2019 9:13:41 AM

Thought you'd find this interesting <https://www.petfoodprocessing.net/articles/13194-midwestern-manufacturer-debuts-legume-free-dog-foods>

From: Rotstein, David
Sent: Monday, June 24, 2019 12:57 PM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>; DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Subject: RE: Calls Complete

Martine deserves a lot of credit. Some of the firms were challenging.

From: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Date: June 24, 2019 at 12:22:40 PM EDT
To: Norris, Anne <Anne.Norris@fda.hhs.gov>, DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: Calls Complete

No problem. It was an interesting morning!

Martine

From: Norris, Anne
Sent: Monday, June 24, 2019 12:15 PM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: Calls Complete

Understood, just confirming. Thanks for handling this, I know it was an unpleasant situation.

From: Hartogenesis, Martine
Sent: Monday, June 24, 2019 12:13 PM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>; DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: Calls Complete

No – I thought the plan was (b) (5). We had several firms that were pretty upset BTW.

From: Norris, Anne
Sent: Monday, June 24, 2019 12:11 PM
To: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: Calls Complete

Thanks! Did you speak to PFI?

From: DeLancey, Siobhan
Sent: Monday, June 24, 2019 12:06 PM
To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: Calls Complete

Thanks!

From: Hartogensis, Martine
Sent: Monday, June 24, 2019 12:04 PM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>; DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: Calls Complete

From: [Norris, Anne](#)
To: [Hartogenesis, Martine](#)
Cc: [DeLancey, Siobhan](#)
Subject: RE: Clearing DCM Comms
Date: Tuesday, June 18, 2019 9:26:34 AM
Attachments: [DCM Project Plan.docx](#)
[image001.png](#)
[image002.jpg](#)
[image003.jpg](#)
[image004.jpg](#)
[image005.jpg](#)
[image006.jpg](#)

Thanks! I think these look good. I believe Tracey and Nadine started a script – see attached. I think Dave knows best about what the current marching orders are with the divisions so you may want to touch base with him directly. Let me know what I can do to help.

Anne

From: Hartogenesis, Martine
Sent: Tuesday, June 18, 2019 7:59 AM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Subject: RE: Clearing DCM Comms

Hi Anne!

I reviewed and these are cleared by me. I added some suggested language (3rd paragraph CVM Update and question 3 in Q&A) (b) (5). See what you think!

(b) (5)

I missed the meeting last week, so I am not sure where we are on that and the script. I am happy to write a script if there isn't one yet.


Lastly, I added in a few places ..labeled as “grain-free” so we are consistent. Not a huge deal, but it might help some of our readers recall the issue.

Thanks again!

Martine

From: Norris, Anne
Sent: Monday, June 17, 2019 4:48 PM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Subject: Clearing DCM Comms

When you go through and clear the comms, do you want to enter in the language that you feel most comfortable with regarding (b) (5)



Is tomorrow late morning a reasonable timeline to get your clearance? Happy to discuss.

Thanks!

Anne Norris
Strategic Initiatives

Office of the Director
Center for Veterinary Medicine
U.S. Food & Drug Administration
O: 240-402-0132
M: (b) (6)
Anne.Norris@fda.hhs.gov



DCM Plan

Activity	Lead	Notes	Timing
<p>(b) (5)</p>			

(b) (5)

(b) (5)	

From: [Norris, Anne](#)
To: [Palmer, Lee Anne](#); [Carey, Lauren](#); [Rotstein, David](#); [Jones, Jennifer L](#); [Peloquin, Sarah](#); [Reimschuessel, Renate](#); [Hartogensis, Martine](#); [Burkholder, William](#); [DeLancey, Siobhan](#)
Subject: RE: DCM paper - Darcy Adin, 2019 Vet Cardiology
Date: Thursday, February 21, 2019 12:12:01 PM
Attachments: [sky488.pdf](#)
[image001.png](#)
[image002.jpg](#)
[image003.jpg](#)
[image004.jpg](#)
[image005.jpg](#)
[image006.jpg](#)

I've lost track of whether we circulated this paper internally, but sharing because it caught the eye of Phyllis Entis from Food Safety News. She hasn't written about it (at least not yet). One of the authors is Greg Aldrich.

From: Norris, Anne
Sent: Tuesday, February 19, 2019 9:09 AM
To: Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>; Reimschuessel, Renate <Renate.Reimschuessel@fda.hhs.gov>; Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Burkholder, William <William.Burkholder@fda.hhs.gov>
Subject: RE: DCM paper - Darcy Adin, 2019 Vet Cardiology

Thanks!

From: Palmer, Lee Anne
Sent: Tuesday, February 19, 2019 9:05 AM
To: Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>; Reimschuessel, Renate <Renate.Reimschuessel@fda.hhs.gov>; Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>; Burkholder, William <William.Burkholder@fda.hhs.gov>
Subject: DCM paper - Darcy Adin, 2019 Vet Cardiology

Hi – please forgive me if we have this already, but I think this just came out.

I haven't read it yet.

Thanks, lee Anne

Lee Anne M. Palmer, VMD, MPH
Team Leader HFV-242, Supervisory VMO

Center for Veterinary Medicine
OSC, Division of Veterinary Product Safety

U.S. Food and Drug Administration

Tel: 240-402-5767

Leeanne.palmer@fda.hhs.gov



Special topic: The association between pulse ingredients and canine dilated cardiomyopathy: addressing the knowledge gaps before establishing causation¹

Wilfredo D. Mansilla,[†] Christopher P.F. Marinangeli,[‡] Kari J. Ekenstedt,^{||} Jennifer A. Larsen,[§] Greg Aldrich,[¶] Daniel A. Columbus,^{**} Lynn Weber,^{**} Sarah K. Abood,^{|||} and Anna K. Shoveller^{†,2}

[†]Department of Animal Biosciences, University of Guelph, Guelph, ON N1G 2W1, Canada; and [‡]Pulse Canada, Winnipeg, Manitoba, Canada, R3C 0A5; ^{||}Department of Basic Medical Sciences, College of Veterinary Medicine, Purdue University, West Lafayette, IN 47907; [§]Department of Molecular Biosciences, School of Veterinary Medicine, University of California, Davis, CA 95616; [¶]Department of Grain Science and Industry, Kansas State University, Manhattan, KS 66506; ^{**}Prairie Swine Centre, Saskatoon, SK S7H 5N9, Canada; ^{**}Department of Veterinary Biomedical Sciences, University of Saskatchewan, 52 Campus Drive, Saskatoon, SK S7N 5B4, Canada; ^{|||}Department of Clinical Studies, University of Guelph, Guelph, ON N1G 2W1, Canada

ABSTRACT: In July 2018, the Food and Drug Administration warned about a possible relationship between dilated cardiomyopathy (DCM) in dogs and the consumption of dog food formulated with potatoes and pulse ingredients. This issue may impede utilization of pulse ingredients in dog food or consideration of alternative proteins. Pulse ingredients have been used in the pet food industry for over 2 decades and represent a valuable source of protein to compliment animal-based ingredients. Moreover, individual ingredients used in commercial foods do not represent the final nutrient concentration of the complete diet. Thus, nutritionists formulating dog food must balance complementary ingredients to fulfill the animal's nutrient needs in the final diet. There are multiple factors that should be considered, including differences in nutrient digestibility and overall bioavailability, the fermentability and quantity of fiber, and interactions among food constituents that can increase the risk of DCM development.

Taurine is a dispensable amino acid that has been linked to DCM in dogs. As such, adequate supply of taurine and/or precursors for taurine synthesis plays an important role in preventing DCM. However, requirements of amino acids in dogs are not well investigated and are presented in total dietary content basis which does not account for bioavailability or digestibility. Similarly, any nutrient (e.g., soluble and fermentable fiber) or physiological condition (e.g., size of the dog, sex, and age) that increases the requirement for taurine will also augment the possibility for DCM development. Dog food formulators should have a deep knowledge of processing methodologies and nutrient interactions beyond meeting the Association of American Feed Control Officials nutrient profiles and should not carelessly follow unsubstantiated market trends. Vegetable ingredients, including pulses, are nutritious and can be used in combination with complementary ingredients to meet the nutritional needs of the dog.

Key words: dilated cardiomyopathy, dogs, feed formulation, grain-free, nutrition, pulse ingredients

© The Author(s) 2019. Published by Oxford University Press on behalf of the American Society of Animal Science.

¹Funding for this project was provided by Pulse Canada. C.P.F.M. works for Pulse Canada and is a former employee of Kellogg Canada. W.D.M., A.K.S., K.J.E., G.A., J.A.L., D.A.C., L.W., and S.K.A. have no conflicts of interest. All authors contributed to the content of this paper. We would

like to acknowledge the contribution of James Templeman, Sarah Dodd, and Emma Thornton.

²Corresponding author: ashovell@uoguelph.ca

Received November 23, 2018.

Accepted January 4, 2019.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

J. Anim. Sci. 2019.XX:XX–XX
doi: 10.1093/jas/sky488

INTRODUCTION

In July 2018, the Food and Drug Administration (FDA) issued a statement relating dilated cardiomyopathy (DCM) in dogs to the consumption of foods that have potatoes and/or pulse ingredients, such as peas and lentils or their coproducts, as main ingredients (FDA, 2018). The FDA's statement, as well as media attention, has raised concern in some pet owners, veterinarians, nutritionists, and the pet food manufacturing and retail industry. The underlying cause for concern with pet food and DCM is that there is a link between nutrition that was previously tied to DCM and insufficient circulating taurine (Fascetti et al., 2003; Backus et al., 2006). The result was an increased need for dietary taurine or its precursor methionine due to higher fermentation of taurine and greater fecal excretion with dietary fermentable fiber (Kim et al., 1996a, 1996b). Whether this has any link to dietary pulses or the greater inclusion of pulses in grain-free dog food has yet to be directly demonstrated and mechanistic research is warranted.

Pulses are a subset of legumes, harvested as a dry crop, with low concentrations of lipid. They include peas, lentils, chickpeas, and dry beans (Marinangeli et al. 2017) which have been used as ingredients in dog food for their protein and fiber for more than 2 decades (Butterwick et al., 1994; Rice and Ihle, 1994). As a source of protein, the amino acid (AA) profile in peas, lentils, chickpeas, and beans is generally high in lysine and low in methionine (NRC, 2006) and serves as a complementary protein to both animal and plant-derived ingredients. As an example, soybean meal is derived from defatted soybeans and has an AA profile similar to pulses. In a 24-wk study that evaluated graded concentrations of soybean meal up to 17% (as-fed basis) in dog foods, soybean meal inclusion did not affect the nutrient status of dogs as indicated by serum biochemistry analysis (Menniti et al., 2014). However, Yamka et al. (2003) demonstrated that using soybean meal at more than 15% inclusion on a dry matter basis decreased crude protein digestibility. Based on the authors' assessment of current formulas in the market, there is a high likelihood that legume seed use in some foods may be greater

than 40%. This inclusion exceeds concentration of legumes previously investigated in dogs. When used to complement the nutritional profile of other ingredients, pulses can be used as nutrient-rich vehicles to meet the nutritional requirements of dogs and other companion animals. Given that companion animals most often consume static diets for long periods of time, overuse of any ingredient could facilitate higher risk of certain nutrient deficiencies if nutrient balance is not considered in the formulation. Thus, the formulation of static diets that use significant concentrations of a single ingredient, relative to other ingredients in the formulation, requires an in-depth knowledge of nutrient interactions, animal physiology, and effects of processing, beyond that of simply meeting minimum nutrient profiles stipulated in the Official Publication of The Association of American Feed Control Officials (AAFCO, 2018).

The present commentary discusses the following: 1) The limited data being used to support linkages between DCM and pulse ingredients; 2) The nutritional factors and physiological mechanisms that should be explored to establish causation between nutritional deficiencies and incidence of DCM; 3) The factors that nutritionists should consider when formulating complete diets destined for long-term consumption; and 4) The disadvantages of formulating protein and minimal AA recommendations rather than a balanced indispensable AA profile.

The Development of Canine DCM, Historical Linkages to Taurine Deficiency, and Pulses

Dilated cardiomyopathy is a disease of the myocardium that results in both mechanical dysfunction (enlarged heart cavities and congestion) and/or electrical dysfunction (arrhythmias and sudden death) (Sisson et al., 2000; Maron et al., 2006; Dutton and López-Alvarez, 2018). Development of DCM is slow and few clinical signs manifest over time. As DCM progresses, signs include lethargy, anorexia, shallow breathing, sudden fainting, and potential death. In some cases, animals may die from irregular heart rhythm without previous signs of the disease. In dogs, DCM can be

caused by various factors. Genetic predisposition is thought to play the most important role in the development of DCM in several dog breeds, mostly large and giant breeds. Genetic mutations associated with DCM have been discovered in American lines of Doberman and Boxer dogs (Meurs et al., 2012; Meurs et al., 2013). However, the Doberman variant's association was not upheld in a European population of Dobermans (Owczarek-Lipska et al., 2013). Similarly, a United Kingdom population of Boxers did not uphold their published DCM-associated variant (Cattanach et al., 2015). It is becoming increasingly clear that the genetic basis for DCM in dogs is not monogenic, but complex and polygenic. Breeds with the highest prevalence of DCM include Dobermans, Boxers, Great Danes, Newfoundlands, Irish Wolfhounds, English Cocker Spaniels, and Portuguese Water Dogs (Monnet et al., 1995; Borgarelli et al., 2006; Werner et al., 2008; Martin et al., 2009), and the genetic basis of DCM in each of these breeds has been investigated (Dutton and López-Alvarez, 2018). In addition, Golden Retrievers and American Cocker Spaniels appear to have breed predispositions to taurine deficiency (Kramer et al., 1995; Bélanger et al., 2005). When dogs are not genetically predisposed for developing DCM, diet and physiology are other factors that may be associated with the disease.

The first link between taurine deficiency and DCM was demonstrated in cats in 1987. Cats diagnosed with DCM recovered after taurine supplementation (Pion et al., 1987). Similarly, an inverse association between dietary taurine and the incidence of DCM in a population of foxes was documented by Moise et al. (1991) and

established the importance of taurine in the family Canidae. In dogs, DCM diagnoses related to low whole blood taurine concentrations have been reported in Cocker Spaniels, Dalmatians, Boxers, Newfoundlands, Portuguese Water Dogs, English Setters, Alaskan Malamutes, and Scottish Terriers (Freeman et al., 1996; Kittleson et al., 1997; Pion et al., 1998; Alroy et al., 2000; Fascetti et al., 2003; Backus et al., 2006). In all these cases, taurine supplementation improved cardiac function. However, dogs, in contrast to cats, can endogenously synthesize taurine from methionine and cysteine (Figure 1). Therefore, the above-mentioned data do not unequivocally establish taurine intake as the underlying mechanism for the development of DCM in dogs, whether they are genetically predisposed. Dietary supply of precursor AAs necessary for taurine synthesis (i.e., methionine and cysteine), metabolic intermediates, and cofactors (such as methyl donors) cannot be ruled out as factors that contribute to the susceptibility of dogs to developing genetic and diet-related DCM. When DCM is diet-related, the formulation and the provision of all nutrients, including indispensable AAs, to facilitate optimum health and wellbeing of dogs should be considered.

Recent reports, including the statement by the FDA (2018), have implicated that lentils, peas, and other legumes seeds could be responsible for the development of DCM in dogs not genetically predisposed to this disease. Such statements and associations between pulse ingredients and incidence of DCM are, at the present time, premature. Animals, including dogs, have no minimum or maximum requirements for ingredients. Ingredients serve




Figure 1. Metabolism of sulfur amino acids. DMG = dimethylglycine; SAH = S-adenosylhomocysteine; SAM = S-adenosylmethionine.

as the vehicle to providing nutrients to animals. As such, animals have nutrient requirements, not ingredient requirements. In diets that have nutrient deficits, imbalances, or exceed maximums, the final nutrient composition of the diet, not the ingredients, should be critiqued. In addition, animal nutritionists should consider that the nutrient concentration of ingredients can vary, nutrient availability is not 100%, and diets formulated to marginally meet requirements could actually be deficient. Overall, it is the responsibility of nutritionists to use different ingredients to formulate diets that can be produced and safely meet the nutritional needs of animals.

Taurine Deficiency and the Development of Canine DCM

For dogs, taurine is a dispensable AA synthesized from methionine and cysteine primarily in the liver (Figure 1). Taurine is not incorporated into proteins. Instead, it is used as a mediator for various biological processes and is the most abundant free AA intracellularly (Huxtable, 1992). In the heart, taurine represents ~60% of the total AA free pool (Huxtable, 1992). The high concentration of taurine in cardiac cells may explain the role of a taurine deficiency in the development of DCM. It has been speculated that taurine contributes to the reabsorption of calcium by the sarcoplasmic reticulum and increases the sensitivity of the myofilaments to calcium (Bakker and Berg, 2002). Thus, low dietary taurine intake and/or reduced synthesis of taurine from methionine and cysteine can deplete calcium pools in the cardiac cells and impede proper contraction of the cardiac muscle tissue, resulting in DCM in dogs.

For diagnosing DCM in dogs and cats, among other diagnostic methods including electrocardiograms and echocardiography, it is common to measure taurine concentration in whole blood. Whole blood samples, and not plasma samples, should be used to assess circulating taurine concentrations. In plasma, free taurine concentrations are much lower compared with intracellular taurine. This suggests that the plasma pool is not representative of taurine in other pools (Schaffer et al., 2010). In platelets, taurine concentration is high and is considered a marker of taurine status. Taurine concentration in platelets is captured when whole blood is analyzed (Huxtable, 1992). However, platelet count can vary depending on the immune status of the animal and whole blood taurine concentration can be affected. In this scenario, whole blood taurine may not represent concentrations of

taurine in muscle cells, including cardiac muscle. These additional variables related to the measurement of taurine status may explain why some dogs diagnosed with DCM have normal whole blood taurine concentrations.

As taurine can be synthesized endogenously in dogs, taurine is not considered an indispensable AA for the species Canidae. Thus, there are no recommendations on minimum dietary concentrations of taurine for dogs reported by the National Research Council (NRC, 2006) or AAFCO (2018). The lack of regulation on minimum taurine concentrations in commercial dog foods suggests that endogenous synthesis of taurine can meet the metabolic needs in all dogs and at all life stages. This assumption may not be accurate as studies have determined that synthesis of taurine is related to the size of dog (Ko et al., 2007), and some dietary factors can increase the physiological need for taurine (Story, 1978). Nutritional factors that increase the dietary requirement, reduce the supply, or increase the excretion of taurine in dogs are discussed in subsequent sections of this review and should be considered to avoid taurine deficiency in dogs and the risk of DCM.

Physiological factors can increase taurine utilization in dogs, and endogenous synthesis of taurine could be insufficient for meeting taurine requirements. For example, compared with smaller size dogs, synthesis of taurine in large dog breeds is up to 50% lower per unit of metabolic body weight (Ko et al., 2007). These results demonstrate that larger dogs are at higher risk for insufficient endogenous taurine synthesis, and dietary supplementation or fortification may be required, even when there is no minimum dietary taurine concentration according to current recommendations (AAFCO, 2018). Obesity and diabetes have also been related to lower concentrations of taurine in blood in humans and rats, respectively (Merheb et al., 2007; Nardelli et al., 2011; Ito et al., 2012), and may increase the requirement for sulfur AAs necessary for endogenous taurine synthesis. This is of importance given that approximately half of dogs in North America are obese (Linder and Mueller, 2014). Data from rats and cats suggest that age and sex could also affect whole body taurine status. Hepatic activity of cysteine sulfonate decarboxylase, the enzyme responsible for taurine synthesis, was shown to be 16 times higher in adult male rats vs. female rats. In the same study, the activity of cysteine sulfonate decarboxylase was higher in 5- to 6-wk-old kittens compared with 15-mo-old cats and in 8-wk-old mice compared with 16-wk-old mice; changes of

the enzyme activity in dogs have not been tested (Worden and Stipanuk, 1985). Overall, these studies suggest that, despite some capacity for endogenous synthesis, physiological need of taurine can be heavily dependent on breed, age, sex, and physiological status. These physiological factors could help us to predict the risk for developing DCM when genotypic and environmental factors, such as diet, are simultaneously considered to ensure that dogs maintain adequate concentrations of taurine and other sulfur AAs.

Given that there are no recommendations for the minimum concentration of taurine in dog food, the concentration of taurine in dog foods can vary substantially depending on the ingredients used. Taurine is very low in plant-based ingredients (Table 1) but is higher in some algae and fungi species and is ubiquitously found in animal tissues, especially in the heart, brain, and white blood cells (Huxtable, 1992). This is relevant, as many grain-free and/or high legume dog foods attempt to limit the use of animal byproducts, which can substantially decrease the levels of dietary taurine. In the context of providing adequate and preventive nutrition, dog foods should include organ meat

or animal byproducts or be fortified with taurine and/or its precursors (methionine and/or cysteine) to ensure the delivery of sufficient levels of taurine.

Effect of Dietary Fiber on Taurine Status and Risk of Canine DCM

Dietary fiber has been shown to affect the taurine status in dogs. For example, commercial diets formulated with lamb meal and rice bran were shown to cause taurine deficiency in part because of low bioavailable cysteine from lamb meal and possibly more importantly due to the effects of rice bran fiber on gastrointestinal metabolism of taurine (Johnson et al., 1998; Tôrres et al., 2003). It has been hypothesized that high-fiber diets can increase susceptibility to taurine deficiency by 2 mechanisms of action linked to obligatory bile acid conjugation with taurine in dogs (O'Mádille et al., 1965) and reliance on enterohepatic circulation for the reabsorption of bile acids and taurine. First, high-fiber diets may increase fecal output and losses of taurine-conjugated bile. This would require higher synthesis rates of bile in the liver, and consequently, higher utilization of taurine

Table 1. Crude protein (CP), fiber, selected amino acids, and carnitine contents in the principal legumes, cereals, and animal-derived ingredients used in dog food formulation

Ingredients	CP, %	Crude fiber, ¹ %	α -amino acids, mg/g protein ¹			Tau, mg/kg ²	Carnitine, mg/kg ³	
			Lys	Met	Cys			
Legumes	Fava beans	27.2	8.55	23.9	7.0	12.5	–	–
	Phaseolus beans	22.9	NR	72.9	12.7	12.7	–	–
	Kidney beans	20.0	6.40	26.5	14.0	12.0	–	–
	Lentils	26.0	NR	65.8	6.9	10.4	–	–
	Lupins	32.4	14.25	48.7	6.5	14.2	–	–
	Chick peas	20.3	6.16	69.4	14.8	21.6	–	–
	Soybean meal	47.7	3.89	62.0	13.8	14.7	–	–
Grains	Barley	11.3	3.90	35.3	17.7	22.9	–	–
	Corn, yellow dent	8.2	1.98	30.3	21.8	23.1	–	–
	Oats	11.2	2.20	43.9	60.9	32.3	–	–
	Rice	7.9	0.52	44.5	31.8	22.9	–	–
	Rye	11.7	2.71	36.9	13.7	16.3	–	–
	Sorghum	9.4	2.14	21.4	17.1	19.2	–	–
	Wheat hard, red	14.5	2.57	27.0	15.2	22.8	–	–
Animal-derived ingredients	Beef, meat	15.0	–	77.3	28.7	15.3	296	150
	Chicken, meat and skin	17.6	–	81.3	26.7	13.1	159	57
	Chicken, by product	59.0	–	48.1	17.3	16.8	3049	120
	Lamb, ground	16.6	–	88.0	25.9	12.0	473	282.3
	Rendered meat	54.1	2.50	53.8	14.2	11.3	NR	NR

Cys = cysteine; Lys = lysine; Met = methionine; NR = not reported; Tau = taurine.

Values are presented on as-fed basis.

¹NRC, 2006; NRC, 2012.

²Spitze et al. 2003.

³Arslan, 2006.

(Story, 1978). Second, high consumption of fermentable fibers may increase the abundance of microbial populations that degrade taurine in the intestinal lumen (Kim et al., 1996a, 1996b). Either alone or together, increased excretion or degradation of taurine from high-fiber diets may decrease enterohepatic circulation and recycling of taurine. Given that taurine is the only AA used for bile acid conjugation in dogs, over time, high-fiber diets could increase the risk of taurine insufficiency in dogs and lead to DCM.

This should not be interpreted as dietary fiber being deleterious to the health of dogs. However, there may be a limit to the benefit for soluble fibers. Legume seeds contain an appreciable quantity of oligosaccharides which are known to be fermentable (Tosh and Yada, 2010). Thus, by a similar mechanism as described above, high levels of legume seed oligosaccharides could ostensibly contribute to taurine depletion via excretion in the feces as bile conjugation and degradation by colonic bacteria. In addition to the physiological benefits of high-fiber diets in certain dogs, formulators should also be cognizant of possible nutritional risks associated with high concentrations of fiber in dog foods. Consequently, dog foods with high concentrations of dietary fiber should be accompanied by higher supplies of taurine or sulfur AAs for endogenous taurine synthesis. Overall, the digestibility and bioavailability of taurine in ingredients used and the effect of other nutrients in taurine metabolism should be considered to avoid taurine deficiency and the development of DCM.

Carnitine Deficiency and Risk of Canine DCM

Carnitine is not nutritionally indispensable since it is endogenously produced in the liver and kidneys from lysine and methionine; it can also be attained exogenously from animal-based products. Carnitine is highly abundant in skeletal and cardiac muscles. Together, these represent >95% of the total carnitine in the body. Carnitine is essential for metabolism of fatty acids used for energy production (Hoppel, 2003). In the heart, where 60% of the energy is derived from fatty acid oxidation, carnitine facilitates the uptake of free fatty acids into the mitochondria to produce ATP (Hoppel, 2003). Plant-based ingredients do not contain carnitine (Table 1). Therefore, in commercial dog foods with reduced inclusion of animal-based ingredients, intakes of carnitine could be decreased if diets are not fortified. Reduced dietary carnitine intake

translates into increased reliance on endogenous synthesis to meet physiological requirements.

Given that carnitine is required for sufficient energy production in cardiac muscle, it is not surprising that carnitine deficiency is associated with DCM. In 1991, a family of Boxers diagnosed with DCM were also diagnosed with carnitine deficiency (Keene et al., 1991). In dogs, carnitine deficiency can occur with aberrations of carnitine regulation in disorders such as cardiomyopathy (including DCM), diabetes, sepsis, and malnutrition (Flanagan et al., 2010). However, carnitine deficiency as a causative factor in the development of DCM or a consequence of cardiac malfunction remains as a subject of debate (Freeman and Rush, 2006). Despite the interest in this metabolite, little progress has been made on determining the effect of carnitine supplementation on alleviating risk of DCM. However, both taurine and carnitine are often supplemented in suprphysiological concentrations once DCM is diagnosed. This practice is supported by positive clinical outcomes, albeit without comparison groups (Kittleson et al., 1997; Sanderson et al., 2001). Concentrations of carnitine in the plasma are relatively insensitive to dietary carnitine, and more invasive techniques (biopsies) are required to determine the concentration of carnitine in muscle tissue (Flanagan et al., 2010; Rășanu et al., 2012). The invasive nature of testing for carnitine status is likely the reason why carnitine is rarely explored when investigating possible causes of canine DCM.

Preventing Diet-Mediated DCM in Dogs by Providing Adequate Sulfur AAs and Maximizing Endogenous Taurine Synthesis

Although taurine is considered a dispensable AA in dogs, endogenous taurine synthesis requires an adequate supply of bioavailable sulfur AA precursors cysteine or methionine (Figure 1). Thus, providing marginal concentrations of these 2 sulfur AAs, or providing sources with lower bioavailability, could increase the risk of taurine deficiency and facilitate the development of DCM. Contrary to taurine, methionine cannot be synthesized endogenously in dogs (NRC, 2006). Therefore, dogs depend on the provision of dietary methionine to meet daily sulfur AA requirements, which includes production of taurine. From an ingredient perspective, methionine and lysine are usually the first or second limiting AAs in dog diets formulated with soybean meal and rendered meats (NRC, 2006). In addition, methionine is particularly susceptible to damage, and subsequent reduction in bioavailability,

secondary to heat processing (Marshall et al. 1982; Hurrell et al., 1983). This suggests that the risk of methionine deficiency is more likely than any other indispensable AA in commercial dog diets. Although the primary role for methionine is protein synthesis, in pigs at least 50% of absorbed methionine acts as a methyl donor and a precursor in the production of cysteine, taurine, sulfate, and pyruvate (Robinson et al., 2016a; Figure 1). These functions of methionine become more crucial when dietary intake of cysteine, taurine, and/or dietary methyl donors (e.g., folate, betaine, and their precursors) is limited (Robinson et al., 2016b), and they need to be considered when nutritionists set criteria for delivery of sulfur AAs in pet foods.

Methionine and cysteine both contribute to the total sulfur AA requirements for humans and animals. For adult dogs at maintenance, the latest guidelines from the NRC (2006) recommend that adult dog foods contain 0.33% (on dry matter basis) methionine when cysteine is provided in excess, and 0.65% for methionine + cysteine. These NRC (2006) recommendations are not based on dose–response studies, but on a 4-yr study where adult dogs were fed low-crude protein diets (Sanderson et al., 2001). In that study, the lowest concentration of methionine in the diet that reported no observable deficiencies was used as the recommended requirement. As companion animals are typically fed a single static diet during adulthood, and for most of their lifespan, it is necessary that AA requirements of dogs should be measured empirically (Baker, 1986). In addition to the lack of empirical data corresponding to the AA requirements of dogs, it is equally important to understand how other dietary (e.g., dietary fiber), environmental, other physiological variables, and breed/genotype may alter AA requirements. The lack of recommendations for taurine in commercial dog food puts a higher stress on accurately meeting requirements for sulfur AAs, not only for protein synthesis, but also for the endogenous synthesis of taurine, for support of optimal methyl status, and for the synthesis of secondary metabolites.

Rethinking Indispensable AA Targets in Commercial Dog Foods

Currently, the ingredients permitted in pet foods and the corresponding nutrient targets are guided by recommendations made by AAFCO (2018). These recommendations are based on the peer-reviewed scientific literature and represented in the Nutrient Requirement of Dogs and Cats

(NRC, 2006). However, AA recommendations made by AAFCO correspond to total AA content within the formulation and do not consider the true ileal digestibility of ingredients. True ileal digestibility of AAs is more representative of nutrient absorption capacity and bioavailability compared with fecal digestibility or total AA content in the diet (Columbus and de Lange, 2012). To account for the reduced digestibility and bioavailability of protein-bound AAs in food ingredients, AAFCO arbitrarily increases AA recommendations relative to those from the NRC to ensure that an adequate supply of AAs is provided, regardless of the ingredients and effects of processing (Table 2). However, this increment is only applied to lysine, threonine, and tryptophan and not applied to other indispensable AAs, including methionine (AAFCO, 2018). For example, the recommended allowance for lysine reported in NRC (2006) is 0.35% for adult dogs at maintenance, whereas the minimum content of lysine to meet AAFCO (2018) recommendations is 0.63%. Nonruminant animals, including dogs, absorb AAs from the duodenum to the terminal ileum (Columbus and de Lange, 2012). Hence, feeding diets with lower ileal digestibility coefficients could decrease actual concentrations of available indispensable AAs, even when meeting AAFCO recommendations. This is of special concern for dietary taurine and other sulfur AAs, considering that there is no regulated minimum threshold for taurine in dog foods and that AAFCO (2018) recommendations for sulfur AAs are not increased compared with NRC (2006) recommendations to account for potential ileal digestibility coefficients. There is a dearth of data in this area to justify empirical adjustments based on different dietary variables. As such, future research should pursue how AA requirements change under different dietary variables that can affect small intestinal digestibility and whole body availability.

It is worthwhile to note that minimum dietary nutrient contents for dog foods, as reported in AAFCO (2018), only consider differences between growth/reproduction and adult life stages. This lack of data places the pregnant bitch in the same group as growing animals. Moreover, most studies on nutrient requirements in dogs have been established using Beagles as a proxy for all dogs. Using a single breed creates a homogenous sample and likely does not account for nutritional variability across pure and mixed breeds, or those of different sizes. Unpublished data from Shoveller et al. investigated the minimum methionine (with excess cysteine) requirements of

Table 2. Recommended allowance (RA) and minimum dietary content suggested by AAFCO for crude protein and essential amino acids in dog food, and their physiological roles and potential interactions

Nutrient	NRC RA ¹ , % DM	AAFCO ² , % DM	Important physiological roles and potential interactions
Crude protein	10	18	Necessary for synthesis of nonessential amino acids
Arginine	0.35	–	Competes with lysine absorption, arginine should be increased when high lysine concentrations in the diet
Histidine	0.19	–	
Lysine	0.35	0.63	Highly reactive to reducing sugars during heating (Maillard reaction), reducing bioavailability
Methionine	0.33	0.33	Requirement increases when methyl donors/acceptors and cysteine are reduced in the diet
Methionine + cystine	0.65	0.65	Requirement is increased with low supply of taurine and during immune challenge
Phenylalanine	0.45	0.45	
Phenylalanine + tyrosine	0.74	0.74	
Threonine	0.43	0.48	Abundant in mucosal proteins (mucin), requirement increases when feeding high fermentable fibers
Tryptophan	0.14	0.16	Precursor for serotonin synthesis. Ratio of Trp: LNAA should be considered; lower ratios may deprive appetite
Valine	0.49	0.49	Abnormal Increment of valine, leucine, or isoleucine (BCAA) will cause catabolism of the other BCAA in the muscle
Isoleucine	0.38	–	
Leucine	0.68	0.68	

AAFCO = The Association of American Feed Control Officials; BCAA = branched chain amino acids; DM = dry matter; NRC = National Research Council; RA = recommended allowance; Trp:LNAA = tryptophan to large neutral amino acid ratio.

¹Recommended Allowance requirements for adult dogs at maintenance, Nutrient Requirements of Dogs and Cats (NRC, 2006).

²Minimum dietary content, AAFCO (2018).

Miniature Dachshunds, Beagles, and Labrador Retrievers as proxies for small, medium, and large dog breeds and found that methionine requirements may differ across breeds or size of dogs and be greater than previously estimated. Thus, given the methods of derivation, single indispensable AA requirements for all dog populations, as presented in AAFCO (2018), may not consider variable AA requirements across dog phenotypes. Moreover, it is widely assumed that endogenous synthesis of dispensable AAs, such as taurine in the dog, is sufficient for meeting metabolic demands. However, recent studies suggest that under some metabolic conditions, dispensable AAs may also be required in diets (Hou et al., 2015). Taurine, as described in this commentary, is a clear example of this paradigm shift. Dietary taurine or the capacity for its adequate endogenous synthesis, especially in circumstances where excessive losses might occur, should be considered in the final formulation of dog foods to decrease the risk of canine DCM.

Nutritionists and regulatory agencies should be aware that, in the spectrum of nutrient requirements, dog populations with higher AA requirements relative to energy intake and other factors could be at a higher risk for a taurine deficiency. More precise categorization of requirements among different canine populations would help us to optimize nutritional

adequacy and decrease risk of diseases, such as DCM, that are possibly linked to nutrient deficiencies.

Effect of Processing on Antinutritional Factors in Plant-Based Ingredients

Just as understanding the inherent nutritional characteristics and the interaction between ingredients is important for preventing nutritional imbalances in pet foods, the effects of processing on these factors are equally important. Raw cereals and legumes contain antinutritional factors such as trypsin inhibitors, phytates, hematoglutinins, and polyphenols that can decrease protein digestion, nutrient absorption, and/or cause illness. Some of these antinutritional factors are thermolabile and, under the right conditions, can be effectively destroyed during the extrusion process improving the overall quality of plant-based ingredients and the final diet (Patterson et al., 2017). Recent reviews across a variety of legumes and legume-derived ingredients show that the activities of trypsin inhibitor, chymotrypsin inhibitor, and hemagglutinating activity were decreased by up to 95% across a variety of thermal treatment conditions, including extrusion (Patterson et al., 2017; Avilés-Gaxiola et al., 2018). Extrusion had modest effects on levels of phytate with reductions ranging from 7% to 26% and varied by legume and extrusion conditions (Patterson




Figure 2. Effect of thermal processing methods on trypsin inhibitor levels (mg/g) soybean kernel. ¹Treatment conditions: None = no treatment; dry extrusion for 25 to 30 sec (1 = 100 °C; 2 = 125 °C; 3 = 140 °C; 4 = 150 °C); wet extrusion for 25 to 30 s with 6% to 8% added moisture (1 = 100 °C; 2 = 125 °C; 3 = 140 °C); micronization with near-infrared rays wavelength of 1.8 to 3.4 μ m for 90 s (1 = 100 °C; 2 = 125 °C; 3 = 140 °C; 4 = 150 °C); microwave roasting at 800 W and 2450 MHz (1 = 1 min [kernel temp = 57 °C], 2 = 2 min [kernel temp = 88 °C], 3 = 3 min [kernel temp = 108 °C], 4 = 4 min [kernel temp = 121 °C], 5 = 5 min [kernel temp = 132 °C]); Autoclaving at 120 °C and 1.2 bars (1 = 10 min, 2 = 20 min, 3 = 30 min). Reprinted with permission from Žilić et al. (2012).

et al., 2017). Figure 2 highlights the variability between processing methods and thermic conditions for decreasing antinutritional factors. For example, when soybeans were subjected to extrusion at increasing temperatures that ranged from 100 to 150 °C, trypsin inhibitor levels were incrementally decreased. At 140 °C, dry extrusion was considerably more effective at decreasing trypsin inhibitors (–91%) compared with wet extrusion (–44%). When the dry extrusion temperature was increased to 150 °C, reductions in trypsin inhibitors were further decreased by 94% (Žilić et al., 2012). Other thermal treatments, such as micronisation, microwave roasting, and autoclaving, also facilitated incremental reductions in trypsin inhibitors with increasing temperatures (Žilić et al., 2012). When formulating foods with higher concentrations of plant-based ingredients, consideration should also be given to the processing methods and the parameters used to effectively optimize the nutritional density and decrease antinutritional factors.

It is important to mention that, while temperature and pressure processing can greatly decrease antinutritional factors, they can also negatively affect bioavailability of AAs. The Maillard reaction is a well-known example of heat-damaged protein (Teodorowicz et al., 2017). In this reaction, lysine interacts with reducing sugars present in the diets forming the Maillard product. The

complex formed can be digested and absorbed by the animal but cannot be utilized for metabolic processes (e.g., protein synthesis). Thus, in heat-damaged proteins, digestibility of AAs can greatly overestimate bioavailability (Moehn et al., 2005). Other products of heat damage on proteins include racemization of AAs (alteration from L to D form) and the formation of cross-linked AAs. Such components can decrease bioavailability of AAs and digestibility of proteins, and their effects on protein quality cannot usually be determined using conventional methods of AA analysis. Pet foods with higher levels of plant-based ingredients may also require optimization of processing methods to maximize their nutritional density and nutrient bioavailability.

Recommendations for Formulating Dog Food With Novel Ingredients

Considering the AA profile of dog foods. Feed formulation for agricultural and companion animals should be based on the ideal protein concept (Baker, 1991; Swanson et al., 2013). The ideal protein is defined as that in which all AAs are in perfect balance compared with the animal's AA requirements (mg/g protein). Hence, all indispensable AAs are equally limiting. However, this is impossible to achieve in practical animal feed formulation, and diets should be formulated considering the first limiting indispensable AA. The first limiting indispensable AA refers to the indispensable AA that is present in the lowest proportion compared with the animal's requirement. By meeting the first indispensable limiting AA requirement, requirements for all other indispensable AAs are also inherently satisfied. Moreover, to avoid the formulation of diets with excessive protein concentration or an excess of indispensable AAs relative to the requirements of dogs, animal nutritionists combine multiple ingredients that are complementary in their AA profiles. Commonly, dog foods are formulated with a higher proportion of animal-derived ingredients, and a lower proportion of plant-based ingredients to meet nutrient recommendations. More recently, however, cereal grains have been removed in some diet formulations or the proportion of animal-based ingredients has been reduced. The production of these types of formulations is often driven by consumer perception, rather than scientific evidence. Allowing consumers to direct the ingredient composition of dog foods, or other pet foods, could perpetuate nutrient deficits that affect the health of animals in the long term.

In the formulation of grain-free pet foods, cereal grains are replaced with alternative ingredient(s). Animal-derived ingredients are expensive relative to plant-based ingredients. Thus, pulses, a subset of legumes, are often used as the replacement. In addition to containing substantial fiber, pulses also contain significant concentrations of protein and are used to partly meet indispensable AA requirements. Of interest, soybean meal and pulses contain 48% and 25% crude protein, respectively, which is substantially greater than the average protein concentration for grains (11%; [Table 1](#)). Although the high-protein content in soybean meal and pulses is indicative of higher concentration of AAs compared with grains, it does not imply AA balance. Soybean meal and pulses are high in lysine (mg/g protein) but low in sulfur AAs (mg/g protein), whereas the reverse is true for cereals. Plant-based ingredients tend to have lower ileal digestibility coefficients for protein compared with protein from animal sources ([FAO and WHO, 1991](#)). Thus, dog foods that contain substantial amounts of pulses, lower proportions of animal-based ingredients, and do not address AA imbalances through the addition of alternate ingredients or fortification, may risk AA deficiencies. To mitigate this risk across the pet food industry and ensure the final pet diets are nutritionally adequate and balanced, it is prudent that the digestibility coefficients of all final pet food products be calculated.

Considering the addition of high-fiber ingredients to dog foods. By definition, dietary fiber is carbohydrates that are resistant to digestion by endogenous enzymes in the gastrointestinal tract ([NRC, 2006](#)). Typical fibers include arabinoxylan, raffinose, inulin, β -glucan, cellulose, and pectin ([NRC, 2006](#)). Common ingredients to increase fiber content in companion animal diets include beet pulp, corn fiber, rice bran, whole grains, and pulse fibers ([de Godoy et al., 2013](#)). Achieving an optimal fiber concentration in canine diets has diverse positive physiological effects in the gastrointestinal tract; for example, higher fermentable fiber intake has been shown to slow the transit time of digesta, increasing satiety of the animal ([Haber et al., 1977](#)). Moreover, high-fiber diets generally have lower energy density making them an important nutritional strategy for controlling body weight ([Johnson et al., 2008](#)) and reducing the incidence of diarrhea ([Homann et al., 1994](#)). Gut health is also improved with higher consumption of fiber; fermentable fiber can act as a prebiotic and increase the population of health-promoting microbiota including lactobacilli and

bifidobacteria ([Roberfroid, 2005](#)). Although not required by AAFCO to fulfill the criteria of “complete and balanced,” fiber is an important component of the diet, and depending on the type of fiber and the amount consumed, fiber can increase the gut health status. Adding the necessary amount and type of fiber in the diet is crucial for optimal dog nutrition.

Despite the benefits of fiber in the diet, fiber can also affect enterohepatic recycling of taurine (discussed above). In monogastric species, including humans, high dietary fermentable fiber may also decrease digestibility and availability of dietary AAs ([Blackburn and Southgate, 1981](#); [Degen et al., 2007](#)) and, in some cases, increase the risk of DCM in dogs fed diets that marginally meet requirements for sulfur AAs. Moreover, higher concentrations of dietary fiber increase the size of the gastrointestinal tract in pigs and poultry ([Nyachoti et al., 2000](#)), increasing nutrient utilization in this organ. It has been determined in pigs that on average the gastrointestinal tract catabolizes 30% of dietary indispensable AAs during absorption, and this utilization represents ~50% for sulfur AAs ([Stoll et al., 1998](#); [Mansilla et al., 2018](#)), further reducing precursor availability for taurine synthesis and increasing the risk for taurine deficiency. For some high-fiber diets, fortification of specific nutrients, including taurine and other sulfur AAs, might be beneficial to avoid nutrient deficiencies.

Compared with the pet food industry, in other industries where high-fiber ingredients (coproducts) are routinely used (e.g., swine industry), the effects of fiber on the absorption of nutrients have been given more attention when formulating diets ([NRC, 2012](#)). For example, highly fermentable fiber in swine diets increases the threonine requirement to compensate for the increase in mucus (mucin protein) production in the intestinal cell lining ([Lien et al., 1997](#); [Mathai et al., 2016](#)). This has underpinned the development of “requirement models” ([NRC, 2012](#)) to tailor nutrient requirements for pigs while accounting for the different nutrient interactions. In contrast, in the pet food industry, the only concentrations of nutrients used for comparison are those recommended by [AAFCO \(2018\)](#). Such recommendations are static and may not encompass all the effects of the different nutrient combinations in the final diet. There is a clear need in companion animal nutrition to improve the understanding of the interactions of different ingredients and how these alter nutrient requirements for different breeds, age, and physiological status of dogs.

Other recent publications highlight the need for careful nutrient formulation. Several recent papers, both original research and reviews, likewise highlight the unknowns surrounding grain-free diets (typically legume or pulse-based, but sometimes also with “exotic” ingredients such as kangaroo, bison, or wild boar) and DCM. For example, [Adin et al. \(2019\)](#) examined 48 dogs of many breeds with diagnosed DCM and having a known diet history. Among grain-free diets being consumed in this study, 1 dog was particularly associated with DCM, possibly underscoring the importance of specific diet formulation. Furthermore, 2 dogs switched from that diet to other grain-free diets showed improvement in their DCM; it is unclear if those dogs were taurine deficient or if they also received taurine and/or carnitine supplementation. This suggests that grain-free composition per se may not be the root cause of DCM. Another recently published case series of 24 Golden Retrievers with DCM and known diet histories were evaluated, and an association between grain-free diets and DCM was suggested ([Kaplan et al., 2018](#)). Most dogs (15 of 24) were fed a single diet which was significantly associated with low blood taurine concentrations, again suggesting that specific diet formulation may play an important role. However, as in the previous study, soluble vs. insoluble fiber concentrations were not available for the diets, nor were taurine, methionine, or cysteine concentrations, meaning that the true nutrient profiles of the diets could not be assessed and reinforcing the point that diet formulation for nutrients—not ingredients—is essential. It also suggests that nutrient requirements may vary widely based on breed, diet, and other phenotypic data. Indeed, most of the dogs with DCM in the previously described study were consuming less energy compared with their predicted requirements ([Kaplan et al., 2018](#)). It also bears pointing out that the numbers in both studies were very low (representing less than 100 DCM-affected dogs between them), which surely represents a fraction of the dogs consuming grain-free, pulse-based diets. A recent thoughtful review supports these conclusions by reiterating the crucial need for plant-based diets for dogs to be formulated with sufficient quantities of bioavailable methionine and cysteine to support adequate taurine synthesis ([Dodd et al., 2018](#)). This can be achieved with the addition of purified AAs and other sources that are readily available ([Gloaguen et al., 2014](#)). Finally, a recent commentary carefully concludes that a true cause-and-effect relationship

between grain-free diets and DCM has not been proven, and other factors may ultimately be more important ([Freeman et al., 2018](#)). Taken together, these recent publications may point to faulty nutrient formulation in some, but not all, grain-free diets.

CONCLUSIONS

Recently, it has been suggested that pulse ingredients in commercial dog foods are associated with a limited number of cases of DCM. Although pulse ingredients have been implicated for having negative effects on the taurine status in dogs (deficiency of which is a known cause of canine DCM) based on the available evidence, the relationship between pulses and canine DCM remains undefined. However, the FDA statement may harm consideration of protein alternatives, such as pulses, as quality ingredients in pet foods and undermine attempts to diversify ingredients used across the food chain as the global population continues to grow. Ingredients do not represent the nutritional composition of the diet, and therefore, nutrient deficiencies should not be attributed to individual ingredients. The authors of this commentary recognize the important role of endogenous, and perhaps exogenous, taurine in the prevention of DCM in some dogs. The assurance of appropriate concentrations of all indispensable sulfur AAs, including methionine and cysteine, is crucial for ensuring adequate endogenous synthesis of taurine and to meet the metabolic demands of dogs. Additional dietary factors, such as methyl donors required for sulfur AA metabolism, carnitine for energy production in muscle, and dietary fiber, as well as animal factors, such as breed, size, and health status, should also be investigated when nutrient deficiency-related DCM is suspected.

It is the responsibility of animal nutritionists to formulate balanced diets for dogs, and other animals, by looking beyond the goal of meeting AAFCO recommendations or satisfying unsubstantiated market trends. Pulses and other plant-based ingredients can be used to formulate nutritionally adequate dog foods, and final product formulations should be assessed for nutrient balance and bioavailability, especially when using a limited number of ingredients. Although dietary factors are important in the prevention of sulfur AA deficiency and development of DCM, empirical data and mechanistic studies are required to better understand the indispensable AA requirements of dogs and preventing DCM. In diets that contain high concentrations of dietary fiber, compensative inclusion

of dietary indispensable sulfur AAs, including exogenous taurine, might be required to offset the possibility of increased fecal excretion or microbial assimilation of taurine in the large intestine. Processing conditions may also require adjustments to ensure the presence or effects of antinutritional factors are minimized and nutrient bioavailability is not compromised. Greater awareness of AA balance is crucial for ensuring that AA requirements are met for dogs consuming static diets.

LITERATURE CITED

- AAFCO. 2018. Association of American feed control officials. Official Publication Association of American Feed Control Inc., Oxford.
- Adin, D., T. C. DeFrancesco, B. Keene, S. Tou, K. Meurs, C. Atkins, B. Aona, K. Kurtz, L. Barron, and K. Saker. 2019. Echocardiographic phenotype of canine dilated cardiomyopathy differs based on diet type. *J. Vet. Card.* 21:1–9. doi: 10.1016/j.jvc.2018.11.002.
- Alroy, J., J. E. Rush, L. Freeman, M. S. Amarendhra Kumar, A. Karuri, K. Chase, and S. Sarkar. 2000. Inherited infantile dilated cardiomyopathy in dogs: genetic, clinical, biochemical, and morphologic findings. *Am. J. Med. Genet.* 95:57–66. doi:10.1002/1096-8628(20001106)95:1<57::AID-AJMG12>3.0.CO;2-O
- Arslan, C. 2006. L-Carnitine and its use as a feed additive in poultry feeding a review. *Revue Med Vet.* 157:134–142.
- Avilés-Gaxiola, S., C. Chuck-Hernández, and S. O. Serna Saldívar. 2018. Inactivation methods of trypsin inhibitor in legumes: a review. *J. Food Sci.* 83:17–29. doi: 10.1111/1750-3841.13985
- Backus, R. C., K. S. Ko, A. J. Fascetti, M. D. Kittleson, K. A. Macdonald, D. J. Maggs, J. R. Berg, and Q. R. Rogers. 2006. Low plasma taurine concentration in newfoundland dogs is associated with low plasma methionine and cyst(e)ine concentrations and low taurine synthesis. *J. Nutr.* 136:2525–2533. doi: 10.1093/jn/136.10.2525
- Baker, D. H. 1986. Problems and pitfalls in animal experiments designed to establish dietary requirements for essential nutrients. *J. Nutr.* 116:2339–2349. doi: 10.1093/jn/116.12.2339
- Baker, D. H. 1991. Comparative nutrition of cats and dogs. *Annu. Rev. Nutr.* 11:239–263. doi: 10.1146/annurev.nu.11.070191.001323
- Bakker, A. J., and H. M. Berg. 2002. Effect of taurine on sarcoplasmic reticulum function and force in skinned fast-twitch skeletal muscle fibres of the rat. *J. Physiol.* 538:185–194. doi: 10.1113/jphysiol.2001.012872
- Bélangier, M. C., M. Ouellet, G. Queney, and M. Moreau. 2005. Taurine-deficient dilated cardiomyopathy in a family of golden retrievers. *J. Am. Anim. Hosp. Assoc.* 41:284–291. doi: 10.5326/0410284
- Blackburn, N. A., and Southgate D. A. T. 1981. Protein digestibility and absorption: effects of fibre and the extent of individual variation. Joint FAO/WHO/UNU Expert Consultation on Energy and Protein Requirements Rome; October 5–17.
- Borgarelli, M., R. A. Santilli, D. Chiavegato, G. D’Agnolo, R. Zanatta, A. Mannelli, and A. Tarducci. 2006. Prognostic indicators for dogs with dilated cardiomyopathy. *J. Vet. Intern. Med.* 20:104–110. doi: 10.1111/j.1939-1676.2006.tb02829.x
- Butterwick, R. F., P. J. Markwell, and C. J. Thorne. 1994. Effect of level and source of dietary fiber on food intake in the dog. *J. Nutr.* 124(12 Suppl):2695S–2700S. doi: 10.1093/jn/124.suppl_12.2695S.
- Cattanach, B. M., J. Dukes-McEwan, P. R. Wotton, H. M. Stephenson, and R. M. Hamilton. 2015. A pedigree-based genetic appraisal of boxer ARVC and the role of the striatin mutation. *Vet. Rec.* 176:492. doi: 10.1136/vr.102821.
- Columbus, D., and C. F. de Lange. 2012. Evidence for validity of ileal digestibility coefficients in monogastrics. *Br. J. Nutr.* 108 (Suppl 2):S264–S272. doi: 10.1017/S0007114512002334.
- Degen, L., V. Halas, and L. Babinszky. 2007. Effect of dietary fibre on protein and fat digestibility and its consequences on diet formulation for growing and fattening pigs: a review. *Act. Agr. Scand. A-AN.* 57:1–9. doi: 10.1080/09064700701372038
- Dodd, S. A. S., J. L. Adolphe, and A. Verbrugghe. 2018. Plant-based diets for dogs. *J. Am. Vet. Med. Assoc.* 253:1425–1432. doi: 10.2460/javma.253.11.1425
- Dutton, E., and J. López-Alvarez. 2018. An update on canine cardiomyopathies – is it all in the genes? *J. Small. Anim. Pract.* 59:455–464. doi: 10.1111/jsap.12841
- FAO. 1991. Food and agriculture organization of the United Nations. Protein quality evaluation. Report of Joint FAO/WHO, Expert Consultation, Rome, Italy.
- Fascetti, A. J., J. R. Reed, Q. R. Rogers, and R. C. Backus. 2003. Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997–2001). *J. Am. Vet. Med. Assoc.* 223:1137–1141. doi: 10.2460/javma.2003.223.1137
- FDA, Center for Veterinary Medicine. 2018. FDA investigating potential connection between diet and cases of canine heart disease. <https://www.fda.gov/animalveterinary/new-sevents/cvmupdates/ucm613305.htm> (Accessed 12 July 2018.)
- Flanagan, J. L., P. A. Simmons, J. Vehige, M. D. Willcox, and Q. Garrett. 2010. Role of carnitine in disease. *Nutr. Metab. (Lond).* 7:30. doi: 10.1186/1743-7075-7-30
- Freeman, L. M., and J. E. Rush. 2006. Cardiovascular diseases: nutritional modulation. In: P. Pibot, V. Biourge, and D. Elliott, editors, *Encyclopedia of canine clinical nutrition*. Aniwa SAS, Aimargues. p. 316–347.
- Freeman, L. M., K. E. Michel, D. J. Brown, P. M. Kaplan, M. E. Stamoulis, S. L. Rosenthal, B. W. Keene, and J. E. Rush. 1996. Idiopathic dilated cardiomyopathy in dalmatians: nine cases (1990–1995). *J. Am. Vet. Med. Assoc.* 209:1592–1596.
- Freeman, L. M., J. A. Stern, R. Fries, D. B. Adin, and J. E. Rush. 2018. Diet-associated dilated cardiomyopathy in dogs: what do we know? *J. Am. Vet. Med. Assoc.* 253:1390–1394. doi: 10.2460/javma.253.11.1390
- Glauguen, M., N. Le Floc’h, E. Corrent, Y. Primot, and J. van Milgen. 2014. The use of free amino acids allows formulating very low crude protein diets for piglets. *J. Anim. Sci.* 92:637–644. doi: 10.2527/jas.2013-6514
- de Godoy, M. R., K. R. Kerr, and G. C. Fahey, Jr. 2013. Alternative dietary fiber sources in companion animal nutrition. *Nutrients* 5:3099–3117. doi: 10.3390/nu5083099.

- Haber, G. B., K. W. Heaton, D. Murphy, and L. F. Burroughs. 1977. Depletion and disruption of dietary fibre. Effects on satiety, plasma-glucose, and serum-insulin. *Lancet* 2:679–682. doi: 10.1016/S0140-6736(77)90494-9
- Homann, H. H., M. Kemen, C. Fuessenich, M. Senkal, and V. Zumtobel. 1994. Reduction in diarrhea incidence by soluble fiber in patients receiving total or supplemental enteral nutrition. *JPEN. J. Parenter. Enteral Nutr.* 18:486–490. doi: 10.1177/0148607194018006486
- Hoppel, C. 2003. The role of carnitine in normal and altered fatty acid metabolism. *Am. J. Kidney Dis.* 41:S4–12. doi: 10.1016/S0272-6386(03)00112-4
- Hou, Y., Y. Yin, and G. Wu. 2015. Dietary essentiality of “nutritionally non-essential amino acids” for animals and humans. *Exp. Biol. Med.* (Maywood). 240:997–1007. doi: 10.1177/1535370215587913
- Hurrell, R. F., P. A. Finot, and J. E. Ford. 1983. Storage of milk powders under adverse conditions. I. Losses of lysine and of other essential amino acids as determined by chemical and microbiological methods. *Br. J. Nutr.* 49:343–354. doi: 10.1079/BJN19830043
- Huxtable, R. J. 1992. Physiological actions of taurine. *Physiol. Rev.* 72:101–163. doi: 10.1152/physrev.1992.72.1.101
- Ito, T., S. W. Schaffer, and J. Azuma. 2012. The potential usefulness of taurine on diabetes mellitus and its complications. *Amino Acids* 42:1529–1539. doi: 10.1007/s00726-011-0883-5
- Johnson, L., A. P. Mander, L. R. Jones, P. M. Emmett, and S. A. Jebb. 2008. Energy-dense, low-fiber, high-fat dietary pattern is associated with increased fatness in childhood. *Am. J. Clin. Nutr.* 87:846–854. doi: 10.1093/ajcn/87.4.846
- Johnson, M. L., C. M. Parsons, G. C. Fahey, Jr, N. R. Merchen, and C. G. Aldrich. 1998. Effects of species raw material source, ash content, and processing temperature on amino acid digestibility of animal by-product meals by cecectomized roosters and ileally cannulated dogs. *J. Anim. Sci.* 76:1112–1122. doi: 10.2527/1998.7641112x
- Kaplan, J. L., J. A. Stern, A. J. Fascetti, J. A. Larsen, H. Skolnik, G. D. Peddle, R. D. Kienle, A. Waxman, M. Cocchiario, C. T. Gunther-Harrington, et al. 2018. Taurine deficiency and dilated cardiomyopathy in golden retrievers fed commercial diets. *PLoS One* 13:e0209112. doi: 10.1371/journal.pone.0209112
- Keene, B. W., D. P. Panciera, C. E. Atkins, V. Regitz, M. J. Schmidt, and A. L. Shug. 1991. Myocardial L-carnitine deficiency in a family of dogs with dilated cardiomyopathy. *J. Am. Vet. Med. Assoc.* 198:647–650.
- Kim, S. W., Q. R. Rogers, and J. G. Morris. 1996a. Dietary antibiotics decrease taurine loss in cats fed a canned heat-processed diet. *J. Nutr.* 126:509–515. doi: 10.1093/jn/126.2.509
- Kim, S. W., Q. R. Rogers, and J. G. Morris. 1996b. Maillard reaction products in purified diets induce taurine depletion in cats which is reversed by antibiotics. *J. Nutr.* 126:195–201. doi: 10.1093/jn/126.1.195
- Kittleson, M. D., B. Keene, P. D. Pion, and C. G. Loyer. 1997. Results of the multicenter spaniel trial (MUST): taurine- and carnitine-responsive dilated cardiomyopathy in american cocker spaniels with decreased plasma taurine concentration. *J. Vet. Intern. Med.* 11:204–211. doi: 10.1111/j.1939-1676.1997.tb00092.x
- Ko, K. S., R. C. Backus, J. R. Berg, M. W. Lame, and Q. R. Rogers. 2007. Differences in taurine synthesis rate among dogs relate to differences in their maintenance energy requirement. *J. Nutr.* 137:1171–1175. doi: 10.1093/jn/137.5.1171
- Kramer, G. A., M. D. Kittleson, P. R. Fox, J. Lewis, and P. D. Pion. 1995. Plasma taurine concentrations in normal dogs and in dogs with heart disease. *J. Vet. Intern. Med.* 9:253–258. doi: 10.1111/j.1939-1676.1995.tb01076.x
- Lien, K. A., W. C. Sauer, and M. Fenton. 1997. Mucin output in ileal digesta of pigs fed a protein-free diet. *Z. Ernahrungswiss.* 36:182–190. doi: 10.1007/BF01611398
- Linder, D., and M. Mueller. 2014. Pet obesity management: beyond nutrition. *Vet. Clin. North Am. Small Anim. Pract.* 44:789–806, vii. doi: 10.1016/j.cvsm.2014.03.004
- Mansilla, W. D., K. E. Silva, C. Zhu, C. M. Nyachoti, J. K. Htoo, J. P. Cant, and C. F. M. de Lange. 2018. Ammonia-nitrogen added to low-crude-protein diets deficient in dispensable amino acid-nitrogen increases the net release of alanine, citrulline, and glutamate post-splanchnic organ metabolism in growing pigs. *J. Nutr.* 148:1081–1087. doi: 10.1093/jn/nxy076
- Marinangeli, C. P. F., J. Curran, S. I. Barr, J. Slavin, S. Puri, S. Swaminathan, L. Tapsell, and C. A. Patterson. 2017. Enhancing nutrition with pulses: defining a recommended serving size for adults. *Nutr. Rev.* 75:990–1006. doi: 10.1093/nutrit/nux058
- Maron, B. J., J. A. Towbin, G. Thiene, C. Antzelevitch, D. Corrado, D. Arnett, A. J. Moss, C. E. Seidman, J. B. Young. 2006. Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation.* 113:1807–1816. doi: 10.1161/CIRCULATIONAHA.106.174287
- Marshall, H. F., K. C. Chang, K. S. Miller, and L. D. Satterlee. 1982. Sulfur amino acid stability: effects of processing on legume proteins. *J. Food Sci.* 47:1170–4. doi: 10.1111/j.1365-2621.1982.tb07642.x
- Martin, M. W., M. J. Stafford Johnson, and B. Celona. 2009. Canine dilated cardiomyopathy: a retrospective study of signalment, presentation and clinical findings in 369 cases. *J. Small Anim. Pract.* 50:23–29. doi: 10.1111/j.1748-5827.2008.00659.x
- Mathai, J. K., J. K. Htoo, J. E. Thomson, K. J. Touchette, and H. H. Stein. 2016. Effects of dietary fiber on the ideal standardized ileal digestible threonine:lysine ratio for twenty-five to fifty kilogram growing gilts. *J. Anim. Sci.* 94:4217–4230. doi: 10.2527/jas.2016-0680
- Menniti, M. F., G. M. Davenport, A. K. Shoveller, J. P. Cant, and V. R. Osborne. 2014. Effect of graded inclusion of dietary soybean meal on nutrient digestibility, health, and metabolic indices of adult dogs. *J. Anim. Sci.* 92:2094–2104. doi: 10.2527/jas.2013-7226
- Merheb, M., R. T. Daher, M. Nasrallah, R. Sabra, F. N. Ziyadeh, and K. Barada. 2007. Taurine intestinal absorption and renal excretion test in diabetic patients: a pilot study. *Diabetes Care* 30:2652–2654. doi: 10.2337/dc07-0872
- Meurs, K. M., S. Lahmers, B. W. Keene, S. N. White, M. A. Oyama, E. Mauceli, and K. Lindblad-Toh. 2012. A splice site mutation in a gene encoding for PDK4, a mitochondrial protein, is associated with

- the development of dilated cardiomyopathy in the doberman pinscher. *Hum. Genet.* 131:1319–1325. doi: 10.1007/s00439-012-1158-2
- Meurs, K. M., J. A. Stern, D. D. Sisson, M. D. Kittleson, S. M. Cunningham, M. K. Ames, C. E. Atkins, T. DeFrancesco, T. E. Hodge, B. W. Keene, et al. 2013. Association of dilated cardiomyopathy with the striatin mutation genotype in boxer dogs. *J. Vet. Intern. Med.* 27:1437–1440. doi: 10.1111/jvim.12163
- Moehn, S., R. F. Bertolo, P. B. Pencharz, and R. O. Ball. 2005. Development of the indicator amino acid oxidation technique to determine the availability of amino acids from dietary protein in pigs. *J. Nutr.* 135:2866–2870. doi: 10.1093/jn/135.12.2866
- Moise, N. S., L. M. Pacioretty, F. A. Kallfelz, M. H. Stipanuk, J. M. King, and R. F. Gilmour, Jr. 1991. Dietary taurine deficiency and dilated cardiomyopathy in the fox. *Am. Heart J.* 121:541–547. doi: 10.1016/0002-8703(91)90724-V
- Monnet, E., E. C. Orton, M. Salman, and J. Boon. 1995. Idiopathic dilated cardiomyopathy in dogs: survival and prognostic indicators. *J. Vet. Intern. Med.* 9:12–17. doi: 10.1111/j.1939-1676.1995.tb03266.x
- Nardelli, T. R., R. A. Ribeiro, S. L. Balbo, E. C. Vanzela, E. M. Carneiro, A. C. Boschero, and M. L. Bonfleur. 2011. Taurine prevents fat deposition and ameliorates plasma lipid profile in monosodium glutamate-obese rats. *Amino Acids* 41:901–908. doi: 10.1007/s00726-010-0789-7
- NRC, National Research Council. 2006. Nutrient requirements of dogs and cats. 10th ed. Natl. Acad. Press, Washington, DC.
- NRC, National Research Council. 2012. Nutrient requirements of swine. 11th ed. Natl. Acad. Press, Washington, DC.
- Nyachoti, C. M., C. F. M. de Lange, B. W. McBride, S. Leeson, and H. Schulze. 2000. Dietary influence on organ size and in vitro oxygen consumption by visceral organs of growing pigs. *Livest Prod Sci.* 65:229–237. doi: 10.1016/S0301-6226(00)00157-3
- O'Máille, E. R., T. G. Richards, and A. H. Short. 1965. Acute taurine depletion and maximal rates of hepatic conjugation and secretion of cholic acid in the dog. *J. Physiol.* 180:67–79.
- Owczarek-Lipska, M., T. B. Mausberg, H. Stephenson, J. Dukes-McEwan, G. Wess, and T. Leeb. 2013. A 16-bp deletion in the canine PDK4 gene is not associated with dilated cardiomyopathy in a European cohort of doberman pinschers. *Anim. Genet.* 44:239. doi: 10.1111/j.1365-2052.2012.02396.x
- Patterson, C. A., J. Curran, and T. Der. 2017. Effect of processing on antinutrient compounds in pulses. *Cereal Chemistry.* 94:2–10. doi: 10.1094/CCHEM-05-16-0144-FI
- Pion, P. D., M. D. Kittleson, Q. R. Rogers, and J. G. Morris. 1987. Myocardial failure in cats associated with low plasma taurine: a reversible cardiomyopathy. *Science* 237:764–768. doi: 10.1126/science.3616607
- Pion, P. D., S. L. Sanderson, and M. D. Kittleson. 1998. The effectiveness of taurine and levocarnitine in dogs with heart disease. *Vet. Clin. North Am. Small Anim. Pract.* 28:1495–514, ix. doi: 10.1016/S0195-5616(98)50134-9
- Rășanu, T., M. Mehedinți-Hâncu, M. Alexianu, T. Mehedinți, E. Gheorghe, and I. Damian. 2012. Carnitine deficiency. *Rom. J. Morphol. Embryol.* 53:203–206.
- Rice, J. E., and S. L. Ihle. 1994. Effects of diet on fecal occult blood testing in healthy dogs. *Can. J. Vet. Res.* 58:134–137.
- Roberfroid, M. B. 2005. Introducing inulin-type fructans. *Br. J. Nutr.* 93 (Suppl 1):S13–S25. doi: 10.1079/BJN20041350
- Robinson, J. L., S. V. Harding, J. A. Brunton, and R. F. Bertolo. 2016a. Dietary methyl donors contribute to whole-body protein turnover and protein synthesis in skeletal muscle and the jejunum in neonatal piglets. *J. Nutr.* 146:2007–2012. doi: 10.3945/jn.115.226035
- Robinson, J. L., L. E. McBreaity, E. W. Randell, J. A. Brunton, and R. F. Bertolo. 2016b. Restriction of dietary methyl donors limits methionine availability and affects the partitioning of dietary methionine for creatine and phosphatidylcholine synthesis in the neonatal piglet. *J. Nutr. Biochem.* 35:81–86. doi: 10.1016/j.jnutbio.2016.07.001
- Sanderson, S. L., K. L. Gross, P. N. Ogburn, C. Calvert, G. Jacobs, S. R. Lowry, K. A. Bird, L. A. Koehler, and L. L. Swanson. 2001. Effects of dietary fat and L-carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am. J. Vet. Res.* 62:1616–1623. doi:10.2460/ajvr.2001.62.1616
- Schaffer, S. W., C. J. Jong, K. C. Ramila, and J. Azuma. 2010. Physiological roles of taurine in heart and muscle. *J. Biomed. Sci.* 17 (Suppl 1):S2. doi: 10.1186/1423-0127-17-S1-S2
- Sisson, D. D., W. P. Thomas, and B. W. Keene. 2000. Primary myocardial disease in the dog. In: S. J. Ettinger, and E. C. Feldman, editors, *Textbook of veterinary internal medicine. Diseases of the dog and cat.* 5th ed. WB Saunders Co., Philadelphia. p. 874–895.
- Spitze, A. R., D. L. Wong, Q. R. Rogers, and A. J. Fascetti. 2003. Taurine concentrations in animal feed ingredients; cooking influences taurine content. *J. Anim. Physiol. Anim. Nutr. (Berl.)* 87:251–262. doi: 10.1046/j.1439-0396.2003.00434.x
- Stoll, B., J. Henry, P. J. Reeds, H. Yu, F. Jahoor, and D. G. Burrin. 1998. Catabolism dominates the first-pass intestinal metabolism of dietary essential amino acids in milk protein-fed piglets. *J. Nutr.* 128:606–614. doi: 10.1093/jn/128.3.606
- Story, J. A., and D. Kritchevsky. 1978. Bile acid metabolism and fiber. *Am. J. Clin. Nutr.* 31 (10 Suppl):S199–S202. doi: 10.1093/ajcn/31.10.S199
- Swanson, K. S., R. A. Carter, T. P. Yount, J. Aretz, and P. R. Buff. 2013. Nutritional sustainability of pet foods. *Adv. Nutr.* 4:141–150. doi: 10.3945/an.112.003335
- Teodorowicz, M., J. van Neerven, and H. Savelkoul. 2017. Food processing: the influence of the Maillard reaction on immunogenicity and allergenicity of food proteins. *Nutrition* 9:835. doi: 10.3390/nu9080835
- Tôrres, C. L., R. C. Backus, A. J. Fascetti, and Q. R. Rogers. 2003. Taurine status in normal dogs fed a commercial diet associated with taurine deficiency and dilated cardiomyopathy. *J. Anim. Physiol. Anim. Nutr. (Berl.)* 87:359–372. doi:10.1046/j.1439-0396.2003.00446.x
- Tosh, S.M., and S. Yada. 2010. Dietary fibres in pulse seeds and fractions: characterization, functional attributes, and applications. *Food Res. Int.* 43:450–460. doi: 10.1016/j.foodres.2009.09.005
- Werner, P., M. G. Raducha, U. Prociuk, M. M. Sleeper, T. J. Van Winkle, and P. S. Henthorn. 2008. A novel locus for dilated cardiomyopathy maps to canine

- chromosome 8. *Genomics* 91:517–521. doi: 10.1016/j.ygeno.2008.03.007
- Worden, J. A., and M. H. Stipanuk. 1985. A comparison by species, age and sex of cysteinesulfinatase activity and taurine concentration in liver and brain of animals. *Comp. Biochem. Physiol.* 82B:233–239. doi: 10.1016/0305-0491(85)90232-9
- Yamka, R. M., U. Jamikorn, A. D. True, and D. L. Harmon. 2003. Evaluation of soybean meal as a protein source in canine foods. *Anim. Feed Sci. Technol.* 109:121–132. doi: 10.1016/S0377-8401(03)00203-7
- Žilić, S., I. Bozović, and V. H. T. Šukalović. 2012. Thermal inactivation of soybean bioactive proteins. *Int. J. Food Eng.* 8:1556–3758. doi:10.1515/1556-3758.2521

From: [Palmer, Lee Anne](#)
To: [Murphy, Jeanette](#); [Norris, Anne](#); [Hartogenesis, Martine](#)
Cc: [Forfa, Tracey](#)
Subject: RE: DCM roll-out timing question?
Date: Thursday, June 20, 2019 10:44:49 AM

(b) (5)

I don't have slides on research gaps, and I think the other slides I would have are going to be shared with the update. Others may have thoughts...(good luck with this).

From: Murphy, Jeanette
Sent: Thursday, June 20, 2019 10:37 AM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Cc: Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>
Subject: DCM roll-out timing question?

Greetings Ladies

I am at a training this week and am with one of the ORA division reps who got Dave Rotstein's email about contacting firms and timing.

I took from Dave's email that the roll-out (b) (5).

I am speaking at a conference next Friday (conference is Thurs/Fri) about FSMA updates. The general focus of the conference is research in rendering and pet food for food safety. I anticipate some of the firms that are listed in the complaint release will be there in addition to members of PFI's nutrition sub-committee who has been working on the DCM issue.

Do you have a slide or two I can incorporate on research gaps in DCM or about current science you want shared. And any side bar talking points you want me to have if asked questions.

(b) (6)

From: [Edwards, David](#)
To: [Hartogenesis, Martine](#)
Cc: [Norris, Anne](#); [Murphy, Jeanette](#)
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference
Date: Friday, November 30, 2018 9:12:00 AM

Bill is willing to be on the panel. He is going to help with the AAFCO labeling workshop as well.

Thanks,
Dave

From: Hartogenesis, Martine
Sent: Thursday, November 29, 2018 9:14 PM
To: Edwards, David <David.Edwards@fda.hhs.gov>
Cc: Norris, Anne <Anne.Norris@fda.hhs.gov>; Murphy, Jeanette <Jenny.Murphy@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Dave,

My apologies as I missed your reply message back (somehow I am not getting all my emails).
Anyway, they are still interested in a CVM rep and I think Bill B would be an excellent choice.

If that works, do you want to ask him? I can ask, but let me know what works best for you.

Thanks again!

Martine

From: Edwards, David <David.Edwards@fda.hhs.gov>
Date: November 9, 2018 at 1:45:59 PM EST
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

I know that AAFCO is putting on a pet food labeling workshop at the Forum as well, so there will likely be some from FDA going to participate in that (Bill, maybe). Should we try to get a 2 for 1 and have him go? Unless you are wanting to go.

Dave

From: Hartogenesis, Martine
Sent: Friday, November 09, 2018 1:40 PM
To: Edwards, David <David.Edwards@fda.hhs.gov>

Subject: FW: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Dave,

See the request below. Looks like the conference is in Kansas City, MO: <https://www.petfoodforumevents.com/>. I could probably attend, but others may want to do it (Ok with me).

We used to bring speaking requests to CEB, but I haven't seen any of those discussions recently. Let me know your thoughts and I am happy to send to Susan DeWitt if you like.

Martine

From: Debbie Phillips <DPhillips@wattglobal.com>
Sent: Friday, November 09, 2018 12:32 PM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Cc: Norris, Anne <Anne.Norris@fda.hhs.gov>
Subject: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Dr. Hartogenesis:

Thank you again for participating in our webinar in September on the atypical cases of canine DCM and their possible link to certain pet diets. This continues to be a topic of conversation and concern in the industry, as you probably are aware, so we are planning a follow-up/update during our annual conference, Petfood Forum, in late April/early May. Could you or one of your colleagues please consider serving as a panel member for this discussion?

The session is currently scheduled for the afternoon of Wednesday, May 1. I know that seems a long way off, but we prefer to issue invitations in advance, as opposed to the hasty invitation for the webinar!

Other panel members will likely include Dr. Greg Aldrich as moderator (who also participated in the webinar, as you know), plus Dr. Jennifer Adolphe, nutrition manager for Petcurean Pet Food; Dr. Kate Shoveller, assistant professor at the University of Guelph; and Dr. Chris Marinangeli, director of nutrition, scientific and regulatory affairs for Pulse Canada.

In case you are not familiar with Petfood Forum, we just held our 26th edition this past April. It is the only event of its kind for the global pet food industry, drawing more than 3,000 people each year from pet food companies around the world, plus from retailers and related businesses, academia and regulatory organizations, such as AAFCO. In addition to education (concurrent scientific tracks plus panel discussions, general sessions, keynotes and other), it includes a trade show featuring the industry's leading suppliers of ingredients, equipment, packaging materials, testing and other services. This year's show had over 400 booths with more than 250 exhibiting companies.

We offer an honorarium to our speakers and panel members, cover their hotel costs and conference registration and reimburse all other travel expenses.

Please let me know if you have questions about Petfood Forum or this panel discussion. Thank you in advance for considering the request!

Sincerely
Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: + (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13

Petfood Forum China 2019: August 20

Petfood R&D Showcase: October 2019

Visit www.PetfoodForumEvents.com

From: [Rotstein, David](#)
To: [Hartogensis, Martine](#)
Subject: RE: logistics-DCM Firm Contact -Quick List 6-18-2019.xlsx
Date: Tuesday, June 18, 2019 2:10:52 PM
Attachments: [image001.png](#)
[image002.jpg](#)
[image003.jpg](#)
[image004.jpg](#)
[image005.jpg](#)
[image006.jpg](#)

Martine,

I'm still confused on that part—it seems like [REDACTED] (b) (5)

I think we can figure out the logistics for the calls. I think having a two day window—one day to hopefully finish and a snow day for firms that we can't get ahold of.

As for the Divisions, [REDACTED] (b) (5)

I do have one Division that has not responded to me with a second email request. If I don't hear from them by tomorrow, I'll go the person above them about it.

I do have thoughts on the call scheduling including a call-in line for the whole day (but I'm worried if we go over or a firm calls in early).

Dave

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/CERT
7519 Standish Place
[REDACTED] (b) (6) (BB)



This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: Hartogensis, Martine
Sent: Tuesday, June 18, 2019 1:49 PM
To: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: logistics-DCM Firm Contact -Quick List 6-18-2019.xlsx

Thanks Dave! [REDACTED] (b) (5)
[REDACTED]
[REDACTED]

Hopefully we can clarify today.

Thanks again!

Martine

From: Rotstein, David
Sent: Tuesday, June 18, 2019 11:39 AM
To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Cc: Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: logistics-DCM Firm Contact -Quick List 6-18-2019.xlsx

Martine,

I'm still awaiting word from CA about two of the firms, but here is the breakdown.

As you can see HAF5E has a the bulk of firms.

Here are some thoughts for discussion:

[REDACTED] (b) (5)
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]

Dave

From: [Palmer, Lee Anne](#)
To: [Murphy, Jeanette](#); [Hartogenesis, Martine](#)
Subject: RE: Meeting with PFI - thoughts, invitees?
Date: Monday, April 08, 2019 12:42:56 PM

One more piece to this – Jen Jones is giving an FDA-wide presentation to the White Oak Veterinarians group on 4/17/19 at noon. Just a heads up, since a wider presentation sometimes grows legs.

From: Palmer, Lee Anne
Sent: Monday, April 8, 2019 11:58 AM
To: Murphy, Jeanette <Jenny.Murphy@fda.hhs.gov>; Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Subject: RE: Meeting with PFI - thoughts, invitees?

Hi Jenny – I defer to Martine, [REDACTED] (b) (5)
[REDACTED] - a collaboration would be good to pursue. I think an initial meeting to explore ways we can collaborate would be a nice way to begin. I this is going to take industry (PFI nutrition sub-committee group), FDA and our academic partners to really get to the bottom of it.

[REDACTED] (b) (5)
[REDACTED]
[REDACTED]
[REDACTED]
[REDACTED]

It may take some discussion and planning first with them before we can provide anything. I could do either week, but hoping on right on the 22nd, will just be getting back from leave.

Thanks, Lee Anne

From: Murphy, Jeanette
Sent: Monday, April 8, 2019 11:39 AM
To: Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Subject: Meeting with PFI - thoughts, invitees?

LeeAnne and Martine,

My apologies for getting this to you so late.

I met with PFI 2 weeks ago to talk about the work of their nutrition sub-committee and there request to get together with our DCM team.

Their nutrition sub-committee is

(b) (4), (b) (5)

(b) (4), (b) (5)

I told them that I was not sure

(b) (5), (b) (4)

(b) (5), (b) (4)

I am happy to help set things up or we can potentially get Mia to help schedule/coordinate.

I don't know who all from CVM should be invited but know there are a lot of folks working on the issue.

Thoughts?

Jenny

Jenny Murphy
Office of Surveillance and Compliance
Center for Veterinary Medicine
7519 Standish Place
Rockville, MD 20855
Telephone: [240-402-6246](tel:240-402-6246)
Email: jenny.murphy@fda.hhs.gov

From: [Conway, Charlotte](#)
To: [Murphy, Jeanette](#)
Subject: RE: SME assistance for meeting with PFI and Tim next week
Date: Thursday, April 11, 2019 8:44:00 AM

10-4

From: Murphy, Jeanette
Sent: Wednesday, April 10, 2019 9:26 PM
To: Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>; Lambkin, Sonya <Sonya.Lambkin@fda.hhs.gov>
Cc: Large, Mabelle <Mabelle.Large@fda.hhs.gov>
Subject: SME assistance for meeting with PFI and Tim next week

Greetings LeeAnne, Charlotte, and Sonya,

PFI is coming in next week, Wed 4/17 (2-3:00 p.m.), to have their initial meeting with Tim. This will serve as both a meet and greet and a chance to discuss some issues.

PFI is only bringing in 3 people (b) (6). PFI has proposed the following agenda topics, and Tim has recommend each of you attend to help discuss topics below. While I know there are a lot of people who work on all of these, probably best to keep the invite list small to not totally outnumber the PFI staff. (FYI...I will be on leave next week and will not be in attendance)

In person attendance likely preferred. Mia will provide you with the calendar invite.

Please let me or Mia know if you have any concerns/conflicts and will not be able to make it.

Thanks,
Jenny

Agenda Items:

(b) (4), (b) (5)

Jenny Murphy
Office of Surveillance and Compliance
Center for Veterinary Medicine
7519 Standish Place
Rockville, MD 20855
Telephone: **240-402-6246**
Email: jenny.murphy@fda.hhs.gov

High Potential for Selenium Biofortification of Lentils (*Lens culinaris* L.)

DIL THAVARAJAH, JAMIE RUSZKOWSKI, AND ALBERT VANDENBERG*

Crop Development Centre, College of Agriculture and Bioresources, University of Saskatchewan,
 51 Campus Drive, Saskatoon, Saskatchewan S7N 5A8, Canada

Beneficial forms of selenium (Se) and their impact on human health are a global topic of interest in public health. We are studying the genetic potential for Se biofortification of pulse crops to improve human nutrition. Lentils (*Lens culinaris* L.) are an important protein and carbohydrate food and are a valuable source of essential dietary components and trace elements. We analyzed the total Se concentration of 19 lentil genotypes grown at eight locations for two years in Saskatchewan, Canada. We observed significant genotypic and environmental variation in total Se concentration in lentils and that total Se concentration in lentils ranged between 425 and 673 $\mu\text{g kg}^{-1}$, providing 77–122% of the recommended daily intake in 100 g of dry lentils. Over 70% of the Se was present as selenomethionine (SeMet) with a smaller fraction (<20%) as inorganic Se and very small amounts as selenocysteine (SeCys). We found that soils from the locations where the lentils were grown were rich in Se (37–301 $\mu\text{g kg}^{-1}$) and that lentils grown in Saskatchewan have the potential to provide an excellent natural source of this essential element. Our analyses gave us a preliminary understanding of the genetic basis of Se uptake in lentil and indicated that any potential strategy for micronutrient biofortification in lentil will require choice of field locations that minimize the spatial variability of soil Se content.

KEYWORDS: Selenium; selenomethionine (SeMet); selenocysteine (SeCys); lentils

INTRODUCTION

Selenium (Se) is an essential micronutrient in human nutrition and is involved in important regulatory and protective mechanisms. The nutritional benefits of Se were first published in 1957 (1). Se or, more specifically, selenocysteine (SeCys) is a key component of certain enzymes, for example, in the Se-dependent iodothyronine deiodinases involved in activating thyroid hormone. It also forms the integral parts of glutathione peroxidases and selenoprotein P (2), containing one or more atoms of Se per protein molecule. Essential Se-based roles in enzymes, antioxidants, and protective pathways have been discovered and have recently gained importance in cancer suppression, HIV treatment, free radical induced diseases, and protection from toxic heavy metals (3–5).

Selenium content in the human diet has increasing importance, as the effect of Se deficiency on human health is becoming a topic of interest in public health systems around the world. A recommended dietary allowance (RDA) of 55 μg of Se day^{-1} has been established for regular adults in the United States (6), and 60–75 μg of Se day^{-1} has been recommended for regular adults in the United Kingdom (7). This requirement is generally met by North Americans; however, large numbers of people in Europe, Asia, Australia, and parts of Africa have intakes of less

than the RDA level. Selenium-enriched commercial fertilizers have been recommended in Finland since 1984 to increase Se content in their major food crops. The recommended fertilizer rate for cereals and other crops was 16 mg of Se as sodium selenate (Na_2SeO_4) per kilogram of fertilizer. Since then, the daily Se intake of the Finnish population has increased from 39 μg of Se day^{-1} in 1984 to 110 μg of Se day^{-1} in 1998 (8). Low intake of Se (<25 $\mu\text{g day}^{-1}$) is linked to specific diseases such as arsenicosis in Bangladesh and fatal juvenile cardiomyopathy (Keshan disease) in China. Deficiency is also linked to specific diseases such as poor skeletal muscle strength in older adults (9), and even a slight deficiency has now been associated with other disorders including chronic heart failure and prostate and bladder cancers (10–14). A dietary intake of 55–200 μg of Se per day is now recommended as safe and adequate to reduce the risks of several types of cancer (15, 16). Recently, several clinical studies examined the relationship between serum Se levels and the prevalence of diabetes among U.S. adults and suggested that the adverse effects of a high intake of Se may increase primary or secondary diabetes (12, 17, 18).

The Se content of the soil from which foods are derived is the major influence on dietary intake of Se. Soil Se is highly variable in distribution and chemical availability. Most soils around the world contain 0.1–2 μg of Se kg^{-1} (19). Deficient soils in New Zealand, Australia, Denmark, central Siberia, northeast to south central China, parts of India, and Bangladesh

* Corresponding author [telephone (306) 966-8786; fax (306) 966-5015; e-mail bert.vandenberg@usask.ca].

Table 1. Market Class, Major Consuming Countries, Protein Content and Cotyledon Color of 19 Lentil Genotypes Grown in Saskatchewan, Canada

market class	seed wt (mg)	major consuming countries	genotype	protein ^a (%)	cotyledon color
extra small red	<30	Bangladesh, Pakistan, Egypt	CDC Robin	26	red
			CDC Rosetown	27	red
			CDC Imperial	27	red
small red	30–50	England, Middle East, Sri Lanka, India, Pakistan	CDC Blaze	27	red
			CDC Impact	27	red
			CDC Redberry	25	red
			CDC Rouleau	22	red
			Red Chief	25	red
large red large green	>65	USA, Dubai, Sri Lanka	Laird	24	yellow
	>65	Spain, Turkey, Iran, Germany, Algeria	CDC Grandora	23	yellow
			CDC Greenland	24	yellow
			CDC Plato	22	yellow
			CDC Sedley	22	yellow
			CDC Sovereign	24	yellow
			CDC Richlea	20	yellow
			CDC Meteor	23	yellow
medium green	50–60	Latin America, Europe	CDC Viceroy	24	yellow
			CDC Milestone	25	yellow
small green	<40	Italy, Morocco, Greece, Mexico	Eston	25	yellow

^a Protein (%) was calculated on the basis of the total seed nitrogen content ($n = 57$).

produce crops that are very low in Se (20). On the other hand, soils of Ireland, Colombia, and Venezuela and of the Great Plains of the United States and Canada are naturally rich in Se (16). Our initial research showed that Saskatchewan soils have abundant Se and that lentils grown in Saskatchewan may have the potential to provide a significant natural source of this essential element (21). However, we have limited understanding of the potential for genetic improvement of Se uptake in Saskatchewan grown lentils.

Biofortification by enrichment of the nutritional contribution of staple food crops through plant breeding is one option that is now widely discussed in the fields of nutrition, public health, and agriculture at national and international levels. Development of an effective biofortification strategy requires the application of genetics and agronomy to provide a solution to wide-scale nutrition problems (22). Studies have demonstrated that there is significant genetic variation for Se uptake in soybean, wheat, and *Brassica* vegetables (23–25).

Pulses combined with cereals are central to the diets of billions of people, and the potential for Se biofortification of pulses is high because of their relatively high protein content. World lentil production on an annual basis is approximately 4 million metric tonnes, and about 20–25% is grown in Saskatchewan, Canada (26). Saskatchewan supplies lentils to consumers in more than 100 countries with concentrated regions of consumption in Europe, the Middle East, and most notably South Asia (Table 1). Health problems due to Se deficiency affect over 100 million people around the world, many of them in lentil-consuming countries. Progress has been made in controlling Se deficiency through dietary supplementation, food fortification, and agronomic fertilization, but new approaches such as biofortification of basic foodstuffs are needed. Supplying essential Se through widely consumed meals such as lentils and rice could help increase the intake of dietary Se in regions where Se intake is insufficient (7). Research is needed to determine whether significant genetic variation exists in pulse crops for Se uptake to develop appropriate breeding strategies in the future. This also requires an understanding of the Se content of soils. We investigated the potential for biofortification of Se content for Saskatchewan-grown lentils as a means of improving human nutrition. The dual objectives of this study were to (1) measure the total Se content of seeds of 19 lentil genotypes grown in 8 key lentil-growing regions in Saskatchewan, Canada,

Table 2. Experimental Design and Sample Protocol

year	2005, 2006
no. of study locations per year	8
study locations (soil zones)	1. Saskatoon (moist dark brown) 2. Kyle (brown) 3. Swift Current (brown) 4. Wilkie (dark brown) 5. Melfort (black) 6. Hodgeville (brown) 7. Rosthern (thin black) 8. Rouleau (moist dark brown)
no. of soil samples per location	4 ($n = 32$)
no. of lentil genotypes per location	19
no. of replications	3 per genotype
no. of lentil seed samples per location (total Se analysis)	114
total no. of lentil samples tested for total Se content	912

in 2005 and 2006 and (2) identify the chemical forms of Se in extra small red lentil cultivar CDC Robin grown in Saskatoon in 2005.

EXPERIMENTAL METHODS

Materials. Se standards and chemicals used for digestion and for total Se measurements were purchased from VWR International (Canada) and Sigma-Aldrich Co. (Canada). High-purity chemicals and solvents for HPLC analysis were purchased from Sigma-Aldrich Co. and were used without further purification.

Soil Samples. Locations of the field research sites in Saskatchewan and sample protocol are listed in Table 2. These locations cover the major lentil-growing areas in Saskatchewan. Four soil cores were collected at each site from the 0–30 cm soil layer. They were air-dried (≤ 40 °C), passed through a 2 mm sieve, homogenized into one composite sample, and stored in plastic vials at -20 °C until analysis. The soil samples were collected in October 2005, about 1 month after the lentil plots were harvested.

Approximately 1 g of soil underwent primary organic digestion in 3 mL of HNO_3 (70%) at 90 °C followed by 1 mL of 30% H_2O_2 and further digestion in 3 mL of 70% HNO_3 and 9 mL of 35% HCl at 90 °C over several hours (24 h). The resulting slurry was filtered and made up to 50 mL in deionized water. Measurements of total Se using this modified method were validated using NIST standard reference material 2586 (soils; $[\text{Se}] = 0.6 \pm 0.005$ mg kg^{-1}). Soils from the South Saskatchewan River basin (Saskatoon, Saskatchewan) were used as the Se

hyperaccumulator *Astragalus bisulcatus* grows naturally, were used as a laboratory reference material and measured periodically to ensure consistency in the method. The total Se concentrations of different soils are indicated as the mean of three replicates with standard error.

Lentil Seed Samples. Lentil seeds were obtained from regional variety trials conducted in 2005 and 2006 by the Crop Development Centre (CDC), University of Saskatchewan, Canada. The selected lentil genotypes, market class, and major consuming countries are listed in **Table 1**. For the genotype \times environment study, samples of between 10 and 20 g of dry lentil seeds (14% moisture) were collected from each location with three replicates. Each replicated seed sample was prepared by standard HNO_3 H_2O_2 digestion as described previously (21). Measurements of total Se concentration using this modified method were validated using NIST standard reference material 1573a (tomato leaves; $[\text{Se}] = 0.054 \pm 0.003 \text{ mg kg}^{-1}$). Total Se was measured by hydride generation flame atomic absorption spectroscopy (HGAAS) on a Varian SpectraAA150 equipped with a hydride generation apparatus (Varian Canada Inc., Mississauga, ON, Canada). Measurements were made on the digested sample solutions outlined above.

Se Speciation. For the Se speciation study, seed samples of 250–500 g were obtained from three replicated plots of the variety CDC Robin (27) grown at the Saskatoon location in 2005. The seeds were dehulled in a Satake TM-05 grain-testing mill (Satake Engineering Co. Ltd. Japan) and then carefully separated by hand into seed coat, embryo, and cotyledon fractions. Se species were separated on a BioCAD Sprint perfusion chromatograph fitted with a 100 μL sample loop using an anion-exchange column (Hamilton PRP-X100, Reno, Nevada, NV) and a reverse-phase C18 column (Varian, Lake Forest, CA) using previously developed and reported methods (28). Anionic exchange was carried out with 10 mM citric acid, 1 mM potassium hydrogen phthalate (KHP), and 1 μM rubidium nitrate made to pH 4.5 with ammonium hydroxide in water and 2% v/v methanol. Other Se compounds were confirmed using the C18 column with 10 mM triethylamine and 1 mM KHP at pH 9 in water and 2% v/v methanol. Relative concentrations of Se species in natural samples were determined by ICP-MS (Saskatchewan Research Council, Saskatoon, SK, Canada) normalized to rubidium-spiked HPLC solvent.

Lentil samples were ground to a fine powder, and a 250 mg subsample was suspended in 4 mL of Millipore water. Samples were digested by 10 mg of protease XIV (*Streptomyces griseus*) at 38 °C for 90 min, centrifuged, filtered through a 0.5 μm PTFE membrane, and mixed with 3 equiv of HPLC solvent. Standards (SeMet, S-methylselenocysteine, selenate, selenite) were used after simple dilution to 40 ng Se mL^{-1} . SeCys was prepared from CysSeSeCys by dissolution at pH 11, followed by sodium borohydride reduction. CysSeSeCys was dissolved with 6 M HCl before dilution with solvent.

Statistical Analysis. The experimental design was a randomized complete block design with 3 replicates, at 8 locations for 19 genotypes over 2 years. Subsamples of lentil seeds for the determination of total Se were randomly taken from the entire harvested sample of each of the field plots. Data from both years and 8 locations were combined, and data error variances were tested for homogeneity. Locations, replications, years, and genotypes were considered as random factors. Class variables were year, location, replication, and genotype. Mixed-model analysis of variance was performed using the PROC GLM procedure of SAS version 8.2. Means were separated by Fisher's protected LSD at $P < 0.05$ (29). For each location–year data were analyzed separately using the General Linear Model procedure (PROC GLM) of SAS version 8.2 (29). Means were separated by Fisher's protected least significant difference (LSD) at $P < 0.05$. The broad sense heritability (H^2) of Se concentration in lentil seeds was calculated from the error mean squares from PROC GLM of SAS version 8.2 (30).

RESULTS

Soil Se Concentrations and Conditions. Se availability in soils depends upon soil pH, aeration, organic carbon, and iron levels. In acidic soils, Se is relatively unavailable to plants and occurs mainly as insoluble selenides and elemental forms. This

Table 3. Total Soil Se Concentration, Soil Texture, and pH from Various Locations in Saskatchewan, Canada

location	soil texture	soil pH	total soil Se ($\mu\text{g kg}^{-1}$)	SE ^a
Saskatoon	clay loam	6.3	301	5
Wilkie	clay loam	5.9	262	5
Melfort	clay loam	7.3	213	7
Kyle	clay loam	6.3	75	3
Rosthern	silt loam	6.5	71	5
Hodgeville	clay loam	7.1	70	5
Swift Current	clay loam	6.4	45	5
Rouleau	heavy clay	7.9	37	3

^a Standard error ($n = 4$).

Table 4. Summary of Combined Analysis of Variance for Total Se Concentration of 19 Lentil Genotypes Grown at Different Locations in Saskatchewan, Canada

source	df	mean square ^a
year	1	1845855*
location	7	19770258*
genotype	18	156536**
replication (year, location)	32	306369*
year \times location	7	8191592*
genotype \times year	18	101913
genotype \times location	126	72551
error	576	32043

^a Mean square was significantly different at *, $P < 0.05$, and **, $P < 0.1$, respectively.

study examined soils from eight different locations in Saskatchewan covering major soil zones where the lentil crop is grown. The total soil Se concentration ranged from 37 to 301 μg of Se kg^{-1} , or equivalent to 0.5–3.8 μmol of Se g^{-1} (**Table 3**). The Saskatoon location showed significantly higher total soil Se concentration than Rouleau or Swift Current. The Wilkie and Melfort locations were the second highest in total soil Se concentration. The most predominant soil texture at these locations was clay loam, and soil pH ranged from 5.9 to 7.9 (**Table 3**). The soil at Rouleau was more alkaline, and at Wilkie it was slightly acidic and poorly aerated.

Total Se Concentration and Se Species (Beneficial Forms) in Lentil Seeds. Combined statistical analysis (mixed model) over the years and locations showed that variation in total Se concentration in lentil seeds was significant ($P < 0.05$ and $P < 0.1$) for years, locations, genotypes, and the interaction between location and year (**Table 4**). As expected with most quantitative traits, the interaction between the genotype and location shows that most of the variation in the total Se concentration in the lentil seeds may have been due to environmental variation such as soil Se content, soil moisture, and other crop management practices. Therefore, data were analyzed and presented in this paper separately for each location–year (**Table 5**). Significant genotypic differences in total Se in lentil seeds were observed at all but two locations: Rosthern and Wilkie in 2005 (**Table 5**). Lentils grown at Saskatoon and Kyle had the greatest mean total Se concentration (643–1884 μg of Se kg^{-1}) compared to those from Swift Current (139–233 μg of Se kg^{-1}), Rosthern (87–305 μg of Se kg^{-1}), and Wilkie (206–392 μg of Se kg^{-1}) (**Table 5**). Furthermore, it was found that the Se concentration of lentil seeds from Wilkie and Melfort was not influenced by soil Se concentration. High levels of seed Se were not observed, despite high soil Se concentration at these locations. The soil moisture conditions, weather patterns, and soil Se available to plants might explain these differences (**Table 3**). Lentils grown at Wilkie may have had a lower Se concentration due to soil

Table 5. Mean Total Se Concentration of 19 Lentil Genotypes Grown at Different Locations in Saskatchewan, Canada in 2005 and 2006

year	location	total Se concn in lentils ($\mu\text{g kg}^{-1}$)			
		min	max	mean (SE) ^a	genotype effect ^b
2005	Saskatoon	900	2104	1324(7)	*
	Kyle	553	851	643(2)	*
	Hodgeville	232	1403	536(3)	*
	Rosthern	162	560	305(3)	NS
	Melfort	108	619	298(3)	*
	Rouleau	149	403	269(1)	*
	Swift Current	137	327	233(1)	*
	Wilkie	76	505	206(3)	NS
	2006	Kyle	1236	2609	1884(5)
Saskatoon		510	1662	885(4)	*
Rouleau		442	990	633(1)	*
Melfort		160	507	308(2)	*
Hodgeville		128	380	220(1)	*
Wilkie		121	614	392(3)	*
Swift Current		61	254	139(1)	*
Rosthern		32	185	87(1)	*

^a SE, pooled standard error of mean calculated from mean square of ANOVA for each location ($n = 57$). ^b Genotype effect was significantly different at $P < 0.05$. NS, not significant at $P < 0.05$.

acidity, lower soil aeration, and high soil iron concentrations. Comparison of Se concentrations of Saskatoon and Kyle reflected the influence of soil moisture. In 2005, Saskatoon had higher precipitation than in 2006, whereas Kyle experienced greater precipitation in 2006 compared to 2005 (31). Between years, seed Se levels at a particular location varied up to 3-fold. The year-to-year variation in Se levels at any specific location can be explained by both soil and weather factors that influence the uptake of Se during grain filling. At each location, the trial fields follow a particular crop rotation and soil properties from field to field can vary substantially in soils derived from glacial till. The weather patterns, particularly temperature and precipitation, are extremely variable in a continental climate and can have a large influence on the availability of soil Se at any time during the growing season.

Se concentration in the lentil seeds varied 4–5-fold across the locations, and on average, seeds of some genotypes had 40–50% more than others (Table 6). The extra small red lentil genotype (CDC Robin) and two of the large green lentil genotypes (CDC Sedley and CDC Grandora) had the highest total Se concentrations (612–672 μg of Se kg^{-1}). The small green lentil genotype, Eston, had the lowest (Table 6). We calculated that a 35 g serving of CDC Robin lentil (95th percentile of lentil intake per person) grown in Saskatchewan could supply 42% of the current RDA in the United States (55 μg of Se day^{-1}).

Our elemental analysis from seed fractions of CDC Robin lentil from a high Se location (Saskatoon) had mean total Se concentrations as follows: embryo axis, 3600 μg of Se kg^{-1} (45.6 μmol of Se kg^{-1}); cotyledon, 2800 μg of Se kg^{-1} (35.5 μmol of Se kg^{-1}); and seed coat, 2600 μg of Se kg^{-1} (32.9 μmol of Se kg^{-1}). Our previous experiments (21) demonstrated that whole seeds of CDC Robin from the Saskatoon location (0.72 μg of Se kg^{-1} ; 9.1 nmol of Se kg^{-1}) and those from Swift Current (0.16 μg of Se kg^{-1} ; 2 nmol of Se kg^{-1}) showed the greatest range of total Se concentration of the locations tested.

The relative content of Se chemical forms in the lentil seeds was determined by various HPLC-ICP-MS techniques. CDC Robin is a commercially grown cultivar in Saskatchewan because of its early maturity, disease resistance, high yield, and

Table 6. Comparison of Total Se Concentration in 19 Lentil Genotypes Grown in Saskatchewan, Canada, in 2005 and 2006

genotype	total Se concn ($\mu\text{g kg}^{-1}$)			%RDA ^a (100 g of lentil)	
	Saskatoon (2005)	Kyle (2006)	mean ^b (8 locations, 2 years)	North America (55 $\mu\text{g day}^{-1}$)	Europe (65 $\mu\text{g day}^{-1}$)
CDC Robin	2104 a	2012 bcd	672 a	122	103
CDC Sedley	1446 abcd	2127 abc	612 a	111	94
CDC Grandora	1694 abc	2351 ab	612 a	111	94
Laird	1232 cd	1970 bcde	593 abc	108	91
CDC Greenland	1064 cd	2609 a	544 abcd	99	84
CDC Imperial	1246 cd	1884 bcdef	538 bcde	98	83
CDC Redberry	1947 ab	1583 defg	533 cde	97	82
CDC Sovereign	1503 abcd	2364 ab	533 cde	97	82
CDC Plato	1178 cd	2035 bcd	532 cde	97	82
CDC Meteor	1483 abcd	1470 efg	510 def	93	78
CDC Blaze	1413 bcd	2119 abc	509 def	93	78
CDC Rosetown	1005 d	1942 bcdef	505 def	92	78
CDC Richlea	900 d	2008 bcd	498 defg	91	77
CDC Impact	1136 cd	1844 bcdef	491 defgh	89	76
Red Chief	1429 abcd	1421 fg	472 efgh	86	73
CDC Viceroy	1009 cd	1685 cdefg	471 efgh	86	72
CDC Milestone	1186 cd	1236 g	457 fgh	83	70
CDC Rouleau	1271 cd	1585 defg	431 gh	78	66
Eston	901 d	1555 defg	425 h	77	65
SE ^c	7	5	6		

^a %RDA was calculated on the basis of the mean total Se concentration across eight locations ($n = 912$) in Saskatchewan. ^b Means within a column followed by different letters are significantly different at $P < 0.05$. ^c SE, pooled standard error of mean calculated from mean square of ANOVA for each location ($n = 57$) and mean of eight locations ($n = 912$).

consumer preference. In addition, it was found that lentil seeds from CDC Robin had the highest Se concentration compared to a wide range of lentil genotypes grown in Saskatchewan. Furthermore, many South Asian consumers (specifically Bangladesh) prefer red cotyledon, extra small seed size (> 30 mg) cultivars such as CDC Robin. On the basis of these factors, we chose CDC Robin to study the Se speciation. More than 70% of the Se in the whole lentil sample was present as organic Se with a small fraction ($< 20\%$) as inorganic Se (Table 7). Small fractions (7%) of SeCys and γ -glutamylselenocysteine were present in the whole lentil seeds, and the concentrations of the other Se species (selenomethionine, dimethylselenoxide, and S-methylselenocysteine) were not significant. In the embryonic axis, $> 80\%$ of the Se was present as organic Se with a small fraction (20%) as inorganic Se. SeMet (73%) and selenate (27%) were the major chemical forms of Se present in CDC Robin cotyledon, and inorganic Se (94%) was the major chemical form of Se present in the lentil seed coat (Table 7). Our results clearly indicated that the field-grown CDC Robin lentils contained predominately organic Se (80%) as SeMet and SeCys with a minor component of inorganic Se (20%).

DISCUSSION

The biological importance of Se and its roles in human health have recently become of great interest in the international community. There is a great necessity for food systems to provide at least 55 μg per day for maximal expression of Se enzymes, and large populations in some parts of the world are Se deficient. Se deficiency compromises the health of developing children and reduces the ability to combat the effects of heavy metals in the human diet (32). As a common, universal, and quick-cooking nutritious food source, lentils can help to deliver

Table 7. Total Se Concentration and Percentage Composition of Se Species for CDC Robin (Saskatoon, 2005)

seed fraction ^b	contribution to total seed wt (%)	total Se (SE) ^c ($\mu\text{g kg}^{-1}$)	percentage of Se species present in lentil seeds as ^a				
			organic Se			inorganic Se	
			SeMet (\pm SE)	selenocysteines (\pm SE) ^d	γ -glutamylselenocysteine (\pm SE)	selenate (\pm SE)	selenite (\pm SE)
CDC Robin whole seed	100	2104	69 \pm 2	7 \pm 1	2 \pm 1	10 \pm 1	9 \pm 1
CDC Robin embryo	5	3600	19 \pm 1	53 \pm 2	8 \pm 1	3 \pm 1	17 \pm 1
CDC Robin cotyledon	88	2800	73 \pm 2	nd ^e	nd	27 \pm 2	nd
CDC Robin seed coat	7	2600	nd	nd	6 \pm 1	80 \pm 2	14 \pm 1

^a Se speciation as determined using LC and ICP-MS. ^b Lentils were collected from Saskatoon location, 2005 ($n = 12$). ^c SE, standard error. ^d Selenocysteine and selenocystine. ^e nd, not detected within the limits of quantitation (1.6 parts per trillion in HPLC fraction).

beneficial Se to those who need it. Lentil-growing regions with adequate soil Se play a fundamental role in this mass distribution. We have shown that Saskatchewan-grown lentils contain 425–673 μg of Se kg^{-1} depending upon location, soil characteristics, and growing conditions. This potentially provides 80–120% of the minimum recommended daily Se intake in only 100 g of dry lentils. Our data are derived from small-plot field trials. The Se concentration available in commercial lentil shipments would likely reflect a blended average across many fields in multiple locations.

There is unique potential for Se-rich lentil and other pulse crops to be grown in western Canada without soil supplementation. We conducted a preliminary analysis of the Se content of lentil seeds grown in some other regions of the world (U.S. Pacific Northwest, >50 samples; Australia, >40 samples; Syria, 7 samples; Bangladesh, 12 samples; India, 10 samples; and Nepal, 5 samples). All samples had very low Se concentration, on average, <5% of the Se content of the lowest Se content lentils from Saskatchewan (Swift Current). Many samples from Syria, Bangladesh, India, and Nepal had no detectable Se (<20 ppb) (data not shown).

The chemical species distribution in seeds of Se is important in terms of nutritional benefits. It has long been understood that certain forms of Se are critical to development and self-regulation, whereas others are potential poisons. The biological fate of Se is also determined by the original form and the transformation that occurs during digestion and absorption. The amino acid SeMet is readily incorporated into protein masses, but SeCys, which is found in key regulatory proteins, is tightly controlled and is catabolized into hydrogen selenide. Inorganic forms, such as selenate and selenite, have been studied for their involvement in the treatment of arsenicosis and excretion of mercury (32).

The presence of Se in plants grown on soils containing available Se has been reported in many studies. Seleniferous green onion (*Allium cepa* L.) predominantly contained SeMet and small amounts of SeCys (33). SeMet and SeCys were the major organic selenides found in sour clover (*Melilotus indica* L.) and alfalfa (*Medicago sativa* L.) grown in seleniferous soils in California (34). SeMet is the major organic form of Se found in wheat, common bean, mushroom, and yeast (35). Our findings for Se in lentils seed are similar to those reported for seeds of seleniferous wheat (*Triticum aestivum* L.), common bean (*Phaseolus vulgaris* L.), alfalfa, and sour clover, which contain mainly SeMet with a smaller fraction of SeCys (34, 35).

HPLC-ICP-MS analysis of the Se species in whole lentils revealed that most of Se was present as SeMet with small amounts of selenate and very small amounts of selenocysteines, selenite, and other selenooligopeptides such as γ -glutamylselenocysteine (gGSeCys) as outlined in **Table 7**. This supports our previous experiments using synchrotron X-ray spectroscopy

to identify Se species in lentil seeds and seed tissues (21). Synchrotron techniques offer a unique advantage in that samples can be run intact with no pretreatment, but it is difficult to differentiate between chemically similar species such as SeMet and Se-methylselenocysteine (Se-MeSeCys) or to reliably detect smaller components when one type is in great excess. HPLC-ICP-MS methods for Se quantitation can be used to differentiate the forms that X-ray techniques cannot. Conversely, the overlap of other chemical species in the HPLC methods can be differentiated in the related synchrotron experiments. Both methods used in conjunction are sufficient to determine the complete set of Se forms present in seeds.

Our analysis of Se speciation in CDC Robin lentil provided an indication that Se species may vary according to the seed component. The seed coat has a unique Se species profile, largely inorganic. The embryonic axis is enriched for SeCys in comparison to cotyledon tissue. Red lentils are usually decorticated prior to cooking in whole or split form. In terms of Se speciation, split lentils may have lower SeCys content because the embryo fraction is often collected as a byproduct for use in animal feed. In some countries, for example, in Bangladesh, consumers have a distinct preference for decorticated unsplit lentils, which may be beneficial for human nutrition.

Other factors, such as cooking, grinding, and digestion that may affect or transform Se speciation, have been investigated. We found that cooking the lentils in boiling water did not change the total Se content (data not shown). There is a migration of Se from the lentils to the liquid broth, but provided the lentils and broth are consumed as a whole food source, the Se concentration and speciation remain intact. However, we would expect a nearly 50% reduction in Se for lentils that are thermally processed in brine (canned) and consumed after the canning brine is discarded (36).

In many parts of the world, lentils with adequate beneficial Se concentration could be considered a natural, whole food source for Se, and a possible solution to Se deficiency-related arsenicosis in Bangladesh and juvenile cardiomyopathy (Keshan disease) in China (32). Supplementation of 200 μg per day may help to prevent certain cancers, such as bladder, prostate, liver, colorectal, and lung cancers (16). Efforts to optimize the Se in food sources must consider not only the overall concentration but the amounts of the various beneficial forms.

Quantitative traits generally depend on the collective interaction of many genes. The expression of quantitative genes is also influenced by the environment. The phenotypic variance calculation is influenced by the number of years, locations, and replicates used in the experiment, therefore, plant breeders commonly use heritability estimates to distinguish the proportion of total phenotypic variation due to genotype and environmental influences. This estimate is then used to design appropriate genetic improvement programs (14, 15, 17, 18, 19, 20, 21).

genetic variance was 1135 and that of phenotypic variance was 2877. The broad sense heritability estimate was 0.4, which indicates that Se content in lentil is in the midrange of heritability. An appropriate genetic improvement strategy for increasing Se content lentil would require that environmental influence be kept to a minimum by careful selection of environments with low spatial variability for soil Se content combined with appropriate replication.

Breeding for enhanced Se accumulation and selective speciation may be an effective strategy to help overcome global Se deficiencies. Some studies have suggested the possibility of genetic improvement for Se uptake in *Brassica* vegetables, wheat and soybean (23–25). By specifically controlling the Se variability in soil Se content, it would be possible to reduce environmental effects as part of our biofortification approach on lentil genetic improvement. It may be possible to screen lentil genotypes for increased Se uptake ability using atomic absorption spectroscopy techniques or possibly using marker-assisted genetic selection. On the basis of our results, we suggest that it may be possible to cost effectively breed lentil cultivars for enhanced Se uptake for specific regions of the world with soils that have lower levels of Se. In regions where Se is highly deficient, it may be necessary to combine this approach with agronomic biofortification using fertilizer with Se additives. This may be particularly important for regions where the rapidly increasing cost of rice may induce further reduction in the land area devoted to pulse production. Further studies are being performed on diverse genotypes, including wild relatives of cultivated lentil, modern commercial cultivars, and genotypes adapted to different geographic locations in Europe, Asia, Africa, and North America.

Se must be available in the soil for uptake and transformation. In general, soil Se is unevenly distributed and varied in availability, ranging from <0.1 to >100 ppm, and most commonly from 1 to 1.5 ppm (19). Ultimately, the total Se in the soil depends on the minerals in the rocks from which the soil was derived. Soils of the Northwest, Southeast, and Great Lakes regions of the United States were derived from volcanic deposits and have low soil Se content (<0.05 ppm) (37). Soils originating from cretaceous shales, such as those found in South Dakota and Montana, tend to have concentrations upward of 10 ppm (37). However, the availability of the Se is greatly dependent on aeration, water availability, pH, and soil texture and composition. In poorly aerated soils, Se is relatively unavailable to plants and occurs mainly as insoluble selenides and elemental forms. Furthermore, wetter soils with alkaline pH have lower Se concentration due to leaching of mobile selenate (20).

Our study indicates that uptake of Se in lentil seeds is affected by soil and environmental conditions such as moisture, soil texture, aeration, and soil fertility and irrigation (7). The higher Se concentration observed from the Saskatoon area may be due to higher spatial variability in the soil combined with wet weather conditions, which would increase availability of soil Se. Soils at Rouleau had the lowest Se concentration (Table 3) and the highest pH. The Rouleau soil was a moist heavy clay with poor aeration, thus reducing the amount of Se that is available to plants. We found that the total soil Se concentration was not the best indicator of plant Se availability, although it is the most commonly used method of reporting Se availability in the literature (38). A complete understanding of the biochemistry of Se in soil and lentil plants will require more in-depth studies of plant biochemistry, agronomy, and physiology.

In summary, our present study shows that Saskatchewan soils are naturally rich in Se and that lentils grown in them have

great potential as a quick-cooking, Se-rich, natural food product. Significant genotypic differences for Se concentration were observed across the locations. In addition, genotype \times environment analysis of the concentration of Se in the lentils indicated that good potential exists for genetic improvement of the concentration of this essential element in lentil. The Se content and the chemical forms of Se within the seed may be altered by conventional plant breeding approaches or by optimizing agricultural production conditions.

ACKNOWLEDGMENT

We thank B. Barlow, D. de Silva, and Dr. A. Sarker (ICARDA) for providing the lentil seeds and soils from the Crop Development Centre, University of Saskatchewan, Canada, and Drs. Gerald F Combs, Jr., Kofi Agblor, and Curtis Pozniak for reviewing the manuscript.

LITERATURE CITED

- (1) Schwarz, K.; Foltz, C. M. Selenium as an integral part of factor 3 against dietary necrotic liver degeneration. *J. Am. Chem. Soc.* **1957**, *79*, 3292–3293.
- (2) Rotruck, J. T.; Pope, A. L.; Ganther, H. E.; Swanson, A. B.; Hafeman, D. G.; Hoekstra, W. G. Selenium. Biochemical role as a component of glutathione peroxidase. *Science* **1973**, *179*, 588–590.
- (3) Bordoni, A.; Danesi, F.; Malaguti, M.; Di Nunzio, M.; Pasqui, F.; Maranesi, M.; Biagi, P. L. Dietary selenium for the counteraction of oxidative damage: fortified foods or supplements? *Br. J. Nutr.* **2008**, *99*, 191–197.
- (4) Gailer, J.; Madden, S.; Burke, M. F.; Denton, M. B.; Aposhian, H. V. Simultaneous multielement-specific detection of a novel glutathione-arsenic-selenium ion [(GS)2AsSe]- by ICP AES after micellar size-exclusion chromatography. *Appl. Organomet. Chem.* **2000**, *14*, 355–363.
- (5) Jansson, B. The role of selenium as a cancer-protecting trace element. *Met. Ions Biol. Syst.* **1980**, *10*, 281–311.
- (6) Monsen, E. R. Dietary reference intakes for the antioxidant nutrients: vitamin C, vitamin E, selenium, and carotenoids. *J. Am. Diet. Assoc.* **2000**, *100*, 637–640.
- (7) Broadley, M. R.; White, P. J.; Bryson, R. J.; Meacham, M. C.; Bowen, H. C.; Johnson, S. E.; Hawkesford, M. J.; McGrath, S. P.; Zhao, F. J.; Breward, N.; Harriman, M.; Tucker, M. Biofortification of UK food crops with selenium. *Proc. Nutr. Soc.* **2006**, *65*, 169–181.
- (8) Wang, W. C.; Naentoe, V.; Maekelae, A. L.; Maekelae, P. Effect of nationwide selenium supplementation in Finland on selenium status in children with juvenile rheumatoid arthritis. a ten-year follow-up study. *Analyst* **1995**, *120*, 955–958.
- (9) Lauretani, F.; Semba, R. D.; Bandinelli, S.; Ray, A. L.; Guralnik, J. M.; Ferrucci, L. Association of low plasma selenium concentrations with poor muscle strength in older community-dwelling adults: the InCHIANTI Study. *Am. J. Clin. Nutr.* **2007**, *86*, 347–352.
- (10) Brinkman, M.; Buntinx, F.; Muls, E.; Zeegers, M. P. Use of selenium in chemoprevention of bladder cancer. *Lancet Oncol.* **2006**, *7*, 766–774.
- (11) de, L. M.; Salen, P. Selenium and antioxidant defenses as major mediators in the development of chronic heart failure. *Heart Fail. Rev.* **2006**, *11*, 13–17.
- (12) Hawkes, W. C.; Alkan, Z.; Lang, K.; King, J. C. Plasma selenium decrease during pregnancy is associated with glucose intolerance. *Biol. Trace Elem. Res.* **2004**, *100*, 19–29.
- (13) Karunasinghe, N.; Ryan, J.; Tuckey, J.; Masters, J.; Jamieson, M.; Clarke, L. C.; Marshall, J. R.; Ferguson, L. R. DNA stability and serum selenium levels in a high-risk group for prostate cancer. *Cancer Epidemiol. Biomarkers Prevent.* **2004**, *13*, 391–397.
- (14) Kellen, E.; Zeegers, M.; Buntinx, F. Selenium is inversely associated with blood C-reactive protein in a Belgian population. *FASEB J.* **2009**, *23*, 1002–1008.

- case-control study on bladder cancer. *Int. J. Urol.* **2006**, *13*, 1180–1184.
- (15) Clark, L. C.; Dalkin, B.; Krongrad, A.; Combs, G. F., Jr.; Turnbull, B. W.; Slate, E. H.; Witherington, R.; Herlong, J. H.; Janosko, E.; Carpenter, D.; Borosso, C.; Falk, S.; Rounder, J. Decreased incidence of prostate cancer with selenium supplementation: results of a double-blind cancer prevention trial. *Br. J. Urol.* **1998**, *81*, 730–734.
- (16) Combs, G. F., Jr.; Lu, J. Selenium as a cancer preventive agent. *Selenium* **2001**, 205–217.
- (17) Bleys, J.; Navas-Acien, A.; Guallar, E. Serum selenium and diabetes in U.S. adults. *Diabetes Care.* **2007**, *30*, 829–834.
- (18) Bleys, J.; Navas-Acien, A.; Guallar, E. Selenium and diabetes: more bad news for supplements. *Ann. Intern. Med.* **2007**, *147*, 271–272.
- (19) Berrow, M. L.; Ure, A. M. Geological materials and soils In *Occurrence and Distribution of Selenium*; CRC Press: Boca Raton, FL, 1989; pp 226–228.
- (20) Combs, G. F., Jr. Selenium in global food systems. *Br. J. Nutr.* **2001**, *85*, 517–547.
- (21) Thavarajah, D.; Vandenberg, A.; George, G. N.; Pickering, I. J. Chemical form of selenium in naturally selenium-rich lentils (*Lens culinaris* L.) from Saskatchewan. *J. Agric. Food Chem.* **2007**, *55*, 7337–7341.
- (22) <http://www.harvestplus.org/>.
- (23) Finley, J. W. Reduction of cancer risk by consumption of selenium-enriched plants: enrichment of broccoli with selenium increases the anticarcinogenic properties of broccoli. *J. Med. Food* **2003**, *6*, 19–26.
- (24) Lyons, G.; Stangoulis, J.; Graham, R. High-selenium wheat: biofortification for better health. *Nutr. Res. Rev.* **2003**, *16*, 45–60.
- (25) Wei, A. Soybean sprout. *Zhengzhou Liangshi Xueyuan Xuebao.* **1996**, *17*, 67–70.
- (26) <http://www.agriculture.gov.sk.ca/Statistics>.
- (27) Vandenberg, A.; Kiehn, F. A.; Vera, C.; Gaudiel, R.; Buchwaldt, L.; Dueck, S.; Wahab, J.; Slinkard, A. E. CDC Robin lentil. *Can. J. Plant Sci.* **2002**, *82*, 111–112.
- (28) Pedrero, Z.; Encinar, J. R.; Madrid, Y.; Camara, C. Identification of selenium species in selenium-enriched *Lens esculenta* plants by using two-dimensional liquid chromatography–inductively coupled plasma mass spectrometry and [⁷⁷Se]selenomethionine selenium oxide spikes. *J. Chromatogr.* **2007**, *1139*, 247–253.
- (29) *SAS Institute SAS User's Guide: Statistics*; 9th ed.; SAS Institute: Cary, NC, 2005.
- (30) Falconer, D. S.; MacKay, T. F. C. Introduction to quantitative genetics. 1996.
- (31) <http://www.weatheroffice.gc.ca>.
- (32) Spallholz, J. E.; Mallory Boylan, L.; Rhaman, M. M. Environmental hypothesis: is poor dietary selenium intake an underlying factor for arsenicosis and cancer in Bangladesh and West Bengal, India? *Sci. Total Environ.* **2004**, *323*, 21–32.
- (33) Hamilton, J. W.; Beath, O. A. Selenium uptake and conversion by certain crop plants. *Agron. J.* **1963**, *55*, 528–531.
- (34) Wu, L.; Guo, X.; Banuelos, G. S. Accumulation of seleno-amino acids in legume and grass plant species grown in selenium-laden soils. *Environ. Toxicol. Chem.* **1997**, *16*, 491–497.
- (35) Li, G. X.; Lee, H. J.; Wang, Z.; Hu, H.; Liao, J. D.; Watts, J. C.; Combs, G. F., Jr.; Lue, J. Superior in vivo inhibitory efficacy of methylseleninic acid against human prostate cancer over selenomethionine or selenite. *Carcinogenesis* **2008**, *29*, 1005–1012.
- (36) Sager, M. Selenium in agriculture, food, and nutrition. *Pure Appl. Chem.* **2006**, *78*, 111–133.
- (37) Kubota, J.; Allaway, W. H.; Carter, D. L.; Gary, E. E.; Lazar, V. A. Selenium in crops in the United States in relation to selenium-responsive diseases of animals. *J. Agric. Food Chem.* **1967**, *15*, 448–453.
- (38) Lyons, G. H.; Genc, Y.; Stangoulis, J. C. R.; Palmer, L. T.; Graham, R. D. Selenium distribution in wheat grain, and the effect of postharvest processing on wheat selenium content. *Biol. Trace Elem. Res.* **2005**, *103*, 155–168.

Received for review July 25, 2008. Revised manuscript received September 8, 2008. Accepted September 18, 2008. Support for this research was provided by the Saskatchewan Pulse Growers (Project PRO0717), Saskatoon, Saskatchewan.

JF802307H

800.218-sub 1 (b) (6)

		Case Sample	Label	AAFCO
		California Naturals Kangaroo & Lentil	Product Nutrient Analysis (website label)	AAFCO-Adult Maint
(b) (4)	Ca	1.30%	0.83%	0.5 to 2.5%
	Mg	0.13%	0.17%	0.06%
	P	0.74%	0.71%	0.4 to 1.6 %
	Fe	30 mg/kg	305 mg/kg	40 mg/kg
	Co	0.12 mg/kg	n/a	25 mg/kg-chicks/rats/sheep max
	Cu	21 mg/kg	13.61 mg/kg	7.3 mg/kg
	Zn	240 mg/kg	193.37 mg/kg	80 mg/kg
	Se	0.7 mg/kg	0.08 mg/kg	0.35 to 2 mg/kg
	Ca:P	1.76:1		1:1 to 2:1
	Cu:Zn	0.09:1		0.09:1-not AAFCO
(b) (4)	Tau	~0.26%		0.1% in Cats
	Cystine	2.32 mg/g = ~0.23%		n/a
	Met	5.78 mg/g = ~0.58%	0.61%	0.33%
	Met-Cys	~0.81%	0.97%	0.65%

800.218-sub 5				
Case Sample		Label	AAFCO	
California Naturals Chicken Meal		Product Nutrient Analysis (website label)	AAFCO-Adult Maint	
(b) (6)	Ca	1.80%	1.98%	0.5 to 2.5%
	Mg	0.14%	0.12%	0.06%
	P	1.30%	1.34%	0.4 to 1.6 %
	Fe	39 mg/kg	156 mg/kg	40 mg/kg
	Co	0.14 mg/kg	n/a	25 mg/kg-chicks/rats/sheep max
	Cu	19 mg/kg	18 mg/kg	7.3 mg/kg
	Zn	330 mg/kg	229 mg/kg	80 mg/kg
	Se	0.66 mg/kg	0.78 mg/kg	0.35 to 2 mg/kg
	Ca:P	1.38:1		1:1 to 2:1
	Cu:Zn	0.06:1		0.09:1-not AAFCO
(b) (6)	Tau	1.08 mg/g = ~0.11%		0.1% in Cats
	Cystine	3.2 mg/g = ~0.32%		n/a
	Met	6.2 mg/g = ~0.62%	0.65%	0.33%
	Met-Cys	~0.94%	0.98%	0.65%

		800.218-sub 4		
		Case Sample		
		Fromm Heartland Gold Grain Free Large Breed Adult	Product Typical Analysis (website label)	AAFCO Growth & Maint
(b) (6)	Ca	1.20%	1.14%	1.2 to 1.8%
	Mg	0.14%	0.17%	0.06%
	P	1%	1.08%	1 to 1.6%
	Fe	30 mg/kg	258.26 mg/kg	88 mg/kg
	Co	0.37 mg/kg	n/a	25 mg/kg-chicks/rats/sheep max
	Cu	25 mg/kg	25.83 mg/kg	12.4 mg/kg
	Zn	170 mg/kg	217.37 mg/kg	100 mg/kg
	Se	0.85 mg/kg	n/a	0.35 to 2 mg/kg
	Ca:P	1.2:1		1:1 to 2:1
	Cu:Zn	0.15:1		0.09:1-not AAFCO
(b) (4)	Tau	1.84 mg/g = ~0.18%	n/a	0.1% in Cats
	Cystine	3.15 mg/g = ~0.32%	n/a	n/a
	Met	4.75 mg/g = ~0.48%	n/a	0.35%
	Met-Cys	~0.79%	n/a	0.70%



FAU Institutional Repository

<http://purl.fcla.edu/fau/fauir>

This paper was submitted by the faculty of FAU's Harbor Branch Oceanographic Institute

Notice: This article was published by the European Association for Aquatic Mammals <http://www.aquaticmammalsjournal.org/> and may be cited as Bossart, Gregory D. , George Hensley, Juli D. Goldstein, Kenny Kroell, Charles A. Manire, R. H. Defran, and John S. Reif (2007) Cardiomyopathy and Myocardial Degeneration in Stranded Pygmy (*Kogia breviceps*) and Dwarf (*Kogia sima*) Sperm Whales, *Aquatic Mammals* 33(2) 214-222, DOI: 10.1578/AM.33.2.2007.214

Cardiomyopathy and Myocardial Degeneration in Stranded Pygmy (*Kogia breviceps*) and Dwarf (*Kogia sima*) Sperm Whales

Gregory D. Bossart,¹ George Hensley,¹ Juli D. Goldstein,¹ Kenny Kroell,¹
Charles A. Manire,² R. H. Defran,¹ and John S. Reif^{1, 3}

¹Center for Coastal Research, Marine Mammal Research and Conservation Program, Harbor Branch Oceanographic Institution, 5600 U.S. 1 North, Fort Pierce, FL 34946, USA; E-mail: GBossart@hboi.edu

²Mote Marine Laboratory and Aquarium, 1600 Ken Thompson Parkway, Sarasota, FL 34236, USA

³Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO 80523, USA

Abstract

Cardiomyopathy (CMP) has been documented as a disease associated with stranded pygmy (*Kogia breviceps*) and dwarf (*Kogia sima*) sperm whales in the United States and Asia. In this study, hearts from 27 pygmy and two dwarf sperm whales stranded in the coastal U.S. Atlantic Ocean and Gulf of Mexico from 1999 to 2006 were analyzed. Gross and microscopic examinations were conducted according to a standardized protocol designed to ensure systematic examination of tissue and data recording. Hearts were weighed and specific measurements made for selected tissues. Fourteen (48.3%) pygmy sperm whales had a microscopic diagnosis of CMP, 12 (41.4%) showed evidence of mild myocardial degeneration (MCD), one (3.4%) had moderate myocarditis and two (6.9%) had no pathological lesions. One dwarf sperm whale had CMP, and the other had mild MCD. The majority of stranded *Kogia* spp. with cardiac lesions came from the southeast Atlantic region (19/27, 70.3%). Cardiomyopathy and MCD lesions were found predominantly among adult whales. An excess of males was found for CMP and MCD (approximately 75% of both groups). The predominant histological lesions found in both disorders were anisokaryosis with karyomegaly and nuclear rowing, followed in frequency by interstitial edema. Cardiac weight, ventricular wall thickness, and valve circumference were compared between pygmy sperm whales with CMP and those with MCD. The largest differences were found for heart weight and intraventricular septum wall thickness, but none of the differences were statistically significant. Further adjustment for sex and body length did not alter the results. In the aggregate, these findings suggest that CMP in *Kogia* spp. is a chronic, progressive condition that represents a continuum from MCD to the more severe forms of the disorder. The etiology of this complex disorder remains unknown.

Key Words: pygmy sperm whale, *Kogia breviceps*, dwarf sperm whale, *Kogia sima*, cardiomyopathy, myocardial degeneration, stranding, U.S. Atlantic Ocean and Gulf of Mexico

Introduction

Cardiomyopathy (CMP) was first described in pygmy (*Kogia breviceps*) and dwarf (*Kogia sima*) sperm whales in 1985 in a study group of 29 beached whales (Bossart et al., 1985). The disease in *Kogia* spp. has been described primarily in whales from the southeastern Atlantic Ocean, but it also occurs in Pacific Ocean whales (Chiu et al., 2003). The etiopathogenesis of the *Kogia* CMP is unknown. Distinct clinical, functional, and pathological patterns of CMP occur in domestic animals and humans, however, and each pattern may be associated with distinct pathogenic mechanisms. While controversies exist with CMP classification schemes, the general clinical, functional, and pathological patterns of CMP are the stress, dilated, hypertrophic, and restrictive forms.

Interest in the etiology and pathogenesis of CMP is ongoing as *Kogia* spp. are the second most common single-stranded cetaceans in the southeastern United States (SEUS) after the bottlenose dolphin (*Tursiops truncatus*). Total annual *Kogia* strandings have ranged from 16 to 69 in the SEUS and from 9 to 40 in Florida (Odell et al., 2004). Annual stranding totals have been highly variable and, at least on the east coast of Florida, may be related to chronic disease and local oceanographic conditions, especially the Gulf Stream (Bossart et al., 1985; Odell et al., 2004). *Kogia* are rarely seen at sea and, despite the relatively high frequency of strandings, very little is known about their biology. In fact, prior to 1966, only one species was recognized (Odell et al., 2004).

The purpose of this study was to further characterize the pathological features of cardiac lesions found in pygmy and dwarf sperm whales using a

newly developed standardized protocol designed to ensure systematic examination of tissue and data recording and to explore potential factors in their etiology.

Materials and Methods

Gross and Microscopic Pathology

The analysis reported here was based on gross and microscopic examination of whole hearts from 27 *K. breviceps* (17 adult males [M], 6 adult females [F], 1 subadult [M], 1 subadult [F], 2 calves [F]) and two *K. sima* (adult [M]) that stranded in the coastal U.S. Atlantic Ocean and Gulf of Mexico between 1999 and 2006 and were submitted to our laboratory for evaluation. A *Kogia* heart dissection manual was developed which describes specific protocols for the collection, fixation, and dissection of heart specimens from *Kogia* spp. (Hensley et al., 2005). Procedures were standardized to ensure systematic gross and microscopic examination of tissue and data recording. *In situ* examination of the heart is detailed in the manual, which also emphasizes the importance of accurately determining the heart weights and specific heart measurements.

Briefly, the formalin-fixed heart was divided into five cross sections of approximately the same width. Cross sections were referred to as Levels 1 through 5, from apex to base, respectively. Heart weights and measurements (right and left ventricular wall thickness at Levels 2 and 4; interventricular septum thickness at Levels 2 and 4; valve circumference [tricuspid, mitral, pulmonary, and aortic]) were determined as described in Hensley et al. (2005). Evaluation included the collection of 12 representative heart sections: septal summit (two blocks), dorsal wall of right ventricle at Level 2, ventral wall of right ventricle at Level 2, dorsal wall of left ventricle at Level 2, ventral wall of left ventricle at Level 2, interventricular septum at Level 2, dorsal wall of right ventricle at Level 4, ventral wall of right ventricle at Level 4, dorsal wall of left ventricle at Level 4, ventral wall of left ventricle at Level 4, and interventricular septum at Level 4. After sectioning, samples were placed into labeled tissue cassettes containing fresh neutral buffered 10% formalin. Tissues were routinely processed, embedded in paraffin, sectioned at 5 μ m, and stained with hematoxylin and eosin for examination by light microscopy. Masson trichrome was used as a special stain to demonstrate the presence of collagen.

Myocardial degeneration (MCD) was diagnosed microscopically as (1) mild, multifocal, or diffuse anisokaryosis with karyomegaly, nuclear rowing, and interstitial edema found in all 12 sections described above; and (2) mild, multifocal,

or diffuse eosinophilic homogenization of sarcoplasm and vacuolization, myofiber disarray (architectural disorganization), wavy-attenuated myofibers, and/or loss of cross-striations in six or less sections described above. Fibrosis and inflammation were absent.

A diagnosis of CMP consisted of a suite of microscopic lesions, including multifocal or diffuse: (1) moderate to severe anisokaryosis with karyomegaly and nuclear rowing, (2) moderate to severe interstitial edema, (3) mild to severe eosinophilic homogenization of sarcoplasm and vacuolization, (4) mild to severe myofiber disarray (architectural disorganization), (5) mild to severe wavy-attenuated myofibers, (6) mild to severe loss of cross-striations, and (7) mild to severe fibrosis in 11 or more sections as described above.

Statistical Analyses

Statistical analyses were conducted to determine whether heart weights, wall thicknesses, and valve dimensions differed among *K. breviceps* affected with MCD and CMP. Analyses were limited to *K. breviceps* because the number ($n = 2$) of *K. sima* was inadequate. Further, the morphometrics of these two species would be expected to differ, thus they were not combined. In the first phase of the analysis, all *K. breviceps* with CMP were compared to those with MCD using the *t*-test procedure in SAS, Version 9.1 (SAS, Cary, NC). A two-way analysis of variance (ANOVA) was then conducted for heart weight and Level 2 intraventricular septum thickness with diagnosis and sex in the model. Finally, PROC GLM in SAS was used to compare heart weight and Level 2 intraventricular septum thickness adjusting for body length in sperm whales with MCD and CMP. An alpha of $p < 0.05$ was considered statistically significant.

Results

The cardiac lesions found in this sample of 29 stranded *Kogia* spp. are summarized in Table 1 along with the geographic region where they occurred. Fourteen (48.3%) whales had a diagnosis of CMP, 12 (41.4%) showed evidence of mild MCD, one (3.4%) had moderate myocarditis, and two (6.9%) had no pathological lesions. Hearts from two *K. sima* were examined: one had evidence of CMP, the other of MCD. Nineteen of 27 (70.3%) *Kogia* spp. with evidence of cardiac pathology came from the southeast Atlantic region that included the east coasts of Florida, Georgia, South Carolina, and North Carolina.

The age and sex distribution of cardiac lesions is shown in Table 2. Lesions occurred predominantly among adult animals. Twenty-five of 27 (92.6%) affected animals were adults and two

Table 1. Cardiac pathology and geographic distribution in 27 pygmy (*Kogia breviceps*) and 2 dwarf (*Kogia sima*) sperm whales

Region	Cardiomyopathy	Myocardial degeneration	Myocarditis	No lesions
SE Atlantic ¹	9	9	1	2
NE Atlantic ²	1	--	--	--
Gulf of Mexico ³	4	3	--	--
Totals	14	12	1	2

¹Florida (east coast), Georgia, South Carolina, North Carolina

²Maryland, Delaware, New Jersey, New York, Connecticut, Rhode Island, Massachusetts, New Hampshire, Maine

³Florida (west coast), Alabama, Mississippi, Louisiana, Texas

Table 2. Cardiac pathology in 27 pygmy (*Kogia breviceps*) and two dwarf¹ (*Kogia sima*) sperm whales by age group and sex

Age class	Cardiomyopathy		Myocardial degeneration		Myocarditis		No lesions	
	Males	Females	Males	Females	Males	Females	Males	Females
Adults	10	3	8	3	1	--	--	--
Subadults	1	--	--	1	--	--	--	--
Calves	--	--	--	--	--	--	--	2
Totals	11	3	8	4	1	0	0	2

¹Includes one *K. sima* male with CMP and one *K. sima* male with MCD

(7.4%) were subadults. No pathological findings were observed in the hearts of the two stranded female calves that were examined. Male whales accounted for the majority of myocardial lesions. Twenty of the 27 (74%) affected whales were male; the excess of males occurred for both CMP (78.6% male) and MCD (66.7% male). Lesions of CMP and MCD were further categorized by severity. Eight of the 14 cases of CMP were categorized as severe, five as moderate, and one as mild. All forms of moderate and severe CMP were observed among adult animals. All cases of MCD were classified as mild.

Among *Kogia* with CMP, anisokaryosis with karyomegaly and nuclear rowing was the most common histological finding (Figures 1, 2 & 3) and was observed in 133 of the 168 (79.2%) tissue blocks examined in the 14 affected whales. In descending frequency, the next most common histological lesions were interstitial edema (61.3%), myofiber disarray (26.2%) (Figure 4), loss of cross-striations (20.2%) (Figure 1), eosinophilic homogenization of sarcoplasm (20.2%), and interstitial edema with fibrosis (17.3%) (Figures 3 & 5). In most cases, the anatomic distribution of each lesion was relatively similar across the 12 sampling sites. There were some exceptions to this observation, however. For example, the frequency of detection of anisokaryosis varied between interventricular septal sites. The distribution of

histopathological lesions of CMP in *Kogia* spp. was analyzed according to their anatomical location; the results are shown in Appendix 1.

Anisokaryosis with karyomegaly and nuclear rowing was also the predominant histological lesion in the 12 *Kogia* with MCD. This lesion was observed in 106 of the 144 (73.6%) tissue blocks examined. The frequency of other histological lesions generally followed the pattern observed in whales with CMP, with interstitial edema (37.5%) the next most commonly found lesion. The frequency of myofiber disarray (5.6%), loss of cross-striations (6.9%), eosinophilic homogenization of sarcoplasm (1.4%), and interstitial edema with fibrosis (2.1%) were substantially lower than those observed in cases of CMP. As was found for CMP, the anatomical distribution of each lesion of MCD was relatively similar across the 12 sampling sites. The distribution of MCD lesions by anatomical location is shown in Appendix 2.

A single case of moderate, multifocal chronic myocarditis without concurrent CMP or MCD was also present. The predominant inflammatory cell type was mature lymphocytes. No infectious agents were observed, and the etiology was not determined.

No statistically significant differences were found between pygmy sperm whales with CMP and those with MCD for any of the 11 parameters tested. Differences in heart weight; right and left




Figure 1. Photomicrograph of heart of a pygmy sperm whale with cardiomyopathy; note the anisokaryosis with karyomegaly (small arrow), loss of cross-striations (arrowhead), and interstitial edema (large arrow). H&E stain; bar = 80 microns.




Figure 2. Photomicrograph of heart of a pygmy sperm whale with cardiomyopathy; note the nuclear rowing of cardiomyocytes (arrows). H&E stain; bar = 80 microns.




Figure 3. Photomicrograph of heart of a pygmy sperm whale with cardiomyopathy; note the interstitial edema (arrow) and presumptive fibrosis (arrowhead). H&E stain; bar = 150 microns




Figure 4. Photomicrograph of heart of a pygmy sperm whale with cardiomyopathy; note the myofiber disarray and architectural disorganization of cardiomyocytes (arrows). H&E stain; bar = 150 microns.




Figure 5. Photomicrograph of heart of a pygmy sperm whale with cardiomyopathy; note the diffuse thickened blue fibrillar deposits of collagen. Masson trichrome stain; bar = 120 microns

ventricular wall thickness at two levels; intraventricular septum thickness at two levels; and circumference of the aortic, pulmonary, mitral, and tricuspid valves were evaluated. The largest differences were found for heart weight and intraventricular septum wall thickness, but none were statistically significant (Table 3). An ANOVA with sex in the model was run to determine whether sex could be acting as a confounder since males would be expected to be larger and were selectively affected. The mean heart weight for *K. breviceps* with CMP was greater than that for MCD for both males and females, but the differences were not statistically significant ($p = 0.14$) (Table 4). Finally, body length was used as a surrogate for body weight to adjust the heart weight data in PROC GLM. The adjusted least square mean heart weight for *K. breviceps* with CMP (1.63 kg)

Table 3. Heart weight, wall thickness, and valve circumference in *Kogia breviceps* with cardiomyopathy and myocardial degeneration

Cardiac parameter	Cardiomyopathy		Myocardial degeneration		P value
	Mean	SD	Mean	SD	
Heart weight (kg)	1.7	0.27	1.4	0.53	0.12
L2 RV wall (cm)	0.5	0.18	0.6	0.25	0.92
L2 LV wall (cm)	1.0	0.47	1.1	0.61	0.54
L2 IV wall (cm)	1.7	0.68	1.3	0.44	0.20
L4 RV wall (cm)	0.7	0.28	0.9	0.60	0.45
L4 LV wall (cm)	1.2	0.58	1.2	0.49	0.86
L4 IV septum (cm)	2.1	0.80	1.8	0.45	0.37
Pulmonary valve (cm)	12.6	1.84	11.8	2.71	0.47
Aortic valve (cm)	11.9	2.34	10.8	2.31	0.38
Tricuspid valve (cm)	19.6	5.24	16.9	2.63	0.26
Mitral valve (cm)	16.3	3.72	14.5	3.28	0.33

Table 4. Heart weight by sex in *Kogia breviceps* with cardiomyopathy and myocardial degeneration

Sex	Number	Cardiomyopathy		Myocardial degeneration		
		Mean weight (kg)	SD	Number	Mean weight (kg)	SD
Males	9	1.7	0.26	8	1.4	0.50
Females	2	1.8	0.42	3	1.4	0.71
Totals	11			11		

*p = 0.14

was greater than that for MCD (1.52 kg), but the difference was not statistically significant (data not shown).

Discussion

The pathological data presented in this report suggest that CMP is a common lesion in *Kogia* spp. that stranded in the coastal waters of the U.S. Atlantic and Gulf of Mexico from 1999 to 2006. These data support and extend a past report of *Kogia* spp. that stranded in the same geographic region from 1980 to 1984 (Bossart et al., 1985). All of the whales in the present study represent single strandings. The consistency, extent, and severity of the pathological lesions suggest that many of these whales were in a state of myocardial decompensation at the time of stranding. Furthermore, this study provides new evidence indicating that *Kogia* CMP is a chronic progressive condition rather than an acute terminal event. Specifically, the similarity in type and distribution of pathological lesions of CMP and MCD implies a continuum of the same process. Thus, MCD may precede and lead to the development of CMP as the whales age. The apparent progressive nature of CMP ultimately would result in a debilitating condition in adults; and the beaching

event of the ailing whales would be dependent on the active movement (or lack of purposeful directional movement) and prevailing environmental conditions such as ocean steering currents and weather conditions. The general trend of greater heart weights in *K. breviceps* with CMP compared to those with MCD supports the hypothesis that the disorder is a progressive condition since heart weight may be increased with CMP in other species (see below). It is important to note that none of the heart weight, wall thickness, or valve circumference differences were statistically significant and that the statistical analyses compared whales with CMP to whales with MCD; an unaffected comparison group was not available.

In terrestrial mammals and humans, the general types of CMP include stress, hypertrophic, dilated (congestive), and restrictive forms. The myocardial lesions of the stress, dilated, and hypertrophic CMP types described in other species were observed in the present study.

The stress form represents an acute process mediated by catecholamines, which may lead to sudden death in humans and animals, usually without a history of pre-existing heart disease (Cebelin & Hirsch, 1980; Liu et al., 1982). Elevated endogenous catecholamines produce distinctive focal hypercontraction and lysis of contractile filaments

in small groups of cardiomyocytes that is termed “contraction band necrosis” (Turnbull & Cowan, 1998). Contraction band necrosis, including loss of cross-striations, interstitial edema, myofiber cytoplasmic hypereosinophilia, and wavy fibers, have been reported previously in the *Kogia* CMP and in other stranded cetaceans (Bossart et al., 1985; Turnbull & Cowan, 1998). Additionally, a more chronic stress hormone component involving cortisol has recently been described in dogs with dilated CMP (Tidholm et al., 2005).

Hypertrophic CMP is well characterized in humans and domestic animals (Liu et al., 1993, 1994; Maron, 1997). The demonstration of specific genetic abnormalities in cardiac energy metabolism or structural and contractile proteins results in approximately half the human cases of hypertrophic CMP (Hughes, 2004). The diagnosis of this form is based on macroscopic enlargement of the heart usually supported by microscopic lesions consisting of myofiber disarray (architectural disorganization). In *Kogia*, heart enlargement is difficult to assess because normal heart weights have not been determined; however, the microscopic lesion of myofiber disarray associated with hypertrophic CMP in other species was seen in 26% of *Kogia* with CMP.

In about half of the human patients with hypertrophic CMP, the disease is familial and is one of the most common causes of a sudden, unexplained death in young male athletes (Schoen, 1999). The preponderance of male whales with myocardial pathology in the current study was a new and interesting finding. In contrast, an earlier study did not find an unusual sex distribution (Bossart et al., 1985). The significance of the excess of CMP among male *Kogia* is unknown but may suggest a sex-linked genetic etiology.

Dilated CMP differs from the hypertrophic form in that the capacity of the ventricle(s) is actually increased, which can impart a “globular” appearance to the heart. In the fresh unfixed heart, the ventricle(s) may feel flabby. Dilated CMP in humans has varied etiologies that may involve complex mechanisms, including postinfectious, autoimmune, and idiopathic factors (Richardson et al., 1996). Dilated CMP also has been associated with L-carnitine and taurine deficiencies in humans, rodents, and domestic animals (Levitan et al., 1987; Keene, 1991; Fascetti et al., 2003; Zaugg et al., 2003). Dilated CMP was recently reported in southern California sea otters (*Enhydra lutris nereis*) and postulated to be associated with domoic acid toxicosis and depletion of myocardial L-carnitine (Kreuder et al., 2005).

The first report of CMP in *Kogia* was a dilated form, which included a grossly dilated flabby right ventricle, generalized myocardial pallor, and

chronic passive congestion of the liver (Bossart et al., 1985). The etiology of this case of dilated CMP could not be determined, but nutritional etiologies, including a thiamine deficiency, were postulated. Thiamine deficiency has been reported in captive marine mammals, and myocardial lesions consistent with thiamine deficiency were seen in captive sea lions fed a diet presumably containing high concentrations of thiaminase (Rigdon & Drager, 1955; Worthy, 2001). In this study, it was difficult to assess the occurrence of gross myocardial changes of dilated CMP as all of the examined hearts had already been fixed, thus distorting normal gross morphology. Previously described microscopic heart lesions of the dilated form were found, however, and these consisted of cardiomyocyte degeneration, loss of cross-striations, interstitial edema, and fibrosis (Bossart et al., 1985).

Although each form of CMP is fundamentally different, they are not necessarily mutually exclusive in a given case. Moreover, transitions from one type to another may occur in humans, reflecting chronicity and/or severity of the basic disease process (Hughes & McKenna, 2005). Specifically, hypertrophic CMP may progress to a dilated phase in human patients and resemble dilated CMP (Maron, 2002). Therefore, it appears that the *Kogia* CMP may be best defined as a “mixed form,” having microscopic components of all three types. Lesions seen uniformly in all sections included eosinophilic homogenization of sarcoplasm, loss of cross-striations, interstitial edema and fibrosis, anisokaryosis with karyomegaly, myofiber disarray (architectural disorganization), and wavy-attenuated myofibers. Thus, the etiology of CMP in *Kogia* is likely complex and multifactorial. Etiologic components may include metabolic factors, such as excessive repeated sublethal episodes of catecholamine release (repeated acute “stress” reactions) and endogenous glucocorticoid release (chronic “stress” response); nutritional deficiencies; and postinfectious, genetic, and toxic factors (e.g., biotoxins). Further studies may help confirm these hypotheses.

Acknowledgments

This work was supported by Prescott Grant #NA05NMF4391182 from the National Marine Fisheries Service and Harbor Branch Oceanographic Institution’s “Protect Florida Whales” program. Marine mammal tissues were collected, analyzed, and archived in accordance with the NOAA Fisheries Permits to the Marine Mammal Health and Stranding Response Program (No. 932-1489-01). We thank Wayne McFee (NOAA/NOS/NCCOS/CCEHBR, Charleston, South Carolina, USA)

and Megan K. Stolen and Wendy Noke Durden (Hubbs-Sea World Research Institute, Orlando, Florida, USA) for heart acquisition. Special thanks go to Dr. Dan Odell for natural history information and Dr. Ruth Ewing for initial assistance in the heart dissection technique. Additionally, we gratefully acknowledge the volunteer members of the Southeastern Marine Mammal Stranding Network and Harbor Branch marine mammal volunteers for their tireless efforts in advancing the science of marine mammal medicine and pathology.

Literature Cited

- Bossart, G. D., Odell, D. K., & Altman, N. H. (1985). Cardiomyopathy in stranded pygmy and dwarf sperm whales. *Journal of the American Veterinary Medical Association*, *187*, 1137-1140.
- Cebilin, M. S., & Hirsch, C. S. (1980). Human stress cardiomyopathy: Myocardial lesions in victims of homicidal assaults without internal injuries. *Human Pathology*, *11*, 123-132.
- Chiu, J., Chiou, T., & Chou, L. (2003). Pathological examinations on the stranded *Kogia* sp. from Taiwan waters, 1998-2002. *Proceedings of the 34th Annual Conference of the International Association for Aquatic Animal Medicine*, Waikoloa, HI. 223 pp.
- Fascetti, A. J., Reed, J. R., Rogers, Q. R., & Backus, R. C. (2003). Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997-2001). *Journal of the American Veterinary Medical Association*, *223*, 1137-1141.
- Hensley, G., Bossart, G. D., Ewing, R., Varela, R., Murdoch, E., Heym, K., et al. (2005). *Kogia heart dissection manual* (Harbor Branch Oceanographic Institution Technical Report 90). Fort Pierce, FL: HBOI.
- Hughes, S. E. (2004). The pathology of hypertrophic cardiomyopathy. *Histopathology*, *44*, 412-427.
- Hughes, S. E., & McKenna, W. J. (2005). New insights into the pathology of inherited cardiomyopathy. *Heart*, *91*, 257-264.
- Keene, B. W. (1991). Myocardial L-carnitine deficiency in a family of dogs with dilated cardiomyopathy. *Journal of the American Veterinary Medical Association*, *198*, 647-650.
- Kreuder, C., Miller, M. A., Lowenstine, L. J., Conrad, P. A., Carpenter, T. E., Jessup, D. A., et al. (2005). Evaluation of cardiac lesions and risk factors associated with myocarditis and dilated cardiomyopathy in southern sea otters (*Enhydra lutris nereis*). *American Journal of Veterinary Research*, *66*, 289-299.
- Levitan, M., Murphy, J., & Sherwood, W. (1987). Adult onset systemic carnitine deficiency: Favorable response to L-carnitine supplementation. *Canadian Journal of Neurological Science*, *14*, 5054.
- Liu, S-K., Chiu, Y. T., & Shyu, J. J. (1994). Hypertrophic cardiomyopathy in pigs: Quantitative pathologic features in 55 cases. *Cardiovascular Pathology*, *3*, 261-268.
- Liu, S-K., Dolensik, E. P., & Herron, A. J. (1982). Myopathy in a nyala. *Journal of the American Veterinary Medical Association*, *181*, 1232-1236.
- Liu, S-K., Roberts, W. C., & Maron, B. J. (1993). Comparison of morphologic findings in spontaneously occurring hypertrophic cardiomyopathy in humans, cats and dogs. *The American Journal of Cardiology*, *72*, 944-951.
- Maron, B. J. (1997). Hypertrophic cardiomyopathy. *Lancet*, *350*, 127-133.
- Maron, B. J. (2002). Hypertrophic cardiomyopathy: A systematic review. *Journal of the American Medical Association*, *287*, 1308-1320.
- Odell, D. K., Barros, N. B., & Stolen, M. K. (2004). *Dwarf and pygmy sperm whale (genus Kogia) stranding patterns in the southeastern United States*. 84th Annual meeting of the American Society of Mammalogists, June 12-16, Arcata, CA. Retrieved 14 May 2007 from <http://abstracts.co.allenpress.com/pweb/asm2004/document/?ID=39074>.
- Pickrell, J. (2003). Whale beachings linked to mysterious heart defect. Retrieved 20 September 2006 from http://news.nationalgeographic.com/news/2003/08/0806_030806_-whaleheart.html.
- Richardson, P., McKenna, W., Bristow, M., Maisch, B., Mautner, B., O'Connell, J., et al. (1996). Report of the 1995 World Health Organization/International Society and Federation of Cardiology Task Force on the definition and classification of cardiomyopathies. *Circulation*, *93*, 841-842.
- Rigdon, R. H., & Drager, G. A. (1955). Thiamine deficiency in sea lions (*Otaria californiana*) fed only frozen fish. *Journal of the American Veterinary Medical Association*, *127*(944), 453-455.
- Schoen, F. J. (1999). The heart. In R. S. Cotran, V. Kumar, & T. Collins (Eds.), *Robbins pathologic basis of disease* (pp. 543-599). Philadelphia: W. B. Saunders.
- Tidholm, A., Haggstrom, J., & Hansson, K. (2005). Vasopressin, cortisol, and catecholamine concentrations in dogs with dilated cardiomyopathy. *American Journal of Veterinary Research*, *66*(10), 1709-1717.
- Turnbull, B. S., & Cowan, D. F. (1998). Myocardial contraction band necrosis in stranded cetaceans. *Journal of Comparative Pathology*, *118*, 317-327.
- Worthy, G. A. (2001). Nutrition and energetics. In L. A. Dierauf & F. M. D. Gulland (Eds.), *Marine mammal medicine* (pp. 791-827). Boca Raton, FL: CRC Press. 1,063 pp.
- Zaugg, C. E., Spaniol, M., Kaufmann, P., Bellahcene, M., Barbosa, V., Tolnay, M., et al. (2003). Myocardial function and energy metabolism in carnitine deficient rats. *Cellular and Molecular Life Sciences*, *60*, 767-775.

Appendix 1. Distribution of histopathological lesions of cardiomyopathy by anatomic site (*n* = 14)

Histological lesion	SS-1	SS-2	DRV-2	VRV-2	DLV-2	VLV-2	IV-2	DRV-4	VRV-4	DLV-4	VLV-4	IV-4	Total
Anisokaryosis with karyomegaly and nuclear rowing	9 (64%)	9 (64%)	11 (79%)	12 (86%)	12 (86%)	11 (79%)	7 (50%)	13 (93%)	12 (86%)	12 (86%)	11 (79%)	14 (100%)	133 (79.20%)
Interstitial edema	8 (57%)	5 (36%)	9 (64%)	9 (64%)	8 (57%)	9 (64%)	6 (43%)	10 (71%)	9 (64%)	9 (64%)	9 (64%)	12 (86%)	103 (61.30%)
Myofiber disarray (architectural disorganization)	1 (7%)	2 (14%)	4 (29%)	4 (29%)	4 (29%)	5 (36%)	3 (21%)	3 (21%)	3 (21%)	3 (21%)	5 (36%)	7 (50%)	44 (26.20%)
Loss of cross-striations	1 (7%)	2 (14%)	1 (7%)	3 (21%)	5 (36%)	4 (29%)	1 (7%)	2 (14%)	2 (14%)	5 (36%)	4 (29%)	4 (29%)	34 (20.20%)
Myofiber cytoplasmic hypereosinophilia	2 (14%)	2 (14%)	1 (7%)	2 (14%)	2 (14%)	5 (36%)	2 (14%)	3 (21%)	4 (29%)	4 (29%)	4 (29%)	3 (21%)	34 (20.20%)
Interstitial edema and fibrosis	0 (0%)	5 (36%)	2 (14%)	1 (7%)	4 (29%)	3 (21%)	1 (7%)	3 (21%)	2 (14%)	3 (21%)	3 (21%)	2 (14%)	29 (17.30%)
Wavy-attenuated myofibers	0 (0%)	1 (7%)	1 (7%)	2 (14%)	1 (7%)	1 (7%)	2 (14%)	1 (7%)	2 (14%)	2 (14%)	2 (14%)	2 (14%)	17 (10.10%)

SS-1 = Septum summit to the right of the midline
 SS-2 = Septum summit to the left of the midline
 DRV-2 = Dorsal right ventricle at Level 2
 VRV-2 = Ventral right ventricle at Level 2
 DLV-2 = Dorsal left ventricle at Level 2
 VLV-2 = Ventral left ventricle at Level 2
 IV-2 = Interventricular septum at Level 2
 DRV-4 = Dorsal right ventricle at Level 4
 VRV-4 = Ventral right ventricle at Level 4
 DLV-4 = Dorsal left ventricle at Level 4
 VLV-4 = Ventral left ventricle at Level 4
 IV-4 = Interventricular septum at Level 4

Appendix 2. Distribution of histopathological lesions of myocardial degeneration by anatomical site (*n* = 12)

Histological lesion	SS-1	SS-2	DRV-2	VRV-2	DLV-2	VLV-2	IV-2	DRV-4	VRV-4	DLV-4	VLV-4	IV-4	Total
Anisokaryosis with karyomegaly and nuclear rowing	9 (75%)	9 (75%)	6 (50%)	7 (58%)	11 (92%)	11 (92%)	10 (83%)	5 (42%)	8 (67%)	9 (75%)	9 (75%)	12 (100%)	106 (73.6%)
Interstitial edema	4 (33%)	3 (25%)	4 (33%)	4 (33%)	5 (42%)	6 (50%)	5 (42%)	2 (17%)	2 (17%)	5 (42%)	6 (50%)	8 (67%)	54 (37.5%)
Myofiber disarray (architectural disorganization)	1 (8%)	0 (0%)	2 (17%)	0 (0%)	1 (8%)	0 (0%)	2 (17%)	0 (0%)	0 (0%)	1 (8%)	1 (8%)	0 (0%)	8 (5.60%)
Loss of cross-striations	1 (8%)	1 (8%)	0 (0%)	0 (0%)	1 (8%)	2 (17%)	2 (17%)	0 (0%)	0 (0%)	1 (8%)	0 (0%)	2 (17%)	10 (6.9%)
Myofiber cytoplasmic hypereosinophilia	1 (8%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (8%)	0 (0%)	0 (0%)	2 (1.40%)
Interstitial edema and fibrosis	1 (8%)	1 (8%)	0 (0%)	0 (0%)	0 (0%)	1 (8%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	3 (2.10%)
Wavy-attenuated myofibers	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (8%)	0 (0%)	1 (0.7%)

SS-1 = Septum summit to the right of the midline

SS-2 = Septum summit to the left of the midline

DRV-2 = Dorsal right ventricle at Level 2

VRV-2 = Ventral right ventricle at Level 2

DLV-2 = Dorsal left ventricle at Level 2

VLV-2 = Ventral left ventricle at Level 2

IV-2 = Interventricular septum at Level 2

DRV-4 = Dorsal right ventricle at Level 4

VRV-4 = Ventral right ventricle at Level 4

DLV-4 = Dorsal left ventricle at Level 4

VLV-4 = Ventral left ventricle at Level 4

IV-4 = Interventricular septum at Level 4

IV-2 = Interventricular septum at Level 2

DRV-4 = Dorsal right ventricle at Level 4

VRV-4 = Ventral right ventricle at Level 4

DLV-4 = Dorsal left ventricle at Level 4

VLV-4 = Ventral left ventricle at Level 4

IV-4 = Interventricular septum at Level 4

IV-4 = Interventricular septum at Level 4

From: [Edwards, David](#)
To: [Hartogenesis, Martine](#); [Murphy, Jeanette](#)
Subject: talk to AFIA Pet Food Conference
Date: Wednesday, January 30, 2019 11:25:00 AM
Attachments: [Final Agenda - Pet Food Conference 092718.docx](#)

Hi Martine and Jenny,

I am slated to talk to the AFIA Pet Food Conference on February 12 in Atlanta. They would like an update on DCM and on FSMA inspections. Do you have a recent presentation that I can update to provide the latest? I have 45 minutes total for talking and questions. Greg Aldrich is speaking after me on DCM, and AAFCO is also on the program to provide updates (see attached).

If I need to ping others, please let me know.

Thank you,
Dave

David Edwards, PhD | Director, Division of Animal Feeds
(240)402-6205 | david.edwards@fda.hhs.gov



2019 Pet Food Conference
Tuesday, February 12, 2019
Atlanta, Ga.
7:30 a.m. to 4:00 p.m.

As of 9/27/18

7:30-8:30 a.m.	Breakfast
8:30-8:40 a.m.	Welcome and Overview of the Conference
8:40-9:30 a.m.	Domestic and Global Industry Trends Jared Koerten, Euromonitor International (http://blog.euromonitor.com/)
9:30-10:00 a.m.	Trade Policy and Outlook Gina Tumbarello, American Feed Industry Association
10:00-10:30 a.m.	Break
10:30-11:15 a.m.	Blockchain Management of Pet Food: A Legal Perspective John Dillard, Olsson Frank Weeda Terman Matz PC
11:15-12:00 p.m.	The Healing Power of Man's Best Friend: Opportunities to Save and Enrich Human Lives Robin Ganzert, Ph.D., American Humane
12:00-1:15 p.m.	Networking Lunch (provided)
1:15-2:00 p.m.	FSMA Inspections and Canine Dilated Cardiomyopathy Update David Edwards, Ph.D., FDA Center for Veterinary Medicine
2:00-2:45	Implications of Canine Dilated Cardiomyopathy on Ingredient Categories Greg Aldrich, Ph.D., Kansas State University
2:45-3:15	Break
3:15-4:00	AAFCO Updates on Pet Food Sue Hays, Association of American Feed Control Officials
4:00 p.m.	Wrap up and Adjourn

**** Agenda Subject to Change

Patient Information

Patient: (b) (6) Age: 8 years Referring Veterinarian: (b) (6)
 Patient Number: (b) (6) Weight:(kg) 29.40 Cardiologist: (b) (6) DVM, DACVIM
 (Cardiology)
 Breed: Labrador Retriever Sex: F Client Number: (b) (6)
 Exam Date: (b) (6) 08:22 BSA: 0.96

History: (b) (6) was presented to the (b) (6) last night for transfer to Cardiology for further evaluation of her heart. She was seen by (b) (6) last week after collapsing last Thursday while playing fetch with her owner. On presentation at (b) (6) she was found to be in atrial fibrillation with evidence of mild heart failure. She was treated with a diltiazem CRI overnight, then oral diltiazem, as well as Lasix, spironolactone, and enalapril. Her heart rhythm converted back to sinus rhythm as of (b) (6) (Friday). She was presented back to (b) (6) on Saturday after collapsing again on Saturday while playing fetch. She was found to still be in a normal heart rhythm and radiographs showed resolution of heart failure at that time. Bloodwork done at (b) (6) (CBC and chem) was reported as unremarkable.

Physical Examination: Grade 3-4/6 left apical holosystolic murmur. Irregular rhythm consistent with sinus arrhythmia. Clear lungs. Moderate femoral pulses. Normal abdominal palpation. Well hydrated. Normal PLNs. mm pink, CRT normal

Diagnostic Tests: BP: 108mmHg 4cm cuff RR

Echo: see below

Telemetry -(b) (6) heart rhythm was monitored throughout her hospital stay and showed a consistent sinus rhythm/arrhythmia with no significant dysrhythmias.

Renal panel pending

Echocardiographic Report

2D-ECHO

LA Systolic Diameter LX	6.3 cm	Aortic Root Diameter	2.3 cm
-------------------------	--------	----------------------	--------

DOPPLER

AV Peak Velocity	104 cm/s	PV Peak Velocity	72.7 cm/s
AV Peak Gradient	4.3 mmHg	PV Peak Gradient	2.1 mmHg
Mitral E Point Velocity	156 cm/s	TR Peak Velocity	343 cm/s
Mitral E to A Ratio	3	TR Peak Gradient	47.2 mmHg
MR Peak Velocity	521 cm/s		

M-MODE

LV Diastolic Diameter MM	6.8 cm	LVPW Diastolic Thickness MM	0.92 cm
LV Systolic Diameter MM	5.8 cm	LVPW Systolic Thickness MM	1.1 cm
LV Fractional Shortening MM	15.1 %	LVPW Percent Thickening MM	0.16
LV Diastolic Volume Cube	314 cm ³	IVS to PW Ratio MM	1.3
LV Systolic Volume Cube	192 cm ³	LV Mass MM	323 g
LV Ejection Fraction Cube	0.39	LV Mass Normalized MM	336 g/m ²
IVS Diastolic Thickness MM	1.2 cm	LA Systolic Diameter MM	3.8 cm
IVS Systolic Thickness MM	1.4 cm	Aortic Root Diameter MM	2.3 cm
IVS Percent Thickening MM	0.22	MV E Point Septal Separation	1.5 cm

Left Ventricle: Dilated, rounded, and poorly contractile chamber.

Left Atrium: Moderate dilation with marked dilation of right pulmonary vein.

Right Ventricle: Normal.

Right Atrium: Normal.

Mitral Valve: Mildly thickened valve leaflets. 4+ eccentric regurgitation. High inflow velocity with restrictive filling pattern.

Aortic Valve: Normal.

Tricuspid Valve: Thickened valve leaflets with multiple 1+ jets of regurgitation. TR velocity is increased consistent with mild pulmonary hypertension.

Pulmonic Valve: Mild valve thickening. 1+ regurgitation. PI velocity is not suggestive of diastolic pulmonary hypertension.

Aorta: Normal.

Pericardium: Normal.

Diagnosis

Dilated cardiomyopathy - This is a disease characterized by weakening of the heart muscle and dilation of the heart chambers. It is most commonly an inherited disease, but can occur as a consequence of other injuries to the heart. Severe valvular heart disease can sometimes lead to heart muscle failure (cardiomyopathy of overload) and since (b) (6) appears to have severe valve disease as well as heart muscle failure, we cannot be sure whether one led to the other or if there are two completely separate disease processes. As the disease progresses, it can lead to congestive heart failure (fluid in the lungs causing shortness of breath and cough). Abnormal heart rhythms are common and can result in sudden death. Most commonly this is an inherited disease, though it can occur secondary to a deficiency in an amino acid called taurine.

Chronic degenerative valve disease - Degenerative changes in one or more heart valves have caused leaking across these valves. This is the source of the heart murmur. As this disease progresses, the heart enlarges. Eventually this can lead to symptoms of cough and shortness of breath (airway compression and/or congestive heart failure).

Atrial fibrillation on presentation at (b) (6), converted back to sinus rhythm (b) (6) - This is a chaotic and rapid heart rhythm from the upper heart chambers. It most commonly occurs secondary to severe underlying heart diseases, though it can occur in isolation in some giant breed dogs. Our goal medically in treating this arrhythmia is to control the heart rate, but (b) (6) has returned to a normal heart rhythm so no specific medication is indicated for the heart rhythm at this time.

Exertional collapse - I suspect the first episode was likely caused by the new onset of the atrial fibrillation in (b) (6), but the second episode is a little harder to explain. We did not find any evidence while monitoring her in the hospital of other arrhythmia, and she had a normal heart rhythm at the emergency visit after her second collapse as well. It is possible that she collapsed as a result of her severe structural heart disease, though this is a little surprising to see recurrent collapse after starting on medications that had been effective in resolving her heart failure.

Recommendations

Please DISCONTINUE:

Diltiazem - This is a medication to slow the heart rate in atrial fibrillation, but it has not been shown to be effective in decreasing the risk of recurrent atrial fibrillation once an animal converts back to sinus rhythm. Since this is the case with (b) and she is no longer in atrial fibrillation, we do not need to keep her on this medication at this time.

Please CONTINUE:

Furosemide (Lasix) 50mg tablets - Give 1 tablet by mouth once every 12 hours. Furosemide (Lasix, Salix) - This is a diuretic (water pill), that prevents the body from retaining excessive sodium and water. It will cause your pet to drink and urinate more frequently. It is important that fresh water is always available.

Spirolactone (Aldactone) 50mg tablets - Give 1 tablet by mouth once daily. This is a diuretic (water pill) that also blocks a hormone that can injure the heart muscle. It works well in combination with the furosemide and enalapril.

Please INCREASE:

Enalapril (Enacard, Vasotec) 10mg tablets - Give 1 and 1/2 tablets by mouth every 12 hours. This medication is a strong drug that dilates blood vessels, permitting the heart to pump blood more efficiently. It can lower blood pressure (hypotension) and cause changes in kidney function and electrolyte values. If your pet develops weakness or depression, decrease the drug dose by 1/2 and call. A kidney panel and blood pressure should be reevaluated 7-10 days after beginning this medication.

Please ADD:

Pimobendan (Vetmedin) 5mg tablets - Give 1 and 1/2 tablets by mouth every 12 hours. This is a drug that is approved for the treatment of congestive heart failure secondary to dilated cardiomyopathy or chronic valve disease (endocardiosis). Studies have shown improved quality of life and increase survival time when this drug is added to other standard cardiac medications. In our experience, side effects are uncommon, but it is important that you advise us if you feel your pet is having any potential adverse effects from this medication. The reported potential side effects listed for this medication are increased heart rate, vomiting, diarrhea, inappetance, uneasiness, incoordination, convulsions, increased drinking and increase urinating.

**Please NOTE: When 10mg tablets of Vetmedin become available again, it would be OK to switch to one 10mg tablet in the morning and 1/2 - 10mg tablet at night for the same total daily dose and this would be more cost effective. Currently, however, the 10mg tablets are on backorder and we do not have a confirmed release date.

With advanced heart disease, our biggest dietary concerns are adequate calorie content and low sodium content. We aim for less than 80mg sodium per 100 kilocalories (kcal) in patients that have developed congestive heart failure. We do not advise protein restriction unless there is concurrent kidney disease (i.e. kidney diets are not advised unless there is concurrent kidney disease). Please refer to our diet handouts with a list of currently adequate diets and treats, though this list is not exclusive. If you wish to feed a diet that is not on these lists, you will need to call the manufacturer of the diet to obtain a sodium content.

One thing that can be very helpful for home monitoring is checking sleeping or resting respiratory rates. A recent study showed that even dogs with severe heart disease rarely have resting respiratory rates greater than 30 breaths per minute unless they are starting to decompensate for that disease. Elevated respiratory rates at home may be even more sensitive than chest radiographs at picking up early decompensation. Count your pet's respiratory rate when he/she is at rest or sleeping (not within 20 minutes of being active). If his/her respiratory rate is greater than 30 breaths per minute, recheck again in a couple of hours. If persistently elevated above this level, call.

Exercise is also a concern in advanced heart disease. While cage rest is ideal with active heart failure, some exercise is permissible in asymptomatic disease. However, vigorous or extended exercise should be avoided.

Please call if you have any concerns about (b), if she develops an increase in respiratory rate or effort, has a persistent cough, or has any further collapse episodes. As long as she is doing well, we will plan to recheck her again in another month and will recheck her heart rhythm, chest radiographs, and kidney panel at that time.

(b) (6) (Cardiology)

(Electronically Signed)

Final Date: (b) (6) 16:50

Amended: (b) (6) 17:16

Like us on Facebook!

www.facebook.com/ (b) (6)

Notes to our clients

- Please bring all medications to your pet's scheduled appointments.
- We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).
- Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.
- If an emergency arises with your pet, (b) (6) is a 24 hour facility.

Left Atrium: Moderate dilation, minimal decrease since initial study.
Right Ventricle: Normal.
Right Atrium: Normal.
Mitral Valve: Mildly thickened valve leaflets. 3-4+ regurgitation.
Aortic Valve: Normal.
Tricuspid Valve: I+ regurgitation. TR velocity consistent with normal pulmonary pressures.
Pulmonic Valve: I+ regurgitation. Normal PI velocity.
Aorta: Normal.
Pericardium: Normal.

Diagnosis

Dilated Cardiomyopathy
Chronic Degenerative Valve Disease
Historical atrial fibrillation with collapse (b) (6) continues to be in a normal sinus rhythm today
Historical congestive heart failure - no evidence of heart failure today

(b) (6) echo today looks stable to slightly improved from his initial echo in January, though his heart is a little larger today than on the radiographs in February. He is showing no signs of recurrent heart failure and his heart rhythm is still normal. Overall, I am happy with where we are overall.

Recommendations

Please continue to give medications as directed:

Furosemide 40mg tablets- Give 1 tablet by mouth once every 12 hours.
Enalapril 10mg tablets- Give 1 and 1/2 tablets by mouth once every 12 hours.
Spironolactone 50mg tablets- Give 1 tablet by mouth once every 24 hours.
Vetmedin 5mg tablets- Give 1 and 1/2 tablets by mouth once every 12 hours.

ADD:

Trazodone 150mg tablets- Give 1 tablet by mouth up to once every 8 hours as needed for storm anxiety.

As long as (b) (6) continues to do well, we will continue to recheck her every 3-4 months with chest radiographs, renal panel, and blood pressure with periodic echocardiograms. Please call, however, if she develops any new or recurrent clinical symptoms.

(b) (6) (Cardiology)

(Electronically Signed)

Final Date: 31 May 2017 15:11

Like us on Facebook!

www.facebook.com/ (b) (6)

Notes to our clients

- Please bring all medications to your pet's scheduled appointments.
- We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).
- Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.
- If an emergency arises with your pet, (b) (6) is a 24 hour facility.

Patient Information

Patient:	(b) (6)	Age: 9	years	Referring Veterinarian:	(b) (6)
Patient Number:	(b) (6)	Weight:(kg)	32.10	Cardiologist:	(b) (6) (Cardiology)
Breed:	Lab	Sex:	F	Client Number:	(b) (6)
Exam Date:	12/11/2017 08:17	BSA:	1.02		

History: Reevaluation of dilated cardiomyopathy with chronic degenerative valve disease, historical atrial fibrillation with collapse, historical congestive heart failure, and urinary incontinence. (b) (6) is doing well at home. Owners do report a new cough with him since his last visit. It is not frequent and is seen at rest and with excitement/activity. She is breathing comfortably. She has a normal appetite and good activity level as well. Owners are transitioning her to a new brand of venison food.

Physical Examination: Temp 102.4. Heart rate 128bpm. RR pant. Grade 3/6 left apical systolic murmur with wide radiation. Regular rhythm. Clear lungs. Fair to moderate femoral pulses. Normal abdominal palpation and PLNs. Well hydrated.

Diagnostic Tests: Chest radiographs: progressive cardiomegaly with VHS 13.5 versus 13 on radiographs in September, normal pulmonary vessels, unchanged lung pattern with no evidence of active heart failure

Renal panel: within normal limits

Echo: see below. ECG during echo showed sinus rhythm.

Echocardiographic Report

2D ECHO

LA Systolic Diameter Lx	6.5 cm	Aortic Root Diameter	2 cm
-------------------------	--------	----------------------	------

DOPPLER

AV Peak Velocity	161 cm/s	PV Peak Velocity	70.7 cm/s
AV Peak Gradient	10.4 mmHg	PV Peak Gradient	2 mmHg
Mitral E Point Velocity	111 cm/s	TR Peak Velocity	269 cm/s
Mitral E to A Ratio	2	TR Peak Gradient	28.9 mmHg
MR Peak Velocity	487 cm/s		

M-MODE

LV Diastolic Diameter MM	6.6 cm	LVPW Systolic Thickness MM	1.2 cm
LV Systolic Diameter MM	5.9 cm	LVPW Percent Thickening MM	0.049
LV Fractional Shortening MM	11.3 %	IVS to PW Ratio MM	1
LV Diastolic Volume Cube	287 cm ³	LV Mass MM	348 g
LV Systolic Volume Cube	200 cm ³	LV Mass Normalized MM	341 g/m ²
LV Ejection Fraction Cube	0.3	RV Diastolic Diameter MM	0.68 cm
IVS Diastolic Thickness MM	1.2 cm	LA Systolic Diameter MM	3.5 cm
IVS Systolic Thickness MM	1.5 cm	Aortic Root Diameter MM	2.3 cm
IVS Percent Thickening MM	0.25	MV E Point Septal Separation	1.7 cm
LVPW Diastolic Thickness MM	1.1 cm		

Left Ventricle: Stable diastolic dimension with progressive increase in systolic dimension and decline in myocardial function.

Left Atrium: Progressive dilation.

Right Ventricle: Mild dilation.

Right Atrium: Mild dilation.

Mitral Valve: Unchanged mild thickening with 3-4+ regurgitation.

Aortic Valve: Normal. Acceleration slope is decreased.

Tricuspid Valve: Two jets of 2+ regurgitation. TR velocity consistent with normal pulmonary pressures.

Pulmonic Valve: Normal. 1+ physiologic regurgitation.

Aorta: Normal.

Pericardium: Normal.

Diagnosis

Dilated cardiomyopathy with chronic degenerative valve disease - (b) (6) heart is bigger and does not contract as well as it did at her last two rechecks. However, she is showing no signs of decompensation at this time.

Historical atrial fibrillation with collapse

Historical congestive heart failure

Urinary incontinence

Recommendations

Please give the following medications as directed:

Glucosamine and coat supplement

Incurin - give 2 tablets by mouth once daily in the mornings

Furosemide 40mg tablets - give 1 tablet by mouth every 12 hours

Enalapril 10mg tablets - give 1 and 1/2 tablets by mouth every 12 hours

Spironolactone 50mg tablets - give 1 tablet by mouth once daily in the mornings

Vetmedin 5mg tablets - INCREASE to 1 and 1/2 tablets by mouth EVERY 8 HOURS. (We discussed that we can go to 1/2 of a 10mg tablet + 1/2 of a 5mg tablet to save some in cost for this medication).

Please call if you have any questions or concerns about (b) (6). As long as she continues to do well, we will recheck her again in another 3-4 months. We will do a brief echo and recheck kidney values and blood pressure at that visit +/- chest radiographs (if she is having any respiratory symptoms).

(b) (6), DVM, DACVIM (Cardiology)

(Electronically Signed)

Final Date: 11 December 2017 14:48

Amended: 11 December 2017 14:49

Like us on Facebook!

www.facebook.com/ (b) (6)

Notes to our clients

- Please bring all medications to your pet's scheduled appointments.
- We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).
- Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.
- If an emergency arises with your pet, (b) (6) is a 24 hour facility.

From: Palmer, Lee Anne
To: Rotstein, David; Jones, Jennifer L; Carey, Lauren; Queen, Jackie L
Subject: RE: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls): (b) (6) EON-345965
Date: Thursday, January 25, 2018 1:23:37 PM

Woops, that's 2 from (b) (6) rather than (b) (6) Same household. thx!

From: Rotstein, David
Sent: Thursday, January 25, 2018 1:00 PM
To: Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Queen, Jackie L <Jackie.Queen@fda.hhs.gov>
Subject: RE: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls): (b) (6) - EON-345965

Fantastic!

Hope it goes well!

From: Palmer, Lee Anne
Sent: Thursday, January 25, 2018 12:57 PM
To: Rotstein, David <David.Rotstein@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Queen, Jackie L <Jackie.Queen@fda.hhs.gov>
Subject: RE: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls): (b) (6) - EON-345965

Hi – trying out the epi summary in EON to give it a dry run (Dave, I've temporarily assigned it to myself). FYI – this makes 6 cases now, 5 from (b) (6) 1 from (b) (6) (involving 2 dogs, 1 death) all eating LID type diet with Kangaroo and Red Lentil, onset dates between 1/20/2017 and 12/30/2017). I'll double check that info as I read through the reports and summarize it in the incident. Just had to try out the new functionality, needed some fun today.

From: Rotstein, David
Sent: Thursday, January 25, 2018 12:27 PM
To: Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Queen, Jackie L <Jackie.Queen@fda.hhs.gov>
Subject: Fwd: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls): (b) (6) - EON-345965

Another one from (b) (6)

From: PFR Event <pfpreventcreation@fda.hhs.gov>
Date: January 25, 2018 at 12:25:12 PM EST
To: Cleary, Michael * <Michael.Cleary@fda.hhs.gov>, HQ Pet Food Report Notification <HQPetFoodReportNotification@fda.hhs.gov>, (b) (5) (b) (5) >
Subject: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls): (b) (6) - EON-345965

A PFR Report has been received and PFR Event [EON-345965] has been created in the EON System

A "PDF" report by name "2040808-report.pdf" is attached to this email notification for your reference Please note that all documents received in the report are compressed into a zip file by name "2040808-attachments.zip" and is attached to this email notification

Below is the summary of the report:

EON Key: EON-345965

ICSR #: 2040808

EON Title: PFR Event created for limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls); 2040808

AE Date	04/24/2017	Number Fed/Exposed	5
Best By Date		Number Reacted	5
Animal Species	Dog	Outcome to Date	Unknown
Breed	Shih Tzu		
Age	8 Years		
District Involved	(b) (6)		

Product information.

Individual Case Safety Report Number: 2040808

Product Group: Pet Food

Product Name: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls)

Description: At his scheduled visit to my clinic, thoracic radiographs showed generalized cardiomegaly which had been progressive compared to prior chest radiographs from his regular veterinarian but there was no evidence of cardiogenic edema. Echocardiogram was performed which showed dilated cardiomyopathy. Fundic exam was abnormal with a suspected partial retinal detachment OS. Diet history revealed that (b)(6) was eating a kangaroo based diet. At this time the patient was continued on the Cough-tabs, Lasix was discontinued, and Vetmedin (2.5mg a.m., 1.25mg p.m.), enalapril (1.25mg BID), and taurine (500mg BID) were started. Taurine was discontinued after a normal taurine level was received. Cough persisted despite these changes and a course of doxycycline was prescribed (50mg BID x 10days). The cough improved significantly but did not completely resolve so the doxycycline was continued an additional 14 days. The dog has since been lost to follow-up. I have attempted to contact the owner and am waiting for a response. I did contact the referring veterinarian and to their knowledge the dog is still alive.

Submission Type: Initial

Report Type: Adverse Event (a symptom, reaction or disease associated with the product)

Outcome of reaction/event at the time of last observation: Unknown

Number of Animals Treated With Product: 5

Number of Animals Reacted With Product: 5

Product Name	Lot Number or ID	Best By Date
limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls)		

Sender information

(b)(6)

Owner information

(b)(6)

To view this PFR Event, please click the link below:

<https://eon.fda.gov/eon/browse/EON-345965>

To view the PFR Event Report, please click the link below:

(b)(6)

This email and attached document are being provided to you in your capacity as a Commissioned Official with the U.S. Department of Health and Human Services as authorized by law. You are being provided with this information pursuant to your signed Acceptance of Commission.

This email message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential. Any dissemination, distribution, or copying is strictly prohibited.

The information is provided as part of the Federal-State Integration initiative. As a Commissioned Official and state government official, you are reminded of your obligation to protect non-public information, including trade secret and confidential commercial information that you receive from the U.S. Food and Drug Administration from further disclosure. The information in the report is intended for situational awareness and should not be shared or acted upon independently. Any and all actions regarding this information should be coordinated through your local district FDA office.

Failure to adhere to the above provisions could result in removal from the approved distribution list. If you think you received this email in error, please send an email to FDAREportableFoods@fda.hhs.gov immediately.

From: [Rotstein, David](#)
To: [Hartogenesis, Martine](#); [DeLancey, Siobhan](#)
Cc: [Norris, Anne](#)
Subject: RE: List of firms with city/state?
Date: Friday, June 21, 2019 5:32:39 AM
Attachments: [image002.png](#)
[image004.jpg](#)
[image006.jpg](#)
[image008.jpg](#)
[image010.jpg](#)
[image012.jpg](#)
[image013.png](#)
[image014.jpg](#)
[image015.jpg](#)
[image016.jpg](#)
[image017.jpg](#)
[image018.jpg](#)

I'd consider this to be the final list. We may learn of co-packers, but it won't change the actual firm marketing the product.

Internally for FDA, the Divisions (HAF) is aware of the update and the FDA National Consumer Complaint Coordinator is also aware. Hopefully having all of this internal awareness will help should firms or consumers contact FDA.

D.

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/CERRT
7519 Standish Place
(b) (6) **(BB)**



This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: Hartogenesis, Martine
Sent: Thursday, June 20, 2019 10:06 PM
To: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Cc: Norris, Anne <Anne.Norris@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: List of firms with city/state?

Hi Siobhan,

Yes, I am happy to be the SME. Sending you the latest list from Dave.

Thanks again!

Martine

From: DeLancey, Siobhan
Sent: Thursday, June 20, 2019 3:18 PM
To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Cc: Norris, Anne <Anne.Norris@fda.hhs.gov>
Subject: List of firms with city/state?

Hi Martine,

[Redacted] (b) (5)
[Redacted]
[Redacted]
[Redacted]
[Redacted]
[Redacted]

[Redacted]
[Redacted]
[Redacted]

Thanks!

Siobhan DeLancey, RVT, MPH
Senior Advisor for Strategic Initiatives
Center for Veterinary Medicine
U.S. Food and Drug Administration
O: 240-402-9973
M: [Redacted] (b) (6)
Siobhan.DeLancey@fda.hhs.gov



Owner: (b) (6)

Accession Number: (b) (6)
Reference Number:
Case Coordinator: (b) (6)
Received: (b) (6) Finalized: 04/18/2017
Sampled:

To: (b) (6)
(b) (6) VETERINARY HEALTH COMPLEX

Phone # (b) (6)

Addended Report

ANATOMIC PATHOLOGY RESULTS

SMALLANIMAL NECROPSY

ANIMAL ID (b) (6)
REF CASE NO (b) (6)
SPECIES Canine
BREED Schnauzer
SEX Mc
AGE 3y
SPECIMEN DESC Body

GROSS An 8.2 kg, 2.5-year-old, castrated male miniature schnauzer dog is presented for postmortem examination. The animal was euthanized and the body is in fair postmortem condition with a euthanasia?to?necropsy interval of approximately 16 hours. The body appears moderately dehydrated. The hair coat is thick and shiny, and no ectoparasites are seen. Throughout the body, subcutaneous and visceral fat stores are adequate and the body condition score is 4/9. Throughout the dentition, a mild amount of dental calculus is present. Multifocally over the left lateral brachium, left cranio-lateral tibia, and left temporalis muscle, there are multiple approximately 4-6 cm long linear skin incisions that have been closed with suture. The left antebrachium is shaved and there are a few ecchymotic hemorrhages in this region. The right antebrachium, dorsal metatarsals, and ventral neck are also shaved. All skeletal muscle is grossly unremarkable.

Free within the thoracic cavity are approximately 15 mL of red, watery fluid. The lungs diffusely fail to collapse and contain multiple rib impressions along the pulmonary pleural surface. They are diffusely pink to red. On cut surface, these ooze a moderate to large amount of frothy, white fluid. The heart is subjectively enlarged. The heart measures as follows: right atrium circumference – 6 cm; right ventricular free wall thickness – 0.5 cm; left atrium circumference – 7 cm; left ventricular free wall thickness – 1 cm; interventricular wall thickness – 0.6 cm; pulmonic valve circumference – 4.5 cm; aortic valve circumference – 3 cm; total weight – 99 g (1.2% of the total body weight); right and left atria weight –

Addended Report

Accession Number: (b) (6)

ANATOMIC PATHOLOGY RESULTS

15 g; right ventricle weigh 22 g; left ventricle and interventricular septum weight – 62 g. Multifocally expanding the free edge of the leaflets of the mitral valve, there are a few small, less than 2 mm diameter, smooth, shiny, white nodules.

Focally within the peripheral left limb of the pancreas, there is a small, 2 mm diameter, firm, white nodule. Multifocally along the capsular surface of the spleen are a few siderotic plaques.

GROSS DIAGNOSIS

1. Lungs: moderate to marked, diffuse pulmonary edema
2. Heart: mild cardiomegaly with mild mitral valve endocardiosis and mild left ventricular hypertrophy and left atrial dilation
3. Thorax: mild pleural effusion
4. Pancreas: focal nodular hyperplasia

**REPORT STATUS
COMMENTS**

PRELIMINARY REPORT-HISTOLOGY PENDING

Except for gross evidence of moderate to marked pulmonary edema and mild cardiac changes, gross examination is otherwise mostly unremarkable. The heart is mildly enlarged with mild mitral valve endocardiosis and associated mild left atrial dilation and left ventricular hypertrophy. This mild degree of cardiac changes does not fit well with the moderate to severe degree of pulmonary edema present. In addition to the mild structural changes, a functional cardiac abnormality may have been contributing to clinical disease in this patient. Samples of lung and heart were rushed and examined histologically. In addition to the pulmonary edema, there is also histologic evidence of diffuse, acute alveolar injury. These pulmonary changes can explain this patient's signs of respiratory distress. However, the heart is histologically unremarkable and the skeletal muscle was grossly unremarkable; as such, a cause for the markedly elevated CK is not yet identified. Histology of skeletal muscle is pending.

**PATH RESIDENT
SENIOR PATH
DATE**

(b) (6) DVM
(b) (6) DVM PHD DACVP
(b) (6)

Final Necropsy Report

MICROSCOPIC

Lung (slide 2): Diffusely throughout all sections, there is evidence of alveolar injury characterized by frequent deposition of bright eosinophilic fibrin within alveolar lumina. This fibrin is often forming hyaline membranes lining alveolar septal walls. There is also frequent deposition of fibrin within the alveolar walls and sometimes effacing and replacing the septa. There are also high numbers of scattered foamy, alveolar macrophages. Multifocally scattered throughout multiple sections, there is occasional type II pneumocyte hyperplasia.

Liver (slide 4): There is diffuse mild to moderate, acute congestion of hepatic sinusoids. Low numbers of individually scattered hemosiderin-

Addended Report

Accession Number: (b) (6)

ANATOMIC PATHOLOGY RESULTS

laden Kupffer cells are present.

Heart (slide 1), Skeletal muscle (slide 3), Kidneys (slide 3), Spleen (slide 3), Stomach (slide 4), Small intestine (slide 4), Colon (slide 4), Pancreas (slide 4), Adrenal gland (slide 4): No remarkable histologic abnormalities are appreciated throughout the tissue sections examined. PAS stains with and without diastase treatment are applied to sections of heart. No evidence of glycogen-storage is appreciated.

FINAL DIAGNOSIS

1. Lungs:
 - a. Severe, diffuse alveolar injury with marked fibrin deposition (hyaline membranes) and marked alveolar histiocytosis and multifocal type II pneumocyte hyperplasia
 - b. Moderate to marked, diffuse pulmonary edema
2. Heart: mild cardiomegaly with mild mitral valve endocardiosis and mild left ventricular hypertrophy and left atrial dilation
3. Thorax: mild pleural effusion

COMMENTS

Histology reveals severe, diffuse alveolar injury and alveolar histiocytosis throughout all examined sections of lung. This pattern of injury is often seen in cases of acute respiratory distress syndrome (ARDS). However, ARDS is a clinical diagnosis. The degree of injury present is severe enough to explain this dog's reported clinical disease of respiratory distress. However, an underlying etiology for this alveolar injury is not identified. The heart and skeletal muscle are histologically unremarkable and, as such, a primary noncardiogenic etiology is suspected. However, if there was clinical evidence of cardiac dysfunction then a functional cardiac abnormality in the absence of a correlating structural change cannot be ruled-out based on postmortem examination. Survey sections of other major organs are also histologically unremarkable. Common causes of ARDS in dogs include aspiration of sterile stomach contents, smoke or other toxin inhalation, near drowning, infection and systemic inflammation (sepsis), drug reaction, etc. In this case, there is no overt evidence of sepsis throughout the other tissues examined.

PATH RESIDENT
SENIOR PATH
FINALIZED DATE

(b) (6) DVM
(b) (6) DVM PHD DACVP
(b) (6)

Report Details - EON-323515	
ICSR:	2023228
Type Of Submission:	Initial
Report Version:	FPSR.FDA.PETF.V.V1
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)
Reporting Type:	Voluntary
Report Submission Date:	2017-07-11 17:06:59 EDT
Reported Problem:	Problem Description: Please note: Dr. Jennifer Jones was consulted prior to submission of this report. She would like to be involved in the case review 3 week history of cough treated unsuccessfully with doxycycline and prednisone. 3 day history of inappetence and vomiting prior to presentation to NCSU emergency service for dyspnea. Radiographs showed severe pulmonary edema and echocardiogram showed severe Dilated Cardiomyopathy. There was an initial response to diuretic therapy however, he declined and was placed on the ventilator for respiratory support and continued CHF treatment. Attempts to wean off the ventilator were unsuccessful and aquaphoresis was performed. He continued to decline despite aggressive therapy and was euthanized. Infectious disease testing was negative and taurine and carnitine analysis showed adequate levels. Necropsy initially did not reveal a cause for DCM and supported alveolar injury (possibly ventilator related). A re-review of the myocardial histopathology by one of our pathologist showed myofiber vacuoles reminiscent of the changes seen in doxorubicin toxicity. Since the dog had not received doxorubicin, the pathologist recommended culturing the food for Streptomyces peucetius - the bacterium which produces doxorubicin. He also recommended testing for Fusarium spp. a fungus which produces Fumonisin B1, a toxin that produces heart failure in pigs. (b) (6) had been fed California Naturals Adult - both kangaroo with lentils and venison with lentils along with Milo's kitchen treats. We have samples of these foods from 6/17 but not the original bags from when he was presented 2/17. These samples were provided at the time his housemate, (b) (6) (unrelated, older miniature schnauzer) also presented with severe DCM and CHF. I will enter this dog as a separate affected patient. Both dogs had extensive infectious disease testing which was negative and nutritional amino acid deficiencies were ruled out. Because of this, their unrelated lineages (although the same breed, they were from different lines), different ages but similar time of presentation ((b) (6) had clinical signs at the time (b) (6) was treated, but didn't present with CHF for several months), we are considering common environmental factors which could precipitate DCM, including food contamination or toxin exposure.
	Date Problem Started: (b) (6)
	Concurrent Medical Problem: No
	Outcome to Date: Died Euthanized
	Date of Death: (b) (6)
Product Information:	Product Name: Alternated between: -California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils and Kangaroo & Red Lentils Recipe
	Product Type: Pet Food
	Lot Number:
	UPC: not available
	Package Type: BAG
	Package Size: 26 Pound
	Purchase Date: 06/01/2017
	Possess Unopened Product: No
	Possess Opened Product: Yes
	Storage Conditions: In a cabinet, in the original bag
Product Use Information:	Description: twice daily feeding The sample we have is from 6/17, however, we do not have food samples from 2/17 when both dogs started with clinical signs.
	Time Interval between Product 2 Years

FDA-CVM-FOIA-2019-1704-000332

	Use and Adverse Event:		
	Product Use Stopped After the Onset of the Adverse Event:	No	
	Perceived Relatedness to Adverse Event:	Possibly related	
	Other Foods or Products Given to the Animal During This Time Period:	Yes	
Manufacturer /Distributor Information:			
Purchase Location Information:	Address:	United States	
Animal Information:	Name:	(b) (6)	
	Type Of Species:	Dog	
	Type Of Breed:	Schnauzer - Miniature	
	Gender:	Male	
	Reproductive Status:	Neutered	
	Weight:	8.2 Kilogram	
	Age:	2.5 Years	
	Assessment of Prior Health:	Excellent	
	Number of Animals Given the Product:	2	
	Number of Animals Reacted:	2	
	Owner Information:	Owner Information provided:	Yes
		Contact:	Name: (b) (6)
			Phone: (b) (6)
			Email: (b) (6)
	Address:	(b) (6) United States	
	Healthcare Professional Information:	Practice Name:	North Carolina State University, College of Veterinary Medicine
		Contact:	Name: Darcy Adin
Phone: (919) 513-6694			
Other Phone: 6145829798			
Email: dbadin@ncsu.edu			
Address:	1060 William Moore Dr Raleigh New York 27607 United States		
Practice Name:	North Carolina State University College of Veterinary Medici		
	Contact:	Name: (b) (6)	
Phone:		9195136694	

		Other Phone: 6145829798
	Practice Name:	North Carolina state University, College of Veterinary Medic
	Contact:	Name: (b) (6)
		Phone: 919 513-6694
		Other Phone: 6145829798

Sender Information:	Name:	Darcy Adin		
	Address:	1060 William Moore Dr Raleigh New York 27607 United States		
	Contact:	Phone:	9195136694	
		Other Phone:	6145829798	
		Email:	dbadin@ncsu.edu	
	Permission To Contact Sender:	Yes		
Preferred Method Of Contact:	Email			
Reported to Other Parties:	Manufacturer			

Additional Documents:	
------------------------------	--

Report Details - EON-323519					
ICSR:	2023230				
Type Of Submission:	Initial				
Report Version:	FPSR.FDA.PETF.V.V1				
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)				
Reporting Type:	Voluntary				
Report Submission Date:	2017-07-11 17:32:14 EDT				
Reported Problem:	Problem Description: Please note: Dr. Jennifer Jones was consulted prior to submission of this report. She would like to be involved in the case review. (b) (6) housemate, (b) (6) (separate report submitted) was diagnosed with DCM and CHF (b) (6) and was euthanized after aggressive treatment of CHF. At that time (b) (6) had 2 syncopal events closely related to each other. His appetite for dog food declined but he would eat it if tempted with treats mixed in. He was presented 6/22/17 for more syncopal events and was similarly diagnosed with severe DCM and CHF. He was able to be successfully treated however and is clinically doing well on CHF medications as of 7/10/17. A re-review of the myocardial histopathology for (b) (6) housemate (b) (6) was requested at this time because of the unusual diagnosis of DCM in a small breed dog living in the same house as another dog similarly diagnosed a few months ago. This re-review by one of our pathologists showed myofiber vacuoles reminiscent of the changes seen in doxorubicin toxicity. Since the dog had not received doxorubicin, the pathologist recommended culturing the food for Streptomyces peucetius - the bacterium which produces doxorubicin. He also recommended testing for Fusarium spp. a fungus which produces Fumonisin B1, a toxin that produces heart failure in pigs. Like (b) (4) (unrelated, younger miniature schnauzer), (b) (6) had been fed California Naturals Adult - both kangaroo with lentils and venison with lentils along with Milo's kitchen treats. We have samples of these foods from 6/17 but not the original bags from when he was presented 2/17. These samples were provided at the time (b) (6) also presented with severe DCM and CHF. Like (b) (6) had extensive infectious disease testing which was negative and nutritional amino acid deficiencies were ruled out. Because of this, their unrelated lineages (although the same breed, they were from different lines), different ages but similar time of presentation, we are considering common environmental factors which could precipitate DCM, including food contamination or toxin exposure. We have plasma, serum, urine and myocardial tissue samples (latter only for (b) (6)) stored at -80 Celsius in addition to food and treat samples.				
	Date Problem Started: (b) (6)				
	Date of Recovery: 07/10/2017				
	Concurrent Medical Problem: No				
	Outcome to Date: Stable				
Product Information:	Product Name: Alternated feedings between: California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food and Kangaroo and Lentils				
	Product Type: Pet Food				
	Lot Number:				
	Package Type: BAG				
	Package Size: 26 Pound				
	Storage Conditions: In the bag in a cabinet				
	<table border="1"> <tr> <td rowspan="2">Product Use Information:</td> <td>Description:</td> <td>the Venison and Kangaroo types were alternated depending on store availability and were fed twice daily along with Milo's kitchen treats</td> </tr> <tr> <td>Perceived Relatedness to Adverse Event:</td> <td>Possibly related</td> </tr> </table>	Product Use Information:	Description:	the Venison and Kangaroo types were alternated depending on store availability and were fed twice daily along with Milo's kitchen treats	Perceived Relatedness to Adverse Event:
Product Use Information:	Description:		the Venison and Kangaroo types were alternated depending on store availability and were fed twice daily along with Milo's kitchen treats		
	Perceived Relatedness to Adverse Event:	Possibly related			
Manufacturer /Distributor Information:					
Purchase Location Information:					
Animal Information:	Name: (b) (6)				

FDA-CVM-FOIA-2019-1704-000335

	Type Of Species:	Dog								
	Type Of Breed:	Schnauzer - Miniature								
	Gender:	Male								
	Reproductive Status:	Neutered								
	Weight:	9.6 Kilogram								
	Age:	7 Years								
	Assessment of Prior Health:	Excellent								
	Number of Animals Given the Product:	2								
	Number of Animals Reacted:	2								
	Owner Information:	<table border="1"> <tr> <td>Owner Information provided:</td> <td>Yes</td> </tr> <tr> <td rowspan="3">Contact:</td> <td>Name: (b) (6)</td> </tr> <tr> <td>Phone: (b) (6)</td> </tr> <tr> <td>Email: (b) (6)</td> </tr> <tr> <td>Address:</td> <td>(b) (6) United States</td> </tr> </table>	Owner Information provided:	Yes	Contact:	Name: (b) (6)	Phone: (b) (6)	Email: (b) (6)	Address:	(b) (6) United States
	Owner Information provided:	Yes								
	Contact:	Name: (b) (6)								
		Phone: (b) (6)								
Email: (b) (6)										
Address:	(b) (6) United States									
Healthcare Professional Information:	Practice Name:	North Carolina State University, College of Veterinary Medicine								
	Contact:	Name: Darcy Adin								
		Phone: (919) 513-6694								
		Other Phone: 6145829798								
		Email: dbadin@ncsu.edu								
	Address:	1060 William Moore Dr Raleigh New York 27607 United States								
	Practice Name:	North Carolina State University, College of Veterinary Medic								
	Contact:	Name: (b) (6)								
		Phone: 919-513-6694								
Other Phone: 6145829798										
Practice Name:	North Carolina State University, College of Veterinary Medic									
Contact:	Name: (b) (6)									
	Phone: 919-513-6694									
	Other Phone: 6145829798									

Sender Information:	Name:	Darcy Adin
	Address:	1060 William Moore Dr Raleigh New York 27607 United States
	Contact:	Phone: 9195136694
		Other Phone: 6145829798
		Email: dbadin@ncsu.edu
Permission To Contact Sender:	Yes	

FDA-CVM-FOIA-2019-1704-000336

	Preferred Method Of Contact:	Email
	Reported to Other Parties:	Manufacturer

Additional Documents:

EON #	Last Name	Product	Lot	BB Date	Signalment	Clinical Signs	Open Product?	Closed Product?	MedRec?	Vet-LIRN Open Product Testing
EON-345822	(b) (6)	California Naturals Kangaroo & Lentil	unknown	unknown	(b) (6) -6 yr FS Lab	cough, dyspnea, collapse, CHF; hx atopy	Yes	No	DCM with left CHF, WB Tau wnl; mild inc ALT/AST/CK	collection in progress
EON-345831	(b) (6)	California Naturals Kangaroo & Lentil; began new Venison Food (Brand?)	unknown	unknown	(b) (6) -8 yr F Lab	collapse, cough, urinary incontinence	No	No	DCM with Chronic Degen Valve Disease, Atrial Fib (resolved), Exertional Collapse; WB Tau: wnl	n/a
EON-345833	(b) (6)	California Naturals Kangaroo & Lentil; began new Venison Food (Brand?)	unknown	unknown	(b) (6) -5 yr FS Lab **euthanized**	dyspnea, cough-productive, urinary incontinence	No	No	DCM with Endocardiosis, CHF, inc SDMA	n/a
EON-345835	(b) (6)	Zignature Kangaroo Limited Ingredient	unknown	unknown	(b) (6) -4 yr MC Cocker Spaniel	cough	No	No	DCM, CHF, WB Tau: LOW, low blood Alb with proteinuria	n/a
EON-345965	(b) (6)	Kangaroo	unknown	unknown	(b) (6) -8 yr MC Shih Tzu	lethargy, cough	unknown	unknown	DCM, WB Tau: wnl	n/a

Vet-LIRN Case Summary Document

Vet-LIRN Case Number:	800.218
EON/CC #:	Multiple
Owner LAST Name:	(b) (6)
Vet LAST Name:	(b) (7)(A)
Vet-LIRN Initiation Date:	1/23/2018
MedRec: Requested:	already received
MedRec: Received:	
MedRec: Significant finding:	All dogs w/ DCM, 2 Tau wnl, 1 not measured, 1 Tau low
Vet-LIRN Tests (planned):	<ul style="list-style-type: none"> • MRx Review • Open Product (b) (6) for Metals and Cys/Met • (b) (6) Blood Fe Panel
Vet-LIRN Test Results:	Low product Fe results
Result Interpretation:	Fe Panel: all wnl Iodine: wnl
IF NFA, justification:	

COMPLAINT Narrative: see xls and multiple PFRs

1/23/2018

JJ-MRx summaries.

The listserv BLUF:

- Diet:
 - Kangaroo: low Tau & Met, high Carn
 - Lentil: Low Cys, Met
- In Cats, plasma Taurine (Tau)
 - falsely increases with damaged WBC and Platelets, short half-life
 - Whole blood Taurine a better measurement b/c longer half-life
 - Skeletal muscle even longer half-life and thought to parallel cardiac muscle Tau content
- From one vet dealing with Golden Retrievers eating designer diets (high quality meats) with pea or lentils-most dogs with low or Low Normal Whole Blood Tau → most recovered with Tau supplementation → Hypothesis was low Cys & Met plus Golden Retriever Tau handling may cause the illness

(b) (6)-6 yr FS Lab

Presenting complaint 1/10/2018: cough, dyspnea, multiple episodes of collapse, cardiomegaly, suspected CHF; seen by rDVM 12/30 for cough → by 1/4/2018: collapse episode-circled, fell over, flopped ~1 min, 2nd less severe episode 1/5; → continued shortness of breath, easily tires; heartworm negative (9/9/2017), eats California Naturals Kangaroo and Lentil with veggies and 2 tbsp canned pumpkin; on Apoquel BID and SLIT;

PE 1/10: T 102.7, P 208, R 150, Gr III/VI left apical systolic murmur with gallop, regular tachycardia, quiet heart sounds, localized fine crackles left cranial hilar region with dry cough, poor femoral pulses, mm pale pink; → hospitalized → occ short paroxysms of slow ventricular tachycardia

Labs: 1/5/2018 Chem: mild inc ALT, AST, and CK

1/10 T4: low normal

BP: 152 mmHg

Echo: sinus tachycardia, DCM with left sided CHF

Whole Blood Tau: 292 (200-350), >150 not known risk of Tau Deficiency;

ECG: HR 189 bpm, sinus tachycardia, atrial enlargement, LVE

1/11 T-Rads: mild decrease in severity of cardiomegaly, resolving cardiogenic edema

(b) (6): 8 yr F Lab

Presenting (b) (6): ER for heart evaluation, seen week prior (Thurs) for collapse while playing fetch; at vet-atrial fibrillation with mild heart failure; normal sinus rhythm (nsr) on 1/20 → 1/21 collapse while fetching → nsr with rads showing resolution of HF, CBC/Chem wnl → recheck on 5/31: no episodes of weakness/collapse, last 2-3 anxious and less social b/c storm anxiety; on daily glucosamine/chondroitin → recheck 12/11: new cough, not frequent, at rest and with excitement/activity; owner transitioning to a new venison food, urinary incontinence

PE (b) (6): Gr III to IV/VI left apical holosystolic murmur, sinus arrhythmia, moderate femoral pulses

5/31: T 103.1, RR 120, Gr III-IV/VI left apical holosystolic murmur

12/11: HR 128, pant, Gr III/VI left apical systolic murmur with wide radiation, fair-moderate

femoral pulse

Labs:

(b) (6)

Whole Blood Taur: 236 (200-350) (>150 = no risk for Tau deficiency)

BP: 108 mmHg

-5/31: 90 mmHg

Echo: DCM, Chronic degenerative valve disease, atrial fibrillation, exertional collapse

-5/31: slight improvement though heart a little larger

-12/11: heart bigger with decreased contractility,

Telemetry: consistent sinus rhythm/arrhythmia

Renal panel: no results listed

-5/31: SDMA 18, normal BUN/Ct, mild low Cl

-12/11: wnl

5/31 T Rads: mild progression of cardiac enlargement

-12/11: progressive cardiomegaly

(b) (6): 5 yr FS Lab → genetic niece of (b) (6)

Presenting complaint (b) (6): new heart murmur, diagnosed with CHF on (b) (6) seen by rDVM for heavy breathing and cough; rads and labs showed enlarged heart, CHF → (b) (6) to ER, increased meds with some improvement → panting and on low sodium Kangaroo and Lentil diet; other dog (b) (6) is (b) (6) aunt, urinary incontinence began prior to CHF treatment/diagnosis; less social and active; on Apoquel → (b) (6) tachypneic, coughed up pink tinged fluid, episode of collapse, dry heaving on way to ER → on Venison and Lentil diet → respiratory rate continue to increase despite Lasix CRI → euthanized PE (b) (6): HR 150, pant, Gr IV/VI left apical systolic murmur with radiation to right, adequate femoral pulse

Labs:

(b) (6) HR 180, pant, pale pk, Gr IV/VI murmur, fine crackles right dorsal lung fields

(b) (6) BP: 110

(b) (6) Renal Panel: mild inc SDMA (19 ug/dL)

(b) (6) Echo: endocardiosis, DCM

(b) (6) ECG: nsr

(b) (6) T Rads: persistent cardiomegaly with mild decreased severity

-9/9-9/10: generally enlarged, unstructured interstitial pulmonary pattern within right middle and cd lobes, mild enlargement cranial lobar pulmonary veins

(b) (6) -4 yr MC Cocker Spaniel

Presenting complaint 9/5/2017: evaluation of cardiomegaly and CHF, diagnosed by rads 8/22; to rDVM 8/22 for week long progressive cough-2x/day with progressive severity; t-rads and labs: CHF; treated with antibiotics and heart meds (concern initially for pneumonia) → improved cough but still occurring; PE 9/5: T 104.4, HR 150, pant, quiet/distant heart sounds with gallop, Gr I/VI systolic murmur on left, fine bilateral crackles, patient shivering so femoral pulse difficult to assess but suspect decreased, palpable jugular pulsation, suspected oral epulis;

-9/19: HR 178-180, T 102.4, quiet heart sounds, gallop, fair femoral pulses, palpable hepatomegaly, epidermal collarettes with exudative crust on ventral abdomen

-11/2: T 103.2, HR 168, RR 110

Labs: 9/5 BP 160 mmHg

-9/19: 152 mmHg

9/5 Chem: Alb 2.5 (inc from 2.1)

-9/19 or 11/2: Renal Panel: BUN 32

9/5 UA: 1.015, pH 8, pro tr

Echo: DCM, CHF

ECG: sinus tachycardia, HR ~200

9/5 Whole Blood Tau: 10 (200-350)

T Rads (9/19 or 11/2?): decreased heart size

PLAN: I Recommend collecting open product from (b) (6) case (only one available) and testing for metals with the other cases. Also based on message board-recommend testing cysteine and methionine from all the DCM cases. If we find something "Low" in the foods for amino acids, we can look at the case from last summer and measure cardiac muscle-Cys, Met, Tau. For (b) (6) case (EON-345822) if dog dies-will ask to save heart tissue for potential testing and histopath. Vet-LIRN can purchase storebought Zignature Essentials for testing based on other results.

DR agrees-If testing confirms an issue with the Amino acids-DAF could have a chat with the firm.

JJ-Contacting the vet to request open product for testing and tissue if dog dies.

1/24/2018

JJ-Vet is collecting the leftover food from (b) (6). The vet mentions additional cases:

For your information, we have pulled all our cases of dilated cardiomyopathy that we diagnosed in 2017 and identified a fifth animal on kangaroo and lentil diet (out of 23 total) - I will submit an FDA case report for this-, and another one on a lentil-containing Vegan diet. There are some cases in which the diet is not specified that we still need to contact, but at least 11 of the cases are on grain-free diets, including all of the cases in "atypical" breeds. As you may have seen from my listserve query to other cardiologists, not only have

Cardiologists been seeing cases in dogs fed kangaroo and lentil, but there were a number of cases in dogs of "atypical" breeds on lentil-based Vegan diets, and a series of Golden retrievers with taurine deficiency on grain-free diets. Grain-free diets are a fad right now and lots of clients are feeding them, so the incidence of grain-free diets may simply be coincidence. Interestingly however, (b) (6), who I saw recently and re-echoed after several months on another diet with no documented improvement in echo findings is still being fed a grain-free diet. I have asked her owner to switch her diet once again to something not "grain-free" and we will see how things go over the next several months.

Emailed the vet-we're sending a kit to collect the food.

1/30/2018

JJ-Vet submitted an additional PFR but it was unclear which brand kangaroo diet was fed.

LAP did an epi summary: FYI – this makes 6 cases now, 5 from Ohio, 1 from NC (involving 2 dogs, 1 death) all eating LID type diet with Kangaroo and Red Lentil, onset dates between 1/20/2017 and 12/30/2017).

MRx summary:

(b) (6)-8 yr MC Shih Tzu

Presenting complaint 6/19/2017: lethargy for past few weeks and progressive coughing that began 6 months ago; started on cough suppressant and Lasix without improvement; RR/effort normal, no episodes weakness/collapse; on limited ingredient diet with Kangaroo as sole protein source; monthly Sentinel, HW negative February; → recheck 7/6-frequently travelling, dog not liking car rides as much (RR 80's), forceful cough-significant increase; bad night-unable to sleep comfortably; 3 urinary accidents in house since starting new meds

PE 6/19: P 128, Gr III-IV/VI left apical systolic murmur with radiation to the right base, fair femoral pulses, excessive bruising after jugular venipuncture; thin body condition, mm light pink/slightly tacky; fundic-suspect partial retinal detachment on nasal aspect of retina OS w/ hyper-reflectivity around edge;

-7/6: Grade IV/VI murmur, 99.6F, HR 165, RR 56

Labs: 6/19 BP: 102 mmHg

-7/6: 138 mmHg

6/19 Rads: generalized cardiomegaly (progressive), mild diffuse bronchointerstitial pattern

-7/6: unchanged cardiomegaly

6/19 CBC: mild non-pre-regenerative anemia, Retic 115 (10-110)

6/19 Chem: Na 154 (142-152), K 5.4 (4-5.4), Cl 107 (108-119), Bicarb 30 (13-27),
SDMA 14 (0-14), Alb 4.1 (2.7-3.9), AST 83 (16-55), CK 1162 (10-200)

-7/6 Renal panel: nsf

6/19 Whole Blood Taurine: 276 (200-350); >150 no known risk for Tau deficiency

6/19 Echo: DCM, mild PV insufficiency, mild tricuspid regurgitation & thickening,
mild MV thickening with +3 regurgitation

1/30/18

JG – Shipped food sample collection kit to Vet and informed Vet with tracking information

Note: Vet hospital's zip code was incorrect. updated.

2/2/2018

JJ-Emailed (b)(4) requesting testing. Make lab submission forms.

JG – Received treats (kibble) in the provided pink zip bag. Also received original package (empty). Seal the original package with tape.

Log in the treat as 800.218-sub6

2/5/2018

JJ-prepared the sample and packaged box to send to (b) (4).

2/22/2018

JJ-We received the results from (b) (4). The only significant finding was all products contained inadequate Fe content, below the AAFCO claim. Fe has been associated with idiopathic DCM (Marinescu and McCullough 2011) in people. According to the article, anemia was found in 12 to 55% of patients in heart failure, and the anemia's severity correlated to the degree of heart failure.

Looking at our cases, a CBC was done for 3 dogs according to their discharge summaries. One dog had a non/pre-regenerative anemia, and the other two had no mention of RBC abnormalities. It's possible that chronically feeding a diet containing low Fe could cause a heart issue, but I would also expect to see anemias in these dogs.

Dr. Adin at NCSU emailed additional theories. She continues to see more cases.

- (b) (5)

- Just thought I would forward this email chain to you as the growing concern about grain-free diets in the veterinary cardiology community continues to expand. (b) (5)

(b) (5)

- (b) (5)

o

(b) (5)

The Vet Cardiology Community is discussing this quite a bit.

2/23/2018

JJ-DR replied-

(b) (5)

Send feed test results to NCSU and (b) (6). I'll ask Dr. Adin if an iron panel has been done on any of the dogs. If not, we can request a full iron panel for one dog at NCSU and one from (b) (6). Will get estimates for the test from each lab. If the dogs are low Fe, we can test a myocardial sample from the (b) (6) case.

2/27/2018

JJ-The vet will perform a blood iron panel. I received an estimate and submitted a PO. This will be an unfasted blood sample.

Per the vet: "We had not done a CBC as one was completed through the referring vet, prior to her initial referral, and was completely normal dated 1/5/18 through (b) (6)

3/13/2018

JJ-The veterinarian sent the iron panel results. Iron 179 (73-245), TIBC 396 (270-530), Ferritin 443 (89-489). This was an unfasted sample and may be after the dog was switched to a new food??

From the vet: I would love to discuss our findings on dilated cardiomyopathy and dietary relationships in our clinic over the past year. In particular, 75% of our DCM cases in 2017 for which we have adequate dietary histories were on grain free diets. We have started to do a survey of other cardiac patients during the same time period to see if we can get an idea of what percentage of our referral population were feeding these type of diets so we can see if this is truly significant. As we have been looking harder at these cases, we have found some other interesting things. For instance, two patients that were found to be taurine deficient were being fed Zignature diets. One was on the kangaroo variety and the other the pork variety. There continues to be a lot of discussion about diets and dilated cardiomyopathy on our list serve, but I can see it degenerating into I diagnosed DCM and the dog is on this diet so therefore this is a problem, which may or may not be true evidence to support a causative role of the diet. I am not sure what information the FDA would want on all this at this point. We are certainly concerned that there may be a wider concern than simply the kangaroo and lentil diets that we first identified as a potential problem.

3/20/2018

JJ-The invoice is still pending.

4/9/2018

JJ-I'll ask the vet if they received the lab invoice.

4/12/2018

JJ-The vet sent the invoice, and I submitted.

I prepared the food samples and they should ship to MSU for iodine screening.

4/23/2018

JJ-We received the MSU results. All products contained less than 10 ppm. NFA.

From: [Forfa, Tracey](#)
To: [Hartogenesis, Martine](#)
Subject: FW: DCM Plan
Date: Tuesday, June 18, 2019 10:07:15 AM
Attachments: [DCM Project Plan.docx](#)

Here you go! The "script" is embedded in the chart. Let me know if you have any questions....and please feel free to tweak it.

From: Forfa, Tracey
Sent: Tuesday, June 4, 2019 12:22 PM
To: Steinberg, Nadine (Nadine.Steinberg@fda.hhs.gov) <Nadine.Steinberg@fda.hhs.gov>; DeLancey, Siobhan (Siobhan.Delancey@fda.hhs.gov) <Siobhan.Delancey@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>; Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>
Cc: Dewitt, Susan J (Susan.Dewitt@fda.hhs.gov) <Susan.Dewitt@fda.hhs.gov>; Cepeda, Sandra <Sandra.Cepeda@fda.hhs.gov>; Murphy, Jeanette (Jenny.Murphy@fda.hhs.gov) <Jenny.Murphy@fda.hhs.gov>; Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>
Subject: DCM Plan

Hi All – As discussed [REDACTED] (b) (5)

[REDACTED] (b) (5)

(b) (5)

Thank you!!!

DCM Plan

Activity	Lead	Notes	Timing
(b) (5)	(b) (5)	(b) (5)	
(b) (5)	(b) (5)		
(b) (5)			

(b) (5)

(b) (5)

From: Rotstein, David
To: Reimschuessel, Renate; Queen, Jackie L; Palmer, Lee Anne; Jones, Jennifer L; Ceric, Olgica; Carey, Lauren
Subject: Fwd: Alternated feedings between-California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food and Kangaroo and Lentils: Darcy Adin - EON-323519
Date: Tuesday, July 11, 2017 5:45:51 PM
Attachments: 2023230-report.pdf

David Rotstein, DVM, MPVM, Dipl.ACVP
 CVM Vet-LIRN Liaison
 CVM OSC/DC/CERT
 7519 Standish Place
 (b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: PFR Event <pfreventcreation@fda.hhs.gov>
Date: July 11, 2017 at 5:36:16 PM EDT
To: HQ Pet Food Report Notification <HQPetFoodReportNotification@fda.hhs.gov>, (b) (6) Cleary, Michael *
 <Michael.Cleary@fda.hhs.gov>
Subject: Alternated feedings between:-California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food and Kangaroo and Lentils: Darcy Adin - EON-323519

A PFR Report has been received and PFR Event [EON-323519] has been created in the EON System

A "PDF" report by name "2023230-report pdf" is attached to this email notification for your reference

Below is the summary of the report:

EON Key: EON-323519
ICSR #: 2023230
EON Title: PFR Event created for Alternated feedings between: California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food and Kangaroo and Lentils; 2023230

AE Date	(b) (6)	Number Fed/Exposed	2
Best By Date		Number Reacted	2
Animal Species	Dog	Outcome to Date	Stable
Breed	Schnauzer - Miniature		
Age	7 Years		
District Involved	PFR-New York DO		

Product information

Individual Case Safety Report Number: 2023230

Product Group: Pet Food

Product Name: Alternated feedings between: California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food and Kangaroo and Lentils

Description: Please note: Dr Jennifer Jones was consulted prior to submission of this report She would like to be involved in the case review (b) (6) housemate, (b) (6) (separate report submitted) was diagnosed with DCM and CHF (b) (6) and was euthanized after aggressive treatment of CHF At that time (b) (6) had 2 syncopal events closely related to each other His appetite for dog food declined but he would eat it if tempted with treats mixed in He was presented (b) (6) for more syncopal events and was similarly diagnosed with severe DCM and CHF He was able to be successfully treated however and is clinically doing well on CHF medications as of 7/10/17 A re-review of the myocardial histopathology for (b) (6) housemate (b) (6) was requested at this time because of the unusual diagnosis of DCM in a small breed dog living in the same house as another dog similarly diagnosed a few months ago This re-review by one of our pathologists showed myofiber vacuoles reminiscent of the changes seen in doxorubicin toxicity Since the dog had not received doxorubicin, the pathologist recommended culturing the food for Streptomyces peucetius - the bacterium which produces doxorubicin He also recommended testing for Fusarium spp a fungus which produces Fumonisin B1, a toxin that produces heart failure in pigs Like (b) (6) (unrelated, younger miniature schnauzer), (b) (6) had been fed California Naturals Adult - both kangaroo with lentils and venison with lentils along with Milo's kitchen treats We have samples of these foods from 6/17 but not the original bags from when he was presented (b) (6) These samples were provided at the time (b) (6) also presented with severe DCM and CHF Like (b) (6) had extensive infectious disease testing which was negative and nutritional amino acid deficiencies were ruled out Because of this, their unrelated lineages (although the same breed, they were from different lines), different ages but similar time of presentation, we are considering common environmental factors which could precipitate DCM, including food contamination or toxin exposure We have plasma, serum, urine and myocardial tissue samples (latter only for (b) (6)) stored at -80 Celsius in addition to food and treat samples

Submission Type: Initial

Report Type: Adverse Event (a symptom, reaction or disease associated with the product)

Outcome of reaction/event at the time of last observation: Stable

Number of Animals Treated With Product: 2

Number of Animals Reacted With Product: 2

--	--	--

Product Name	Lot Number or ID	Best By Date
Alternated feedings between: California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food and Kangaroo and Lentils		

Sender information

Darcy Adin
 1060 William Moore Dr
 Raleigh, NY 27607
 USA

Owner information

(b) (6)

To view this PFR Event, please click the link below:
<https://eon.fda.gov/eon/browse/EON-323519>

To view the PFR Event Report, please click the link below:
<https://eon.fda.gov/eon/> (b) (6)

This email and attached document are being provided to you in your capacity as a Commissioned Official with the U S Department of Health and Human Services as authorized by law You are being provided with this information pursuant to your signed Acceptance of Commission

This email message is intended for the exclusive use of the recipient(s) named above It may contain information that is protected, privileged, or confidential Any dissemination, distribution, or copying is strictly prohibited

The information is provided as part of the Federal-State Integration initiative As a Commissioned Official and state government official, you are reminded of your obligation to protect non-public information, including trade secret and confidential commercial information that you receive from the U S Food and Drug Administration from further disclosure The information in the report is intended for situational awareness and should not be shared or acted upon independently Any and all actions regarding this information should be coordinated through your local district FDA office

Failure to adhere to the above provisions could result in removal from the approved distribution list If you think you received this email in error, please send an email to FDAREportableFoods@fda.hhs.gov immediately

From: Rotstein, David
To: Olesen, Jackie L; Jones, Jennifer L; Reimschuessel, Renate; Gerig, Olga; Palmer, Lee Anne; Carey, Lauren
Subject: Fwd: Alternated between: -California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils and Kangaroo & Red Lentils Recipe: Darcy Adin - EON-323515
Date: Tuesday, July 11, 2017 5:17:06 PM
Attachments: 2023228-report.pdf

David Rotstein, DVM, MPVM, Dipl.ACVP
 CVM Vet-LIRN Liaison
 CVM OSC/DC/CERT
 7519 Standish Place
 (b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: PFR Event <pfrventcreation@fda.hhs.gov>
Date: July 11, 2017 at 5:16:15 PM EDT
To: Cleary, Michael * <Michael.Cleary@fda.hhs.gov>, HQ Pet Food Report Notification <HQPetFoodReportNotification@fda.hhs.gov>,
 (b) (6)
Subject: Alternated between: --California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils and Kangaroo & Red Lentils Recipe: Darcy Adin - EON-323515

A PFR Report has been received and PFR Event [EON-323515] has been created in the EON System

A "PDF" report by name "2023228-report pdf" is attached to this email notification for your reference

Below is the summary of the report:

EON Key: EON-323515
ICSR #: 2023228
EON Title: PFR Event created for Alternated between: -California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils and Kangaroo & Red Lentils Recipe; 2023228

AE Date	(b) (6)	Number Fed/Exposed	2
Best By Date		Number Reacted	2
Animal Species	Dog	Outcome to Date	Died Euthanized
Breed	Schnauzer - Miniature		
Age	2 5 Years		
District Involved	PFR-New York DO		

Product information

Individual Case Safety Report Number: 2023228

Product Group: Pet Food

Product Name: Alternated between: -California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils and Kangaroo & Red Lentils Recipe

Description: Please note: Dr Jennifer Jones was consulted prior to submission of this report. She would like to be involved in the case review. 3 week history of cough treated unsuccessfully with doxycycline and prednisone. 3 day history of inappetence and vomiting prior to presentation to NCSU emergency service for dyspnea. Radiographs showed severe pulmonary edema and echocardiogram showed severe Dilated Cardiomyopathy. There was an initial response to diuretic therapy however, he declined and was placed on the ventilator for respiratory support and continued CHF treatment. Attempts to wean off the ventilator were unsuccessful and aquaphoresis was performed. He continued to decline despite aggressive therapy and was euthanized. Infectious disease testing was negative and taurine and carnitine analysis showed adequate levels. Necropsy initially did not reveal a cause for DCM and supported alveolar injury (possibly ventilator related). A re-review of the myocardial histopathology by one of our pathologist showed myofiber vacuoles reminiscent of the changes seen in doxorubicin toxicity. Since the dog had not received doxorubicin, the pathologist recommended culturing the food for Streptomyces peucetius - the bacterium which produces doxorubicin. He also recommended testing for Fusarium spp. a fungus which produces Fumonisin B1, a toxin that produces heart failure in pigs. (b) (6) had been fed California Naturals Adult - both kangaroo with lentils and venison with lentils along with Milo's kitchen treats. We have samples of these foods from 6/17 but not the original bags from when he was presented. (b) (6). These samples were provided at the time his housemate (b) (6) (unrelated, older miniature schnauzer) also presented with severe DCM and CHF. I will enter this dog as a separate affected patient. Both dogs had extensive infectious disease testing which was negative and nutritional amino acid deficiencies were ruled out. Because of this, their unrelated lineages (although the same breed, they were from different lines), different ages but similar time of presentation. (b) (6) had clinical signs at the time. (b) (6) was treated, but didn't present with CHF for several months), we are considering common environmental factors which could precipitate DCM, including food contamination or toxin exposure.

Submission Type: Initial

Report Type: Adverse Event (a symptom, reaction or disease associated with the product)

Outcome of reaction/event at the time of last observation: Died Euthanized

Number of Animals Treated With Product: 2

Number of Animals Reacted With Product: 2

Product Name	Lot Number or	Best By
---------------------	----------------------	----------------

	ID	Date
Alternated between: -California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils and Kangaroo & Red Lentils Recipe		

Sender information

Darcy Adin
1060 William Moore Dr
Raleigh, NY 27607
USA

Owner information

(b) (6)

To view this PFR Event, please click the link below:

<https://eon.fda.gov/eon/browse/EON-323515>

To view the PFR Event Report, please click the link below:

<https://eon.fda.gov/> (b) (6)

This email and attached document are being provided to you in your capacity as a Commissioned Official with the U S Department of Health and Human Services as authorized by law You are being provided with this information pursuant to your signed Acceptance of Commission

This email message is intended for the exclusive use of the recipient(s) named above It may contain information that is protected, privileged, or confidential Any dissemination, distribution, or copying is strictly prohibited

The information is provided as part of the Federal-State Integration initiative As a Commissioned Official and state government official, you are reminded of your obligation to protect non-public information, including trade secret and confidential commercial information that you receive from the U S Food and Drug Administration from further disclosure The information in the report is intended for situational awareness and should not be shared or acted upon independently Any and all actions regarding this information should be coordinated through your local district FDA office

Failure to adhere to the above provisions could result in removal from the approved distribution list If you think you received this email in error, please send an email to FDAReportableFoods@fda.hhs.gov immediately

From: [Jones, Jennifer L](#)
To: [Rotstein, David](#); [Palmer, Lee Anne](#); [Carey, Lauren](#); [Queen, Jackie L](#)
Cc: [Ceric, Olgica](#); [Reimschuessel, Renate](#)
Subject: Head"s up-potential DCM case-Dr. Adin-NCSU-2 Schnauzers
Date: Tuesday, July 11, 2017 11:38:00 AM
Attachments: [image001.png](#)
[image002.png](#)

Vet will submit PFR online →
2 dogs-unrelated miniature schnauzers


Dog 1: 2 yr → presented 2/2017 with fulminant CHF → severe DCM on echo, taurine/carnitine normal, infectious disease testing negative, died on the ventilator, necropsy done-myocardial changes were subtle but could be similar to moldy corn toxicity in pigs → plasma, urine, serum, and myocardial tissue available

Dog 2: 7 yr, had a syncopal episode ~2/2017 but presented to vet for progressive frequency of syncopal episodes → 6/2017 for CHF, diagnosed with DCM similar to housemate, nearly same image on Echo, taurine/carnitine normal, infectious disease testing negative, they have changed the diet (Hill's) and dog is responding to treatment; plasma, urine, and serum available

Dogs were eating California Naturals (different bag than from 2/2017) and treats (Milo's Kitchen);
Vet has samples of food and treats

Jennifer L. A. Jones, DVM

Veterinary Medical Officer
U.S. Food & Drug Administration
Center for Veterinary Medicine
Office of Research
Veterinary Laboratory Investigation and Response Network (Vet-LIRN)
8401 Muirkirk Road, G704
Laurel, Maryland 20708
new tel: 240-402-5421
fax: 301-210-4685
e-mail: jennifer.jones@fda.hhs.gov
Web: <http://www.fda.gov/AnimalVeterinary/ScienceResearch/ucm247334.htm>



**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958501

This is an accurate reproduction of the original electronic record as of 07/27/2016

Flag	Flag Remarks				
Episode Number	Origin	Basis	Sample Type	FIS Smpl Num	Status
	Domestic	Surveillance	Official	16260362	Completed
FEI	Date Collected	Product Code	Responsible Firm	PAC	Hours
3004211953	06/23/2016	72AYT02	Manufacturer	71R801	11
Compliance Num	Country of Origin				
	United States				
Related Smpl Num	Position Class	Sampling District	NDC Number	Permit Number	Storage Rqrmnt.
958500	INV	NWJ-DO			Ambient
Dealer is Consumer	Crx/DEA Schedule	Recall Num	Consumer Compl. Num	Brand Name	
No			146048	Merrick	

Product Description

See Remarks

Product Label

See continuation.

Reason for Collection

Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

MFG Codes

"16025DL1 38310 14133"

Expiration Date

07/26/17

Firm Legal Name

Merrick Pet Care, Inc.

Address

3275 Tierra Blanca Rd Hereford, TX
79045-7823 US

Type of Firm

Manufacturer

Firm FEI

3004211953

FCE

02944

(b) (6)

(b) (6)

Dealer

(b) (6)

Size of Lot

One paper bag weighing 5.4kg

Est. Value

\$.00

Rcpt Type

FDA484

Carrier Name

Date Shipped

Description of Sample

One unopened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe weighing 5.4kg

Method of Collection

See continuation.

How Prepared

See continuation.

Collector's Identification on Package and/or Label

958501 06/23/2016 EB

Collector's Identification on Seal

958501 06/23/2016 Esteban Beltran Investigator

Sample Delivered To

SRL-ACNA

Date Delivered

06/28/2016

Orig C/R & Records To

NWJ-DO

Lab w/Split Sample

0

Lab

SRL

Document Number

1
2
3

Document Date

06/23/2016
06/23/2016
06/23/2016

Document Type

Other
Other
Other

Document Remarks

Copy FDA 484, Receipt for Sample, 1 pg.
Copy FDA 484, Receipt for Sample Amend, 1 pg.
Photos of Product labeling, 3 pages

Date: 07/27/2016

Page: 1 of 3

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958501

This is an accurate reproduction of the original electronic record as of 07/27/2016

Remarks

See continuation.

Payment Amount	Payment Method	704(d) Sample	702(b) Portion	Collector's Name
\$27.99	Cash	No	No	Esteban Beltran
Name of Signer	Date & Time of Signature			Meaning
Esteban Beltran	07/07/2016 08:55 AM ET			Collector

**Food and Drug Administration Office of Regulatory Affairs
Collection Report**

For Sample Number: 958501

This is an accurate reproduction of the original electronic record as of 07/27/2016

Continuation:

Product Label

Finished Product: Label on bag read in parts: "****Lot #:16025DL1 38310 14133****Merrick Whole Health Made Right Purrfect Bistro Grain Free *** REAL CHICKEN RECIPE FISH-FREE *** Net Wt 12Lb (5.4kg) *** WHOLE HEALTH MADE RIGHT! *** SQF INSTITUTE CERTIFIED *** MERRICK PET CARE, INC. P.O. BOX 9800 AMARILLO, TX 79105 USA WWW.MERRICKCARE.COM *** Best By: July 26, 2017****"

Method of Collection

On 06/23/2016, I collected a sample from the storage area of a retail store. The sample was placed in a clear plastic bag. I officially sealed the clear plastic bag containing the sample with a FDA415a on site. The sample was transported via GOV to the NBRP sample prep room.

How Prepared

On 06/28/16 I placed the sample into a shipping box with bubble wrap. I secured the shipping box for shipment to SRL-ACNA via UPS (Tracking #: 1Z A47 51E 01 9888 4498). I delivered the sample inside to the NBRP reception area for UPS pickup.

Remarks

Product Description: Poultry based dry cat food packed in a dark orange paper bag with brown letters and an image of a cat on the front.

An amendment to the original FDA 484 was done in order to further describe what each sample number consists of and to identify what lot number of the product pertains to the sample number. CSO Gobiga Vanniyasingam assisted during the assignment. Related samples include 958500, 958502, 958503, 958504.

Research Paper

Mannose Binding Lectin and Macrophage Migration Inhibitory Factor Gene Polymorphisms in Turkish Children with Cardiomyopathy: No Association with MBL2 Codon 54 A/B Genotype, but an Association between MIF -173 CC Genotype

Nilgun Col-Araz¹, Sibel Oguzkan-Balci²✉, Osman Baspinar³, Tugce Sever², Ayse Balat⁴, Sacide Pehlivan²,

1. University of Gaziantep, Faculty of Medicine, Department of Pediatrics, Gaziantep, Turkey.
2. University of Gaziantep, Faculty of Medicine, Department of Medical Biology and Genetics, Gaziantep, Turkey.
3. University of Gaziantep, Faculty of Medicine, Department of Pediatric Cardiology, Gaziantep, Turkey.
4. University of Gaziantep, Faculty of Medicine, Department of Pediatric Nephrology, Gaziantep, Turkey.

✉ Corresponding author: Assos. Prof. Dr. Sibel Oguzkan Balci, University of Gaziantep, Faculty of Medicine, Department of Medical Biology and Genetics, 27310 Gaziantep, Turkey. E-mail: oguzkan@gantep.edu.tr Tel: +90 342 3606060, Fax: +90 342 3601617.

© Ivyspring International Publisher This is an open-access article distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by-nc-nd/3.0/>) Reproduction is permitted for personal, noncommercial use, provided that the article is in whole, unmodified, and properly cited

Received: 2012.06.27; Accepted: 2012.08.19; Published: 2012.08.22

Abstract

Myocardial inflammation is one of the commonest mechanisms in cardiomyopathy (CMP). Mannose binding lectin (MBL) is a key molecule in innate immunity, while macrophage migration inhibitory factor (MIF) is a constitutive element of the host defenses. We investigated the possible association between polymorphisms of MBL2 and MIF genes and CMP in Turkish children. Twenty-children with CMP and 30 healthy controls were analyzed for codon 54 A/B polymorphism in MBL, and -173 G/C polymorphism in MIF genes by using PCR-RFLP methods. No significant difference was found between genotypes and alleles of MBL2 gene codon 54 A/B polymorphism in patients and controls ($p>0.05$). However, serum uric acid levels was found higher in dilated CMP patients with AA genotype. Frequency of MIF -173 CC genotype was significantly higher in patients ($p<0.05$), and sodium levels were higher in patients with MIF -173 CC genotype. This study is the first to investigate the MBL and MIF gene polymorphisms in Turkish children with CMP. We conclude that CC genotype of MIF (-173) polymorphism may be a risk factor for CMP patients. However, further studies with larger samples are needed to address the exact role of this polymorphism in CMP.

Key words: Cardiomyopathy, Children, Macrophage migration inhibitory factor, Mannose binding lectin, polymorphism.

Introduction

Cardiomyopathy (CMP) is defined as “diseases of the myocardium associated with cardiac dysfunction” by World Health Organization (WHO), and it is an important cause of chronic congestive cardiac failure in children. The reported incidence for cardiomyopathies is 1,13-1,24 per 100,000 children [1, 2].

Although the pathogenesis of disease is not fully understood, disturbances of the cellular and humoral immune system are frequently observed in CMPs, and myocardial inflammation is one of the commonest mechanisms in cardiomyopathy [3].

Mannose binding lectin (MBL) and Macrophage

migration inhibitory factor (MIF) play substantial roles in the pathogenesis of several inflammatory and autoimmune disorders [4, 5]. Mannose binding lectin is a key molecule in innate immunity with the capacity to bind to microorganisms and kill them by initiating the lectin pathway of complement activation [5]. Furthermore, MBL has a major role in the modulation of inflammation but the mechanisms responsible for MBL interactions with inflammatory pathways is remain unclear [6]. Several studies suggest that, there is a modulatory role of MBL in autoimmune disease such as rheumatoid arthritis and systemic lupus erythematosus [6-8]. Previous studies show that the absence of MBL may affect occurrence of cardiovascular complications and myocardial ischemia/reperfusion injury, and CMP in MBL null animal models (9-11). Mannose Binding Lectin deficiency has been reported by three single nucleotide polymorphisms (SNPs) in codon 52, 54 and 57 of exon 1 in the MBL2 gene [6]. These SNPs are frequently referred to as variants B, C, and D (B, C, and D, denoting the substitution of aspartic acid for glycine codon 54, the substitution of glutamic acid for glycine codon 57, and the substitution of cysteine for arginine codon 52, respectively). Each of these variant alleles affect the stability of the final protein product, resulting in decreased serum levels and a dysfunctional MBL variant with a lower molecular weight than the normal MBL [6, 12].

Macrophage migration inhibitory factor is a constitutive element of the host antimicrobial defences and stress response that promotes proinflammatory function of the innate and acquired immune system. Macrophage migration inhibitory factor plays a regulator role in the immune response system and promotes proinflammatory biological activities. MIF is constitutively expressed in variety types of tissue and cells, including innate immune cells such as monocytes and macrophages [4]. Recently, it has been shown that MIF gene expression is higher in the heart with impaired glucose tolerance with cardiac dysfunction in rats, and elevated levels of MIF were associated with cardiac dysfunction in diabetic patients [13]. Mutations of the human MIF gene would predispose affected hosts to altered to sensibility or severity of inflammatory diseases such as juvenile idiopathic rheumatoid arthritis, and glomerulonephritis [4, 14, 15].

To our knowledge, no studies have investigated the possible roles of MBL2 and MIF gene polymorphisms in children with CMP. The aim of the present study is to investigate any possible association between polymorphisms of MBL2 and MIF genes and CMP in a group of Turkish children, and to investigate the association between the identified genotypes

and their clinical features.

Materials and Methods

Patients and controls: Twenty unrelated Turkish children with CMP, followed up in the Paediatric Cardiology Clinic of the Gaziantep University, Medical Faculty, were compared with 30 age- and sex-matched healthy controls. Relatives of CMP patients did not included as healthy controls. The diagnosis of CMP were made by signs and symptoms (irritability, feeding difficulties, weakness, fatigue, dizziness, syncope, tachypnea, tachycardia, hepatomegaly, and evidence of fluid retention), chest X ray (cardiomegaly, pulmonary venous congestion, pulmonary oedema), electrocardiography (hypertrophy of left ventricle with strain, low voltage complexes) and echocardiographic signs. Cardiomyopathies were classified according to their structural and functional abnormalities such as dilated, in the setting of reduced left ventricular systolic function; hypertrophic, in the presence of unexplained septal hypertrophy of the left ventricle; restrictive, when impaired diastolic filling with preserved systolic function and normal ventricular wall thickness [1]. The study was approved by the Local Ethics Committee of the Faculty of Medicine, and informed consents were obtained from the parents of children. The medical records of all children with CMP were reviewed for information about age, sex, and to document clinical presentation including symptoms, family history, laboratory and echocardiographic findings.

Genotyping: All patients and controls were analyzed for codon 54 A/B (gly54asp) variation in exon 1 of MBL2 gene and -173 G/C polymorphism in MIF gene. Genomic DNA was extracted from peripheral blood samples using the salting out procedure [16].

Genotyping of MBL2 gene codon 54 A/B: Polymerase Chain reaction (PCR) was performed using a forward (5'-TAGGACAGAGGGCATGCTC-3') and a reverse (5'-CAGGCAGTTTCTCTGGAAGG-3') primers in a 25 µl volume containing 50 ng DNA, 2 mM dNTPs, 2 nmol of each primer, 1.5 mM MgCl₂ and 3U Taq polymerase. The product 349 bp was digested with restriction enzyme *BanI* (Fermentas) identify codon 54 polymorphism, respectively. *BanI* digestion was performed at 50 °C for 60 minutes with 5 U enzyme. After enzyme digestion, products were visualized by electrophoresis on 3% agarose gel. The *BanI* restriction site is present on wild type allele A and absent on variant allele B [17].

Genotyping of MIF gene -173 G/C: PCR was performed using a forward (5'-ACTAAGAAAGACCCGAGGC-3') and reverse (5'-GGGGCACGTTGGTGTTC-3') primers. For

MIF (-173), a 330 bp fragment was amplified, which was then digested with AluI restriction enzyme (Fermentas), overnight at 37 °C. The products were then separated on 3% agarose gel. The PCR product contains two restriction site for allele C and one of these sites is destroyed when the presence of allele G [18].

Statistical Analysis: All statistical analyses were performed with the Statistical Package for the Social Science for Windows (version 18.0; SPSS Inc, Chicago, IL, U.S.A.). Results are given as mean±SD, while allele frequencies and the distribution of genotype are given as %. Clinical features and MBL/MIF gene polymorphisms were compared using the chi-square and the Fisher's exact tests. Differences between groups were compared by Kruskal-Wallis variant analysis and the Mann-Whitney U-test. Statistical significance was considered at $p < 0.05$. Hardy-Weinberg equilibrium (HWE) was calculated using De-finetti program [19]. Differences in allele and genotype distributions were assessed using odds ratios (ORs) and 95% confidence intervals. Sample size was estimated using a power calculation based on other studies [20]. The minimum sample size was determined as 44 person in each group at the 80% power level with an α error of 5%.

Results

Clinical features: The age ranged from 3 months to 13 years (mean: 3.47 ± 3.38 years; median: 2.00-IR:3.00) in patients with CMP (n=20, 10 females/10 males). According to the echocardiographic evaluation, 80% (n=16) of the patients had dilated cardiomyopathy, 15% (n=3) hypertrophic cardiomyopathy and 5% (n=1) restrictive cardiomyopathy. All patients had clinical findings of CMP. Echocardiographic findings of CMP patients were shown in Table 1.

Genotype frequencies of MBL2 and MIF genes

The distribution of AA, AB, and BB genotypes for MBL codon 54 were 65 % (13), 25 % (5) and 10 % (2) in CMP compared with 76.7 % (23), 23.3 % (7) and 0 % (0) in the controls. The allele frequency of A/B in MBL was 77.5 % (31), and 22.5 % (9) in CMP compared with 88.3 % (53), and 11.7 % (7) in the controls. No significant difference was found between genotypes and alleles of MBL2 gene in patients and controls ($p > 0.05$).

The distribution of GG, GC, and CC genotypes

for MIF (-173) were 50 %, 30 %, and 20 % in CMP compared with 56.7 %, 43.3 % and 0 % in the controls (Table 2). CC genotype was significantly higher in patient group ($p = 0.0210$, Table 2). The allele frequency of G/C in MIF was 65 %, and 35 % in CMP compared with 78.3 %, and 21.7 % in the controls. The observed genotype counts were not deviated significantly from those expected according to the Hardy-Weinberg Equilibrium for MBL and MIF gene polymorphisms ($p > 0.05$).

Genotype frequencies of MBL2 and MIF genes

The distribution of AA, AB, and BB genotypes for MBL codon 54 were 65 % (13), 25 % (5) and 10 % (2) in CMP compared with 76.7 % (23), 23.3 % (7) and 0 % (0) in the controls. The allele frequency of A/B in MBL was 77.5 % (31), and 22.5 % (9) in CMP compared with 88.3 % (53), and 11.7 % (7) in the controls. No significant difference was found between genotypes and alleles of MBL2 gene in patients and controls ($p > 0.05$).

The distribution of GG, GC, and CC genotypes for MIF (-173) were 50 %, 30 %, and 20 % in CMP compared with 56.7 %, 43.3 % and 0 % in the controls (Table 2). CC genotype was significantly higher in patient group ($p = 0.0210$, Table 2). The allele frequency of G/C in MIF was 65 %, and 35 % in CMP compared with 78.3 %, and 21.7 % in the controls. The observed genotype counts were not deviated significantly from those expected according to the Hardy-Weinberg Equilibrium for MBL and MIF gene polymorphisms ($p > 0.05$).

Association between the identified genotypes and patients clinical/laboratory characteristics: We investigated correlations of MBL/MIF genotypes with clinical and laboratory findings of patients such as duration of symptoms, ejection fraction (EF), fractional shortening (FS), left ventricle end diastolic diameter (LVEDD), left ventricle end systolic diameter (LVESD), left ventricle end diastolic volume (LVEDV), and left ventricle end systolic volume (LVESV). No relationship was found between MBL/MIF genotypes and these parameters (data not shown).

In children with dilated CMP (n=16), serum uric acid levels were higher in patients with MBL AA genotype ($p = 0.033$, Table 3), while plasma sodium (Na) levels were higher in patients with MIF CC genotype ($p = 0.042$, Table 4).

Table 1. Echocardiographic signs of children with cardiomyopathy (CMP).

	Dilated CMP Mean \pm SD (min-max) (n=16)	Hypertrophic CMP Mean \pm SD (min-max) (n=3)	Restrictive CMP (n=1)
Ejection fraction (EF) (%)	33.75 \pm 11.13 (19-58)	77.00 \pm 3.60 (74-81)	58.00
Fractional shortening (FS) (%)	15.07 \pm 5.89 (8-29)	35.67 \pm 6.02 (30-42)	29.00
Left ventricle end diastolic diameter (LVEDD) (cm)	4.75 \pm 1.04 (2.40-5.90)	2.83 \pm 0.72 (2.00-3.30)	2.40
Left ventricle end systolic diameter (LVESD) (cm)	3.98 \pm 1.07 (1.70-5.40)	1.63 \pm 0.40 (1.20-2.00)	1.70
Left ventricle end diastolic volume (LVEDV) (ml)	119.94 \pm 42.08 (29.60-173.00)	32.600 \pm 17.30 (12.70-44.10)	45.00
Left ventricle end systolic volume (LVESV) (ml)	82.15 \pm 37.53 (18.10-141.00)	8.15 \pm 4.67 (3.36-12.70)	22.00

Table 2. Genotype and allele frequencies of Macrophage Migration Inhibitory Factor (MIF) gene -173 G/C polymorphism in children with cardiomyopathy (CMP).

MIF Genotype	Control n(%)	CMP Patients n(%)	Odds Ratio (95% C.I.)	p
GG	17 (56.7)	10 (50)	0.823 (0.266 - 2.541)	0.4793 ^a
GC	13 (43.3)	6 (30)	0.560 (0.169 - 1.858)	0.2578 ^a
CC	0 (0)	4 (20)	16.636 (0.842 - 328.60)	0.0210 ^a
MIF Allele				
G	47 (78.3)	26 (65)	0.513 (0.210 - 1.256)	0.1072
C	13 (21.7)	14 (35)	1.947 (0.796 - 4.761)	0.1072
HWE (p)	0.129	0.127		

^aFisher exact test, HWE: Hardy-Weinberg Equilibrium.

Table 3. Association with serum uric acid levels and Mannose Binding Lectin (MBL2) gene codon 54 A/B polymorphism in children with dilated cardiomyopathy.

MBL genotypes	Serum uric acid level (mg/dL) Mean \pm SD (min-max)	95% CI	p
AA	6,139 \pm 1,508 (4,5-8,4)	4,979- 7,299	0.033
AB	3,000 \pm 0,989 (2,3-3,7)	- 5,894- 11, 894	

Table 4. Association with plasma sodium (Na) levels and Macrophage Migration Inhibitory Factor (MIF) gene -173 G/C polymorphism in children with dilated cardiomyopathy.

MIF genotypes	Plasma Na level (mEq/L) Mean \pm SD (min-max)	95% CI	p
CC	137,50 \pm 2,121 (136-139)	118,44-156,56	0.042
GC	131,00 \pm 1,414 (130-132)	118,29- 143,71	

Discussion

Although the pathogenesis of CMP is not fully understood, cellular as well as humoral autoimmune responses are critically associated with the pathogenesis and progression of the disease. Furthermore, disturbances of the cellular and humoral immune system are frequently observed, and myocardial inflammation is one of the commonest mechanisms in cardiomyopathy [3]. Two of the postulated factors are; firstly, myocardial inflammation mediated by the effector cells of the immune system; and secondly local effect of inflammatory mediators, released by the infiltrating lymphocytes, macrophages or endothelial cells [21]. Both MIF and MBL play several roles in innate and adaptive immune responses, and changes in levels of MBL and MIF are implicated as playing causative role in many disease states [4, 6, 22].

Better understanding of the molecular genetics underlying CMP may provide a means of early diagnosis, genotype-based therapy, and even prevention of the disease.

Mannose binding lectin deficiency is associated with susceptibility to infectious and autoimmune diseases and serum MBL levels vary substantially because of the variant alleles in exon 1 of the MBL2 gene, located on chromosome 10 in the humans [22]. In the present study, homozygosity for MBL variant allele was observed only two patients with dilated CMP and the A/A genotype was higher than the variant alleles in CMP patients, but these results were not statistically significant.

Messias-Reason et al, suggested that wild type variants of MBL2 gene was significantly higher in rheumatic heart diseases [23]. Ramasawmy et al. reported that subjects homozygous for the wild type allele had a higher concentration of MBL than heterozygous subjects and than homozygous for the variant MBL2 alleles [24]. Schafranski et al showed that, genotypes associated with a higher level of MBL seem to represent a risk factor for the evolution of rheumatic carditis, and MBL play a substantial role in the progression of the disease to chronic form [25]. However, some studies indicated that there was either no association of MBL gene polymorphisms and systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA), or even an increased risk for RA has been demonstrated in MBL insufficiency [8, 26]. In the present study, homozygosity for MBL variant allele was observed only two patients with dilated CMP and the A/A genotype was higher than the variant alleles in CMP patients, but these results were not statistically significant. These different results in literature highlight the need for further detailed studies

to understand the exact role of MBL2 gene polymorphism in several diseases.

We investigated correlations of MBL genotypes with clinical and laboratory findings of the disease, and found that uric acid levels were higher in patients with MBL AA genotype than the other genotypes. Uric acid is a useful marker for the decompensation trigger in chronic heart failure, and might be related to inflammatory responses [27]. It has been shown that the left ventricular hypertrophy has an important potential to increase uric acid level [28]. Gullu et al. measured uric acid levels in idiopathic dilated cardiomyopathy, and they observed that serum uric acid levels are significantly higher in the lower coronary flow reverse group than in the higher coronary flow reverse group [29]. Interestingly, Garred et al reported that homozygous for wild type alleles in exon 1 of MBL2 gene were more likely to show evidence of persistent inflammation [30]. We suggest that the reason of increased uric acid level in our patients may be both inflammation and chronic heart failure.

Macrophage migration inhibitory factor plays an important role in the control of innate immune responses and promotes proinflammatory biological activities. Four polymorphisms of the human MIF gene (-794, -173, +254, +656) have been reported, and this polymorphisms would predispose affected hosts to altered susceptibility to or severity of inflammatory or infectious disease [4]. Patients with -173 C allele (that is, guanine-to-cytosine transition at position -173) had increased levels of MIF, and increased MIF concentrations had been associated with severe clinical manifestations, high severity scores, and poor outcome of inflammatory disease [4, 31, 32]. In the other studies, no association were found in genotype distributions of MIF -173 G/C polymorphism between ulcerative colitis, juvenile rheumatoid arthritis and healthy controls [14, 33]. However, Donn et al showed that MIF -173 C allele was associated with juvenile idiopathic arthritis [31, 32]. Moreover, MIF-173 C allele had a significantly greater number of joints with active arthritis and was associated with a poor response to glucocorticoids in patients with juvenile idiopathic arthritis [31]. Berdeli et al showed that, the MIF -173 C allele was a poor outcome predictor in JRA [14].

Miller et al. demonstrated that MIF released from ischemic cardiomyocytes stimulates adenosine monophosphate-activated protein kinase (AMPK) activation and promotes glucose uptake, and thereby protects the heart against ischaemia reperfusion injury [34]. Jian et al found that MIF protein is constitutively expressed by cardiomyocytes in vivo and is increased in the myocardium of infants with cyanotic

cardiac defects in myocardial biopsy materials [35]. Tereshchenko et al did not reveal an association of the myocardial infarction with the MIF-173 C allele polymorphism [36]. In the present study, homozygosity for MIF-173 C allele was observed only four patients with dilated CMP. Recently, it has been shown that presence of -173C allele indicates higher MIF levels [37], and cardiac inflammation (autoimmune, viral or post viral) has an important component in the pathogenesis of dilated CMP [38]. Therefore we suggest that, CC genotype in our patients may be partially responsible from inflammation in dilated CMP, and MIF polymorphism may contribute to MIF release from cardiomyocytes in children with CMP. However, we could not find any relationship between MIF genotypes and cardiac functions. Considering the limited number of our patients, we cannot say that MIF polymorphism does not modulate cardiac functions. Further detailed studies with large patient numbers are needed for this suggestion.

It has been shown that, plasma brain natriuretic peptide (BNP) concentrations were increased in various forms of heart disease with impaired left ventricular systolic function including cardiomyopathy [39]. Natriuretic peptides inhibit the transport of sodium and water in proximal tubules and block reabsorption of sodium [40]. In this study, plasma sodium levels were higher in patients with MIF CC genotype than the other genotypes. However, we could not conclude whether CC genotype of MIF has any effect on BNP with this study. This hypothesis needs further evaluation.

Conclusion

This study is the first to investigate the MBL and MIF gene polymorphisms in Turkish children with CMP. We conclude that CC genotype of MIF (-173) gene may be a risk factor for CMP patients. However, further studies with larger samples are needed to address the exact role of this polymorphism in CMP.

Competing Interests

The authors have declared that no competing interest exists.

References

- Nugent AW, Daubeney DE, Chandras P, et al National Australian Childhood Cardiomyopathy Study The epidemiology of childhood cardiomyopathy in Australia *N Engl J Med* 2003; 348 1639-46
- Lipshultz SE, Sleeper TA, Towbin JA, et al The incidence of pediatric cardiomyopathy in two regions of the United States *N Engl J Med* 2003; 348 1647-55
- Maisch B, Richter A, Sandmoller A, BMBF – Herth Failure Network Inflammatory dilated cardiomyopathy (DCMI) *Herz* 2005; 30 535-44
- Renner P, Roger T, Calandra T Macrophage migration inhibitory factor Gene polymorphisms and susceptibility to inflammatory diseases *Clin Infect Dis* 2005; 41 513-9
- Ruskamp JM, Hoekstra MO, Rovers MM, et al Mannose-binding lectin and upper respiratory tract infections in children and adolescents *Arch Otolaryngol Head Neck Surg* 2006;132 482-6
- Turner MW The role of mannose-binding lectin in health and disease *Mol Immunol* 2003; 40 423-9
- Garred P, Madsen HO, Hallberg P, et al Mannose-binding lectin polymorphisms and susceptibility to infection in systemic lupus erythematosus *Arthritis Rheum* 1999; 42 2145-52
- Ip WK, Lau L, Chan SY, et al Mannose-binding lectin and rheumatoid arthritis in Southern Chinese *Arthritis Rheum* 2000; 43 1679-87
- Zou C, La Bonte LR, Pavlov VI, et al Murine hyperglycemic vasculopathy and cardiomyopathy whole-genome gene expression analysis predicts cellular targets and regulatory networks influenced by mannose binding lectin *Front Immunol* 2012; 3 15
- Pavlov VI, La Bonte LR, Baldwin WM, et al Absence of mannose-binding lectin prevents hyperglycemic cardiovascular complications *Am J Pathol* 2012; 180 104-12
- Busche MN, Walsh MC, McMullen ME, et al Mannose-binding lectin plays a critical role in myocardial ischaemia and reperfusion injury in a mouse model of diabetes *Diabetologia* 2008; 51 1544-51
- Schafrański MD, Ferrari LP, Scherner D, et al High-producing MBL2 genotypes increase the risk of acute and chronic carditis in patients with history of rheumatic fever *Mol Immunol* 2008; 45 3827-31
- Yu XY, Chen HM, Liang JL, et al Hyperglycemic myocardial damage is mediated by proinflammatory cytokine macrophage migration inhibitory factor *PLoS One* 2011; 25 e16239
- Berdeli A, Ozyürek AY, Ulger Z, et al Association of macrophage migration inhibitory factor gene -173G/C polymorphism with prognosis in Turkish children with juvenile rheumatoid arthritis *Rheumatol Int* 2006; 26 726-31
- Lan HY, Yang N, Nikolic- Paterson DJ, et al Expression of macrophage migration inhibitory factor in human glomerulonephritis *Kidney Int* 2000; 57 499-509
- Miller SA, Dykes DD, Polesky HF A simple salting out procedure for extracting DNA from human nucleated cells *Nucleic Acids Res* 1988; 16 1215
- Vardar F, Pehlivan S, Onay H, et al Association between mannose binding lectin polymorphisms and predisposition to bacterial meningitis *Turk J Pediatr* 2007; 49 270-3
- Akcali A, Pehlivan S, Pehlivan M, et al Association of macrophage migration inhibitory factor gene promoter polymorphisms with multiple sclerosis in Turkish patients *J Int Med Res* 2010; 38 69-77
- [Internet] Strom TM, Wienker TF Hardy-Weinberg equilibrium online analysis program <http://ihg.gsf.de/cgi-bin/hw/hwa2.pl>
- Col-Araz N, Pehlivan S, Baspinar O, et al Association of macrophage migration inhibitory factor and mannose-binding lectin-2 gene polymorphisms in acute rheumatic fever *Cardiol Young* 2012; [Epub ahead of print]
- Pankuweit S, Ruppert V, Maisch B Inflammation in dilated cardiomyopathy *Herz* 2004; 29 788-93
- Worthley DL, Barty PG, Mullighan CG Mannose-binding lectin biology and clinical implications *Intern Med J* 2005; 35 548-55
- Messias Reason LJ, Schafrański MD, Jensenius JC, et al The association between mannose-binding lectin gene polymorphism and rheumatic heart disease *Hum Immunol* 2006; 67 991-8
- Ramasawmy R, Spina SG, Fae KC, et al Association of mannose-binding lectin gene polymorphism but not of mannose-binding serine protease 2 with chronic severe aortic regurgitation of rheumatic etiology *Clin Vaccine Immunol* 2008; 15 932-6
- Schafrański MD, Ferrari LP, Scherner D, et al High-producing MBL2 genotypes increase the risk of acute and chronic carditis in patients with history of rheumatic fever *Mol Immunol* 2008; 45 3827-31
- Horiuchi T, Tsukamoto H, Morita C, et al Mannose-binding lectin (MBL) gene mutation is not a risk factor for systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA) in Japanese *Genes Immun* 2000; 1 464-6
- Leyva F, Anker SD, Godsland IF, et al Uric acid in chronic heart failure A marker of chronic inflammation *Eur Hearth J* 1998; 19 1814-22
- Kurata A, Shigematsu Y, Higaki J Sex-related differences in relations of uric acid to left ventricular hypertrophy and remodeling in Japanese hypertensive patients *Hypertens Res* 2005; 28 133-9
- Gullu H, Erdogan D, Caliskan M, et al Elevated serum uric acid levels impair coronary microvascular function in patients with idiopathic cardiomyopathy *Eur J Hearth Fail* 2007; 9 466-8
- Garred P, Madsen HO, Marquarth H, et al Two edged role of mannose binding lectin in rheumatoid arthritis a cross sectional study *J Rheumatol* 2000; 27 26-34

- 31 De Benedetti FD, Meazza C, Vivarelli M, et al The British Paediatric Rheumatology Study Group Functional and prognostic relevance of the -173 polymorphism of the macrophage migration inhibitory factor gene in systemic-onset juvenile idiopathic arthritis *Arthritis Rheum* 2003; 48 1398-407
- 32 Donn R, Alourfi Z, De Benedetti F, et al The British Paediatric Rheumatology Study Group Mutation screening of the macrophage migration inhibitory factor gene Positive association of a functional polymorphism of macrophage migration inhibitory factor with juvenile idiopathic arthritis *Arthritis Rheum* 2002; 46 2402-9
- 33 Nohara H, Okayama N, Inoune N, et al Association of the 173G/C polymorphism of the macrophage migration inhibitory factor gene with ulcerative colitis *J Gastroenterol* 2004; 39 242-6
- 34 Miller EJ, Li J, Leng L, et al Macrophage migration inhibitory factor stimulates AMP-activated protein kinase in the ischaemic hearth *Nature* 2008; 451 578-82
- 35 Jian Z, Li JB, Ma RY, et al Increase of macrophage migration inhibitory factor (MIF) expression in cardiomyocytes during chronic hypoxia *Clin Chim Acta* 2009; 405 132-8
- 36 Tereshchenko IP, Petrakova J, Mrazek F, et al The macrophage migration inhibitory factor (MIF) gene polymorphism in Czech and Russian patients with myocardial infarction *Clin Chim Acta* 2009; 402 199-202
- 37 Xie Q, Wang SC, Bian G, et al Association of MIF -173G/C and MBL2 codon 54 gene polymorphisms with rheumatoid arthritis a meta-analysis *Hum Immunol* 2012; [Epub ahead of print]
- 38 Pankuweit S, Ruppert V, Maisch B Inflammation in dilated cardiomyopathy *Herz* 2004; 29 788-93
- 39 Bielecka-Dabrowa A, Wierzbicka M, Dabrowa M, et al New methods in laboratory diagnostics of dilated cardiomyopathy *Cardiol J* 2008; 15 388-95
- 40 Bettencourt PM Clinical usefulness of B-type natriuretic peptide measurement present and future perspectives *Hearth* 2005; 91 1489-94

For reprint orders, please contact reprints@expert-reviews.com

EXPERT
REVIEWS

Nutritional and micronutrient determinants of idiopathic dilated cardiomyopathy: diagnostic and therapeutic implications

Expert Rev. Cardiovasc. Ther. 9(9), 1161–1170 (2011)

Victor Marinescu¹ and Peter A McCullough²

¹Department of Medicine, William Beaumont Hospital, Royal Oak, MI 48073, USA

²St John Providence Health System, Providence Park Heart Institute, Novi, MI 48374, USA

*Author for correspondence:

Tel.: +1 248 885 4197

Fax: +1 248 453 5879

victor.marinescu@beaumont.edu

Idiopathic dilated cardiomyopathy (IDCM) is the term used to describe a group of myocardial diseases of unknown cause whose common clinical presentation is heart failure. The prevalence of IDCM is estimated to be between 7 and 13% of patients with systolic heart failure. Throughout medical history, several nutrient-deficient states have been identified as the root cause of IDCMs, Keshan's disease being one such example, where selenium deficiency-induced heart failure is now well documented. This raises the question of whether a micro- or macro-nutrient imbalance can provide the milieu for inefficient energy expenditure and cardiac metabolism in the context of IDCMs, either causing or exacerbating the condition. To date, there is insufficient evidence in the literature to support this theory, although numerous studies suggest a link between nutrient deficiencies, inefficient energy expenditure and subsequent heart failure. Given the unique metabolic needs of the failing heart, the role of micronutrient testing and supplementation in IDCMs warrants further well-designed studies.

KEYWORDS: heart failure • idiopathic dilated cardiomyopathy • macrominerals • metabolic cardiology • micronutrients • multivitamin supplementation • vitamins

(b) (4)

(b) (4)



(b) (4)



(b) (4)



(b) (4)



(b) (4)



(b) (4)




(b) (4)



(b) (4)



(b) (4)



Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.

From: [Edwards, David](#)
To: [Burkholder, William](#)
Cc: [Hartogenesis, Martine](#)
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference
Date: Tuesday, February 26, 2019 1:53:50 PM
Attachments: [Mansilla 2019 J Anim Sci.pdf](#)

Also, most of the panel are authors on the attached paper that is to come out in March.

Dave

From: Hartogenesis, Martine
Sent: Tuesday, February 26, 2019 1:47 PM
To: Burkholder, William <William.Burkholder@fda.hhs.gov>
Cc: Edwards, David <David.Edwards@fda.hhs.gov>
Subject: FW: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Bill,

[REDACTED] (b) (5)
[REDACTED] Jen and Lee Anne can help you with slides.

Does that work for you? Thanks again!

Martine

From: Norris, Anne
Sent: Tuesday, February 26, 2019 1:12 PM
To: Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>; Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

From Petfood Forum web page:

[Update on canine DCM investigation: lessons learned, insights for the future](#)

In July 2018, the U.S. Food and Drug Administration (FDA) announced it was investigating cases of canine dilated cardiomyopathy (DCM) in breeds not typically associated with the disease, and that many of the dogs had been fed grain-free pet foods high in pulses, legumes or potatoes. Is there really a link between those ingredients or these pet food formulations and the cases of DCM? This panel will provide the latest updates on the FDA and other investigations, and discuss any lessons learned and insights for the industry going forward.

Greg Aldrich, Ph.D. (moderator), research associate professor at Kansas State University and president of Pet Food Ingredients & Technology

Jennifer Adolphe, R.D., Ph.D., nutrition manager for Petcurean Pet Nutrition

William Burkholder, D.V.M., Ph.D., veterinary medical officer with the Center for Veterinary Medicine, Food and Drug Administration

Chris Marinangeli, Ph.D., director of nutrition, scientific and regulatory affairs for Pulse Canada

Anna Kate Shoveller, Ph.D., assistant professor, Department of Animal Biosciences, at University of Guelph

From: Forfa, Tracey

Sent: Thursday, February 21, 2019 1:42 PM

To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Thanks! I will set up a time where we can touch base and decide.

From: Hartogenesis, Martine

Sent: Thursday, February 21, 2019 1:41 PM

To: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>

Cc: Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Ok, thanks!

Martine

From: DeLancey, Siobhan

Sent: Thursday, February 21, 2019 12:19 PM

To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>

Cc: Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

I'm going to defer to Tracey, but I think it may warrant further discussion. Folding her in here.

From: Hartogenesis, Martine

Sent: Thursday, February 21, 2019 12:03 PM

To: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Siobhan,

Thank you so much for weighing in. [REDACTED] (b) (5)

[REDACTED]

[REDACTED] (b) (5).

Martine

From: DeLancey, Siobhan

Sent: Thursday, February 21, 2019 10:55 AM

To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hmmm. [REDACTED] (b) (5)

[REDACTED] (b) (5). Also, wondering who the audience would be for this? I assume anyone can attend, whether or not they are members of this organization or PFI? It was a little unfortunate that AFIA got the update overview before our communication went out, and so I think that the other advocates are even more sensitive to the appearance that we are only talking to industry. [REDACTED] (b) (5)

From: Hartogensis, Martine

Sent: Thursday, February 21, 2019 9:54 AM

To: Norris, Anne <Anne.Norris@fda.hhs.gov>

Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Just circling back on this....Siobhan, any thoughts?

Martine

From: Norris, Anne

Sent: Tuesday, February 19, 2019 10:50 AM

To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>

Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>

Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

It seems duplicative to do a [REDACTED] (b) (5)

[REDACTED] but defer to Siobhan for her thoughts.

From: Hartogenesis, Martine
Sent: Tuesday, February 19, 2019 10:40 AM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Yes, correct! I think this request is for a similar webinar as we did last summer/fall. I believe Bill is still scheduled to go to the Forum.

Let me know your thoughts, of if you would prefer to coordinate.

Thanks again!

Martine

From: Norris, Anne
Sent: Tuesday, February 19, 2019 10:21 AM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

This is separate from the request for Bill to present at the Petfood Forum (the same publication's annual conference)? Last I knew, Bill was going to do that because he was doing a labeling workshop at that event as well.

Just making sure I'm not getting my wires crossed!

From: Hartogenesis, Martine
Sent: Tuesday, February 19, 2019 9:29 AM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>; Burkholder, William <William.Burkholder@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Edwards, David <David.Edwards@fda.hhs.gov>; Schell, Timothy <Timothy.Schell@fda.hhs.gov>
Subject: FW: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi DCM Team!

See the request below from Watt Global Media. You may recall we participated in a webinar to talk about DCM late last summer (following our update).

My thought was to give our group an opportunity to present updated information [REDACTED] (b) (5) [REDACTED] (similar to Dave's AFIA slides).

They are looking for a late March/early April timeframe, so please let me know if that would work for you.

Jen and Lee Anne, I would be looking to you for some slides if we need to [REDACTED] (b) (5) [REDACTED]

Thank you all in advance!

Martine

From: Debbie Phillips <DPhillips@wattglobal.com>

Sent: Monday, February 18, 2019 9:41 AM

To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>

Subject: FW: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hello Martine:

I saw your colleague, Dr. Dave Edwards, present last week at the AFIA pet food conference in Atlanta, and it reminded me that I still need to determine if this webinar is a "go." Can you please let me know if you're still available to participate and if you have time during the last week of March or first two weeks of April – which week would be best for you?

As I explained in my email below, we are broadening the topic of the webinar somewhat; it will still include a discussion of the DCM investigation but also cover other, related topics, especially those relevant to novel ingredients and their relative safety in pet foods.

During his presentation, Dr. Edwards gave a brief list of pet food-related issues that CVM is investigating, and I believe your contribution to this webinar could be a brief update on several or all of these:

- *L. monocytogenes, Clostridium, Salmonella* (including recalls in raw pet food)
- DCM
- Elevated thyroid levels in pet foods
- Vitamin D-related recalls
- Pentobarbital in pet foods

Please let me know if that makes sense to you and your availability. Thank you!

Sincerely,
Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13

Petfood Forum China 2019: August 20

Petfood R&D Showcase 2019: October 15-17

Visit www.PetfoodForumEvents.com

From: Debbie Phillips
Sent: Friday, February 1, 2019 9:23 AM
To: 'Hartogenesis, Martine' <Martine.Hartogenesis@fda.hhs.gov>
Subject: FW: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Martine:

Happy 2019, a month late! I imagine you're very busy considering everything that's been happening (or not happening) in DC recently, but I wanted to follow up on the potential webinar we had corresponded about late last year.

For various reasons, we have decided to broaden the topic of this webinar and cover other aspects of so-called novel pet food ingredients, including benefits, challenges, investigations, etc. That would include the update on the canine DCM investigation and situation –and possibly other situations related to pet food that CVM is involved with now?

Could you please let me know if this is something you could still participate in? Right now we are looking at having the webinar in late March (last week of the month) or first couple of weeks of April. Is there a timeframe that's best for you?

Fyi, I am also following up with Dr. Burkholder about his participating in the DCM panel discussion at our Petfood Forum conference in late April/early May.

Thanks very much,
Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13

Petfood Forum China 2019: August 20

Petfood R&D Showcase: October 2019

Visit www.PetfoodForumEvents.com

From: Debbie Phillips
Sent: Friday, November 30, 2018 9:48 AM
To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Martine,

Sounds good all around! And yes for the webinar; registration for our webinars is free so you can have as many sign up as you wish.

Thanks again,
Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13
Petfood Forum China 2019: August 20
Petfood R&D Showcase: October 2019
Visit www.PetfoodForumEvents.com

From: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Sent: Friday, November 30, 2018 8:40 AM
To: Debbie Phillips <DPhillips@wattglobal.com>
Cc: Edwards, David <David.Edwards@fda.hhs.gov>; Murphy, Jeanette <Jenny.Murphy@fda.hhs.gov>; Burkholder, William <William.Burkholder@fda.hhs.gov>; Norris, Anne <Anne.Norris@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Debbie,

Dr. Bill Burkholder is willing to be on the panel in St. Louis.

I am willing to be on the webinar in March as the lead for CVM. We have a wonderful team, so would we be able to include them similar to the last webinar?

Thanks again!

Martine

From: Debbie Phillips <DPhillips@wattglobal.com>
Date: November 30, 2018 at 9:35:42 AM EST
To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Subject: RE: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Martine,

That's great, thank you! And don't worry about the timing; I know it's a very busy time of year.

Not to push my luck, but do you think someone from CVM could also participate in a webinar in March on this? It would be a follow-up/update to September's webinar and seek to inform those in the industry who cannot attend Petfood Forum. We don't know the exact timing in March yet, other than it probably wouldn't be the week of March 18, as there is a major pet show that week.

Thanks again, please let me know if I can answer any questions.

Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: +1 (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13

Petfood Forum China 2019: August 20

Petfood R&D Showcase: October 2019

Visit www.PetfoodForumEvents.com

From: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Sent: Thursday, November 29, 2018 8:17 PM
To: Debbie Phillips <DPhillips@wattglobal.com>
Cc: Norris, Anne <Anne.Norris@fda.hhs.gov>
Subject: Re: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Debbie!

My sincere apologies for not following up sooner. We did discuss internally and think it would be a wonderful opportunity for CVM.

I am working on finding the right person to represent CVM and will get back to you asap.

Thank you so much for your patience and sending my apologies again!

Martine

From: Debbie Phillips <DPhillips@wattglobal.com>
Date: November 29, 2018 at 4:23:54 PM EST
To: Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>
Cc: Norris, Anne <Anne.Norris@fda.hhs.gov>
Subject: FW: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Dr. Hartogensis and Anne:

I hope you both had nice Thanksgiving celebrations last week! And I'm following up on my message below to see if you have had an opportunity to consider this invitation?

Also, please note that we are tentatively planning another webinar about the investigation and situation, probably in March. That would provide a six-month update since the first webinar and also sort of tease the upcoming session at Petfood Forum.

Please let me know if you have any questions. Thank you again for your consideration.

Sincerely,
Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: +1 (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13

Petfood Forum China 2019: August 20

Petfood R&D Showcase: October 2019

Visit www.PetfoodForumEvents.com

From: Debbie Phillips
Sent: Friday, November 9, 2018 11:32 AM
To: Martine.Hartogensis@fda.hhs.gov
Cc: Anne.Norris@fda.hhs.gov
Subject: Invitation to participate in panel discussion on DCM investigation at Petfood Forum conference

Hi Dr. Hartogensis:

Thank you again for participating in our webinar in September on the atypical cases of canine DCM and their possible link to certain pet diets. This continues to be a topic of conversation and concern in the industry, as you probably are aware, so we are planning a follow-up/update during our annual conference, Petfood Forum, in late April/early May. Could you or one of your colleagues please consider serving as a panel member for this discussion?

The session is currently scheduled for the afternoon of Wednesday, May 1. I know that seems a long way off, but we prefer to issue invitations in advance, as opposed to the hasty invitation for the webinar!

Other panel members will likely include Dr. Greg Aldrich as moderator (who also participated in the webinar, as you know), plus Dr. Jennifer Adolphe, nutrition manager for Petcurean Pet Food; Dr. Kate Shoveller, assistant professor at the University of Guelph; and Dr. Chris Marinangeli, director of nutrition, scientific and regulatory affairs for Pulse Canada.

In case you are not familiar with Petfood Forum, we just held our 26th edition this past April. It is the only event of its kind for the global pet food industry, drawing more than 3,000 people each year from pet food companies around the world, plus from retailers and related businesses, academia and regulatory organizations, such as AAFCO. In addition to education (concurrent scientific tracks plus panel discussions, general sessions, keynotes and other), it includes a trade show featuring the industry's leading suppliers of ingredients, equipment, packaging materials, testing and other services. This year's show had over 400 booths with more than 250 exhibiting companies.

We offer an honorarium to our speakers and panel members, cover their hotel costs and conference registration and reimburse all other travel expenses.

Please let me know if you have questions about Petfood Forum or this panel discussion. Thank you in advance for considering the request!

Sincerely
Debbie

DEBBIE PHILLIPS-DONALDSON
Editor-in-Chief, Petfood Industry/Petfood Forum
www.PetfoodIndustry.com
Watt Global Media
Office: +1.815.966.5424
Mobile: + (b) (6)
dphillips@wattglobal.com
Skype: wattdebbie.phillips
www.gotomeet.me/DebbiePhillipsDonaldson

Check out these upcoming Petfood Forum conferences!

Petfood Forum and Petfood Innovation Workshop 2019: April 29-May 1

Petfood Forum Europe 2019: June 13

Petfood Forum China 2019: August 20

Petfood R&D Showcase: October 2019

Visit www.PetfoodForumEvents.com

The association between pulse ingredients and canine dilated cardiomyopathy: addressing the knowledge gaps before establishing causation

Wilfredo D. Mansilla^{*}, Christopher P.F. Marinangeli[†], Kari J. Ekenstedt[‡], Jennifer A. Larsen[#], Greg Aldrich[°], Daniel A. Columbus[§], Lynn Weber[¶], Sarah K. Abood[‡], Anna K. Shoveller^{*†}

^{*}Department of Animal Biosciences, University of Guelph, 50 Stone Road East, Guelph, Ontario, Canada, N1G 2W1

[†]Pulse Canada, 920-220 Portage Avenue, Winnipeg, Manitoba, Canada, R3C 0A5

[‡]Department of Basic Medical Sciences, College of Veterinary Medicine, Purdue University, 625 Harrison Street, West Lafayette, Indiana, USA, 47907

[#]Department of Molecular Biosciences, School of Veterinary Medicine, University of California, Davis, 1 Shields Ave, Davis, California, USA, 95616

[°]Department of Grain Science and Industry, Kansas State University, Manhattan, Kansas, USA, 66506

[§]Prairie Swine Centre, 2105 – 8th Street East, Saskatoon, Saskatchewan, Canada, S7H 5N9

[¶]Department of Veterinary Biomedical Sciences, University of Saskatchewan, 52 Campus Drive, Saskatoon, SK, Canada, S7N 5B4

[‡]Dept. of Clinical Studies, University of Guelph, 50 Stone Road East, Guelph, Ontario, Canada, N1G 2W1

[†]**Corresponding author:** Anna K. Shoveller; ashovell@uoguelph.ca Current address: 50 Stone Road East, Department of Animal Biosciences, University of Guelph, Guelph, Ontario, N1G 2W1, Canada.

Author disclosure: Funding for this project was provided by Pulse Canada. C.P.F.M. works for Pulse Canada and is a former employee of Kellogg Canada. W.D.M., A.K.S., K.J.E., G.A., J.A.L., D.A.C., L.W., and S.K.A. have no conflicts of interest. All authors contributed to the content of this paper.

Acknowledgments: The authors would like to acknowledge the contribution of James Templeman, Sarah Dodd, and Emma Thornton.

© The Author(s) 2019. Published by Oxford University Press on behalf of the American Society of Animal Science.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

ABSTRACT

In July 2018, the Food and Drug Administration (FDA) warned about a possible relationship between dilated cardiomyopathy (DCM) in dogs and the consumption of dog food formulated with potatoes and pulse ingredients. This issue may impede utilization of pulse ingredients in dog food or consideration of alternative proteins. Pulse ingredients have been used in the pet food industry for over 2 decades and represent a valuable source of protein to compliment animal-based ingredients. Moreover, individual ingredients used in commercial foods do not represent the final nutrient concentration of the complete diet. Thus, nutritionists formulating dog food must balance complementary ingredients to fulfill the animal's nutrient needs in the final diet. There are multiple factors that should be considered, including differences in nutrient digestibility and overall bioavailability, the fermentability and quantity of fiber, and interactions among food constituents that can increase the risk of DCM development. Taurine is a dispensable amino acid that has been linked to DCM in dogs. As such, adequate supply of taurine and/or precursors for taurine synthesis play an important role in preventing DCM. However, requirements of amino acids in dogs are not well investigated and are presented in total dietary content basis which does not account for bioavailability or digestibility. Similarly, any nutrient (e.g. soluble and fermentable fiber) or physiological condition (e.g. size of the dog, sex, age) that increases the requirement for taurine will also augment the possibility for DCM development. Dog food formulators should have a deep knowledge of processing methodologies and nutrient interactions beyond meeting AAFCO nutrient profiles and should not carelessly follow unsubstantiated market trends. Vegetable ingredients, including pulses, are nutritious and can be used in combination with complementary ingredients to meet the nutritional needs of the dog.

Key words: dilated cardiomyopathy, dogs, feed formulation, grain-free, nutrition, pulse ingredients

INTRODUCTION

In July 2018, the Food and Drug Administration (FDA) issued a statement relating dilated-cardiomyopathy (DCM) in dogs to the consumption of foods that have potatoes and/or pulse ingredients, such as peas and lentils or their co-products, as main ingredients (FDA, 2018). The FDA's statement, as well as media attention, has raised concern in some pet owners, veterinarians, nutritionists, and the pet food manufacturing and retail industry. The underlying cause for concern with pet food and DCM is that there is a link between nutrition that was previously tied to DCM and insufficient circulating taurine (Fascetti et al., 2003; Backus et al., 2006). The result, was an increased need for dietary taurine or its precursor methionine due to higher fermentation of taurine and greater fecal excretion with dietary fermentable fiber (Kim et al., 1996ab). Whether this has any link to dietary pulses or the greater inclusion of pulses in grain-free dog food has yet to be directly demonstrated and mechanistic research is warranted.

Pulses are a subset of legumes, harvested as a dry crop, with low concentrations of lipid. They include peas, lentils, chickpeas, and dry beans (Marinangeli et al. 2017) which have been used as ingredients in dog food for their protein and fiber for more than 2 decades (Butterwick et al., 1994; Rice and Ihle, 1994). As a source of protein, the amino acid (AA) profile in peas, lentils, chickpeas, and beans are generally high in lysine and low in methionine (NRC, 2006) and serve as a complementary protein to both animal and plant-derived ingredients. As an example, soybean meal is derived from defatted soybeans and has an amino acid profile similar to pulses. In a 24-week study that evaluated graded concentrations of soybean meal up to 17 % (as-fed basis) in dog foods, soybean meal inclusion did not affect the nutrient status of dogs as indicated by serum biochemistry analysis (Menniti et al., 2014). However, Yamka et al. (2003) demonstrated that using soybean meal at more than 15 % inclusion on a dry matter basis decreased crude protein digestibility. Based on the authors assessment of current formulas in the market, there is a high likelihood that legume seed use in some foods may be greater than 40 %. This inclusion exceeds concentration of legumes previously investigated in dogs. When used to complement the nutritional profile of other ingredients, pulses can be used as nutrient-rich vehicles to meet the nutritional requirements of dogs and other companion animals. Given that companion animals most often consume static diets for long periods of time, overuse of any ingredient could facilitate higher risk of certain nutrient deficiencies if nutrient balance is not considered in the formulation. Thus, the formulation of static diets that use significant concentrations of a single ingredient, relative to other ingredients in the formulation, requires an in-depth knowledge of nutrient interactions, animal physiology, and effects of processing, beyond that of simply meeting minimum nutrient profiles stipulated in the Official Publication of The Association of American Feed Control Officials (AAFCO, 2018).

The present commentary discusses: 1. The limited data being used to support linkages between DCM and pulse ingredients; 2. The nutritional factors and physiological mechanisms that should

be explored to establish causation between nutritional deficiencies and incidence of DCM; 3. The factors that nutritionists should consider when formulating complete diets destined for long term consumption; and 4. The disadvantages of formulating to protein and minimal AA recommendations rather than to a balanced indispensable AA profile.

The development of canine DCM, historical linkages to taurine deficiency and pulses

Dilated cardiomyopathy is a disease of the myocardium that results in both mechanical dysfunction (enlarged heart cavities and congestion) and/or electrical dysfunction (arrhythmias and sudden death) (Sisson et al., 2000; Maron et al., 2006; Dutton and Alvarez, 2018). Development of DCM is slow and few clinical signs manifest over time. As DCM progresses, signs include lethargy, anorexia, shallow breathing, sudden fainting, and potential death. In some cases, animals may die from irregular heart rhythm without previous signs of the disease. In dogs, DCM can be caused by various factors. Genetic predisposition is thought to play the most important role in the development of DCM in several dog breeds, mostly large and giant breeds. Genetic mutations associated with DCM have been discovered in American lines of Doberman and Boxer dogs (Meurs et al., 2012; Meurs et al., 2013). However, the Doberman variant's association was not upheld in a European population of Dobermans (Owczarek-Lipska et al., 2013). Similarly, a UK population of Boxers did not uphold their published DCM-associated variant (Cattanach et al., 2015). It is becoming increasingly clear that the genetic basis for DCM in dogs is not monogenic, but complex and polygenic. Breeds with the highest prevalence of DCM include Dobermans, Boxers, Great Danes, Newfoundlands, Irish Wolfhounds, English Cocker Spaniels, and Portuguese Water Dogs (Monnet et al., 1995; Borgarelli et al., 2006; Werner et al., 2008; Martin et al., 2009), and the genetic basis of DCM in each of these breeds has been investigated (Dutton and Alvarez, 2018). In addition, Golden Retrievers and American Cocker Spaniels appear to have breed predispositions to taurine deficiency (Kramer et al., 1995; Bélanger et al., 2005). When dogs are not genetically predisposed for developing DCM, diet and physiology are other factors that may be associated with the disease.

The first link between taurine deficiency and DCM was demonstrated in cats in 1987. Cats diagnosed with DCM recovered after taurine supplementation (Pion et al., 1987). Similarly, an inverse association between dietary taurine and the incidence of DCM in a population of foxes was documented by Moise et al. (1991) and established the importance of taurine in the family Canidae. In dogs, DCM diagnoses related to low whole blood taurine concentrations have been reported in Cocker Spaniels, Dalmatians, Boxers, Newfoundlands, Portuguese Water Dogs, English Setters, Alaskan Malamutes, and Scottish Terriers (Freeman et al., 1996; Kittleson et al., 1997; Pion et al., 1998; Alroy et al., 2000; Fascetti et al., 2003; Backus et al., 2006). In all these cases, taurine supplementation improved cardiac function. However, dogs, in contrast to cats, can endogenously synthesize taurine from methionine and cysteine (Figure 1). Therefore, the

abovementioned data does not unequivocally establish taurine intake as the underlying mechanism for the development of DCM in dogs, whether or not they are genetically predisposed. Dietary supply of precursor AAs necessary for taurine synthesis (i.e. methionine and cysteine), metabolic intermediates, and co-factors (such as methyl donors) cannot be ruled out as factors that contribute to the susceptibility of dogs to developing genetic and diet-related DCM. When DCM is diet-related, the formulation and the provision of all nutrients, including indispensable AAs, to facilitate optimum health and wellbeing of dogs should be considered.

Recent reports, including the statement by the FDA (2018), have implicated that lentils, peas and other legumes seeds could be responsible for the development of DCM in dogs not genetically predisposed to this disease. Such statements and associations between pulse ingredients and incidence of DCM are, at the present time, premature. Animals, including dogs, have no minimum or maximum requirements for ingredients. Ingredients serve as the vehicle to providing nutrients to animals. As such, animals have nutrient requirements, not ingredient requirements. In diets that have nutrient deficits, imbalances, or exceed maximums, the final nutrient composition of the diet, not the ingredients, should be critiqued. In addition, animal nutritionists should consider that the nutrient concentration of ingredients can vary, nutrient availability is not 100 %, and diets formulated to marginally meet requirements could actually be deficient. Overall, it is the responsibility of nutritionists to use different ingredients to formulate diets that can be produced and safely meet the nutritional needs of animals.

Taurine deficiency and the development of canine DCM

For dogs, taurine is a dispensable AA synthesized from methionine and cysteine primarily in the liver (Figure 1). Taurine is not incorporated into proteins. Instead, it is used as a mediator for various biological processes and is the most abundant free AA intracellularly (Huxtable, 1992). In the heart, taurine represents ~60% of the total AA free pool (Huxtable, 1992). The high concentration of taurine in cardiac cells may explain the role of a taurine deficiency in the development of DCM. It has been speculated that taurine contributes to the reabsorption of calcium by the sarcoplasmic reticulum and increases the sensitivity of the myofilaments to calcium (Bakker and Berg, 2002). Thus, low dietary taurine intake and/or reduced synthesis of taurine from methionine and cysteine can deplete calcium pools in the cardiac cells and impede proper contraction of the cardiac muscle tissue, resulting in DCM in dogs.

For diagnosing DCM in dogs and cats, among other diagnostic methods including electrocardiograms and echocardiography, it is common to measure taurine concentration in whole blood. Whole blood samples, and not plasma samples, should be used to assess circulating taurine concentrations. In plasma, free taurine concentrations are much lower compared to intracellular taurine. This suggests that the plasma pool is not representative of taurine in other

pools (Schaffer et al., 2010). In platelets, taurine concentration is high and is considered a marker of taurine status. Taurine concentration in platelets is captured when whole blood is analyzed (Huxtable, 1992). However, platelet count can vary depending on the immune status of the animal and whole blood taurine concentration can be affected. In this scenario, whole blood taurine may not represent concentrations of taurine in muscle cells, including cardiac muscle. These additional variables related to the measurement of taurine status may explain why some dogs diagnosed with DCM have normal whole blood taurine concentrations.

As taurine can be synthesized endogenously in dogs, taurine is not considered an indispensable AA for the species Canidae. Thus, there are no recommendations on minimum dietary concentrations of taurine for dogs reported by the National Research Council (NRC, 2006) or AAFCO (2018). The lack of regulation on minimum taurine concentrations in commercial dog foods suggests that endogenous synthesis of taurine can meet the metabolic needs in all dogs and at all life stages. This assumption may not be accurate as studies have determined that synthesis of taurine is related to the size of dog (Ko et al., 2007), and some dietary factors can increase the physiological need for taurine (Story, 1978). Nutritional factors that increase the dietary requirement, reduce the supply, or increase the excretion of taurine in dogs are discussed in subsequent sections of this review and should be considered to avoid taurine deficiency in dogs and the risk of DCM.

Physiological factors can increase taurine utilization in dogs, and endogenous synthesis of taurine could be insufficient for meeting taurine requirements. For example, compared to smaller size dogs, synthesis of taurine in large dog breeds is up to 50% lower per unit of metabolic body weight (Ko et al., 2007). These results demonstrate that larger dogs are at higher risk for insufficient endogenous taurine synthesis, and dietary supplementation or fortification may be required, even when there is no minimum dietary taurine concentration according to current recommendations (AAFCO, 2018). Obesity and diabetes have also been related to lower concentrations of taurine in blood in humans and rats, respectively, (Merheb et al., 2007; Nardelli et al., 2011; Ito et al., 2012) and may increase the requirement for sulfur AAs necessary for endogenous taurine synthesis. This is of importance given that approximately half of dogs in North America are obese (Linder and Mueller, 2014). Data from rats and cats suggests that age and sex could also affect whole body taurine status. Hepatic activity of cysteine sulfonate decarboxylase, the enzyme responsible for taurine synthesis, was shown to be 16× higher in adult male rats versus female rats. In the same study, the activity of cysteine sulfonate decarboxylase was higher in 5-6-week-old kittens compared to 15-month-old cats and in 8-week-old mice compared to 16-week old mice; changes of the enzyme activity in dogs have not been tested (Worden and Stipanuk, 1985). Overall, these studies suggest that, despite some capacity for endogenous synthesis, physiological need of taurine can be heavily dependent on breed, age, sex, and physiological status. These physiological factors could help to predict the risk for developing

DCM when genotypic and environmental factors, such as diet, are simultaneously considered to ensure dogs maintain adequate concentrations of taurine and other sulfur AAs.

Given that there are no recommendations for the minimum concentration of taurine in dog food, the concentration of taurine in dog foods can vary substantially depending on the ingredients used. Taurine is very low in plant-based ingredients (Table 1) but is higher in some algae and fungi species and is ubiquitously found in animal tissues, especially in the heart, brain, and white blood cells (Huxtable, 1992). This is relevant, as many grain-free and/or high legume dog foods attempt to limit the use of animal by-products, which can substantially decrease the levels of dietary taurine. In the context of providing adequate and preventive nutrition, dog foods should include organ meat or animal by-products or be fortified with taurine and/or its precursors (methionine and/or cysteine) to ensure the delivery of sufficient levels of taurine.

Effect of dietary fibre on taurine status and risk of canine DCM

Dietary fiber has been shown to affect the taurine status in dogs. For example, commercial diets formulated with lamb meal and rice bran were shown to cause taurine deficiency in part because of low bioavailable cysteine from lamb meal and possibly more importantly due to the effects of rice bran fiber on gastrointestinal metabolism of taurine (Johnson et al., 1998; Tôrres et al., 2003). It has been hypothesized that high fiber diets can increase susceptibility to taurine deficiency by 2 mechanisms of action linked to obligatory bile acid conjugation with taurine in dogs (O'Mádille et al., 1965) and reliance on enterohepatic circulation for the reabsorption of bile acids and taurine. First, high fiber diets may increase fecal output and losses of taurine-conjugated bile. This would require higher synthesis rates of bile in the liver, and consequently, higher utilization of taurine (Story, 1978). Second, high consumption of fermentable fibres may increase the abundance of microbial populations that degrade taurine in the intestinal lumen (Kim et al., 1996ab). Either alone or together, increased excretion or degradation of taurine from high fibre diets may decrease enterohepatic circulation and recycling of taurine. Given that taurine is the only AA used for bile acid conjugation in dogs, over time, high fiber diets could increase the risk of taurine insufficiency in dogs and lead to DCM.

This should not be interpreted as dietary fiber being deleterious to the health of dogs. However, there may be a limit to the benefit for soluble fibers. Legume seeds contain an appreciable quantity of oligosaccharides which are known to be fermentable (Tosh and Yada. 2010). Thus, by a similar mechanism as described above, high levels of legume seed oligosaccharides could ostensibly contribute to taurine depletion via excretion in the feces as bile conjugation and degradation by colonic bacteria. In addition to the physiological benefits of high fiber diets in certain dogs, formulators should also be cognizant of possible nutritional risks associated with

high concentrations of fiber in dog foods. Consequently, dog foods with high concentrations of dietary fiber should be accompanied with higher supplies of taurine or sulfur AAs for endogenous taurine synthesis. Overall, the digestibility and bioavailability of taurine in ingredients used and the effect of other nutrients in taurine metabolism should be considered to avoid taurine deficiency and the development of DCM.

Carnitine deficiency and risk of canine DCM

Carnitine is not nutritionally indispensable since it is endogenously produced in the liver and kidneys from lysine and methionine; it can also be attained exogenously from animal-based products. Carnitine is highly abundant in skeletal and cardiac muscles. Together, these represent > 95% of the total carnitine in the body. Carnitine is essential for metabolism of fatty acids used for energy production (Hoppel, 2003). In the heart, where 60% of the energy is derived from fatty acid oxidation, carnitine facilitates the uptake of free fatty acids into the mitochondria to produce ATP (Hoppel, 2003). Plant-based ingredients do not contain carnitine (Table 1). Therefore, in commercial dog foods with reduced inclusion of animal-based ingredients, intakes of carnitine could be decreased if diets are not fortified. Reduced dietary carnitine intake translates into increased reliance on endogenous synthesis to meet physiological requirements.

Given that carnitine is required for sufficient energy production in cardiac muscle, it is not surprising that carnitine deficiency is associated with DCM. In 1991, a family of Boxers diagnosed with DCM were also diagnosed with carnitine deficiency (Keene et al., 1991). In dogs, carnitine deficiency can occur with aberrations of carnitine regulation in disorders such as cardiomyopathy (including DCM), diabetes, sepsis, and malnutrition (Flanagan et al., 2010). However, carnitine deficiency as a causative factor in the development of DCM or a consequence of cardiac malfunction remains as a subject of debate (Freeman and Rush, 2006). Despite the interest in this metabolite, little progress has been made on determining the effect of carnitine supplementation on alleviating risk of DCM. However, both taurine and carnitine are often supplemented in supraphysiological concentrations once DCM is diagnosed. This practice is supported by positive clinical outcomes, albeit without comparison groups (Kittleson et al. 1997; Sanderson et al. 2001). Concentrations of carnitine in the plasma are relatively insensitive to dietary carnitine, and more invasive techniques (biopsies) are required to determine the concentration of carnitine in muscle tissue (Flanagan et al., 2010; Rășanu et al., 2012). The invasive nature of testing for carnitine status is likely the reason why carnitine is rarely explored when investigating possible causes of canine DCM.

Preventing diet-mediated DCM in dogs by providing adequate sulfur AAs and maximizing endogenous taurine synthesis

Although taurine is considered a dispensable AA in dogs, endogenous taurine synthesis requires an adequate supply of bioavailable sulfur AA precursors cysteine or methionine (Figure 1). Thus, providing marginal concentrations of these 2 sulfur AAs, or providing sources with lower bioavailability, could increase the risk of taurine deficiency and facilitate the development of DCM. Contrary to taurine, methionine cannot be synthesized endogenously in dogs (NRC, 2006). Therefore, dogs depend on the provision of dietary methionine to meet daily sulfur AA requirements, which includes production of taurine. From an ingredient perspective, methionine and lysine are usually the first or second limiting AAs in dog diets formulated with soybean meal and rendered meats (NRC, 2006). In addition, methionine is particularly susceptible to damage, and subsequent reduction in bioavailability, secondary to heat processing (Marshall et al. 1982; Hurrell et al. 1983). This suggests that the risk of methionine deficiency is more likely than any other indispensable AA in commercial dog diets. Although the primary role for methionine is protein synthesis, in pigs at least 50% of absorbed methionine acts as a methyl donor and a precursor in the production of cysteine, taurine, sulfate, and pyruvate (Robinson et al., 2016a) (Figure 1). These functions of methionine become more crucial when dietary intake of cysteine, taurine, and/or dietary methyl donors (e.g. folate, betaine, and their precursors) is limited (Robinson et al., 2016b), and they need to be considered when nutritionists set criteria for delivery of sulfur AAs in pet foods.

Methionine and cysteine both contribute to the total sulfur AA requirements for humans and animals. For adult dogs at maintenance, the latest guidelines from the NRC (2006) recommend that adult dog foods contain 0.33% (on dry matter basis) methionine when cysteine is provided in excess, and 0.65% for methionine + cysteine. These NRC (2006) recommendations are not based on dose-response studies, but on a 4-year study where adult dogs were fed low-crude protein diets (Sanderson et al., 2001). In that study, the lowest concentration of methionine in the diet that reported no observable deficiencies was used as the recommended requirement. As companion animals are typically fed a single static diet during adulthood, and for most of their lifespan, it is necessary that AA requirements of dogs should be measured empirically (Baker, 1986). In addition to the lack of empirical data corresponding to the AA requirements of dogs, it is equally important to understand how other dietary (e.g. dietary fiber), environmental, other physiological variables, and breed/genotype may alter AA requirements. The lack of recommendations for taurine in commercial dog food puts a higher stress on accurately meeting requirements for sulfur AAs, not only for protein synthesis, but also for the endogenous synthesis of taurine, for support of optimal methyl status, and for the synthesis of secondary metabolites.

Rethinking indispensable AA targets in commercial dog foods

Currently, the ingredients permitted in pet foods and the corresponding nutrient targets are guided by recommendations made by AAFCO (2018). These recommendations are based on the

peer-reviewed scientific literature and represented in the Nutrient Requirement of Dogs and Cats (NRC, 2006). However, AA recommendations made by AAFCO correspond to total AA content within the formulation and do not consider the true ileal digestibility of ingredients. True ileal digestibility of AAs is more representative of nutrient absorption capacity and bioavailability compared to fecal digestibility or total AA content in the diet (Columbus and de Lange, 2012). To account for the reduced digestibility and bioavailability of protein-bound AAs in food ingredients, AAFCO arbitrarily increases AA recommendations relative to those from the NRC to ensure that an adequate supply of AAs is provided, regardless of the ingredients and effects of processing (Table 2). However, this increment is only applied to lysine, threonine, and tryptophan and not applied to other indispensable AAs, including methionine (AAFCO, 2018). For example, the recommended allowance for lysine reported in NRC (2006) is 0.35% for adult dogs at maintenance, while the minimum content of lysine to meet AAFCO (2018) recommendations is 0.63%. Non-ruminant animals, including dogs, absorb AAs from the duodenum to the terminal ileum (Columbus and de Lange, 2012). Hence, feeding diets with lower ileal digestibility coefficients could decrease actual concentrations of available indispensable AAs, even when meeting AAFCO recommendations. This is of special concern for dietary taurine and other sulfur AAs, considering that there is no regulated minimum threshold for taurine in dog foods and that AAFCO (2018) recommendations for sulfur AAs are not increased compared to NRC (2006) recommendations to account for potential ileal digestibility coefficients. There is a dearth of data in this area to justify empirical adjustments based on different dietary variables. As such, future research should pursue how amino acid requirements change under different dietary variables that can affect small intestinal digestibility and whole body availability.

It is worthwhile to note that minimum dietary nutrient contents for dog foods, as reported in AAFCO (2018), only considers differences between growth/reproduction and adult life stages. This lack of data places the pregnant bitch in the same group as growing animals. Moreover, most studies on nutrient requirements in dogs have been established using Beagles as a proxy for all dogs. Using a single breed creates a homogenous sample and likely does not account for nutritional variability across pure and mixed breeds, or those of different sizes. Unpublished data from Shoveller et al. investigated the minimum methionine (with excess cysteine) requirements of Miniature Dachshunds, Beagles, and Labrador Retrievers as proxies for small, medium, and large dog breeds and found that methionine requirements may differ across breeds or size of dogs and be greater than previously estimated. Thus, given the methods of derivation, single indispensable AA requirements for all dog populations, as presented in AAFCO (2018), may not consider variable AA requirements across dog phenotypes. Moreover, it is widely assumed that endogenous synthesis of dispensable AAs, such as taurine in the dog, is sufficient for meeting metabolic demands. However, recent studies suggest that under some metabolic conditions, dispensable AAs may also be required in diets (Hou et al., 2015). Taurine, as described in this commentary, is a clear example of this paradigm shift. Dietary taurine or the capacity for its

adequate endogenous synthesis, especially in circumstances where excessive losses might occur, should be considered in the final formulation of dog foods to decrease the risk of canine DCM.

Nutritionists and regulatory agencies should be aware that, in the spectrum of nutrient requirements, dog populations with higher AA requirements relative to energy intake and other factors could be at a higher risk for a taurine deficiency. More precise categorization of requirements among different canine populations would help to optimize nutritional adequacy and decrease risk of diseases, such as DCM, that are possibly linked to nutrient deficiencies.

Effect of processing on anti-nutritional factors in plant-based ingredients.

Just as understanding the inherent nutritional characteristics and the interaction between ingredients is important for preventing nutritional imbalances in pet foods, the effects of processing on these factors are equally important. Raw cereals and legumes contain anti-nutritional factors such as trypsin inhibitors, phytates, hemagglutinins, and polyphenols that can decrease protein digestion, nutrient absorption, and/or cause illness. Some of these anti-nutritional factors are thermolabile and, under the right conditions, can be effectively destroyed during the extrusion process improving the overall quality of plant-based ingredients and the final diet (Patterson, et al., 2017). Recent reviews across a variety of legumes and legume-derived ingredients show that the activities of trypsin inhibitor, chymotrypsin inhibitor, and hemagglutinating activity were decreased by up to 95 % across a variety of thermal treatment conditions, including extrusion (Patterson, et al., 2017; Aviles-Gaxiola et al. 2018). Extrusion had modest effects on levels of phytate with reductions ranging from 7 to 26 % and varied by legume and extrusion conditions (Patterson, et al., 2017). **Figure 2** highlights the variability between processing methods and thermic conditions for decreasing anti-nutritional factors. For example, when soybeans were subjected to extrusion at increasing temperatures that ranged from 100 to 150 °C, trypsin inhibitor levels were incrementally decreased. At 140 °C, dry extrusion was considerably more effective at decreasing trypsin inhibitors (-91 %) compared to wet extrusion (-44 %). When the dry extrusion temperature was increased to 150 °C, reductions in trypsin inhibitors were further decreased by 94 % (Zilic et al., 2012). Other thermal treatments, such as micronisation, microwave roasting, and autoclaving also facilitated incremental reductions in trypsin inhibitors with increasing temperatures (Zilic et al., 2012). When formulating foods with higher concentrations of plant-based ingredients, consideration should also be given to the processing methods and the parameters used to effectively optimize the nutritional density and decrease anti-nutritional factors.

It is important to mention that, while temperature and pressure processing can greatly decrease anti-nutritional factors, they can also negatively impact bioavailability of amino acids. The Maillard reaction is a well-known example of heat damaged-protein (Teodorowicz et al., 2017).

In this reaction, lysine interacts with reducing sugars present in the diets forming the Maillard product. The complex formed can be digested and absorbed by the animal but cannot be utilized for metabolic processes (e.g. protein synthesis). Thus, in heat damaged proteins, digestibility of amino acids can greatly overestimate bioavailability (Moehn et al., 2005). Other products of heat damage on proteins include racemization of amino acids (alteration from L to D form) and the formation of cross-linked amino acids. Such components can decrease bioavailability of amino acids and digestibility of proteins, and their effects on protein quality cannot usually be determined using conventional methods of amino acid analysis. Pet foods with higher levels of plant-based ingredients may also require optimization of processing methods to maximize their nutritional density and nutrient bioavailability.

Recommendations for formulating dog food with novel ingredients

Considering the AA profile of dog foods

Feed formulation for agricultural and companion animals should be based on the ideal protein concept (Baker, 1991; Swanson et al., 2013). The ideal protein is defined as that in which all AAs are in perfect balance compared to the animal's AA requirements (mg/g protein). Hence, all indispensable AAs are equally limiting. However, this is impossible to achieve in practical animal feed formulation, and diets should be formulated considering the first limiting indispensable AA. The first limiting indispensable AA refers to the indispensable AA that is present in the lowest proportion compared to the animal's requirement. By meeting the first indispensable limiting AA requirement, requirements for all other indispensable AAs are also inherently satisfied. Moreover, to avoid the formulation of diets with excessive protein concentration or an excess of indispensable AAs relative to the requirements of dogs, animal nutritionists combine multiple ingredients that are complementary in their AA profiles. Commonly, dog foods are formulated with a higher proportion of animal-derived ingredients, and a lower proportion of plant-based ingredients to meet nutrient recommendations. More recently, however, cereal grains have been removed in some diet formulations or the proportion of animal-based ingredients has been reduced. The production of these types of formulations are often driven by consumer perception, rather than scientific evidence. Allowing consumers to direct the ingredient composition of dog foods, or other pet foods, could perpetuate nutrient deficits that affect the health of animals in the long term.

In the formulation of grain-free pet foods, cereal grains are replaced with alternative ingredient(s). Animal-derived ingredients are expensive relative to plant-based ingredients. Thus, pulses, a subset of legumes, are often used as the replacement. In addition to containing substantial fiber, pulses also contain significant concentrations of protein and are used to partly

meet indispensable AA requirements. Of interest, soybean meal and pulses contain 48% and 25% crude protein, respectively, which is substantially greater than the average protein concentration for grains (11%) (Table 1). While the high protein content in soybean meal and pulses is indicative of higher concentration of AAs compared to grains, it does not imply AA balance. Soybean meal and pulses are high in lysine (mg/g protein) but low in sulfur AAs (mg/g protein), while the reverse is true for cereals. Plant-based ingredients tend to have lower ileal digestibility coefficients for protein compared to protein from animal sources (FAO and WHO 1991). Thus, dog foods that contain substantial amounts of pulses, lower proportions of animal-based ingredients, and do not address AA imbalances through the addition of alternate ingredients or fortification, may risk AA deficiencies. To mitigate this risk across the pet food industry and ensure the final pet diets are nutritionally adequate and balanced, it is prudent that the digestibility coefficients of all final pet food products be calculated.

Considering the addition of high fiber ingredients to dog foods

By definition, dietary fiber is carbohydrates that are resistant to digestion by endogenous enzymes in the gastrointestinal tract (NRC, 2006). Typical fibers include arabinoxylan, raffinose, inulin, β -glucan, cellulose, and pectin (NRC, 2006). Common ingredients to increase fiber content in companion animal diets include beet pulp, corn fiber, rice bran, whole grains, and pulse fibers (de Godoy et al., 2013). Achieving an optimal fiber concentration in canine diets has diverse positive physiological effects in the gastrointestinal tract; for example, higher fermentable fiber intake has been shown to slow the transit time of digesta, increasing satiety of the animal (Haber et al., 1977). Moreover, high fiber diets generally have lower energy density making them an important nutritional strategy for controlling body weight (Johnson et al., 2008) and reducing the incidence of diarrhea (Homan et al., 1994). Gut health is also improved with higher consumption of fiber; fermentable fiber can act as a prebiotic and increase the population of health-promoting microbiota including lactobacilli and bifidobacteria (Roberfroid, 2005). Although not required by AAFCO to fulfill the criteria of “complete and balanced”, fiber is an important component of the diet, and depending on the type of fiber and the amount consumed, fiber can increase the gut health status. Adding the necessary amount and type of fiber in the diet is crucial for optimal dog nutrition.

Despite the benefits of fiber in the diet, fiber can also affect enterohepatic recycling of taurine (discussed above). In monogastric species, including humans, high dietary fermentable fiber may also decrease digestibility and availability of dietary AAs (Blackburn and Southgate, 1981; Degen et al., 2007) and, in some cases, increase the risk of DCM in dogs fed diets that marginally meet requirements for sulfur AAs. Moreover, higher concentrations of dietary fiber increase the size of the gastrointestinal tract in pigs and poultry (Nyachoti et al., 2000) increasing nutrient utilization in this organ. It has been determined in pigs that on average the gastrointestinal tract catabolizes 30% of dietary indispensable AAs during absorption, and this utilization represents ~50% for sulfur AAs (Stoll et al., 1998; Mansilla et al., 2018), further

reducing precursor availability for taurine synthesis and increasing the risk for taurine deficiency. For some high fiber diets, fortification of specific nutrients, including taurine and other sulfur AAs, might be beneficial to avoid nutrient deficiencies.

Compared to the pet food industry, in other industries where high fiber ingredients (co-products) are routinely used (e.g. swine industry), the effects of fiber on the absorption of nutrients have been given more attention when formulating diets (NRC, 2012). For example, highly fermentable fiber in swine diets increases the threonine requirement to compensate for the increase in mucus (mucin protein) production in the intestinal cell lining (Lien et al., 1997; Mathai et al., 2016). This has underpinned the development of “requirement models” (NRC, 2012) to tailor nutrient requirements for pigs while accounting for the different nutrient interactions. In contrast, in the pet food industry, the only concentrations of nutrients used for comparison are those recommended by AAFCO (2018). Such recommendations are static and may not encompass all the effects of the different nutrient combinations in the final diet. There is a clear need in companion animal nutrition to improve the understanding of the interactions of different ingredients and how these alter nutrient requirements for different breeds, age, and physiological status of dogs.

Other recent publications highlight the need for careful nutrient formulation

Several recent papers, both original research and reviews, likewise highlight the unknowns surrounding grain-free diets (typically legume or pulse-based, but sometimes also with “exotic” ingredients such as kangaroo, bison, or wild boar) and DCM. For example, Adin et al. (2019) examined 48 dogs of many breeds with diagnosed DCM and having a known diet history. Among grain-free diets being consumed in this study, 1 was particularly associated with DCM, possibly underscoring the importance of specific diet formulation. Further, 2 dogs switched from that diet to other grain-free diets showed improvement in their DCM; it is unclear if those dogs were taurine deficient or if they also received taurine and/or carnitine supplementation. This suggests that grain-free composition per se may not be the root cause of DCM. Another recently published case series of 24 Golden Retrievers with DCM and known diet histories were evaluated, and an association between grain-free diets and DCM was suggested (Kaplan et al., 2018). Most dogs (15 out of 24) were fed a single diet which was significantly associated with low blood taurine concentrations, again suggesting that specific diet formulation may play an important role. However, as in the previous study, soluble versus insoluble fiber concentrations were not available for the diets, nor were taurine, methionine, or cysteine concentrations, meaning that the true nutrient profiles of the diets could not be assessed and reinforcing the point that diet formulation for nutrients – not ingredients – is essential. It also suggests that nutrient requirements may vary widely based on breed, diet, and other phenotypic data. Indeed, most of the dogs with DCM in the previously described study were consuming less energy compared to their predicted requirements (Kaplan et al., 2018). It also bears pointing out that the numbers in both studies were very low (representing less than 100 DCM-affected dogs between them),

which surely represents a fraction of the dogs consuming grain-free, pulse-based diets. A recent thoughtful review supports these conclusions by reiterating the crucial need for plant-based diets for dogs to be formulated with sufficient quantities of bioavailable methionine and cysteine to support adequate taurine synthesis (Dodd et al., 2018). This can be achieved with the addition of purified amino acids and other sources that are readily available (Gloaguen et al., 2014). Finally, a recent commentary carefully concludes that a true cause-and-effect relationship between grain-free diets and DCM has not been proven, and other factors may ultimately be more important (Freeman et al., 2018). Taken together, these recent publications may point to faulty nutrient formulation in some, but not all, grain-free diets.

CONCLUSIONS

Recently, it has been suggested that pulse ingredients in commercial dog foods are associated with a limited number of cases of DCM. While pulse ingredients have been implicated for having negative effects on the taurine status in dogs (deficiency of which is a known cause of canine DCM) based on the available evidence, the relationship between pulses and canine DCM remains undefined. However, the FDA statement may harm consideration of protein alternatives, such as pulses, as quality ingredients in pet foods and undermine attempts to diversify ingredients used across the food chain as the global population continues to grow. Ingredients do not represent the nutritional composition of the diet, and therefore, nutrient deficiencies should not be attributed to individual ingredients. The authors of this commentary recognize the important role of endogenous, and perhaps exogenous, taurine in the prevention of DCM in some dogs. The assurance of appropriate concentrations of all indispensable sulfur AAs, including methionine and cysteine, is crucial for ensuring adequate endogenous synthesis of taurine and to meet the metabolic demands of dogs. Additional dietary factors, such as methyl donors required for sulfur AA metabolism, carnitine for energy production in muscle, and dietary fiber, as well as animal factors, such as breed, size, and health status, should also be investigated when nutrient deficiency-related DCM is suspected.

It is the responsibility of animal nutritionists to formulate balanced diets for dogs, and other animals, by looking beyond the goal of meeting AAFCO recommendations or satisfying unsubstantiated market trends. Pulses and other plant-based ingredients can be used to formulate nutritionally adequate dog foods, and final product formulations should be assessed for nutrient balance and bioavailability, especially when using a limited number of ingredients. Although dietary factors are important in the prevention of sulfur AA deficiency and development of DCM, empirical data and mechanistic studies are required to better understand the indispensable AA requirements of dogs and preventing DCM. In diets that contain high concentrations of dietary fiber, compensative inclusion of dietary indispensable sulfur AAs, including exogenous taurine, might be required to offset the possibility of increased fecal excretion or microbial

assimilation of taurine in the large intestine. Processing conditions may also require adjustments to ensure the presence or effects of anti-nutritional factors are minimized and nutrient bioavailability is not compromised. Greater awareness of AA balance is crucial for ensuring that AA requirements are met for dogs consuming static diets.

Accepted Manuscript

REFERENCES

- AAFCO. 2018. Association of American Feed Control Officials. Oxford, In: Official Publication Association of American Feed Control Inc.
- Adin, D., T. C. DeFrancesco, B. Keene, S. Tou, K. Meurs, C. Atkins, B. Aona K. Kurtz, L. Barron K. Saker. 2019. Echocardiographic phenotype of canine dilated cardiomyopathy differs based on diet type. *J. Vet. Card.* 21:1-9.
- Alroy, J., J. E. Rush, L. Freeman, M. S. Amarendhra Kumar, A. Karuri, K. Chase, S. Sarkar. 2000. Inherited infantile dilated cardiomyopathy in dogs: genetic, clinical, biochemical, and morphologic findings. *Am J Med Genet* 95:57-66.
- Arslan, C. 2006. L-Carnitine and its use as a feed additive in poultry feeding a review. *Revue Med Vet.* 157:134-142.
- attributes, and applications. *Food Research International* 43 (2010) 450–460
- Avilés-Gaxiola, S., C. Chuck-Hernández, S. O. Serna Saldívar. 2018. Inactivation Methods of Trypsin Inhibitor in Legumes: A Review. *J. Food Sci.* 83:17-29. doi: doi:10.1111/1750-3841.13985.
- Backus, R.C., K. S. Ko, A. J. Fascetti, M. D. Kittleson, K. A. MacDonald, D. J. Maggs, Q. R. Rogers. 2006. Low plasma taurine concentration in Newfoundland dogs is associated with low plasma methionine and cyst(e)ine concentrations and low taurine synthesis. *J Nutr.* 136:2525-2533. doi: 10.1093/jn/136.10.2525
- Baker, D. H. 1986. Problems and pitfalls in animal experiments designed to establish dietary requirements for essential nutrients. *J. Nutr.* 116:2339-2349. doi: 10.1093/jn/116.12.2339
- Baker, D. H. 1991. Review Comparative nutrition of cats and dogs. *Annu Rev Nutr.* 11:239-263. DOI: 10.1146/annurev.nu.11.070191.001323
- Bakker, A. J., H. M. Berg. 2002. Effect of taurine on sarcoplasmic reticulum function and force in skinned fast-twitch skeletal muscle fibres of the rat. *J Physiol.* 538:185–194. doi: 10.1113/jphysiol.2001.012872
- Bélanger, M.C., M. Ouellet, G. Queney, and M. Moreau. 2005. Taurine-deficient dilated cardiomyopathy in a family of golden retrievers. *J. Am. Anim. Hosp. Assoc.* 41(5):284-291.
- Blackburn, N.A., Southgate DAT. 1981. Protein digestibility and absorption: effects of fibre and the extent of individual variation. Joint FAO/WHO/UNU Expert Consultation on Energy and Protein Requirements Rome, 5 to 17 October.

- Borgarelli, M., R. A. Santilli, D. Chiavegato, G. D'Agnolo, R. Zanatta, A. Mannelli, A. Tarducci. 2006. Prognostic indicators for dogs with dilated cardiomyopathy. *J. Vet. Intern. Med.* 20:104-110. <https://doi.org/10.1111/j.1939-1676.2006.tb02829.x>
- Butterwick, R. F., P. J. Markwell, C. J. Thorne. 1994. Effect of level and source of dietary fiber on food intake in the dog. *J. Nutr.* 124:2695S-2700S. https://doi.org/10.1093/jn/124.suppl_12.2695S
- Cattanach, B. M., J. Dukes-McEwan, P. R. Wotton, H. M. Stephenson, R. M. Hamilton. 2015. A pedigree-based genetic appraisal of Boxer ARVC and the role of the Striatin mutation. *Vet. Rec.* 176:492. doi: 10.1136/vr.102821.
- Chemical Hazards Emergency Medical Management. Key Principles of Toxicology and Exposure Washington D.C.: U.S. Department of Health and Human Services,; 2017. Available from: <https://chemm.nlm.nih.gov/toxprinciples.htm>.
- Columbus, D., C. F. M. de Lange. 2012. Evidence for validity of ileal digestibility coefficients in monogastrics. *Br. J. Nutr.* 108:S264-S272. doi: 10.1017/S0007114512002334.
- de Godoy, M. R., K. R. Kerr, J. C. Fahey. 2013. Alternative dietary fiber sources in companion animal nutrition. *Nutrients* 5:3099-3117. doi: 10.3390/nu5083099.
- Degen, L., V. Halas, L. Babinszky. 2007. Effect of dietary fibre on protein and fat digestibility and its consequences on diet formulation for growing and fattening pigs: A review. *Act. Agr. Scand. A-AN.* 57:1-9. <https://doi.org/10.1080/09064700701372038>
- Dodd, S. A. S., J. L. Adolphe, A. Verbrugghe. 2018. Plant-based diets for dogs. *J. Am. Vet. Med. Assoc.* 253:1425-1432. <https://doi.org/10.2460/javma.253.11.1425>
- Dutton, E., J. López-Alvarez. 2018. An update on canine cardiomyopathies – is it all in the genes? *J. Small. Anim. Pract.* 59:455-464. <https://doi.org/10.1111/jsap.12841>
- FAO. 1991. Food and Agriculture Organization of the United Nations. Protein Quality Evaluation. Report of Joint FAO/WHO, Expert Consultation. Rome, Italy.
- Fascetti, A. J., J. R. Teed, Q. E. Rogers, R. C. Backus. 2003. Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997-2001). *J. Am. Vet. Med. Assoc.* 223:1137-1141. doi: 10.2460/javma.2003.223.1137
- FDA, Center for Veterinary Medicine. FDA Investigating Potential Connection Between Diet and Cases of Canine Heart Disease. July 12th, 2018. <https://www.fda.gov/animalveterinary/newsevents/cvmupdates/ucm613305.htm>
- Flanagan, J.L., P. A. Simmons, J. V. Vehige, M. D. P. Willcox, Q. Garrett. 2010. Role of carnitine in disease. *Nutr. Metab. (Lond).* 7:30. doi: 10.1186/1743-7075-7-30.

- Freeman, L. M., J. A. Stern, R. Fries, D. B. Adin, J. E. Rush. 2018. Diet-associated dilated cardiomyopathy in dogs: what do we know? *J. Am. Vet. Med. Assoc.* 253:1390-1394. <https://doi.org/10.2460/javma.253.11.1390>.
- Freeman, L.M., J. E. Rush. 2006. Cardiovascular diseases: nutritional modulation. In: *Encyclopedia of Canine Clinical Nutrition*. Pibot, P., V. Biourge, D. Elliott. Aimargues: Aniwa SAS. Pp. 316-347.
- Freeman, L.M., K. E. Michel, D. J. Brown, P. M. Kaplan, M. E. Stamoulis, S. L. Rosenthal, B. W. Keene, J. E. Rush. 1996. Idiopathic dilated cardiomyopathy in Dalmatians: nine cases (1990-1995). *J. Am. Vet. Med. Assoc.* 209:1592-1596.
- Gloaguen, M., N. Le Floch, E. Corrent, Y. Primot, J. van Milgen. 2014. The use of free amino acids allows formulating very low crude protein diets for piglets. *J. Anim. Sci.* 92(2):637-44. doi: 10.2527/jas.2013-6514.
- Haber, G. B., K. W. Heaton, D. Murphy, L. F. Burroughs. 1977. Depletion and disruption of dietary fibre. Effects on satiety, plasma glucose, and serum-insulin. *Lancet.* 2:679-682.
- Homan, H. H., M. Kemen, C. Fuessenich, M. Senkal, V. Zumtobel. 1994. Reduction in diarrhea incidence by soluble fiber in patients receiving total or supplemental enteral nutrition. *J. Parenter. Enter. Nutr.* 18:486-490. doi: 10.1177/0148607194018006486
- Hoppel, C. 2003. The role of carnitine in normal and altered fatty acid metabolism. *Am. J. Kidney Dis.* 41:S4-12. doi: 10.1016/S0272-6386(03)00112-4
- Hou, Y., Y. Yin, G. Wu. 2015. Dietary essentiality of "nutritionally non-essential amino acids" for animals and humans. *Exp Biol Med.* 240(8):997-1007. doi: 10.1177/1535370215587913
- Hurrell, R. F., P. A. Finot, J. E. Ford. 1983. Storage of milk powders under adverse conditions. I. Losses of lysine and of other essential amino acids as determined by chemical and microbiological methods. *Br. J. Nutr.* 49(3):343-54. doi: 10.1079/BJN19830043
- Huxtable, R.J. 1992. Physiological actions of taurine. *Physiol. Rev.* 72:101-163. doi: 10.1152/physrev.1992.72.1.101
- Ito, T., S. W. Schaffer, J. Azuma. 2012. The potential usefulness of taurine on diabetes mellitus and its complications. *Amino Acids.* 42(5): 1529-1539. doi: 10.1007/s00726-011-0883-5
- Johnson, L., A. P. Mander, L. R. Jones, P. M. Emmett, S. A. Jebb. Energy-dense, low-fiber, high-fat dietary pattern is associated with increased fatness in childhood. *Am. J. Clin. Nutr.* 87:846-854. doi: 10.1093/ajcn/87.4.846
- Johnson, M. L., C. M. Parsons, G. C. Jr Fahey, N. R. Merchen, C. G. Aldrich. 1998. Effects of species raw material source, ash content, and processing temperature on amino acid digestibility of animal by-product meals by cecectomized roosters and ileally cannulated dogs. *J. Anim. Sci.* 76:1112-1122.

- Kaplan, J. L., J. A. Stern, A. J. Fascetti, J. A. Larsen, H. Skolnik, G. D. Peddle, R. D. Kienle, A. Waxman, M. Cocchiaro, C. T. Gunther-Harrington, T. Klose, K. LaFauci, B. Lefbom, M. Machen Lamy, R. Malakoff, S. Nishimura, M. Oldach, S. Rosenthal, C. Stauthammer, L. O'Sullivan, L. C. Visser, R. William, and E. Ontiveros. 2018. Taurine deficiency and dilated cardiomyopathy in golden retrievers fed commercial diets. *PLoS ONE* 13(12):e0209112. <https://doi.org/10.1371/journal.pone.0209112>
- Keene, B.W., D. P. Panciera, C. E. Atkins, V. Regitz, M. J. Schmidt, A. L. Shug. 1991. Myocardial L-carnitine deficiency in a family of dogs with dilated cardiomyopathy. *J. Am. Vet. Med. Assoc.* 198:647-650.
- Kim, S.W., Q. R. Rogers, J. G. Morris. 1996(a). Dietary antibiotics decrease taurine loss in cats fed a canned heat-processed diet. *J. Nutr.* 126:509–515. DOI: 10.1093/jn/126.2.509
- Kim, S.W., Q. R. Rogers, J. G. Morris. 1996(b). Maillard reaction products in purified diets induce taurine depletion in cats which is reversed by antibiotics. *J Nutr.* 126:195–201. doi: 10.1093/jn/126.1.195
- Kittleson, M. D., B. Keene, P. D. Pion, C. G. Loyer. 1997. Results of the multicenter spaniel trial (MUST): taurine- and carnitine-responsive dilated cardiomyopathy in American cocker spaniels with decreased plasma taurine concentration. *J. Vet. Intern. Med.* 11:204-211. doi: 10.1111/j.1939-1676.1997.tb00092.x
- Ko, K.S., R. C. Backus, J. R. Berg, M. W. Lame, Q. R. Rogers. 2007. Differences in taurine synthesis rate among dogs relate to differences in their maintenance energy requirement. *J Nutr.* 137:1171–1175. doi: 10.1093/jn/137.5.1171
- Kramer, G.A., M.D. Kittleson, P.R. Fox, J. Lewis, and P.D. Pion. 1995. Plasma taurine concentrations in normal dogs and in dogs with heart disease. *J. Vet. Intern. Med.* 9(4):253-258.
- Lien, K. A., W. C. Sauer, M. Fenton. 1997. Mucin output in ileal digesta of pigs fed a protein-free diet. *Z Ernährungswiss.* 36:182–190. doi: 10.1007/BF01611398
- Linder, D., M. Mueller. 2014. Pet Obesity Management: Beyond Nutrition. *Vet. Clin. N. Am-Small.* 44:789-806. doi: 10.1016/j.cvsm.2014.03.004
- Mansilla, W.D., K. E. Silva, C. Zhu, C. M. Nyachoti, J. K. Htoo, J. P. Cant, C. F. M. de Lange. 2018. Ammonia-nitrogen added to low-crude-protein diets deficient in dispensable amino acid-nitrogen increases the net release of alanine, citrulline, and glutamate post-splanchnic organ metabolism in growing pigs. *J. Nutr.* 148:1081-1087. doi: 10.1093/jn/nxy076.
- Marinangeli, C. P. F., J. Curran, S. I. Barr, J. Slavin, S. Puri, S. Swaminathan, L. Tapsell, C. A. Patterson. 2017. Enhancing nutrition with pulses: defining a recommended serving size for adults. *Nutr Rev.* 75:990-1006. doi: 10.1093/nutrit/nux058.

- Maron, B. J., J. A. Towbin, G. Thiene, C. Antzelevitch, D. Corrado, D. Arnett, A. J. Moss, C. E. Seidman, J. B. Young. 2006. Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation*. 113:1807-1816. doi: 10.1161/CIRCULATIONAHA.106.174287
- Marshall, H. F., K. C. Chang, K. S. Miller, L. D. Satterlee. 1982. Sulfur amino acid stability: Effects of processing on legume proteins. *J Food Sci*. 47:1170-4. doi: 10.1111/j.1365-2621.1982.tb07642.x
- Martin, M.W., M. J. Stafford Johnson, B. Celona. 2009. Canine dilated cardiomyopathy: a retrospective study of signalment, presentation and clinical findings in 369 cases. *J. Small Anim. Pract*. 50:23-29. doi: 10.1111/j.1748-5827.2008.00659.x
- Mathai, J. K., J. K. Htoo, J. E. Thomson, K. J. Touchette, H. H. Stein. 2016. Effects of dietary fiber on the ideal standardized ileal digestible threonine:lysine ratio for twenty-five to fifty kilogram growing gilts. *J. Anim. Sci*. 94:4217–4230. doi: 10.2527/jas.2016-0680.
- Menniti, M. F., G. M. Davenport, A. K. Shoveller, J. P. Cant, V. R. Osborne. 2014. Effect of graded inclusion of dietary soybean meal on nutrient digestibility, health, and metabolic indices of adult dogs. *J. Anim. Sci*. 92:2094-2104. doi: 10.2527/jas.2013-7226
- Merheb, M., R. T. Daher, M. Nasrallah, R. Sabra, F. N. Ziyadeh, K. Barada. 2007. Taurine intestinal absorption and renal excretion test in diabetic patients: a pilot study. *Diabetes Care* 30:2652–2654. doi: 10.2337/dc07-0872
- Meurs, K. M., J. A. Stern, D. D. Sisson, M. D. Kittleson, S. M. Cunningham, M. K. Ames, C. E. Atkins, T. DeFrancesco, T. E. Hodge, B. W. Keene, Y. Reina Doreste, M. Leuthy, A. A. Motsinger-Reif, S. P. Tou. 2013. Association of dilated cardiomyopathy with the striatin mutation genotype in boxer dogs. *J. Vet. Intern. Med*. 27:1437–1440. doi: 10.1111/jvim.12163
- Meurs, K. M., S. Lahmers, B. W. Keene, S. N. White, M. A. Oyama, E. Mauceli, K. Lindblad-Toh. 2012. A splice site mutation in a gene encoding for PDK4, a mitochondrial protein, is associated with the development of dilated cardiomyopathy in the Doberman pinscher. *Human Genetics* 131:1319–1325. doi: 10.1007/s00439-012-1158-2
- Moehn, S., R. F. Bertolo, P. B. Pencharz, R. O. Ball. 2005. Development of the indicator amino acid oxidation technique to determine the availability of amino acids from dietary protein in pigs. *J Nutr*. 135(12):2866-70.
- Moise, N.S., L. M. Pacioretty, F. A. Kallfelz, M. H. Stipanuk, J. M. King, R. F. Jr Gilmour. 1991. Dietary taurine deficiency and dilated cardiomyopathy in the fox. *Am. Heart J*. 121:541-547. doi: 10.1016/0002-8703(91)90724-V


- Monnet, E., E. C. Orton, M. Salman, J. Boon. 1995. Idiopathic dilated cardiomyopathy in dogs: survival and prognostic indicators. *J. Vet. Intern. Med.* 9:12-17. doi: 10.1111/j.1939-1676.1995.tb03266.x
- Nardelli, T. R., R. A. Ribeiro, S. L. Balbo, E. C. Vanzela, E. M. Carneiro, A. C. Boschero, M. L. Bonfleur. 2011. Taurine prevents fat deposition and ameliorates plasma lipid profile in monosodium glutamate-obese rats. *Amino Acids.* 41(4):901-908. doi: 10.1007/s00726-010-0789-7
- NRC, National Research Council. 2006. *Nutrient Requirements of Dogs and Cats.* 10th ed. The National Academy Press, Washington, DC.
- NRC, National Research Council. 2012. *Nutrient Requirements of Swine.* 11th ed. The National Academic Press, Washington, DC.
- Nyachoti, C. M., C. F. M. de Lange, B. W. McBride, S. Leeson, H. Schulze. 2000. Dietary influence on organ size and in vitro oxygen consumption by visceral organs of growing pigs. *Livest Prod Sci.* 65:229-237. doi: 10.1016/S0301-6226(00)00157-3
- O'Máille, E. R. L., T. G. Richards, A. H. Short. 1965. Acute taurine depletion and maximal rates of hepatic conjugation and secretion of cholic acid in the dog. *J Physiol.* 1965;180:67-79.
- Owczarek-Lipska, M., T. B. Mausberg, H. Stephenson, J. Dukes-McEwan, G. Wess, T. Leeb. 2013. A 16-bp deletion in the canine PDK4 gene is not associated with dilated cardiomyopathy in a European cohort of Doberman Pinschers. *Anim Genet.* 44:239. doi: 10.1111/j.1365-2052.2012.02396.x
- Patterson, C. A., J. Curran, T. Der. 2017. Effect of Processing on Antinutrient Compounds in Pulses. *Cereal Chemistry.* 94(1):2-10. doi: doi:10.1094/CCHEM-05-16-0144-FI.
- Pion, P. D., M. D. Kittleson, Q. R. Rogers, J. G. Morris. 1987. Myocardial failure in cats is associated with low plasma taurine: a reversible cardiomyopathy. *Science* 237:764-768. doi: 10.1126/science.3616607
- Pion, P. D., S. L. Sanderson, M. D. Kittleson. 1998. The effectiveness of taurine and levocarnitine in dogs with heart disease. *Vet. Clin. North Am. Small Anim. Pract.* 28:1495-1514. doi: 10.1016/S0195-5616(98)50134-9
- Rășanu, T., M. Mehedinți-Hâncu, M. Alexianu, T. Mehedinți, E. Gheorghe, I. Damian. Carnitine deficiency. *Rom J Morphol Embryol.* 2012;53:203-206.
- Rice, J. E., S. L. Ihle. 1994. Effects of diet on fecal occult blood testing in healthy dogs. *Can. J. Vet. Res.* 58:134-137.
- Roberfroid, M. B. 2005. Introducing inulin-type fructans. *Br. J. Nutr.* 93:S13-25.

- Robinson, J. L., L. E. McBreairty, E. W. Randell, J. A. Brunton, R. F. Bertolo. 2016(b). Restriction of dietary methyl donors limits methionine availability and affects the partitioning of dietary methionine for creatine and phosphatidylcholine synthesis in the neonatal piglet. *J. Nutr. Biochem.* 35:81-86. doi: 10.1016/j.jnutbio.2016.07.001.
- Robinson, J. L., S. V. Harding, J. A. Brunton, R. F. Bertolo. 2016(a). Dietary methyl donors contribute to whole-body protein turnover and protein synthesis in skeletal muscle and the jejunum in neonatal piglets. *J. Nutr.* 146: 2007–2012. doi: 10.3945/jn.115.226035
- Sanderson, S.L., K. L. Gross, P. N. Ogburn, C. Calvert, G. Jacobs, S. R. Lowry, K. A. Bird, L. A. Koehler, L. L. Swanson. 2001. Effects of dietary fat and L-carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am J Vet Res.* 62:1616-1623. doi: 10.2460/ajvr.2001.62.1616
- Schaffer, S. W., C. J. Jong, K. C. Ramila, J. Azuma. 2010. Physiological roles of taurine in heart and muscle. *J Biomed Sci* 17:S2. doi: 10.1186/1423-0127-17-S1-S2
- Sisson, D. D., W. P. Thomas, B. W. Keene. 2000. Primary myocardial disease in the dog. In: Ettinger SJ, Feldman EC, editors. *Textbook of veterinary internal medicine. Diseases of the dog and cat.* 5th edition. Philadelphia: WB Saunders Co. Pp. 874–895.
- Spitze, A. R., D. L. Wong, Q. R. Rogers, A. J. Fascetti. 2003. Taurine concentrations in animal feed ingredients; cooking influences taurine content. *J. Anim. Physiol.* 87:251-262. OI: 10.1046/j.1439-0396.2003.00434.x
- Stoll, B., J. Henry, P. J. Reeds, H. Yu, F. Jahoor, D. G. Burrin. 1998. Catabolism dominates the first-pass intestinal metabolism of dietary essential amino acids in milk protein-fed piglets. *J Nutr.* 128:606-614. doi: 10.1093/jn/128.3.606
- Story, J.A., D. Kritchevsky. 1978. Bile acid metabolism and fiber. *Am. J. Clin. Nutr.* 31:S199-S202. doi: 10.1093/ajcn/31.10.S199
- Swanson, K. S., R. A. Carter, T. P. Yount, J. Aretz, P. R. Buff. 2013. Nutritional Sustainability of Pet Foods. *Adv Nutr.* 4(2):141–150. doi: 10.3945/an.112.003335
- Teodorowicz, M., J. van Neerven, H. Savelkoul. 2017. Food processing: The influence of the Maillard reaction on immunogenicity and allergenicity of food proteins. *Nutr* 9:835. doi: 10.3390/nu9080835
- Tôrres, C. L., R. C. Backus, A. J. Fascetti, Q. R. Rogers. 2003. Taurine status in normal dogs fed a commercial diet associated with taurine deficiency and dilated cardiomyopathy. *J Anim Physiol Anim Nutr (Berl).* 87:359-372. doi: 10.1046/j.1439-0396.2003.00446.x
- Tosh, S.M. S. Yada. Dietary fibres in pulse seeds and fractions: Characterization, functional
Werner, P., M. G. Raducha, U. Prociuk, M. M. Sleeper, T. J. Van Winkle, P. S. Henthorn. 2008. A novel locus for dilated cardiomyopathy maps to canine chromosome. *Genomics.* 2008; 91(6):517-521. doi: 10.1016/j.ygeno.2008.03.007

- Worden, J. A., M. H. Stipanuk. 1985. A comparison by species, age and sex of cysteinesulfinate decarboxylase activity and taurine concentration in liver and brain of animals. *Comp Biochem Physiol* 82B:233-239. doi: 10.1016/0305-0491(85)90232-9
- Yamka, R. M., U. Jamikorn, A. D. True, D. L. Harmon. 2003. Evaluation of soybean meal as a protein source in canine foods. *Anim. Feed Sci. Technol.* 109:121-132. doi: 10.1016/S0377-8401(03)00203-7
- Žilić, S., I. Bozović, V. H. T. Šukalović. 2012. Thermal Inactivation of Soybean Bioactive Proteins. *International Journal of Food Engineering* 8:1556-3758 doi: <https://doi.org/10.1515/1556-3758.2521>.


Accepted Manuscript

Figure 1. Metabolism of sulfur amino acids. DMG: dimethylglycine, SAH, S-adenosylhomocysteine; SAM, S-adenosylmethionine



Accepted Manuscript

Figure 2. Effect of thermal processing methods on trypsin inhibitor levels (mg/g) soybean kernel. ¹Treatment conditions: None = no treatment; Dry Extrusion for 25 to 30 sec (1=100 °C; 2=125 °C; 3=140 °C; 4=150 °C); Wet Extrusion for 25 to 30 sec with 6 to 8 % added moisture (1=100 °C; 2=125 °C; 3=140 °C); Micronisation with near-infrared rays wavelength of 1.8 to 3.4 μ m for 90 sec (1=100 °C; 2=125 °C; 3=140 °C; 4=150 °C); Microwave roasting at 800 W and 2450 MHz (1 = 1 min (kernel temp = 57 °C), 2 = 2 min (kernel temp = 88 °C), 3 = 3 min (kernel temp = 108 °C), 4 = 4 min (kernel temp = 121 °C), 5 = 5 min (kernel temp = 132 °C)); Autoclaving at 120 °C and 1.2 bars (1 = 10 min, 2 = 20 min, 3 = 30 min). Reprinted with permission from Zilic et al. (2012)



ACCEPTED

Table 1. Crude protein (CP), fiber, selected amino acids, and carnitine contents in the principal legumes, cereals, and animal-derived ingredients used in dog food formulation.¹

Ingredients	CP, %	Crude fiber, ² %	α -amino acids, mg/g protein ²			Tau, mg/kg ³	Carnitine, mg/kg ⁴	
			Lys	Met	Cys			
Legumes	Fava Beans	27.2	8.55	23.9	7.0	12.5	--	
	Phaseolus beans	22.9	NR	72.9	12.7	12.7	--	
	Kidney beans	20.0	6.40	26.5	14.0	12.0	--	
	Lentils	26.0	NR	65.8	6.9	10.4	--	
	Lupins	32.4	14.25	48.7	6.5	14.2	--	
	Chick peas	20.3	6.16	69.4	14.8	21.6	--	
	Soybean meal	47.7	3.89	62.0	13.8	14.7	--	
Grains	Barley	11.3	3.90	35.3	17.7	22.9	--	
	Corn, yellow dent	8.2	1.98	30.3	21.8	23.1	--	
	Oats	11.2	2.20	43.9	60.9	32.3	--	
	Rice	7.9	0.52	44.5	31.8	22.9	--	
	Rye	11.7	2.71	36.9	13.7	16.3	--	
	Sorghum	9.4	2.14	21.4	17.1	19.2	--	
	Wheat hard, red	14.5	2.57	27.0	15.2	22.8	--	
Animal-derived ingredients	Beef, meat	15.0	--	77.3	28.7	15.3	296	150
	Chicken, meat and skin	17.6	--	81.3	26.7	13.1	159	57
	Chicken, by product	59.0	--	48.1	17.3	16.8	3049	120
	Lamb, ground	16.6	--	88.0	25.9	12.0	473	282.3
	Rendered meat	54.1	2.50	53.8	14.2	11.3	NR	NR

Cys: cysteine, Lys: lysine, Met: methionine, NR: not reported, Tau: taurine.

¹Values are presented in as-fed basis.

²NRC, 2006; NRC, 2012

³Spitze et al. 2003

⁴Arslan, 2006

Table 2. Recommended allowance (RA) and minimum dietary content suggested by AAFCO for crude protein and essential amino acids in dog food, and their physiological roles and potential interactions.

Nutrient	NRC RA¹, % DM	AAFCO², % DM	Important physiological roles and potential interactions
Crude protein	10	18	Necessary for synthesis of non-essential amino acids
Arginine	0.35	--	Competes with lysine absorption, arginine should be increased when high lysine concentrations in the diet
Histidine	0.19	--	
Lysine	0.35	0.63	Highly reactive to reducing sugars during heating (Maillard reaction), reducing bioavailability
Methionine	0.33	0.33	Requirement increases when methyl donors/acceptors and cysteine are reduced in the diet
Methionine + cystine	0.65	0.65	Requirement is increased with low supply of taurine and during immune challenge
Phenylalanine	0.45	0.45	
Phenylalanine + tyrosine	0.74	0.74	
Threonine	0.43	0.48	Abundant in mucosal proteins (mucin), requirement increases when feeding high fermentable fibers
Tryptophan	0.14	0.16	Precursor for serotonin synthesis. Ratio of Trp: LNAA should be considered; lower ratios may deprive appetite
Valine	0.49	0.49	Abnormal Increment of valine, leucine, or isoleucine (BCAA) will cause catabolism of the other BCAA in the muscle
Isoleucine	0.38	--	
Leucine	0.68	0.68	

AAFCO: The Association of American Feed Control Officials, BCAA: branched chain amino acids, DM: dry matter, NRC: National Research Council, RA: recommended allowance, Trp: LNAA: tryptophan to large neutral amino acid ratio.

¹Recommended Allowance requirements for adult dogs at maintenance, Nutrient Requirements of Dogs and Cats (NRC, 2006).

²Minimum dietary content, AAFCO (2018).

From: [Norris, Anne](#)
To: [Hartogenesis, Martine](#)
Cc: [DeLancey, Siobhan](#); [Rotstein, David](#)
Subject: RE: DCM Timing
Date: Thursday, June 20, 2019 10:29:48 AM

Just wanted to check – are you planning to [REDACTED] (b) (5)

From: Hartogenesis, Martine
Sent: Wednesday, June 19, 2019 2:22 PM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: DCM Timing

Ok, great. We were planning to start [REDACTED] (b) (5)

Let me know if that works for everyone!

Martine

From: Norris, Anne
Sent: Wednesday, June 19, 2019 2:18 PM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: RE: DCM Timing

Whew, good question! I think we're shooting [REDACTED] (b) (5) I don't see it happening sooner than that. Hope that helps!

From: Hartogenesis, Martine
Sent: Wednesday, June 19, 2019 2:16 PM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Rotstein, David <David.Rotstein@fda.hhs.gov>
Subject: DCM Timing

Hi Anne,

Just checking in on your current DCM comms timing..? [REDACTED] (b) (5)

Thanks very much in advance!

Martine

From: [Forfa, Tracey](#)
To: [Norris, Anne](#); [Rotstein, David](#); [Jones, Jennifer L](#); [Palmer, Lee Anne](#); [Burkholder, William](#); [Carey, Lauren](#); [Steinberg, Nadine](#)
Cc: [DeLancey, Siobhan](#); [Hartogensis, Martine](#); [Peloquin, Sarah](#)
Subject: RE: DCM-firm contacts
Date: Monday, June 03, 2019 3:56:50 PM
Attachments: [image001.png](#)
[image002.jpg](#)
[image003.jpg](#)
[image004.jpg](#)
[image005.jpg](#)
[image006.jpg](#)

Hi – That is correct, I have been tasked with [REDACTED] (b) (5)
[REDACTED] Thanks for checking in.

From: Norris, Anne
Sent: Monday, June 3, 2019 3:36 PM
To: Rotstein, David <David.Rotstein@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Burkholder, William <William.Burkholder@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>; Steinberg, Nadine <Nadine.Steinberg@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>
Subject: RE: DCM-firm contacts

Adding in Tracey and Nadine because [REDACTED] (b) (5)
[REDACTED]

From: Rotstein, David
Sent: Monday, June 3, 2019 3:27 PM
To: Norris, Anne <Anne.Norris@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Burkholder, William <William.Burkholder@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>
Cc: DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>; Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>
Subject: DCM-firm contacts

Everyone,

(b) (5)

(b) (5)

Dave

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/CERT
7519 Standish Place
(b) (6) (BB)



This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

-----Original Appointment-----

From: Solomon, Steven M

Sent: Wednesday, May 29, 2019 11:40 AM

To: Solomon, Steven M; Flynn, William T; Forfa, Tracey; Norris, Anne; Schell, Timothy; Jones, Jennifer L; Palmer, Lee Anne; Burkholder, William; Carey, Lauren

Cc: DeLancey, Siobhan; Hartogensis, Martine; Murphy, Jeanette; Dewitt, Susan J; Cepeda, Sandra; Steinberg, Nadine; Rotstein, David; Reimschuessel, Renate; Ceric, Olgica; Peloquin, Sarah

Subject: FW: Checkpoint on DCM

When: Thursday, May 30, 2019 12:00 PM-1:00 PM (UTC-05:00) Eastern Time (US & Canada).

Where: CVM 7500 Conf E473 and WebEx

-----Original Appointment-----

From: Solomon, Steven M

Sent: Monday, May 20, 2019 10:36 AM

To: Solomon, Steven M; Flynn, William T; Forfa, Tracey; Norris, Anne; Schell, Timothy; Jones, Jennifer L; Palmer, Lee Anne; Burkholder, William; Carey, Lauren

Cc: DeLancey, Siobhan; Hartogensis, Martine; Murphy, Jeanette; Dewitt, Susan J; Cepeda, Sandra; Steinberg, Nadine

Subject: Checkpoint on DCM

When: Thursday, May 30, 2019 12:00 PM-1:00 PM (UTC-05:00) Eastern Time (US & Canada).

Where: CVM 7500 Conf E473 and WebEx

Purpose: [REDACTED] (b) (5)

Meeting materials forthcoming.

Apologies for the lunchtime meeting, but schedules were tight.

[Join Webex meeting](#)

Meeting number (access code): [REDACTED] (b) (6)

Meeting password: [REDACTED] (b) (6)

Join by phone

[REDACTED] (b) (6) US Toll

+ [REDACTED] (b) (6) US Toll Free

[Global call-in numbers](#) | [Toll-free calling restrictions](#)

[Can't join the meeting?](#)

If you are a host, [go here](#) to view host information.

IMPORTANT NOTICE: Please note that this Webex service allows audio and other information sent during the session to be recorded, which may be discoverable in a legal matter. By joining this session, you automatically consent to such recordings. If you do not consent to being recorded, discuss your concerns with the host or do not join the session.

From: [Rotstein, David](#)
To: [Jones, Jennifer L](#)
Subject: RE: EON-266814; EON-266821 and EON-266827-Merrick-FW: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action
Date: Wednesday, August 24, 2016 4:31:23 PM
Attachments: [958504-Taurine.pdf](#)
[958501-Taurine.pdf](#)
[958500-Taurine.pdf](#)

Here you go

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/ICERT
7519 Standish Place, RM 120
240-402-5613 (Office) (**NEW NUMBER**)
(b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david.rotstein@fda.hhs.gov.

From: Jones, Jennifer L
Sent: Wednesday, August 24, 2016 7:23 AM
To: Rotstein, David
Subject: RE: EON-266814; EON-266821 and EON-266827-Merrick-FW: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Sorry to both you Dave, do you also have the official results report? I just need a copy for the final report, so it's not urgent.

Jennifer Jones, DVM
Veterinary Medical Officer

From: Jones, Jennifer L
Sent: Wednesday, August 24, 2016 7:08 AM
To: Rotstein, David
Subject: RE: EON-266814; EON-266821 and EON-266827-Merrick-FW: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Thanks Dave!

Jennifer Jones, DVM
Veterinary Medical Officer

From: Rotstein, David
Sent: Wednesday, August 24, 2016 6:35 AM
To: Jones, Jennifer L
Subject: EON-266814; EON-266821 and EON-266827-Merrick-FW: Moisture Content: RE: Quick

Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Here you go Jen.

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/ICERT
7519 Standish Place, RM 120
240-402-5613 (Office) (NEW NUMBER)

(b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david_rotstein@fda.hhs.gov.

From: Burkholder, William
Sent: Wednesday, July 27, 2016 4:30 PM
To: Rotstein, David; Benjamin, Linda
Cc: Krieger, Darlene; Queen, Jackie L; Hodges, April; Conway, Charlotte
Subject: RE: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

OK Everyone. The product appears to be a dry extruded product, for which the AAFCO Cat Food Nutrient Profiles content for taurine is 0.10% on a dry matter basis. Clearly all three samples were analyzed to contain more than that amount of taurine. **On a dry matter basis the concentration of taurine in the samples was analyzed to be:**

FACTS #	Amount Taurine Found	%Moisture	%Dry Matter	Amount
Taurine on a Dry Matter Basis				
958500	0.183g/100g ≈ 0.18%	2.20%	100 – 2.20 = 97.80%	
	0.183/0.9780 = 0.187%			
958501	0.153g/100g ≈ 0.15%	1.99%	100 – 1.99 = 98.01%	
	0.153/0.9801 = 0.156%			
958504	0.171g/100g ≈ 0.17%	2.79%	100 – 2.79 = 97.21%	
	0.171/0.9721 = 0.176%			

All of the Dry Matter Taurine percentages are above 0.10%. IF any of the samples were canned cat food, they would not be in compliance with the AAFCO Cat Food Nutrient Profiles for the recommended minimum taurine content and IF the label indicated the product was formulated to meet the AAFCO Cat Food Nutrient Profiles the product would be misbranded.

The answer to the question of consequence/causation of the taurine content in the product from which these three samples originated to the cats in the consumer complaint is that this(ese) lot(s) of product are not indicated to be causative. However, dilated cardiomyopathy from taurine deficiency occurs over a long period of exposure to a deficient diet (months to a year or more), so, if these cats were eating the Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry for the 3 years

indicated in the complaint, it is possible that the product was deficient for some long interval of time during that three year period and that a return to "normal" taurine levels in the diet were insufficient to correct the problem in the three cats that developed low blood taurine and the two with dilated cardiomyopathy. Treatment for dilated cardiomyopathy caused by taurine deficiency takes higher daily doses of taurine for several months than normal dietary amounts and is not completely curative.

Recommendations for regulatory steps to consider [REDACTED] (b) (5)

Consider recommending the owner have an ophthalmic exam performed on the cat being treated for low blood taurine to see if there are signs of retinal degeneration due to taurine deficiency.

William J. Burkholder, DVM, PhD, DACVN
Leader, Nutrition and Labeling Team I, HFV-228
Division of Animal Feeds
Center for Veterinary Medicine
United States Food and Drug Administration
7519 Standish Place
Rockville, Maryland 20855
Phone: 240-402-5900
Fax: 240-453-6882
E-mail: william.burkholder@fda.hhs.gov

The opinions and information in this message are those of the author and do not necessarily reflect the views and policies of the U.S. Food and Drug Administration. Because of the nature of electronically transferred information, the integrity or security of this message cannot be guaranteed. This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution or copying is strictly prohibited. If you think you have received this e-mail message in error, please e-mail the sender immediately at William.Burkholder@fda.hhs.gov.

From: Rotstein, David
Sent: Wednesday, July 27, 2016 2:23 PM
To: Benjamin, Linda; Burkholder, William
Cc: Krieger, Darlene; Queen, Jackie L; Hodges, April
Subject: Moisture Content: RE: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Please see the moisture content below:

The moisture content for the samples are as follows:

FACTS #	Amount Taurine Found	%Moisture
958500	0.183g/100g ≈ 0.18%	2.20%

958501	0.153g/100g ≈ 0.15%	1.99%
958504	0.171g/100g ≈ 0.17%	2.79%

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/ICERT
7519 Standish Place, RM 120
240-402-5613 (Office) (NEW NUMBER)
(b) (6) (BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david_rotstein@fda.hhs.gov.

From: Benjamin, Linda
Sent: Wednesday, July 27, 2016 7:54 AM
To: Burkholder, William
Cc: Rotstein, David; Krieger, Darlene; Queen, Jackie L; Hodges, April
Subject: FW: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Hi Bill - Could you please respond to Dave Rotstein.

Dave - As ORS's numbers are very close to the 0.2% guarantee, it might be helpful to know the AV, CV, and/or 95% confidence limit for the analytical method. Additionally, do you know if the numbers below are being reported on a dry matter basis? FYI, the sample description on the collection reports (first 3 attachments) has either "One unopened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe weighing 5.4kg" or "Opened bag of Merrick Purrfect Bistro Grain Free Real Chicken Recipe that only had 0.15kg of product. This sample was used by the consumer" but below **my green highlight** you referenced taurine # for canned products.

Sorry Bill - I just want to make sure you have everything you need.

Thanks for the opportunity to comment,
Linda

From: Rotstein, David
Sent: Wednesday, July 27, 2016 7:22 AM
To: Benjamin, Linda
Cc: Krieger, Darlene; Queen, Jackie L; Hodges, April
Subject: Quick Review/Screening: ORA ORS Cat Food Results--Assessment of Taurine Results for Possible Product Action

Linda,

We received an email from ORS with results for taurine for a cat food. Testing was based on a consumer complaint for 3 cats with cardiac disease and low taurine.

ORS has not finalized the results, but sent on the findings for the DRY cat food and asked whether

CVM considers the results to be low based on the AAFCO requirements for wet cat food.

REQUEST: To answer the following questions:

|

- 1) Is the taurine low for a dry cat food based on AAFCO nutrient profiles?
- 2) If the taurine is low, would it be biologically significant for cats that ate this as their sole/primary diet?

The responses will [REDACTED] (b) (5)

REVIEWERS: Bill Burkholder, Krisztina Atkinson, Randall Lovell.

Date Needed: IDEAL—By our Wednesday Pet Food Outbreak Meeting at 11 AM today (7/27). (I will be out on Thursday and Friday and if no one can respond before the meeting today, please include Jackie Queen on the response).

Email from the ORS Lab:

David I hope you can help us with these findings.

We received three consumer complaint Dry Cat Food products for Amino Acid analysis. We assayed the products for the Amino Acid profile and found only Taurine low.

FACTS #	Amount Taurine Found
958500	0.183g/100g ≈ 0.18%
958501	0.153g/100g ≈ 0.15%
958504	0.171g/100g ≈ 0.17%

The label for all of the samples are the same and Taurine is declared 0.20% minimum. The AAFCO Nutrient Profile from August 2015 states that the minimum limits for Taurine is 0.20% in canned products. Do you consider these product violated?

Attachments:

**Collection Reports
Pet Food Report
Vet-LIRN Summary**

Medical records were collected and evaluated by Vet-LIRN. These can be provided by request.

Thank you

David Rotstein, DVM, MPVM, Dipl. ACVP
CVM Vet-LIRN Liaison
CVM OSC/DC/ICERT
7519 Standish Place, RM 120

240-402-5613 (Office) (**NEW NUMBER**)

(b) (6)(BB)

This e-mail message is intended for the exclusive use of the recipient(s) named above. It may contain information that is protected, privileged, or confidential, and it should not be disseminated, distributed, or copied to persons not authorized to receive such information. If you are not the intended recipient, any dissemination, distribution, or copying is strictly prohibited. If you think you received this e-mail message in error, please e-mail the sender immediately at david_rotstein@fda.hhs.gov.

Food and Drug Administration Office of Regulatory Affairs

Summary Report

For Sample Number: 958500

TD Sample Number:

Import Sample Number

This is an accurate reproduction of the original electronic record as of 08/24/2016

Sample Class: Normal Everyday Sample **Sample Origin:** Domestic **Sample Basis:** Surveillance
Sample Flag: **Sample Type:** Official **Collecting District:** NWJ-DO
Home District: **Orig C/R and Records To:** NWJ-DO **Collection PACs:** 71R801

Product Name: Poultry Prod Pet Cat Food; Not Elsewhere Classified (NEC); Packaged Food (Not Commercially Sterile)

Product Description: See Remarks Section.

Collection Reason: Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

Lab: SRL **Split Num:**0 **Date Received:** 06/29/2016 **Date Out of Lab:** 08/04/2016
District **District Conclusion** **District**
Conclusion: No Action Indicated (NAI) **Made By:** Tweedley, Karen P 08/12/2016
Disposition **Disposition** **Disposition**
Reason: NAI By Examining District **Authorized By:** Tweedley, Karen P **Authorized Date:** 08/12/2016

Performing Org	PAC	LID	PAF	Compliance No	Lab Class-Description	Laboratory Status
ACNA-N	71R801		NAR		1 - In Compliance	Completed

Lab Conclusion

Sample Narrative - Method: AccQTag AAA(Waters) Analysis - Taurine
Amt Found - 0.187% (dry matter basis)
Meets AAFCO minimum requirement of 0.10%

Sample Narrative - Method: Instrument Manual (Denver Instrumentst IR60)/ AOAC 930.15 Analysis - Moisture
Amt Found - 2.20%
Amt Declared - 11.00% max

Lab Conclusion Date **Lab Conclusion Made By**
08/04/2016 Hawes,Brian M

Food and Drug Administration Office of Regulatory Affairs

Summary Report

For Sample Number: 958501

TD Sample Number:

Import Sample Number

This is an accurate reproduction of the original electronic record as of 08/24/2016

Sample Class: Normal Everyday Sample **Sample Origin:** Domestic **Sample Basis:** Surveillance
Sample Flag: **Sample Type:** Official **Collecting District:** NWJ-DO
Home District: **Orig C/R and Records To:** NWJ-DO **Collection PACs:** 71R801

Product Name: Poultry Prod Pet Cat Food; Not Elsewhere Classified (NEC); Packaged Food (Not Commercially Sterile)

Product Description: See Remarks

Collection Reason: Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

Lab: SRL **Split Num:**0 **Date Received:** 06/29/2016 **Date Out of Lab:** 08/04/2016
District **District Conclusion** **District**
Conclusion: No Action Indicated (NAI) **Made By:** Tweedley, Karen P 08/12/2016
Disposition **Disposition** **Disposition**
Reason: NAI By Examining District **Authorized By:** Tweedley, Karen P **Authorized Date:** 08/12/2016

Performing Org	PAC	LID	PAF	Compliance No	Lab Class-Description	Laboratory Status
ACNA-N	71R801		NAR		1 - In Compliance	Completed

Lab Conclusion

Sample Narrative - Method: AccQTag AAA Analysis - Taurine
Amt Found - 0.156% (dry matter basis)
Meets AAFCO minimum requirement of 0.10%

Sample Narrative - Method: Instrument Manual (Denver Instrumentst IR60)/ AOAC 930.15 Analysis - Moisture
Amt Found - 1.99%
Amt Declared - 11.00% max

Lab Conclusion Date **Lab Conclusion Made By**
08/04/2016 Hawes,Brian M

Food and Drug Administration Office of Regulatory Affairs

Summary Report

For Sample Number: 958504

TD Sample Number:

Import Sample Number

This is an accurate reproduction of the original electronic record as of 08/24/2016

Sample Class: Normal Everyday Sample **Sample Origin:** Domestic **Sample Basis:** Surveillance
Sample Flag: **Sample Type:** Official **Collecting District:** NWJ-DO
Home District: **Orig C/R and Records To:** NWJ-DO **Collection PACs:** 71R801

Product Name: Poultry Prod Pet Cat Food; Not Elsewhere Classified (NEC); Packaged Food (Not Commercially Sterile)

Product Description: See Remarks Section.

Collection Reason: Sample collected per FACTS Assignment ID #11650647 and OP ID # 8660426 referencing Consumer Complaint #146048 reporting the illness of multiple cats from the same household. Sample testing request: Taurine.

Lab: SRL **Split Num:**0 **Date Received:** 06/29/2016 **Date Out of Lab:** 08/04/2016
District **District Conclusion** **District**
Conclusion: No Action Indicated (NAI) **Made By:** Ciaccia, Andrew 08/17/2016
Disposition **Disposition** **Disposition**
Reason: NAI By Home District **Authorized By:** Ciaccia, Andrew **Authorized Date:** 08/17/2016

Performing Org	PAC	LID	PAF	Compliance No	Lab Class-Description	Laboratory Status
ACNA-N	71R801		NAR		1 - In Compliance	Completed

Lab Conclusion

Sample Narrative - Method: AccQTag AAA Waters Analysis - Taurine
Amt Found - 0.176% (dry matter basis)
Meets AAFCO minimum requirement of 0.10%

Sample Narrative - Method: Instrument Manual (Denver Instrumentst IR60)/AOAC 930.15 Analysis - Moisture
Amt Found - 2.79%
Amt Declared - 11.00% max

Lab Conclusion Date **Lab Conclusion Made By**
08/04/2016 Hawes,Brian M

From: [Lambkin, Sonya](#)
To: [Hartogenesis, Martine](#); [Wittig, Julianna](#); [Conway, Charlotte](#); [McCoig, Amber](#)
Subject: RE: FYSA - Pet Foods Agenda
Date: Thursday, March 14, 2019 10:03:03 AM

Hello, thanks for the call this morning -

(b) (5)

Thanks,
Sonya

From: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>
Date: March 14, 2019 at 8:18:23 AM EDT
To: Wittig, Julianna <Julianna.Wittig@fda.hhs.gov>, Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>, Carey, Lauren <Lauren.Carey@fda.hhs.gov>, Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>, Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>, Lambkin, Sonya <Sonya.Lambkin@fda.hhs.gov>, Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>
Cc: Ceric, Olgica <Olgica.Ceric@fda.hhs.gov>, Reimschuessel, Renate <Renate.Reimschuessel@fda.hhs.gov>, Duggirala, Heshia Jani <Heshia.Duggirala@fda.hhs.gov>, McCoig, Amber <Amber.McCoig@fda.hhs.gov>, Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>
Subject: RE: FYSA - Pet Foods Agenda

Yes, it shouldn't take up too much time.

Martine

From: Wittig, Julianna
Sent: Thursday, March 14, 2019 8:17 AM
To: Hartogenesis, Martine <Martine.Hartogenesis@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>; Lambkin, Sonya <Sonya.Lambkin@fda.hhs.gov>; Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>
Cc: Ceric, Olgica <Olgica.Ceric@fda.hhs.gov>; Reimschuessel, Renate <Renate.Reimschuessel@fda.hhs.gov>; Duggirala, Heshia Jani <Heshia.Duggirala@fda.hhs.gov>; McCoig, Amber <Amber.McCoig@fda.hhs.gov>; Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>
Subject: RE: FYSA - Pet Foods Agenda

Okay. I was thinking that too based on some feed back re purpose of this meeting. Has it changed at all since we last spoke? More like a reminder?

From: Hartogenesis, Martine

Sent: Thursday, March 14, 2019 8:16 AM

To: Wittig, Julianna <Julianna.Wittig@fda.hhs.gov>; Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>; Lambkin, Sonya <Sonya.Lambkin@fda.hhs.gov>; Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>

Cc: Ceric, Olgica <Olgica.Ceric@fda.hhs.gov>; Reimschuessel, Renate <Renate.Reimschuessel@fda.hhs.gov>; Duggirala, Heshu Jani <Heshu.Duggirala@fda.hhs.gov>; McCoig, Amber <Amber.McCoig@fda.hhs.gov>; Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>

Subject: RE: FYSA - Pet Foods Agenda

Thanks Julianna!

[Redacted] (b) (5)

Thanks in advance!

Martine

From: Wittig, Julianna

Sent: Wednesday, March 13, 2019 5:52 PM

To: Palmer, Lee Anne <LeeAnne.Palmer@fda.hhs.gov>; Carey, Lauren <Lauren.Carey@fda.hhs.gov>; Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov>; Peloquin, Sarah <Sarah.Peloquin@fda.hhs.gov>; Lambkin, Sonya <Sonya.Lambkin@fda.hhs.gov>; Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>

Cc: Ceric, Olgica <Olgica.Ceric@fda.hhs.gov>; Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>; Reimschuessel, Renate <Renate.Reimschuessel@fda.hhs.gov>; Duggirala, Heshu Jani <Heshu.Duggirala@fda.hhs.gov>; McCoig, Amber <Amber.McCoig@fda.hhs.gov>

Subject: FYSA - Pet Foods Agenda

Hi all,

So we are all on the same page, team mbrs shared time frames:

(b) (5)

Thanks,
J

From: [Reimschuessel, Renate](#)
To: [Jones, Jennifer L](#); [Rotstein, David](#); [Palmer, Lee Anne](#); [Carey, Lauren](#); [Queen, Jackie L](#)
Cc: [Ceric, Olgica](#)
Subject: RE: Head's up-potential DCM case-Dr. Adin-NCSU-2 Schnauzers
Date: Tuesday, July 11, 2017 11:50:53 AM
Attachments: [image001.png](#)
[image002.png](#)

(b) (5)

Renate Reimschuessel V.M.D. Ph.D. Vet-LIRN

Phone 1-240-402-5404

Fax 301-210-4685

<http://www.fda.gov/AnimalVeterinary/ScienceResearch/ucm247334.htm>

From: Jones, Jennifer L
Sent: Tuesday, July 11, 2017 11:38 AM
To: Rotstein, David; Palmer, Lee Anne; Carey, Lauren; Queen, Jackie L
Cc: Ceric, Olgica; Reimschuessel, Renate
Subject: Head's up-potential DCM case-Dr. Adin-NCSU-2 Schnauzers

Vet will submit PFR online →

2 dogs-unrelated miniature schnauzers

Dog 1: 2 yr → presented 2/2017 with fulminant CHF → severe DCM on echo, taurine/carnitine normal, infectious disease testing negative, died on the ventilator, necropsy done-myocardial changes were subtle but could be similar to moldy corn toxicity in pigs → plasma, urine, serum, and myocardial tissue available

Dog 2: 7 yr, had a syncopal episode ~2/2017 but presented to vet for progressive frequency of syncopal episodes → 6/2017 for CHF, diagnosed with DCM similar to housemate, nearly same image on Echo, taurine/carnitine normal, infectious disease testing negative, they have changed the diet (Hill's) and dog is responding to treatment; plasma, urine, and serum available

Dogs were eating California Naturals (different bag than from 2/2017) and treats (Milo's Kitchen); Vet has samples of food and treats

Jennifer L. A. Jones, DVM

Veterinary Medical Officer

U.S. Food & Drug Administration

Center for Veterinary Medicine

Office of Research

Veterinary Laboratory Investigation and Response Network (Vet-LIRN)

8401 Muirkirk Road, G704

Laurel, Maryland 20708

new tel: 240-402-5421

fax: 301-210-4685

e-mail: jennifer.jones@fda.hhs.gov

Web: <http://www.fda.gov/AnimalVeterinary/ScienceResearch/ucm247334.htm>



From: [Jones, Jennifer L](#)
To: "Darcy Adin"
Cc: [Ceric, Olgica](#)
Subject: RE: Pet food concern
Date: Wednesday, July 12, 2017 7:02:00 AM
Attachments: [image001.png](#)
[image003.png](#)
[image004.png](#)

Thank you, Darcy.

We received the complaints on our end, and will be in touch about next steps.

Kind regards,

Jennifer

Jennifer Jones, DVM
Veterinary Medical Officer



From: Darcy Adin [mailto:dbadin@ncsu.edu]
Sent: Tuesday, July 11, 2017 5:44 PM
To: Jones, Jennifer L
Subject: Re: Pet food concern

Hi Jennifer,

I've submitted the reports through the portal - one for each dog. The numbers are:

2023230 (I) for (b) (6)
2023228 (I) for [REDACTED]

I've also attached the visit summaries for (b) (6) (2) and (b) (5) (1) as well as (b) (6) necropsy report. I have the biological samples stored at -80 and also have food samples.

Thank you so much for your help and I'll look forward to hearing from you or someone on your team!

Take care

Darcy

On Tue, Jul 11, 2017 at 7:33 AM, Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov> wrote:

Hi Darcy,

I can chat today from [REDACTED] (b) (5)

Jen

Jennifer Jones, DVM
Veterinary Medical Officer



From: Darcy Adin [mailto:dbadin@ncsu.edu]
Sent: Monday, July 10, 2017 6:47 PM
To: Jones, Jennifer L
Cc: Reimschuessel, Renate; (b) (6)

Subject: Re: Pet food concern

Thank you Dr. Jones! I'm sorry I am just reading email now so I have missed you! I'd love to chat with you about the cases. I am on clinics this week but can try to call you if you have another block of time tomorrow (or later this week). My number is [614-582-9798](tel:614-582-9798) or I can be paged from (b) (6).

Thank you!

Darcy

On Jul 10, 2017, at 1:05 PM, Jones, Jennifer L <Jennifer.Jones@fda.hhs.gov> wrote:

Hello Dr. Adin,

Please let me know if you'd like to chat about the case this afternoon. I'll be in the office from 1-3pm (tel: [240-402-5421](tel:240-402-5421)).

If you suspect an animal's illness may be due to the food, you can submit a report at www.safetyreporting.hhs.gov

Please mention Vet-LIRN encouraged you to submit a report. Please email me the ICSR number (similar to a confirmation number), so I can find the report on my end.

Thank you,

Jennifer

Jennifer Jones, DVM

Veterinary Medical Officer

<[image001.png](#)> <[image004.png](#)>

From: Darcy Adin [mailto:dbadin@ncsu.edu]
Sent: Monday, July 10, 2017 11:31 AM
To: (b) (6)
Cc: Jones, Jennifer L; Reimschuessel, Renate
Subject: Re: Pet food concern

Thank you! I will work on this submission later today. I appreciate your help!

On Mon, Jul 10, 2017 at 10:49 AM, (b) (6) <[\(b\) \(6\)@ncsu.edu](mailto:(b) (6)@ncsu.edu)> wrote:

Hi Dr. Adin,

As we discussed this morning, (b) (5) is the FDA program. Here is a website that highlights how you can report a complaint.

<https://www.fda.gov/AnimalVeterinary/SafetyHealth/ReportaProblem/ucm182403.htm>

We work with the FDA Vet-LIRN program on diagnostics from the pet side, but

they agree to include the case in the program and would coordinate with us (or another laboratory). I have copied Dr. Jones and Dr. Reimschuessel here - they can help let us know the process to see if these cases are eligible.

Regards,

(b) (6)

(b) (6)

--

Darcy B. Adin, DVM, DACVIM (Cardiology)
Clinical Assistant Professor of Cardiology
North Carolina State University
NC State Veterinary Hospital
1060 William Moore Drive
Raleigh, NC 27607
[919-513-6032](tel:919-513-6032)

--

Darcy B. Adin, DVM, DACVIM (Cardiology)
Clinical Assistant Professor of Cardiology
North Carolina State University
NC State Veterinary Hospital
1060 William Moore Drive
Raleigh, NC 27607
919-513-6032

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6712303>

Taurine and Carnitine in Canine Cardiomyopathy

Article in *Veterinary Clinics of North America Small Animal Practice* December 2006

DOI: 10.1016/j.cvsm.2006.08.010 Source: PubMed

CITATIONS

13

READS

1,126

1 author:



[Sherry L Sanderson](#)

University of Georgia

26 PUBLICATIONS 309 CITATIONS

SEE PROFILE



Taurine and Carnitine in Canine Cardiomyopathy

Sherry Lynn Sanderson, DVM, PhD

Department of Physiology and Pharmacology, University of Georgia, College of Veterinary Medicine, 501 DW Brooks Drive, Athens, GA 30602, USA

Dilated cardiomyopathy (DCM) is one of the most common acquired cardiovascular diseases in dogs [1–4]. Although few studies of the prevalence of DCM in the overall population of dogs have been reported, estimates range from 0.5% to 1.1% [5,6]. Only degenerative valvular disease and, in some regions of the world, heartworm infection are more common causes of cardiac morbidity and mortality in dogs. DCM is seen most commonly in large and giant breeds of dogs, although its frequency seems to be increasing in medium sized breeds, such as the English and American cocker spaniels [4–8]. It has been reported rarely in small and miniature breeds of dogs [9].

DCM is particularly challenging to veterinarians because the cause is often unknown and can vary among dog breeds [10]. Because most cases of DCM in dogs are classified as idiopathic, most therapies can be classified as “Band Aid therapies” that palliate the effects of this disease for a short duration but do little to address the primary disease process. Therefore, DCM is almost always a progressive disease, and most dogs will eventually succumb to their disease. Survival times in dogs with DCM are variable and can be influenced by several factors, including breed. However, the prognosis for survival of dogs with DCM remains poor, with reported survival rates of 17.5% at 1 year and 7.5% at 2 years [11–13]. Until recently, reported cases of DCM reversal in dogs were very rare.

With advancements in echocardiology, diagnostic capabilities in canine cardiology have improved dramatically over the past 2 decades. Therapeutic advances have made surprisingly little progress. Symptomatic treatment is the standard care and outcome remains poor.

Recently, more promising therapies for dogs with DCM have resulted from a clearer understanding of the importance of biochemistry and nutrition in managing this disease. Nutrition is now widely accepted as an important adjunct to medical therapy in dogs with DCM.

E-mail address: sanderso@vet.uga.edu

The importance of nutrition in managing DCM has changed dramatically in the past 10 to 15 years. Historically, dietary sodium restriction was the most common nutritional recommendation for dogs with DCM. The importance of other nutrients in the origin and management of this disease was largely unknown. More recently, widely accepted beliefs about the role nutrient deficiencies could play in DCM have been proven false, further enhancing the ability to direct therapy at an underlying cause rather than just the symptoms.

This article focuses on two nutrients, taurine and carnitine, that play an important role in the cause and treatment of DCM in some dogs. Known risk factors for developing deficiencies of these nutrients are discussed, along with the use of taurine and carnitine for treating DCM in dogs.

TAURINE

What is Taurine?

Taurine is a sulfur containing amino acid. Unlike most other amino acids, taurine is not incorporated into proteins but rather is one of the most abundant free amino acids in the body. Taurine is found in highest tissue concentrations in cardiac muscle, skeletal muscle, the central nervous system, and platelets [14].

Other than conjugation of bile acids and detoxification of xenobiotics through conjugation and excretion in bile, the function of taurine in mammals is not well understood but is highly diverse [14,15]. Since the mid 1970s, taurine has been known to be essential for normal retinal function in cats [16]. In addition, clinical and experimental evidence collected in the late 1980s documented that taurine is essential for normal myocardial function [17-20].

Taurine is involved with numerous metabolic processes, including antioxidant, retinal photoreceptor activity, development of the nervous systems, stabilization of neural membranes, reduction in platelet aggregation, and reproduction [15,16,21-26]. Although the importance of taurine for normal myocardial function is also well recognized, the mechanisms underlying its effect on the heart remain unknown. Much of the available evidence supports the theory that taurine's major effect on cellular function in the heart is modulating tissue calcium concentrations and availability [14,27,28]. In addition, taurine may inactivate free radicals and protect the heart by changing cellular osmolality [29]. Taurine may also have an effect on osmoregulation in the myocardium. Taurine is a small but highly charged osmotically active molecule, and experts have proposed that alterations in cellular osmolality induced by changes in intracellular taurine concentration are a protective mechanism in nervous tissue and myocardium [29]. Other proposed mechanisms specifically related to myocardial function include *N*-methylation of cell membrane phospholipids [30], direct effects on contractile proteins [31,32], and interactions with the renin-angiotensin-aldosterone system [33]. Taurine is a natural antagonist of angiotension II.

Is Taurine an Essential Amino Acid in Dogs?

Taurine is an essential amino acid in cats, and it is well known that taurine deficiency can cause DCM, retinal degeneration, and reproductive anomalies in this species [18]. However, taurine is not considered an essential amino acid in dogs. One explanation for the differences in taurine requirements between cats and dogs is that the activity of cysteine sulfinic acid decarboxylase (the rate limiting enzyme in the synthesis of taurine from cysteine and methionine) is higher in dogs than cats [34]. However, the difference in activity of this enzyme between dogs and cats does not fully explain the difference in requirements. The activity of this enzyme in humans is even lower than in cats, and taurine is not considered an essential amino acid in healthy adult humans. Therefore, cats and dogs may have additional differences that may explain why taurine is an essential amino acid in cats and not in dogs.

A study in dogs conducted in the 1980s at the University of California at Davis showed that feeding taurine free diets or diets found to be taurine depleting in cats [35] did not result in taurine depletion when fed to a group of eight healthy beagles [36]. In addition, results of an early clinical study in dogs, also conducted at this University soon after the relationship between taurine deficiency and DCM was discovered in cats, were unrewarding. These studies showed that dogs could not become taurine depleted from diet alone, and that taurine did not play a considerable role in the development of DCM in dogs.

Emergence of Taurine Deficiency in Dogs with Dilated Cardiomyopathy

The belief that taurine deficiency could not cause DCM in dogs was challenged in 1989 when taurine deficiency was linked to DCM in foxes [37]. This study reopened taurine's possible role in DCM in dogs, and a collaborative study between the University of California at Davis and the Animal Medical Center in New York City was initiated [38]. In this study, plasma taurine levels were evaluated in dogs with DCM and in those with chronic degenerative mitral valve disease. Surprisingly, results of this study showed that plasma taurine concentration was low in 17% of 75 dogs with DCM, and this deficiency occurred in breeds not commonly afflicted with DCM, such as American cocker spaniels and golden retrievers. However, because the plasma taurine concentration in breeds more commonly affected with DCM were within the reference range, experts concluded that taurine deficiency was unlikely to play an important role in the etiopathogenesis or therapy of DCM in dogs.

Multicenter Spaniel Trial (MUST) Study

Anecdotal reports emerged regarding supplementing American cocker spaniels diagnosed with DCM with taurine; however, initial reports of taurine supplementation were unrewarding. When Kittelson and colleagues [8] gave taurine and L carnitine supplements to two American cocker spaniels with DCM, both dogs experienced response. These findings initiated the Multicenter Spaniel Trial (MUST) study. In this study, baseline plasma taurine concentrations and echocardiograms were collected in 11 American cocker spaniels diagnosed

with DCM. All dogs were found to have low plasma taurine concentrations at baseline (<50 nmol/mL). After baseline information was collected, dogs were randomly assigned to receive supplementation with both taurine (500 mg by mouth every 8 hours) and *L* carnitine (1000 mg by mouth every 8 hours) or a placebo for 4 months, and echocardiograms were reevaluated after 2 and 4 months of therapy. The group supplemented with both taurine and carnitine showed significant echocardiographic improvement, whereas dogs receiving the placebo did not.

After this initial 4 month period, dogs that had received the placebo initially received supplements of both taurine and carnitine, and subsequently showed echocardiographic improvement after 2 to 4 months of therapy. The magnitude of echocardiographic improvement in the American cocker spaniels was not as dramatic as that seen after taurine supplementation in cats with taurine deficiency DCM. Nonetheless, after 4 months of supplementation, the improvement in myocardial function in each dog was significant enough to allow discontinuation of cardiovascular drug therapy. Improvements were seen in not only cardiovascular function but also survival times. The mean survival time for dogs in this study was 28.3 ± 19.1 months, compared with an average life expectancy for dogs treated with conventional drug therapy of approximately 6 months. Based on results from this study, the current recommendation is to supplement American cocker spaniels diagnosed with DCM with both taurine and carnitine at the doses mentioned earlier.

University of Minnesota Study in Urolith-forming Dogs Diagnosed with Dilated Cardiomyopathy

Around the same time the MUST study was initiated, a separate clinical study was initiated at the University of Minnesota. The population of dogs studied consisted of those with either cystine or urate urolithiasis that developed DCM after long term consumption of a protein restricted diet that was being used to manage their stone disease (Sherry L. Sanderson, DVM, PhD, unpublished data, 1998). Dogs in group 1 underwent only conventional drug therapy for their heart disease, whereas those in group 2 underwent taurine and/or carnitine supplementation in addition to conventional drug therapy as needed. Dogs in group 1 that were in Modified New York Heart Association (MNYHA) functional class I and II heart failure received enalapril (0.25 mg/kg by mouth every 12 hours) and digoxin (0.01–0.02 mg/kg by mouth divided twice a day), and dogs in MNYHA functional class III and IV received furosemide (dose varied depending on severity of heart disease) in addition to enalapril and digoxin. The population of dogs in group 1 (N = 6) consisted of five English bulldogs (four with cystine urolithiasis, one with urate urolithiasis) and one Dalmatian with urate urolithiasis. The population of dogs in Group 2 (N = 8) consisted of five English bulldogs (three with cystine urolithiasis, two with urate urolithiasis), two Dalmatians with urate urolithiasis, and one miniature Dachshund with cystine urolithiasis. Because when this study was initiated experts believed that dogs with DCM did not have low plasma taurine

concentrations, none of the dogs in group 1 had these concentrations evaluated at baseline. Plasma taurine concentrations evaluated before supplementation in seven of eight dogs in group 2 ranged from 2 nmol/mL to 45 nmol/mL (mean, 20.9 nmol/mL). These results were below the reference range of 41 nmol/mL to 97 nmol/mL that the investigators established from healthy adult beagles. Echocardiography was performed at baseline and once every 2 months. Details from this study will be published later, but a few interesting and important results were noted:

1. The average life expectancy for dogs in group 1 was 10.5 months, and all dogs were euthanized because of progressive congestive heart failure that became refractory to therapy. The average life expectancy for dogs in group 2 was 47.1 months, and only three of eight dogs were euthanized because of progressive congestive heart failure. In addition, three of five dogs that did not succumb to their heart disease received only taurine and/or carnitine supplementation and no conventional drug therapy for the management of their heart disease.
2. DCM reversed in three of eight dogs in group 2. DCM returned in one dog after the owner discontinued taurine and carnitine supplementation on their own, and in an additional dog when the dose of carnitine was reduced because of diarrhea associated with carnitine supplementation.
3. Dogs consuming a protein-restricted diet long-term could develop taurine deficiency, in contrast to results from previous studies that concluded that a diet could not induce taurine deficiency in dogs. This finding provided an impetus for further examining the effects on plasma and whole blood taurine levels in healthy adult dogs consuming a protein-restricted diet long-term.

Diet-Induced Taurine Deficiency in Healthy Adult Dogs

Previous reports indicated that dogs could not develop diet induced taurine deficiency, even when fed a diet devoid of taurine. However, based on the finding of University of Minnesota study that dogs developed low plasma taurine levels after consuming a protein restricted diet long term, a more controlled study was undertaken to determine the cause of this problem and evaluate the effects of long term taurine deficiency on cardiac function in healthy adult dogs [39].

This study involved 17 healthy adult beagles. Baseline plasma and whole blood taurine levels were evaluated, and echocardiography was performed to assess cardiac function. Once baseline data was collected, dogs were fed one of three protein restricted diets for 48 months. All three diets had similar levels of protein; one diet was also low in fat, a second was high in fat, and a third was high in fat and supplemented with *L* carnitine at 200 mg/kg of diet. All diets contained methionine and cystine concentrations at or above recommended minimum requirements established by the Association of American Feed Control Officials (AAFCO) [40]. After diet assignment, plasma taurine and whole blood taurine concentrations and echocardiography were evaluated every 6 months.

All three dietary treatments caused a significant decrease in whole blood taurine concentration compared with baseline concentrations. Dogs in the high fat

group also experienced a significant decrease in plasma taurine concentration. This study was the first to show that diet could induce taurine deficiency in healthy adult dogs, in contrast to previous studies.

Another important observation was that one dog with taurine deficiency developed DCM, and that taurine supplementation resulted in almost complete reversal of the disease. This study was also the first to clearly document in dogs that taurine deficiency preceded DCM, and that taurine supplementation resulted in substantially improved cardiac function, similar to cats.

Why Did Dogs Develop Taurine Deficiency While Consuming a Protein-Restricted Diet?

The exact mechanism for this problem is unknown. However, this study showed that the AAFCO recommended minimum requirements for amino acids may need to be modified in dogs consuming a protein restricted diet long term. Many therapeutic diets for dogs are now supplemented with taurine.

Additional Examples of Diet-Induced Taurine Deficiency in Dogs

Soybean based diets

Taurine deficiency was identified in two unrelated dogs fed a tofu based diet [41]. Although the diet was low in protein, it met the National Research Council's published requirements for protein and other nutrients in dogs [42]. The authors attributed taurine deficiency to the fact that the primary protein source was soybean curd, which is low in sulfur containing amino acids and devoid of taurine compared with meat proteins [43]. In addition, soybean curd has been shown to accelerate the loss of bile acids in cats [44].

Lamb meal and rice diets

Taurine deficiency was also identified in 12 Newfoundlands consuming two different commercially available lamb meal and rice diets [41]. Echocardiography was performed in six of the dogs, and none were diagnosed with DCM. The taurine deficiency was reversed when the diet was either changed or when the lamb meal and rice diets were supplemented with methionine. This study did not identify the exact mechanism for the development of taurine deficiency in the dogs consuming the lamb meal and rice diets.

In a study by Fascetti and colleagues [45], DCM and taurine deficiency were identified in 12 large and giant breed dogs consuming commercially available diets that contained lamb meal, rice, or both as primary ingredients. All dogs received supplements of with taurine (1000 3000 mg by mouth every 24 hours), and significant echocardiographic improvement occurred in 9 of the 12 dogs that underwent an echocardiogram repeated after taurine supplementation. The authors hypothesized that taurine deficiency caused DCM and was caused by inadequate or unavailable dietary sulfur amino acids, which are essential precursors of taurine synthesis.

In a similar report, five related golden retrievers were diagnosed with taurine deficiency and DCM [46]. Three of five dogs were consuming lamb meal and rice or lamb and rice diets. All showed significant improvement after taurine

supplementation (500 mg by mouth every 12 hours), and all five dogs survived for more than 3 years. The authors attribute the DCM to a suspected autosomal recessive mode of inheritance; however, the potential role diet played in the development of taurine deficiency warrants mentioning.

Potential Causes of Taurine Deficiency in Dogs Consuming Lamb Meal and Rice or Lamb and Rice Diets

Torres and colleagues [47] compared the effects of consuming a lamb meal and rice based diet with effects of consuming a poultry by product based diet in 12 beagles aged 5 to 5.5 months. Although the differences in plasma and whole blood taurine concentrations did not differ among diet groups, dogs consuming the lamb meal and rice based diet excreted less taurine in their urine than dogs consuming the poultry by product based diet. When the lamb meal and rice diet was supplemented with methionine, urinary taurine excretion increased by 54%. Because taurine homeostasis in dogs is achieved primarily through regulating renal taurine excretion, the amount of taurine excreted in urine is a sensitive indicator of the adequacy of either taurine synthesis or absorption of dietary precursor amino acids. The authors concluded that reduced bioavailability of sulfur amino acids in the lamb meal and rice diet is a likely cause of taurine deficiency. This finding is supported by the increase in urine taurine concentrations after supplementation with methionine. Johnson and colleagues [48] showed that ileal digestibility of amino acids in dogs depends on the raw material sources and the temperature used to process feeds and provides a mechanism for these specific dietary effects.

A second potential, although related, cause of taurine deficiency in dogs consuming lamb meal and rice diets was proposed [49,50]. When dietary protein is low in quality, undigested protein reaches the colon, where it serves as a substrate for bacterial growth. Some bacteria produce cholytaurine hydrolase, an enzyme that causes release of taurine from taurocholic and other bile acids that are normally conserved in the enterohepatic circulation, resulting in increased fecal loss of taurine. Studies in dogs [49] and cats [50] have found that diets containing rice bran and whole rice products provide a source of moderately fermentable fiber and high amounts of fat. These fermentable fibers may increase the number of bacteria in the colon and result in a greater loss of taurine in the feces similar to the mechanism for undigested protein. The fat content of the diet can also affect taurine metabolism through altering intestinal bacteria and subsequent changes in the excretion of bile acids.

How Should Samples be Collected to Evaluate Plasma and Whole Blood Taurine Concentrations?

Fasting versus postprandial blood samples

Although fasting has no effect on plasma taurine concentrations in humans [51], food deprivation causes a small but significant reduction in plasma taurine concentrations in cats [52]. In a study by Torres and colleagues [47], plasma taurine concentrations were significantly reduced in food restricted dogs compared with ad libitum fed dogs. Whole blood taurine concentrations were

also reduced, although the whole blood taurine results were not statistically significant between the two groups. Because of the potential for food intake to affect plasma and whole blood taurine concentrations in dogs, withholding food, but not water, is recommended for 8 hours before sampling.

Anticoagulant used for plasma sample collection

Paired analysis of samples comparing taurine concentrations in plasma collected in lithium heparin with those collected in sodium citrate showed that plasma taurine concentrations are higher when lithium heparin is used as the anticoagulant [38]. Because most studies have used heparinized plasma samples to evaluate plasma taurine levels in dogs, these are recommended rather than sodium citrate plasma samples.

Plasma taurine sample collection

Heparinized, nonhemolyzed blood samples should be obtained and stored on ice until they are processed. After centrifuging, the plasma should be separated immediately from the cellular components, and a small amount of plasma should be left above the buffy coat to prevent contamination of the plasma with cells. Hemolysis and platelet or white blood cell contamination falsely elevates plasma taurine concentrations. Samples should be frozen until analyzed for plasma taurine concentrations.

Whole blood taurine sample collection

Heparinized whole blood should be frozen until samples can be analyzed. Because the red blood cells are lysed before analysis, hemolyzed samples do not adversely affect whole blood taurine analysis.

Plasma and whole blood taurine samples can be sent to the Department of Molecular Biosciences at the School of Veterinary Medicine, University of California, Davis, for analysis.

Which is Better: Plasma Taurine Concentrations or Whole Blood Taurine Concentrations

Earlier studies evaluating the relationship between taurine deficiency and DCM in dogs relied primarily on plasma taurine concentrations to predict tissue taurine concentrations. Studies conducted in dogs by this author showed findings similar to those reported in cats [53]. Relying on plasma taurine concentrations alone does not reliably assess tissue taurine concentrations in dogs. Simultaneously evaluating plasma and whole blood taurine concentrations predicts skeletal and cardiac muscle taurine concentrations better than evaluating either test alone. Therefore, when evaluating taurine status in dogs with DCM, plasma and whole blood taurine concentrations should be assessed simultaneously.

Reference Ranges for Plasma and Whole Blood Taurine Concentrations in Dogs

The reference range used in earlier studies evaluating plasma and whole blood taurine concentrations in dogs was extrapolated from the reference range use in

cats. However, reference ranges for plasma and whole blood taurine concentrations in dogs were published recently (Table 1).

Delaney and colleagues [49] have also suggested that plasma taurine concentrations less than 40 nmol/mL are critically low, as are whole blood taurine concentrations less than 150 nmol/mL. In addition, Sanderson and colleagues [53] found that low plasma taurine concentrations can exist without the presence of DCM.

Therefore, results showed that the onset of clinical signs in dogs, just as in cats, was variable when taurine concentrations declined markedly below the normal range [18].

Which Dogs Diagnosed with Dilated Cardiomyopathy Should Receive Taurine Supplementation?

Evaluation of plasma and whole blood taurine concentrations is recommended for all dogs diagnosed with DCM. An association between taurine deficiency and DCM was found in various breeds of dogs, including American cocker spaniels, Newfoundlands, golden retrievers, Labrador retrievers, Dalmatians, English bulldogs, and Portuguese water dogs. Taurine supplementation is highly recommended in any of these breeds that develop DCM.

Not all dogs with DCM will show dramatic improvement with taurine supplementation. However, even if plasma and whole blood taurine concentrations are within the reference range, giving taurine supplements to dogs diagnosed with DCM may still have some benefits. Because taurine is extremely safe and inexpensive, the risks and costs of supplementation are minimal, even if dogs have normal levels of plasma and whole blood. Proposed mechanisms for the beneficial actions of taurine on the myocardium include modulating tissue calcium concentrations and availability in the heart; inactivating free radicals and protecting the heart through altering cellular osmolality; osmoregulating the myocardium; directly affecting contractile proteins; and serving as a natural antagonist of angiotension II. Dogs with DCM that do not have taurine deficiency may still benefit from some of these proposed mechanisms of action for taurine.

Table 1
Normal concentrations of taurine in dogs

Plasma (nmol/mL)	Whole blood (nmol/mL)
41–97 ^a	155–347 ^a
72.8–81.2 ^b	255.8–276.2 ^b

^aReference range established from 18 healthy adult beagles consuming a canned commercial maintenance diet. *Data from* Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein restricted diets. *Am J Vet Res* 2001;62:1616–23.

^bReference range established from 131 healthy adult dogs of various breeds consuming a variety of commercial adult maintenance diets. *Data from* Delaney SJ, Kass PH, Rogers QR, et al. Plasma and whole blood taurine in normal dogs of varying size fed commercially prepared food. *J Anim Physiol* 2003;87:236–44.

Recommended dose for taurine supplementation

This author has successfully used doses of 500 to 1000 mg of taurine administered orally two to three times per day for small dogs (<25 kg), and 1 to 2 g of taurine administered orally two to three times per day for large dogs (25–40 kg). These doses have been shown to normalize plasma and whole blood taurine levels in taurine deficient dogs. Many other doses for taurine are reported in the literature. Whether a smaller or less frequent dose of taurine than what this author recommends can be used successfully remains to be determined. If doses are used that differ from those this author recommends, plasma and whole blood taurine concentrations must be reevaluated after taurine supplementation is initiated to determine if the dose being given is effective and appropriate. Another important point is that echocardiographic improvement in myocardial function is not usually documented before 2 months of supplementation, and often no improvement is documented before 4 months of supplementation. However, the dogs may feel better clinically and be more active before improvement in cardiac function is documented. Owners must not withdraw taurine supplementation prematurely before deciding if their dogs benefit.

Where Can Taurine be Purchased?

Taurine can be purchased through several retail outlets. If taurine is purchased through a health food store, consumers must look for a product that contains a USP certification symbol on the label. This symbol ensures that what is listed on the label is exactly what is found in the product.

LEVOCARNITINE (L-CARNITINE)

What is L-Carnitine?

L carnitine (β hydroxy γ trimethylaminobutyric acid) is a small water soluble molecule with a molecular weight of 160. In dogs, carnitine is obtained either from dietary protein or endogenous synthesis in the liver using the essential precursor amino acids lysine and methionine. Synthesis also requires iron, vitamin C, and vitamin B₆ as cofactors [54]. Although carnitine is classified as an amino acid derivative, it is not an α amino acid and the amino group is not free. Therefore carnitine is not used for protein synthesis [55].

Carnitine is found in the body either as free carnitine, short chain acyl carnitine, or long chain acylcarnitine. Acylcarnitine is carnitine bound to a fatty acid. Total carnitine is the sum of all the individual carnitine fractions. The free carnitine fraction is normally higher than either the short chain acylcarnitine fraction or the long chain acylcarnitine fraction.

Cardiac and skeletal muscles are significant storage sites, containing 95% to 98% of the carnitine in the body [56], and carnitine is concentrated in these tissues through an active membrane transport mechanism. The heart is unable to synthesize carnitine and depends on transport of carnitine from the circulation into cardiac muscle, which results in up to a 100 \times gradient between extracellular and intracellular concentrations.

Only the *L* form of carnitine exists naturally in the body. The *D* form competitively inhibits the actions of the *L* form, thereby inhibiting carnitine enzyme systems. In addition, mammals are unable to convert *D* carnitine to *L* carnitine, and therefore this discussion focuses on *L* carnitine.

Why is *L*-Carnitine Important for Normal Myocardial Function?

The normal heart obtains approximately 60% of its total energy production from oxidation of long chain fatty acids [57]. Long chain fatty acids in the cytosol of myocardial cells combine with coenzyme A (CoA) as the first step toward beta oxidation. However, long chain fatty acids must be transported across the inner mitochondrial membrane to generate energy, and the inner mitochondrial membrane is normally impermeable to such bulky polar molecules. Therefore, transport is accomplished through a “carnitine shuttle.” In the carnitine shuttle, the activated fatty acid in the cytosol reacts with carnitine to form a more permeable molecule. This reaction occurs on the outer surface of the inner mitochondrial membrane and is catalyzed by the enzyme carnitine acyl transferase I. The newly formed long chain acyl carnitine ester molecule is permeable to the inner mitochondrial membrane and is transported across this membrane, where the enzyme acyltransferase II converts the long chain acyl carnitine back to free carnitine and the long chain fatty acid. Therefore, carnitine functions as a cofactor of several important enzymes necessary for transport of long chain fatty acids from the cytosol into the mitochondrial matrix [58,59]. Once inside the mitochondria, fatty acids undergo beta oxidation to generate energy [60].

Another important function of carnitine is its buffering capacity, which modulates the intramitochondrial acyl CoA:CoA ratio [58]. This process is important because acyl CoA is the activated form of fatty acids used for beta oxidation and lipid synthesis. However, buildup of acyl CoA derivatives in the mitochondria results in decreased free CoA, which inhibits oxidative metabolism. Acyl CoA derivatives also act as detergents at high concentrations. Carnitine also facilitates removal of accumulating short and medium chain organic acids from the mitochondria. Therefore carnitine also has a role in detoxification in the mitochondria.

What Causes *L*-Carnitine Deficiency?

Carnitine deficiency can be a primary or secondary disorder. Primary carnitine deficiencies may arise from genetic defects in synthesis, renal transport, intestinal absorption, transmembrane uptake mechanisms, or excessive degradation of carnitine [61]. In humans, primary carnitine deficiencies have been associated with cardiomyopathies that are usually not present at birth but take 3 to 4 years to develop. *L* carnitine therapy can prevent and reverse cardiac dysfunction in some patients.

Secondary carnitine deficiencies are believed to be much more common in humans and can have many causes [61]. In humans, carnitine deficiency can result from inborn errors of metabolism or develop in patients undergoing long term total parenteral nutrition, vegetarians, and infants fed formulas not

supplemented with carnitine. Carnitine deficiencies are recognized in dogs, but the incidence is not known.

What are the Consequences of *L*-Carnitine Deficiency?

Carnitine deficiency has been shown to cause or be associated with DCM in humans [62–64], hamsters [65,66], and dogs [36,67–69]. More widespread studies have not been undertaken in dogs because carnitine status is difficult to thoroughly assess.

What Types of Carnitine Deficiency Exist in Dogs?

Carnitine deficiency in dogs is classified as either (1) plasma carnitine deficiency, characterized by low concentrations of free plasma carnitine; (2) systemic carnitine deficiency, characterized by low concentrations of free plasma and tissue carnitine; or (3) myopathic carnitine deficiency, characterized by low free myocardial carnitine concentrations in the presence of normal and sometimes elevated plasma carnitine concentrations. Plasma carnitine deficiency alone is not a well documented state and is included to account for the fact that plasma carnitine, but not tissue carnitine sampling, is often pursued in veterinary medicine.

For example, if plasma carnitine concentration is used to assess carnitine status of a dog, it can help diagnose carnitine deficiency when it is low. However, if plasma carnitine concentration is normal, it does not rule out the possibility of the myopathic form of carnitine deficiency, and the myopathic form of carnitine deficiency is estimated to occur in 17% to 60% of dogs with DCM. Evaluating cardiac muscle carnitine concentrations requires a fluoroscopy guided endomyocardial biopsy, which is not practical to perform in most private practice situations and is not without risk. Therefore, diagnosing and determining the incidence of myopathic carnitine deficiency in dogs with cardiac disease remains elusive, but may be an underdiagnosed cause of DCM in dogs.

L-Carnitine Deficiency and Associated Myocardial Disease States in Dogs

Carnitine deficiency was associated with DCM in dogs in a limited number of clinical reports [8,9,68–70]. The first reported case of carnitine deficiency was in a family of boxers [69]. The sire, dam, and two littermates were diagnosed with DCM. One offspring had a low plasma carnitine concentration and low myocardial carnitine concentration at DCM diagnosis. After undergoing treatment with high dose *L* carnitine (220 mg/kg/d orally), this dog's fractional shortening (FS) increased from 18% to 28%. This dog's littermate had low myocardial and normal plasma carnitine concentrations and responded similarly to high dose *L* carnitine supplementation, with its FS increasing from 2% to 24%. The latter dog experienced a decline in myocardial function after *L* carnitine therapy was withdrawn. Both parents of these littermates had normal plasma and low myocardial carnitine concentrations. Unfortunately, both parents died soon after beginning *L* carnitine supplementation.

Costa and Labuc [70] presented another case report of two boxers with DCM. One was treated with 250 mg/kg/d of *L* carnitine orally, and the other was not treated. The myocardial concentration of carnitine was found to be low in the dog that did not receive supplementation and elevated in the dog that did.

Concurrent supplementation with carnitine and taurine has shown benefit in American cocker spaniels with DCM [8]. An unpublished study by this author in 1998 showed beneficial effects from carnitine supplementation in urolith forming dogs diagnosed with DCM while consuming a protein restricted diet (Sherry Lynn Sanderson, DVM, PhD, unpublished material). Both studies showed dramatic improvement in myocardial function and survival times in dogs that received supplementation.

Which Came First: Carnitine Deficiency or Dilated Cardiomyopathy?

A common argument made against the role of carnitine deficiency in dogs diagnosed with DCM is that if carnitine deficiency is diagnosed after the onset of DCM, whether carnitine deficiency caused the DCM or DCM caused the carnitine deficiency is unclear. When myocardial cells are damaged, as may occur with DCM, carnitine can leak out of the cells, resulting in low myocardial carnitine levels. In this situation, the DCM caused the carnitine deficiency. Most published studies linking carnitine deficiency to DCM in dogs have shown this scenario when carnitine deficiency was diagnosed after the onset of DCM.

In an unpublished study conducted at the University of Minnesota, this author documented carnitine deficiency before the onset of DCM in three dogs (Sherry Lynn Sanderson, DVM, PhD, unpublished material, 1998). Therefore, the association of carnitine deficiency with DCM at diagnosis may not always imply a cause and effect relationship. However, this study indicates that carnitine deficiency can cause DCM in dogs.

Which Dogs with Dilated Cardiomyopathy Should Receive Carnitine Supplementation?

The importance of carnitine supplementation in the treatment and survival times of some dogs with DCM should not be overlooked. In the first reported study linking carnitine deficiency to DCM in boxers, two of four dogs experienced good response to carnitine supplementation [69]. Considering the generally poor prognosis of this disease in boxers, carnitine supplementation provides owners one additional option for treating this disease, and has made a dramatic difference in the survival times and quality of life of some dogs.

The importance of carnitine supplementation in American cocker spaniels with DCM and urolith forming dogs with DCM should also not be overlooked. Although a few anecdotal reports exist in which American cocker spaniels with DCM experienced good response to taurine supplementation alone, most cases have shown response to combined supplementation with taurine and carnitine. In the above study by this author, a miniature Dachshund diagnosed with carnitine deficiency before the onset of DCM underwent treatment

only with carnitine supplementation, and its heart disease reversed. Although DCM in many dogs is not associated with carnitine deficiency, carnitine and taurine supplementation offer the most promising hope for improved quality of life and survival times in dogs that experience response.

How is Carnitine Deficiency Diagnosed?

Because performing endomyocardial biopsies is impractical for most clinicians in private practice, most screening for carnitine deficiency relies solely on plasma carnitine levels. The method for plasma carnitine sample collection is almost identical to that used for plasma taurine sample collection. Fasting, heparinized, nonhemolyzed blood samples should be obtained and stored on ice until they are processed. The plasma should be immediately separated from the cellular components ideally in a cold centrifuge, and a small amount of plasma should be left above the buffy coat to prevent contamination of the plasma with cells. Samples should be frozen immediately until analyzed for plasma carnitine concentrations.

What is the Recommended Dose for Carnitine Supplementation in Dogs?

The doses of carnitine being administered may contribute to the lack of favorable results with carnitine supplementation that some investigators observed. The recommended doses for carnitine supplementation in dogs with DCM vary widely in the literature. Although most authors recommend a carnitine dose of 50 to 100 mg/kg orally every 8 hours, the effective dose may depend on the form of carnitine deficiency. In a limited number of cases studied at the University of Minnesota, where pre and post carnitine supplemented plasma and cardiac muscle carnitine levels were obtained, this author's clinical impression was that the effective therapeutic dose in dogs with systemic carnitine deficiency was much lower than the effective dose in dogs with myopathic carnitine deficiency.

Some experts speculate that the myopathic form of carnitine deficiency may be caused by a carnitine transport defect in the heart, and much higher plasma levels of carnitine seem to be needed to overcome this defect and achieve normal concentrations of carnitine in the heart than for the systemic form of carnitine deficiency. Based on this work, the dose of carnitine recommended by this author for systemic carnitine deficiency is 100 mg/kg orally every 8 hours. However, if the myopathic form of carnitine deficiency is present or suspected, the author recommends starting carnitine supplementation at 200 mg/kg orally every 8 hours to maximize the chances that carnitine supplementation will improve myocardial function.

Carnitine is a very safe substance. Diarrhea was the only adverse effect of high doses of carnitine, reported in approximately two thirds of dogs. If diarrhea occurs, the highest dose of carnitine that the dog will tolerate without causing diarrhea should be administered. Therefore, like taurine, *L* carnitine is a safe substance to administer, and, except for the expense, few drawbacks exist to supplementing a dog with DCM with carnitine (carnitine is much more expensive than taurine. Another important point is that the time it takes for

improvement in myocardial function to occur is very similar to that for taurine supplementation. Echocardiographic improvement in myocardial function is not usually documented before 2 months of supplementation with carnitine, and often improvement is not documented for up to 4 months. However, dogs may feel better clinically and be more active before improvement in cardiac function is documented. Owners must not withdraw carnitine supplementation prematurely before determining whether their dogs benefit.

Where Can *L*-Carnitine Be Purchased?

Although *L* carnitine can be purchased from health food stores, this source is extremely expensive. Purity of the sample is also of great importance. Therefore, only products that contain the USP certification seal should be purchased from health food stores.

L Carnitine can also be purchased less expensively in bulk. Bulk carnitine can be purchased from Ajinomotousa, Inc (500 Frank W Burr Boulevard; Park Central West; Teaneck, New Jersey). At last check, the company required a minimum purchase of 10 kg at one time. However, the individual expense can be reduced if several owners split an order. If carnitine is purchased in bulk, owners must measure out the carnitine they are giving to their dogs. One teaspoon of carnitine is equivalent to 2 g of carnitine. Therefore, fractions of a teaspoon can be administered if necessary. Owners must be sure to purchase *L* carnitine, not *D* or the *DL* isomers, because *D* carnitine interferes with *L* carnitine use.

Which Dogs with Dilated Cardiomyopathy Should be Supplemented With Carnitine?

Carnitine supplementation should be recommended for boxers, American cocker spaniels, and dogs with cystine or urate urolithiasis that are diagnosed with DCM. Even if carnitine deficiency did not cause DCM, supplementing dogs with carnitine does not hurt them, and supplementation may be beneficial even if carnitine deficiency is not present. The major drawback to supplementing dogs with carnitine is the expense and occasional gastrointestinal upset.

What are the Reference Ranges for Carnitine Concentrations in Dogs?

The reference ranges for carnitine concentrations in dogs are listed in Table 2 [69].

SUMMARY

Some newer more promising therapies for dogs with DCM do not involve drugs but rather nutritional supplements. Two of the more common nutritional supplements administered to dogs with DCM are taurine and carnitine. Deficiencies of these nutrients have been shown to cause DCM in dogs, and some breeds have been shown to experience dramatic improvement in myocardial function after supplementation with one or both nutrients. Although most dogs diagnosed with DCM do not have a documented taurine or carnitine deficiency, they may still benefit from supplementation. Both nutrients are very

Table 2
Normal concentrations of carnitine in dogs

Carnitine fraction	Plasma carnitine (nmol/mL)	Cardiac muscle carnitine (nmol/mg of NCP)
Free	8–36	4–11
Esterified	0–7	0–4
Total	12–38	5–13

Abbreviation: NCP, noncollagenous protein.

safe to administer to dogs. For some owners, the high cost of carnitine is the only deterrent to giving their dogs supplements of both nutrients.

References

- [1] Cobb MA. Idiopathic dilated cardiomyopathy: advances in etiology, pathogenesis and management. *J Small Anim Pract* 1992;33:112–8.
- [2] Keene BW. Canine cardiomyopathy. In: Kirk RW, editor. *Current veterinary therapy X. Small animal practice*. Philadelphia: WB Saunders Co; 1989. p. 240–51.
- [3] Sisson DD, Thomas WP, Keene BW. Primary myocardial disease in the dog. In: Ettinger SJ, Feldman EC, editors. *Textbook of veterinary internal medicine. Diseases of the dog and cat*. 5th edition. Philadelphia: WB Saunders Co; 2000. p. 874–95.
- [4] Buchanan JW. Causes and prevalence of cardiovascular disease. In: Kirk RW, Bonagura JD, editors. *Kirk's current veterinary therapy XI*. Philadelphia: WB Saunders Co; 1992. p. 647–55.
- [5] Sisson D, Thomas WP. Myocardial diseases of dogs and cats. In: Ettinger S, editor. *Textbook of veterinary internal medicine*. 4th edition. Philadelphia: WB Saunders; 1995. p. 995–1032.
- [6] Fioretti M, Delli CE. Epidemiological survey of dilatative cardiomyopathy in dogs [abstract]. *Veterinaria* 1988;2:81.
- [7] Gooding JP, Robinson WF, Wyburn RS, et al. A cardiomyopathy in the English cocker spaniel: a clinico-pathological investigation. *J Small Anim Pract* 1982;23:133–48.
- [8] Kittleson MD, Keene B, Pion PD, et al. Results of the Multicenter Spaniel Trial (MUST): taurine and carnitine-responsive dilated cardiomyopathy in American cocker spaniels with decreased plasma taurine concentration. *J Vet Intern Med* 1997;11:204–11.
- [9] Sanderson S, Osborne C, Ogburn P, et al. Canine cystinuria associated with carnitinuria and carnitine deficiency [abstract]. *J Vet Intern Med* 1995;9:212.
- [10] Calvert CA. Dilated congestive cardiomyopathy in Doberman pinchers. *Compend Contin Educ Pract Vet* 1986;8:417–30.
- [11] Monnet E, Orton C, Salman M, et al. Idiopathic dilated cardiomyopathy in dogs: survival and prognostic indicators. *J Vet Intern Med* 1995;9:12–7.
- [12] Tidholm Am Svensson H, Sylven C. Survival and prognostic factors in 189 dogs with dilated cardiomyopathy. *J Am Anim Hosp Assoc* 1997;33:364–8.
- [13] Tidholm A, Johsson L. A retrospective study of canine dilated cardiomyopathy (189 cases). *J Am Anim Hosp Assoc* 1997;33:544–50.
- [14] Tenaglia A, Cody R. Evidence for a taurine-deficient cardiomyopathy. *Am J Cardiol* 1988;62:136–9.
- [15] Hayes KC. Taurine requirement in primates. *Nutr Rev* 1985;43:65–70.
- [16] Hayes KC, Carey RE, Schmidt SY. Retinal degeneration associated with taurine deficiency in the cat. *Science* 1975;188:949–51.
- [17] Huxtable RJ. From heart to hypothesis: a mechanism for the calcium modulatory actions of taurine. *Adv Exp Med Biol* 1987;217:371–87.

- [18] Pion PD, Kittleson MD, Rogers QR, et al. Myocardial failure in cats associated with low plasma taurine: a reversible cardiomyopathy. *Science* 1987;237:764–8.
- [19] Schaffer SW, Seyed-Mozaffari M, Kramer J, et al. Effect of taurine depletion and treatment on cardiac contractility and metabolism. *Prog Clin Biol Res* 1985;179:167–75.
- [20] Takihara K, Azuma J, Awata N, et al. Beneficial effect of taurine in rabbits with chronic congestive heart failure. *Am Heart J* 1986;112:1278–84.
- [21] Franconi F, Bennardini F, Mattana A, et al. Taurine levels in plasma and platelets in insulin-dependent and non-insulin-dependent diabetes mellitus: correlation with platelet aggregation. *Adv Exp Med Biol* 1994;359:419–23.
- [22] Green TR, Fellman JH. Effect of photolytically generated riboflavin radicals and oxygen on hypotaurine antioxidant free radical scavenging activity. *Adv Exp Med Biol* 1994;359:19–29.
- [23] Rebel G, Petegnief V, Lleu P, et al. New data on the regulation of taurine uptake in cultured nervous cells. *Adv Exp Med Biol* 1994;359:225–32.
- [24] Schmidt SY. Biochemical and functional abnormalities in retinas of taurine-deficient cats. *Fed Proc* 1980;39:2706–8.
- [25] Sturman JA, Hayes KC. The biology of taurine in nutrition and developments. *Adv Nutr Res* 1980;3:231–99.
- [26] Sturman JA. Dietary taurine and feline reproduction and development. *J Nutr* 1991;121:S166–70.
- [27] Huxtable RJ, Chubb J, Asari J. Physiological and experimental regulation of taurine content in the heart. *Fed Proc* 1980;39:2685–90.
- [28] Schaffer SW, Kramer J, Chovan JP. Regulation of calcium homeostasis in the heart by taurine. *Fed Proc* 1980;39:2691–4.
- [29] Huxtable RJ. Physiological actions of taurine. *Physiol Rev* 1992;72:101–63.
- [30] Hamaguchi T, Azuma J, Schaffer S. Interaction of taurine with methionine: inhibition of myocardial phospholipids methyltransferase. *J Cardiovasc Pharmacol* 1991;18:224–30.
- [31] Lake N. Loss of cardiac myofibrils: mechanism of contractile deficits induced by taurine deficiency. *Am J Physiol* 1993;264(4 Part 2):H1323–6.
- [32] Steele DS, Smith GL, Miller DJ. The effects of taurine on Ca^{2++} uptake by the sarcoplasmic reticulum and Ca^{2++} sensitivity of chemically skinned rat heart. *J Physiol* 1990;422:499–511.
- [33] Gentile S, Bologna E, Terracina D, et al. Taurine-induced diuresis and natriuresis in cirrhotic patients with ascites. *Life Sci* 1994;54:1585–93.
- [34] Jacobsen JG, Thomas LL, Smith LH Jr. Properties and distribution of mammalian L-cysteine sulfinic carboxylases. *Biochim Biophys Acta* 1964;85:113–6.
- [35] Pion PD, Kittleson MD, Thomas WP, et al. Clinical findings in cats with dilated cardiomyopathy and relationship to finding taurine deficiency. *J Am Vet Med Assoc* 1992;201:267–74.
- [36] Pion PD, Sanderson SL, Kittleson MD. The effectiveness of taurine and levocarnitine in dogs with heart disease. *Vet Clin North Am Small Anim Pract* 1998;28:1495–514.
- [37] Moise NS. Cardiomyopathy in the fox and association with low dietary taurine. In: *Proceedings of the Seventh American College of Veterinary Internal Medicine Forum*. 1989. p. 834–5.
- [38] Kramer GA, Kittleson MD, Fox PR, et al. Plasma taurine concentration in normal dogs and dogs with heart disease. *J Vet Intern Med* 1995;9:253–8.
- [39] Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L-carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am J Vet Res* 2001;62:1616–23.
- [40] Association of American Feed Control Officials, Inc. Model bill and regulation. AAFCO Official Publication; 2003.
- [41] Backus RC, Cohen G, Pion PD, et al. Taurine deficiency in Newfoundlands fed commercially available complete and balanced diets. *J Am Vet Med Assoc* 2003;223:1130–6.

- [42] National Research Council. Nutrient requirements of dogs, revised 1985. Washington (DC): National Academy Press; 1985.
- [43] Spitze AR, Wong DL, Rogers QR, et al. Taurine concentrations in animal feed ingredients; cooking influences taurine content. *J Anim Physiol* 2003;87:251–62.
- [44] Kim SW, Morris JG, Rogers QR. Dietary soybean protein decreases plasma taurine in cats. *J Nutr* 1995;125:2831–7.
- [45] Fascetti AJ, Reed JR, Rogers QR, et al. Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997–2001). *J Am Vet Med Assoc* 2003;112:1137–41.
- [46] Belanger MC, Quillet M, Queney G, et al. Taurine-deficient dilated cardiomyopathy in a family of Golden Retrievers. *J Am Anim Hosp Assoc* 2005;41:284–91.
- [47] Torres CL, Backus RC, Fascetti AJ, et al. Taurine status in normal dogs fed a commercial diet associated with taurine deficiency and dilated cardiomyopathy. *J Anim Physiol* 2003;87:359–72.
- [48] Johnson ML, Parsons CM, Fahey GC, et al. Effects of species raw material source, ash content and processing temperature on amino acid digestibility of animal by-product meals by cecectomized roosters and ileally cannulated dogs. *J Anim Sci* 1998;76:1112–22.
- [49] Delaney SJ, Kass PH, Rogers QR, et al. Plasma and whole blood taurine in normal dogs of varying size fed commercially prepared food. *J Anim Physiol* 2003;87:236–44.
- [50] Stratton-Phelps M, Backus RB, Rogers QR, et al. Dietary rice bran decreases plasma and whole-blood taurine in cats. *J Nutr* 2002;132:1745S–7S.
- [51] Trautwein EA, Hayes KC. Taurine concentration in plasma and whole blood in humans: estimation of error from intra- and interindividual variation and sampling technique. *Am J Clin Nutr* 1990;52:758–64.
- [52] Pion PD, Lewis J, Greene K, et al. Effects of meal feeding and food deprivation on plasma and whole blood taurine concentration in cats. *J Nutr* 1991;121:S177–8.
- [53] Sanderson SS, Osborne C, Gross K, et al. Reliability of canine plasma and whole blood taurine concentrations as indicators of cardiac and skeletal muscle taurine concentrations. [abstract]. *J Vet Intern Med* 1998;12:224.
- [54] Bremer J. Carnitine-metabolism and function. *Physiol Rev* 1983;63:1420–80.
- [55] Leibowitz BE. Carnitine. *Adv Res Press* 1987;2:1–13.
- [56] Rebouche CJ, Engel AG. Kinetic compartmental analysis of carnitine metabolism in the dog. *Arch Biochem Biophys* 1983;220:60–70.
- [57] Neely JR, Morgan HA. Relationship between carbohydrate metabolism and energy balance of heart muscle. *Annu Rev Physiol* 1974;36:413–59.
- [58] Stumpt DA, Parker WD Jr, Angelini C. Carnitine deficiency, organic acidemias, and Reye's syndrome. *Neurology* 1985;35:1041–5.
- [59] Gilbert EF. Carnitine deficiency. *Pathology* 1985;17:161–9.
- [60] Mayes PA. Oxidation of fatty acids: ketogenesis. In: Murray RK, Granner DK, Mayes PA, et al, editors. *Harper's biochemistry*. 24th edition. Norwalk (CT): Appleton & Lange; 1996. p. 224–35.
- [61] Paulson DJ. Carnitine deficiency-induced cardiomyopathy. *Mol Cell Biochem* 1998;180:33–41.
- [62] Periera RR, Scholte HR, Luyt-Houwen IEM, et al. Cardiomyopathy associated with carnitine loss in kidneys and small intestines. *Eur J Pediatr* 1988;148:193–7.
- [63] Pierpont MEM. Carnitine and myocardial function. In: Carter AL, editor. *Current concepts in carnitine research*. 1st edition. Boca Raton (FL): CRC Press; 1992. p. 197–213.
- [64] Paulson DJ, Sanjak M, Shug AL. Carnitine deficiency and the diabetic heart. In: Carter AL, editor. *Current concepts in carnitine research*. 1st edition. Boca Raton (FL): CRC Press; 1992. p. 215–30.
- [65] Whitmar JT. Energy metabolism and mechanical function in perfused hearts of Syrian hamsters with dilated or hypertrophic cardiomyopathy. *J Mol Cell Cardiol* 1986;18:307–17.

- [66] Whitmar JT. L-Carnitine treatment improves cardiac performance and restores high-energy phosphate pools in cardiomyopathic Syrian hamsters. *Circ Res* 1987;61:396–408.
- [67] Sanderson S, Ogburn P, Osborne C. Heart disease management—Indications for nondrug therapies. *Vet Forum* 1996;13:36–43.
- [68] McEntee K, Clercx C, Snaps F, et al. Clinical, electrocardiographic, and echocardiographic improvements after L-carnitine supplementation in a cardiomyopathic Labrador. *Canine Pract* 1995;20:12–5.
- [69] Keene BW, Panciera DP, Atkins CE, et al. Myocardial L-carnitine deficiency in a family of dogs with dilated cardiomyopathy. *J Am Vet Med Assoc* 1991;198:647–50.
- [70] Costa ND, Labuc RH. Case report: efficacy of oral carnitine therapy for dilated cardiomyopathy in boxer dogs. *J Nutr* 1995;124(supp):2687S–92S.

14770

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/index.cfm>

Vet/Tech Contact: _____
Company Name: _____
Address: _____
(b) (6)

Email: _____
Tel: _____ **(b) (6)** Fax: _____ **(b) (6)**

Billing Contact: _____ **(b) (6)** TAX ID: _____
Email: _____ **(b) (6)** Tel: _____ **(b) (6)**

Patient Name: _____ **(b) (6)**
Species: Feline
Owner's Name: _____ **(b) (6)**

Sample Type: Plasma Whole Blood Urine Food Other: _____
Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 368 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

NC State University
Veterinary Hospital
1052 William Moore Drive
Raleigh, NC 27607
Discharge Comments

Fax: Admin
 Fax: Referral

Small Animal (919) 513-6500
 Large Animal (919) 513-6630

Client (b) (6)	Patient (b) (6) SCHNAUZER MC BLACK CANINE	Case # (b) (6) 8.2 kg	Attending DVM (b) (6) Student (b) (6) Discharging DVM (b) (6) Referring DVM (b) (6)
--------------------------	---	-------------------------------------	--

Admission Date/Time: (b) (6) 09:55 PM **Discharge Date/Time:** (b) (6) 10:57 AM **Discharge Status:**

*****NOTICE OF EUTHANSIA*****

Case Summary

Diagnosis:

- 1) Biventricular congestive heart failure (left significantly worse than right)
- 2) Cardiomyopathy (suspect secondary) vs. myocarditis vs. tachycardia-induced cardiomyopathy vs. other

History:

(b) (6) is a 2 and ½ year old male castrated Miniature Schnauzer who presented the NCSU ER on Thursday, (b) (6) for labored breathing and was subsequently transferred to NCSU Cardiology. (b) (6) initially developed a cough three weeks ago; (b) (6) describes the cough as a wheezing-type cough that occurred more frequently at night. When (b) (6) showed no signs of improvement, (b) (6) presented to his primary veterinarian on Thursday (b) (6). Kennel cough was suspected as the underlying cause of his cough and prednisone and doxycycline were prescribed. On Monday (b) (6) (b) (6) became uninterested in his food and began vomiting. The following day (b) (6) continued vomiting and developed labored breathing and subsequently re-presented to the rDVM for evaluation. Bloodwork and thoracic radiographs were performed. Bloodwork was reportedly unremarkable at this time and there was concern for aspiration pneumonia on his radiographs. Nebulization was performed and subcutaneous fluids, enrofloxacin, unasyn, and gentamicin were administered for treatment. While in-hospital, (b) (6) regurgitated. On Wednesday (b) (6), (b) (6) had improvement in respiratory effort but he still was not eating; subcutaneous fluids, enrofloxacin, unasyn, and gentamicin were performed again on an outpatient basis. The day of presentation (b) (6), (b) (6) syringe fed (b) (6), but as she attempted he immediately regurgitated. (b) (6) developed marked labored breathing following this and was presented to an emergency hospital. Thoracic radiographs were performed (uploaded in eFilm) and revealed cardiomegaly; a diffuse, severe mixed interstitial to alveolar pattern that is most severe caudodorsally; hepatomegaly; and decreased abdominal serosal contrast. (b) (6) was referred to NCSU for further care and ventilation if indicated.

(b) (6) has a history of developing coffee brown urine, sometimes with clumping, after strenuous activity when it is hot outside. The discolored urine typically develops after the activity and lasts 24-36 hours before clearing up. A urinalysis was performed and (b) (6) primary veterinarian detected crystals in the urine. (b) (6) diet was changed to an unknown diet to decrease the amount of crystals prevent stone formation; this diet was ultimately discontinued. Other than this, prior to the coughing that began 3 weeks ago (b) (6) was a normal, healthy dog with no significant medical history. He previously had no respiratory signs or changes in drinking/appetite/urination/defecation. (b) (6) lives with one other dog (not a relative) who is healthy and is currently up to date on his vaccinations (b) (6) is not current on any flea/tick prevention but receives heartworm prevention. (b) (6) is fed California Natural dog food.

Physical Exam Findings (on presentation):

Weight: 8.0 kg BCS: 5/9
 T: 100.2 F
 P: 160 bpm
 R: 64 breaths/min
 MMs/CRT: pale pink / <2 sec
 Attitude: Alert, responsive
 Hydration: Adequate

EENT/Oral: No ocular discharge noted, clear ear canals AU, no crusting/erythema noted; moist/flat nasal planum, no nasal discharge noted

PLN: No peripheral lymphadenopathy noted

CV/R: Grade I-II/VI left apical systolic murmur, femoral pulses hypokinetic but synchronous; jugular venous distention present; normal, albeit tachycardic, rhythm auscultes; dyspneic, inspiratory crackles in all lung fields on bilateral auscultation

GI/GU: Abdomen tense but non-painful on palpation; cranial abdominal organomegaly noted; no obvious fluid wave or masses noted

MSK: Appropriate and symmetrical muscling; ambulatory with no observed lameness; full ortho exam not performed due to patient's status

INTEG: Clean hair coat, no evidence of ectoparasites

NEURO: Mentally appropriate, intact cranial nerve reflexes, limited exam performed due to patient's status

Main Diagnostics (b) (6)

1. Big 4 - Glu: 135, Azo: 15-20, PCV: 40%, TS: 5.0 g/dL
2. Venous blood gas: pH 7.34, PCO2 46, lactate 2.4, HCO3 24.8
3. CBC - WBC 9.4, PCV 45, Seg 7.9, Band 0.18, Plt 157
4. Chemistry - BUN 19, creat 0.4, Phos 6.2, K 4.9, Na 140, TP 4.2
5. Blood Pressure - 90 mmHg systolic via Doppler
6. Urinalysis (post-lasix) - USG 1.011, otherwise unremarkable
7. Cardiac Troponin I - 0.79
8. Bap GM - pending
9. Vector borne panel - pending
10. Taurine levels - pending
11. Carnitine Levels - pending
12. Echocardiogram - Severely dilated and hypocontractile left and right ventricles, severely dilated left and right atria. Changes consistent with DCM (primary vs. secondary) vs. myocarditis vs. pacing-induced vs. other

Main Diagnostics (b) (6)

1. Chemistry - Gluc 225, BUN 29, Creat 0.7, phos 11.7, TP 5.0, ALT 53, CK 709, K 3.3, Cl 95, Na 144
2. Chest radiographs (9:15 AM) - final report pending - Severe generalized cardiomegaly with biventricular heart failure; improved from rDVM radiographs taken prior to presentation
3. Chest radiographs (5:00 PM) - final report pending - Progressive severe diffuse alveolar pattern consistent with worsening cardiogenic pulmonary edema; cannot exclude ventilator - induced lung injury and/or pneumonia

Main Diagnostics (b) (6)

1. CBC - WBC 9.9, Plt (clumping), PCV 44, Seg 8.0, Band 0.7
2. Chem - Gluc 136, BUN 12, Creat 0.7, Phos 4.6, ALT 88, CK 13,621, K 4.3, Na 151, Cl 109, AST 577
3. Coag - PT 9.1, PTT 14, Dimer 189, Fib 539, INR 1.09
4. Chest radiographs (1:30 AM, immediately post-ultrafiltration) - final report pending - markedly improved pulmonary infiltrates consistent with improved cardiogenic edema; residual interstitial to patchy alveolar pattern, decreased caudal cava size consistent with hypovolemia
5. Chest radiographs (11:00 AM) - final report pending - overall improved pulmonary pattern with persistent and in some areas slightly more condensed interstitial to patchy alveolar pattern. Consistent with continued improvement of congestive heart failure with possible concurrent bronchopneumonia; left lateral image shows suspected hiatal hernia

Main Diagnostics (b) (6)

1. CBC - WBC 6.8, PCV 39, Prot 6.9, Seg 4.2, Band 0.54, Toxic Neut Mild, Plt 97
2. Chem - Gluc 165, BUN 37, Creat 0.9, Phos 8.1, ALT 147, AST 1006, CK 35,930, Na 135, K 3.8, Cl 90
3. Urine Creat - 27.9
4. Urine Sodium - pending
5. Chest radiographs (10:00 AM) - final report pending - markedly progressive alveolar pattern consistent with significantly worsened cardiogenic edema; cannot exclude less likely differentials such as ARDS, ALI, hemorrhage, PTE, and pneumonia
6. ECG - suspect atrial tachycardia

Main Therapeutics Throughout Hospitalization (PO medications given through NG tube)

1. Furosemide - 2 boluses given on presentation followed by 1mg/kg/hr CRI with intermittent boluses given as needed
2. Pimobendan - initially 2.5mg PO TID increased to 5mg PO QID throughout hospitalization
3. Dobutamine - titrated between 7.5 mcg/kg/min increasing up to 20mcg/kg/min throughout almost entirety of hospitalization
4. Maropitant - 1mg/kg IV SID
5. Meropenem - 1.06mg/kg/hr CRI
6. Cisapride - 7.4mg PO TID
7. Pantoprazole - 7.4mg IV BID
8. Nitroglycerin Paste
9. Torsemide - 2.5-5mg PO BID
10. Hydrochlorothiazide - 6.25mg PO BID (started (b) (6))
11. Diltiazem (started (b) (6)) - 0.25mg IV (given slowly over 25 minutes) followed by CRI of 2-5mcg/kg/min (titrated PRN)
12. Triple antibiotic OD
13. NG tube feeding - as recommended per NCSU Nutrition service with supplements of Fish oil, taurine, and carnitine

Brief Daily Summary:

(b) (6) (b) (6) presented late in the evening on (b) (6) to the ER and after a TFAST was performed showing severe cardiomegaly with hypocontractility of the ventricles in addition to reviewing the rDVM radiographs, pimobendan and Lasix were given. An

echocardiogram was then performed (performed sternally cage-side given patient status) and a diagnosis of severe cardiomyopathy (primary vs. secondary DCM vs. myocarditis, vs. pacing-induced cardiomyopathy vs. other) and he was quickly given another dose of Lasix.

(b) (6) After the second bolus of Lasix was given he was immediately placed on a Lasix CRI at 1mg/kg/hr and Dobutamine at 5mcg/kg/min quickly uptitrated to 10mcg/kg/min. After 4-6 hours of clinical worsening and the suspicion for respiratory fatigue, mechanical ventilation was recommended to the owner and pursued. Recheck radiographs (first rads performed at NCSU) after being on ventilation and continued CHF treatment showed continued severe pulmonary infiltrates but significantly improved compared to rDVM rads prior to any intervention. Because of this improvement we continued with aggressive CHF management and mechanical ventilation. Throughout the day he started to show some worsening clinically while on the ventilator and recheck radiographs were performed around 4:30PM. These radiographs showed worsening of the pulmonary infiltrates despite aggressive therapy (dobutamine, pimobendan, Lasix, etc.). At that time aquapheresis was discussed with the owner and pursued. This was from approximately 7PM-1AM (including setup and moving the patient. This procedure was performed successfully with no significant complications.

(b) (6) Immediately after aquapheresis therapy recheck radiographs were performed (approximately 1:30AM) which showed a marked improvement in terms of pulmonary infiltrates. An interstitial to alveolar pattern persisted, mostly ventrally distributed, but was significantly improved. He was maintained on mechanical ventilation until approximately 5-6:00 AM when he was slowly weaned off ventilation and extubated. He handled this quite well and while sedated as the medications wore off, he clinically was markedly improved from presentation. His congestive heart failure medications were continued at aggressive doses (dobutamine at 15-20mcg/kg/min, Lasix had continued to be at 1mg/kg/hr CRI since presentation, Pimobendan was approximately 0.52mg/kg PO TID etc.). Recheck radiographs were performed around 2:00PM which showed an overall improvement in the pulmonary pattern, although a ventrally distributed alveolar pattern persisted and in some areas were slightly worse. We continued therapy for CHF and the patient was already being covered for pneumonia with meropenem (this antibiotic was chosen based on the patient's recent history of many antibiotics given including doxycycline, enrofloxacin, unasyn, and gentamycin). He continued to do well until acutely worsening was seen during the owner visit approximately 4:00PM (20 minutes into the visit) where his respiratory rate/effort declined. He was given another bolus of Lasix on top the CRI, butorphanol, and an increase in dobutamine. He was able to relax and showed slight improvement, although was unable to breathe as comfortably as prior to the episode. He remained tachycardic throughout the night but otherwise normal.

(b) (6) He remained tachycardic throughout the night at approximately 180bpm and the following morning he showed signs of atrial tachycardia at a rate of 190bpm. He was weaned off his dobutamine and started on diltiazem boluses followed by a CRI which resolved the atrial tach and a sinus tach at 150bpm persisted. He clinically was worse than the day before and repeat chest radiographs were performed that showed severe worsening of his pulmonary infiltrates. After discussion with the owner, we elected to continue even more aggressive diuretic therapy (multiple boluses of Lasix on top of the 1mg/kg/hr CRI he had been on almost since presentation) but by 3:00 PM after only further worsening the decision for euthanasia was made. The owners elected for necropsy with private cremation.

I am sorry for the loss of your patient. Both (b) (6) were absolutely wonderful to work with. If you have any questions at all, please do not hesitate to call us at 919-513-6694.

(b) (6) DVM

NC State University
Veterinary Hospital
1052 William Moore Drive
Raleigh, NC 27607
Discharge Comments

Fax: Admin
 Fax: Referral

Small Animal (919) 513-6500
 Large Animal (919) 513-6630

Client (b) (6)	Patient (b) (6) SCHNAUZER MC GRAY&WHITE CANINE	Case # (b) (6) 9.9 kg	Attending DVM (b) (6) Student Discharging DVM (b) (6) Referring DVM (b) (6)
--------------------------	--	-------------------------------------	--

Admission Date/Time: (b) (6) 10:46 AM **Discharge Date/Time:** (b) (6) 06:25 PM **Discharge Status:** UNDETERMINED

CASE SUMMARY:

HISTORY:

(b) (6) a 7 year old male castrated Schnauzer, was presented to NCSU Small Animal Emergency on (b) (6) for episodes of collapse. (b) (6) first collapse episode was in Mid February; he was described to fall over for 6 seconds with no loss of consciousness, and he immediately returned to normal after. (b) (6) had another collapse episode two weeks later, which looked the same and lasted the same amount of time as the first. (b) (6) next episode of collapse was 6/3/17, after a two hour hike, when he collapsed and screamed, which lasted for six seconds, and he was normal after. (b) (6) experienced another collapse a week after, he whined, it lasted six seconds, and was normal after. A week after that, (b) (6) had another collapse episode, where a puppy ran towards him, (b) (6) tried to run with him, stopped, got wobbly, stood still, and was normal after a few seconds, but did not fall to the floor. (b) (6) also reports that he is panting more than usual. He had a good appetite for treats, but seems more reluctant to eat his food since February. He does eventually eat his food if mixed in with some treats. (b) (6) is fed California Natural, Kangaroo or Venison flavor. There has been no change with (b) (6) urination or defecation, and is otherwise healthy. (b) (6) is up to day on flea and tick preventative with a three month Bravecto given on 5/2/17, but is not up to day on Heartworm preventative with Heartgard last given on 4/26/17.

PHYSICAL EXAM FINDINGS:

Weight: 9.9 kgs
 T: 100.6 F
 P: 130 bpm
 R: pant
 MMs/CRT: pink/<2 sec
 BSC: 5/9
 Pain Score: 0/4
 Attitude: BAR
 Hydration: euhydrated
 EENT: Unremarkable cornea/sclera OU, ears free of debris, no nasal discharge, no oral ulcers/lesions/foreign bodies detected
 PLN: No lymphadenopathy detected
 CV/R: No murmur or arrhythmia auscultated, femoral pulses strong and synchronous, normal bronchovesicular lung sounds noted on bilateral auscultation
 GI/GU: Normal on palpation, no fluid wave present, no organomegaly detected
 MSK: Ambulatory on all four limbs with no observed lameness.
 INTEG: Clean hair coat, no evidence of ectoparasites
 NEURO: BAR, normal gait on all four limbs, intact reflexes, cranial nerves intact

DIAGNOSTIC TESTS:

1. Big 4: BG 64 (recheck 79); BUN 15-26; PCV 45; TS 7.2
2. Blood pressure (Doppler): 130 mmHg
3. Echocardiogram: Moderately to severely decreased LV ejection fraction, LV cavity size is severely increased, severely dilated LA, moderate to severe mitral valve regurgitation, anterior and posterior leaflets mild thickened due to endocardiosis, mild tricuspid regurgitation, moderately elevated pulmonary artery systolic pressure. Diagnosis: mitral valve endocardiosis with left atrial enlargement and heart failure, decreased left ventricular systolic function, suspect dilated cardiomyopathy
4. ECG: regular sinus rhythm, complexes suggest LV enlargement, tall R waves
5. Thoracic radiographs: Severe left-sided cardiomegaly with moderate left atrial enlargement, unstructured interstitial pattern, and mild lobar venous distention: most consistent with left-sided congestive heart failure. ***Final report pending***
6. Urinalysis (cytocentesis): USG 1.019, protein negative, quiet sediment

Pending Lab Results:

1. Taurine
2. Carnitine
3. Vector borne panel

4. BAPGM
5. Troponin-I
6. T4
7. Toxoplasma
8. Neospora
9. Chagas
10. Complete amino acids

TREATMENTS:

1. Butorphanol 0.2 mg/kg IV
2. Furosemide 2 mg/kg SQ
3. Pimobendan 2.5 mg PO

ASSESSMENT:

Thank you for bringing (b) (6) in to see us! He is a very cute and sweet boy. (b) (6) presented to us today for evaluation of progressive episodes of collapse. On physical exam today, (b) (6) was stable, but he had mildly increased breath sounds noted in all lung fields. During more excitable moments, (b) (6) showed increased respiratory effort with an abdominal component to his breathing pattern. We consulted with the NCSU Cardiology Service, who performed an ECG and echocardiogram. Unfortunately, (b) (6) echocardiogram revealed evidence of mild mitral valve endocardiosis (i.e. chronic mitral valve disease) and suspected dilated cardiomyopathy. Mitral valve endocardiosis is a chronic, progressive condition in which the valve leaflets become progressively thickened and no longer close appropriately allowing mitral regurgitation. Over time, the left atrium will enlarge and this can lead to congestive heart failure. Typically, in patients with mitral valve disease, the systolic heart function/contractility is maintained until late stages of the disease. Unfortunately, (b) (6) systolic function was significantly decreased, and he showed abnormal dilation of his left atrium and ventricle. These findings were most consistent with a condition called dilated cardiomyopathy (DCM). DCM is a disease of unknown etiology affecting the muscle of the heart and is most commonly seen in large breed dogs (such as Dobermans, Great Danes, and Labrador Retrievers), but there is a small case report of this disease occurring in Standard Schnauzers. Although the exact mechanism of DCM is currently unknown, dietary taurine/carnitine deficiencies, genetics, infectious diseases, and toxins have all been linked to DCM. In order to assess for some of these possible causes, we have submitted testing for multiple nutritional deficiencies and infectious diseases. The NCSU Cardiology Service will call you as these tests become available. DCM leads to poor contractility and low cardiac output, and we suspect that (b) (6) episodes of collapse are most likely due to his low cardiac output during exertion. DCM can also lead to fibrosis and remodeling of the myocardium, which can lead to secondary arrhythmias. Fortunately, we saw no evidence of arrhythmias on (b) (6) ECG today. We performed chest radiographs to evaluate (b) (6) heart and lungs, and (b) (6) had evidence of left-sided congestive heart failure on his radiographs and impending right-sided congestive heart failure on his echocardiogram today. We are starting him on three medications to treat his heart disease and congestive heart failure today, and we may consider adding additional medications and supplements in the future. Please monitor (b) (6) for signs of worsening of his heart failure such as increased exercise intolerance, labored breathing, increased coughing or fainting. Call NCSU Cardiology or your referring veterinarian if any of these signs occur. Also, please start taking respiratory rates when (b) (6) is resting. This number should improve on his medications and remain less than 40 at rest. Please contact us or his primary care veterinarian if his respiratory rate or effort are worsening or you have any other concerns.

INSTRUCTIONS FOR CARE OF YOUR PET:**Medications:**

1. Pimobendan 2.5 mg capsules - Please give 1 capsule by mouth every 12 hours as directed. This medication dilates blood vessels (decreasing the workload of the heart) and increases the heart's pumping ability. Possible side effects include vomiting, diarrhea, and arrhythmias. Please begin this medication tonight.
2. Enalapril 5 mg tablets - Please give 1 tablet by mouth every 12 hours as directed. This is an ACE-inhibitor and is used as a cardioprotective medication to decrease remodeling and fibrosis of the heart. This medication will be given lifelong. Possible side effects include decreased blood pressure (hypotension) and decreased blood flow through the kidneys. Blood pressure and kidney bloodwork will need to be periodically monitored while receiving this medication. Please begin this medication tonight.
3. Furosemide 12.5 mg tablets - Please give 1 tablet by mouth every 12 hours as directed. This is a diuretic. Possible side effects include electrolyte abnormalities and decreased blood flow through the kidneys. Kidney bloodwork and electrolytes will need to be periodically monitored while receiving this medication. Please begin this medication tonight.

Activity / Cautions:

Please keep (b) (6) quiet until his recheck exam in several weeks. (b) (6) should be able to regulate his own energy level at home and be taken out for short bathroom breaks.

Diet:

Please continue (b) (6) current diet for the next several weeks. After this, he should be transitioned to a new meat-based commercial diet. We recommend that you consider one of the larger pet food brands (i.e. Royal Canin, Hills, Iams, Purina). Avoid salty treats, such as hot dog or jerky treats.

Monitoring:

1. Please monitor (b) (6) for signs of cardiac decompensation. This would include lethargy, cough, difficulty breathing (increased respiratory rate/effort), worsened abdominal distension, and episodes of collapse or fainting.

2. Please monitor (b) (6) resting respiratory rate. In order to do this, you can count the number of times that (b) (6) breathes in 15 seconds and multiply this number by 4 to get the number of breaths per minute. If you notice that this number is increasing or that it is consistently greater than 35-40 breaths per minute, then you should have (b) (6) evaluated by a veterinarian. A mobile app (Cardalis) is available to record instant rates and trends in respiratory rates.

PLAN FOR RE-EVALUATION OF YOUR PET:

Please call the Cardiology Service tomorrow to schedule an appointment for (b) (6) during the week of July 10th. You can schedule this appointment by calling (b) (6). Please bring a sample of (b) (6) current diet to this appointment for possible testing. He will also likely have a blood pressure, chest radiographs, and a renal panel performed during this appointment to re-evaluate his heart failure.

If you have any questions or concerns before this appointment, please have (b) (6) re-evaluated by your primary care veterinarian or the NCSU Emergency Service. (b) (6) may need recheck bloodwork and chest X-rays prior to this appointment if he is not doing well. He also may need medication adjustments and potentially hospitalization.

In order to help expedite medication refills, please visit us online at www.ncstatevets.org and select Pet Owners, Pharmacy Refills.

(b) (6) (b) (6) (b) (6)

SMALL ANIMAL EMERGENCY SERVICE TEAM:

Faculty: (b) (6)

Residents/Fellows: (b) (6)

Interns: (b) (6)

Supervisor: (b) (6)

Technicians: (b) (6)

Client Services: (b) (6)

Referring Veterinarians - please visit us online at www.ncstatevets.org/veterinarians and fill out our RDVM Feedback Survey!

The following instructions were provided at discharge

CLINICIAN: (b) (6), DVM

Date: (b) (6)

CASE SUMMARY:

Thank you for entrusting us with the care of your companion. The Discharge Summary will be emailed to you and faxed to your RDVM within 24 hours of release/discharge from our facility at (b) (6). If you or your veterinarian do not receive this, please contact the (b) (6) Emergency Service to request a copy. The following will briefly outline the care your companion should receive at home and was explained by our staff at the time of discharge:

DIAGNOSIS (ES):

- 1. Left-sided congestive heart failure
- 2. Mitral valve endocardiosis with left atrial enlargement
- 3. Dilated cardiomyopathy

TREATMENTS:

- 1. Butorphanol 0.2 mg/kg IV
- 2. Furosemide 2 mg/kg SQ

3. Pimobendan 2.5 mg PO

INSTRUCTIONS FOR CARE OF YOUR PET:

Medications:

1. Pimobendan 2.5 mg capsules - Please give 1 capsule by mouth every 12 hours as directed. This medication dilates blood vessels (decreasing the workload of the heart) and increases the heart's pumping ability. Possible side effects include vomiting, diarrhea, and arrhythmias. Please begin this medication tonight.

2. Enalapril 5 mg tablets - Please give 1 tablet by mouth every 12 hours as directed. This is an ACE-inhibitor and is used as a cardioprotective medication to decrease remodeling and fibrosis of the heart. This medication will be given lifelong. Possible side effects include decreased blood pressure (hypotension) and decreased blood flow through the kidneys. Blood pressure and kidney bloodwork will need to be periodically monitored while receiving this medication. Please begin this medication tonight.

3. Furosemide 12.5 mg tablets - Please give 1 tablet by mouth every 12 hours as directed. This is a diuretic. Possible side effects include electrolyte abnormalities and decreased blood flow through the kidneys. Kidney bloodwork and electrolytes will need to be periodically monitored while receiving this medication. Please begin this medication tonight.

Activity / Cautions:

Please keep (b) (6) quiet until his recheck exam in several weeks. (b) (6) should be able to regulate his own energy level at home and be taken out for short bathroom breaks.

Diet:

Please continue (b) (6) current diet for the next several weeks. After this, he should be transitioned to a new meat-based commercial diet. We recommend that you consider one of the larger pet food brands (i.e. Royal Canin, Hills, Iams, Purina). Avoid salty treats, such as hot dog or jerky treats.

Monitoring:

1. Please monitor (b) (6) for signs of cardiac decompensation. This would include lethargy, cough, difficulty breathing (increased respiratory rate/effort), worsened abdominal distension, and episodes of collapse or fainting.

2. Please monitor (b) (6) resting respiratory rate. In order to do this, you can count the number of times that (b) (6) breathes in 15 seconds and multiply this number by 4 to get the number of breaths per minute. If you notice that this number is increasing or that it is consistently greater than 35-40 breaths per minute, then you should have (b) (6) evaluated by a veterinarian. A mobile app (Cardalis) is available to record instant rates and trends in respiratory rates.

PLAN FOR RE-EVALUATION OF YOUR PET:

Please call the Cardiology Service tomorrow to schedule an appointment for (b) (6) during the week of July 10th. You can schedule this appointment by calling (b) (6). Please bring a sample of (b) (6) current diet to this appointment for possible testing. He will also likely have a blood pressure, chest radiographs, and a renal panel performed during this appointment to re-evaluate his heart failure.

If you have any questions or concerns before this appointment, please have (b) (6) re-evaluated by your primary care veterinarian or the NCSU Emergency Service. (b) (6) may need recheck bloodwork and chest X-rays prior to this appointment if he is not doing well. He also may need medication adjustments and potentially hospitalization.

IF YOU HAVE ANY QUESTIONS OR PROBLEMS, PLEASE CALL THE SMALL ANIMAL EMERGENCY SERVICE AT (b) (6).

PLEASE CALL TO MAKE YOUR FOLLOW UP APPOINTMENT AS RECOMMENDED

_____/_____/_____
Owner/Agent Signature Clinician's Signature Student's Signature

Copy to: Owner / Medical Record / Fax to RDVM

NC State University
Veterinary Hospital
1052 William Moore Drive
Raleigh, NC 27607
 Discharge Comments

Fax: Admin
 Fax: Referral

Small Animal (919) 513-6500
 Large Animal (919) 513-6630

Client (b) (6)	Patient (b) (6) SCHNAUZER MC GRAY&WHITE CANINE	Case # (b) (6) 9.9 kg	Attending DVM ADIN, DARCY Student (b) (6) Discharging DVM Referring DVM (b) (6)
--------------------------	--	-------------------------------------	--

Admission Date/Time: (b) (6) 10:59 AM **Discharge Date/Time:** (b) (6) 03:16 PM **Discharge Status:**

CASE SUMMARY

DIAGNOSIS:

Dilated Cardiomyopathy (DCM)
 Mitral valve endocardiosis and regurgitation

HISTORY: (b) (6) a 7 year old neutered male miniature Schnauzer, was presented to the NCSU cardiology service for re-evaluation of dilated cardiomyopathy. At his prior visit, (b) (6) presented to our Small Animal Emergency on (b) (6) for episodes of collapse. (b) (6) had his first two collapse episodes in mid February; he was described to fall over for ~6 seconds with no loss of consciousness, and he immediately returned to normal after. He later has several more collapsing episodes in June 2017, which prompted his presentation to NCSU SAES. Collapse episodes were preceded by excitement or exertion. He was discharged (b) (6) with Pimobendan 2.5mg PO BID, Enalapril 5mg PO BID and Lasix 12.5mg PO BID. (b) (6) house mate, (b) (6), unfortunately passed away from DCM and CHF in February 2017. Infectious disease testing and amino acid testing did not identify a cause for the DCM. Histopathology was relatively unremarkable, however, findings could have been consistent with a toxic insult.

Since (b) (6) (b) (6) reports (b) (6) is doing much better. He has not had any more collapsing episodes, excepting one moment where he stumbled but did not fall when excited 7/4/17. He is tolerating the medications and eating well. No coughing, sneezing, vomiting or diarrhea. His RR has been very normal since starting medications. His diet has been changed to Science Diet adult small breed from California Naturals.

The tests results from (b) (6) (b) (6) visit were normal, including vector borne testing (IFA, PCR) and taurine and carnitine analysis. A full amino acid profile did not reveal significant abnormalities, however, consultation with UCD is pending for a full interpretation.

PHYSICAL EXAM FINDINGS:

T- 99.7* P- 136 bpm R- 36 Wt- 9.6kg
 MM- pink, slightly tacky but no skin tenting
 CRT- <2 seconds
 BCS- 6/9
 Attitude- BAR
 Hydration: equivocal mild dehydration <5%

EENT: Unremarkable cornea/sclera OU, ears free of debris, mild dry crust in both nares, no oral ulcers/lesions/foreign bodies detected

PLN: No lymphadenopathy detected

CV/R: Grade 2/6 left apical systolic murmur, no arrhythmias auscultated, femoral pulses strong and synchronous, normal bronchovesicular lung sounds noted on bilateral auscultation- no crackles or wheezes, eupneic

GI/GU: Soft, non-painful, slightly distended abdomen on palpation, no fluid wave present, no organomegaly detected.

MSK: Ambulatory on all four limbs with no observed lameness.

INTEG: Clean hair coat, no evidence of ectoparasites

NEURO: BAR, normal gait on all four limbs, intact reflexes, cranial nerves intact

RESULTS OF DIAGNOSTIC TESTS:

BLOOD PRESSURE- 110 mmHg systolic

CHEST RADIOGRAPHS- Moderate left sided cardiomegaly with no signs of congestive heart failure. Moderate hepamegaly.

RENAL PANEL: all normal (BUN 16, creat 0.6, K 4.0)

ASSESSMENT:

(b) (6) was presented today for a recheck of his recently diagnosed dilated cardiomyopathy (DCM). DCM is a disease of unknown cause affecting the muscle of the heart and is most commonly seen in large breed dogs (such as Dobermans, Great Danes, and Labrador Retrievers). Although the exact mechanism of DCM is currently unknown, dietary taurine/carnitine deficiencies, genetics and toxins have all been linked to DCM. Infectious and nutritional causes of DCM have been ruled out to the best of our ability to test for (b) (6). The overall effect of DCM is a decrease in the contractility (pumping ability) of the heart. Because the heart is unable to pump

with enough vigor to move blood adequately forward into circulation, a volume overload occurs and the heart dilates to accommodate it. As a result, the chambers of the heart become very large, and the walls of the heart become very thin. Ultimately, the heart is unable to accommodate and dilate further; the result is back-up of blood from the heart and into the lungs, known as congestive heart failure. It is important to know that this disease is progressive, and ultimately those patients affected with it will experience congestive heart failure.

Given the unusual timing of (b) (6) both developing dilated cardiomyopathy within the same time frame, their different ages, unrelatedness, same environment and lack of an identifiable cause, we are continuing to hunt for an environmental explanation for (b) (6) DCM. We will keep you up to date as we pursue toxin testing in the food and treats you have brought us. However, negative toxin testing may not completely rule out a toxin since we do not have samples representative of the onset of cardiac signs in both dogs. We have changed (b) (6) food to address this possibility.

We are glad to see that (b) (6) is doing well clinically. His chest xrays showed resolution of congestive heart failure and his renal panel was normal indicating that he is tolerating the medications well. We would like to add spironolactone to his regime because his potassium, while normal, is at the low end of the range, and because this may have some long term benefit through anti-aldosteronism. We have also listed doses for supplements, that while unproven in their benefit, are not harmful and may help his myocardial function. Taurine and carnitine supplementation are unlikely to be helpful since his plasma and whole blood concentrations are normal.

MEDICATIONS:

Please continue the following medications:

FUROSEMIDE (12.5 mg tablets): Please give 1 tablet by mouth every 12 hours.

PIMOBENDAN (2.5 mg capsules): Please give 1 capsule by mouth every 12 hours. If (b) (6) continues to have near-collapse episodes, you may increase this to 1 capsule 3 times daily. If the pharmacy gives you 2.5 mg tablets in the future you can also give 1.5 of these tablets twice daily.

ENALAPRIL (5 mg tablets): Please give 1 tablet by mouth every 12 hours.

Please start the following medication:

SPIRONOLACTONE 25 mg tablets: Give 1 tablet by mouth every 24 hours. This is a weak diuretic that can also may have cardioprotective effects. Side effects can include electrolyte abnormalities and decreased blood flow through the kidneys. Kidney bloodwork and electrolytes will need to be periodically monitored while receiving this medication.

(b) (6) taurine and carnitine assessments were normal, making supplementation unlikely to be beneficial. Other supplements with theoretical but unstudied benefits are included below. These are not likely to be harmful and no interactions with his medications are anticipated.

1. D-ribose 5 grams (1 scoop) daily.
2. Coenzyme Q10 (as ubiquinol not ubiquinone) 100 mg capsules: give 1 capsule 2-3 times daily
3. Fish oil 450 mg of EPA and DHA per day. This is an approximate dose and can be rounded up or down depending on the fish oil chosen. Nature Made and Nordic Naturals are high quality brands.

ACTIVITY:

Please avoid strenuous exercise or situations which place undue stress on your pet. In general, pets with congestive heart failure will self-regulate their exercise. Please monitor for any change in exercise capability.

Please continue to monitor (b) (6) respiratory rate and call if this increases. Please also call if he begins coughing or collapse episodes recur.

DIET:

A diet that is moderately restricted in salt is ideal for cardiac patients, as excessive salt load can cause fluid accumulation. A commercial "Senior" diet is formulated with an appropriate amount of salt for your pet. Please also avoid salty treats, such as hot dogs or jerky treats.

NEXT APPOINTMENT:

(b) (6) should have a recheck appointment with NCSU Cardiology in 3-4 months to evaluate chest X-rays, blood pressure, kidney values, echocardiogram, and troponin. If he begins to show signs of heart failure prior to your next recheck, please call NCSU Cardiology so that we can recheck your pet sooner.

CLINICIANS:

(b) (5) Dr. Darcy Adin, (b) (6)

RESIDENTS:

(b) (6)

CLINICAL TECHNICIANS:

(b) (6)

RESEARCH TECHNICIAN:

(b) (6)

CLIENT SERVICES:

(b) (6)

In order to help expedite medication refills, please visit us online at www.ncstatevets.org and select Pet Owners, Pharmacy Refills.

NOTE: If your pet is in need of emergency aid and you are not able to get to the NC State Veterinary Hospital quickly, please seek care at the nearest veterinary emergency facility. Take these discharge instructions and current medications with you so that the treating veterinarian will know as much as possible regarding your pet's medical condition.

Owner - (b) (6)

Clinician - Darcy Adin, DVM

Student - (b) (6)

Referring Veterinarians - please visit us online at www.ncstatevets.org/veterinarians and fill out our RDVM Feedback Survey!

**Follow-up Case Information Uniform Data Entry Form
Vet-LIRN**

Date (mm/dd/yy)
EON/CC Number:

PATIENT INFORMATION

Pet Name
 Dog Cat
Breed
Age in years (if < 6 months, put 0.5)
Gender:
 M MN F FS

This form serves as a Uniform Data Entry Form to capture additional case specific information not clear from the Consumer Complaint or Medical Records in a standardized manner. Because each follow-up interview made with owners features questions tailored specifically to the case, each box of information contained in this Uniform Data Entry Form may not be completed.

HISTORY-Additional Comments from Owner

Owner's Description of What Happened:

Any Health Problems Prior to the Event (e.g. allergies, surgeries):

Sensitive GI tract (e.g. stomach upset when switching foods, eats a lot of grass) Yes
Changes to the pet's diet prior to illness Yes
Date Diet Change:

CLINICAL INFORMATION--Additional Comments from Owner on What Happened

Appetite Increased Decreased
Vomiting Yes
Diarrhea Yes
Duration of Diarrhea (days)
Blood in Feces Fresh,Red
 Coffee Ground
 Black,Tarry
Water Consumption Increased Decreased
Urination Increased Decreased
Lethargy Yes
Other:

MEDICATIONS-Taken Prior to the Event and Mentioned by Owner

List medications mentioned by owner (e.g. NSAIDs, steroids, heartworm/flea prevention, antibiotics, etc.)

List probiotics, vitamins, or supplements mentioned by owner:

**Follow-up Case Information Uniform Data Entry Form
Vet-LIRN**

EON/CC Number: 266,814

Owner: (b) (6)

Pet's Name: (b) (6)

DIET-Any other foods the owner mentions were given to the animal during this period. (check all that apply)

Commercial Dry Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name: Merrick Grain Free Bistro Chicken-started in January of 2014, print out of all pet food purchases thru Jan 2014 --> on the Chicken pretty much the entire time; owner has a listing of all purchases from pet

Commercial Wet-Canned Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name: Fancy Feast Wet food given to (b) (6) to stimulate her appetite after her illness onset, but she only licked the gravy. So for ~1 week before the other 4 house cats were tested for taurine, only (b) (6)

Commercial Wet-Pouch Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name:

Commercial-Raw Product Use as Part of Diet: Primary Secondary Occasional

List Product Label Name:

Homemade-Raw Product Use as Part of Diet: Primary Secondary Occasional

Describe Product Type:

Homemade-Cooked Product Use as Part of Diet: Primary Secondary Occasional

Describe Product Type:

Table Scraps/Human Food (as an occasional contribution to diet) Describe Product Type(s): not in past 5 years

Pet Treat Products Product Use as Part of Diet: Primary Secondary Occasional

Commercial Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Rawhides or Pig Ears Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Marrow Bones Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Chicken Jerky Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Duck Jerky Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

Sweet Potato Jerky or Treats Product Label Name/Lot: Date first fed

How Product Administered: Date last fed

**Follow-up Case Information Uniform Data Entry Form
Vet-LIRN**

EON/CC Number:

Owner:

Pet's Name:

DIET-continued-Any other foods the owner mentions were given to the animal during this period. (check all that apply)

Other Treats Product Label Name/Lot: Date first fed
How Product Administered Date last fed

ENVIRONMENTAL EXPOSURES-Environmental Exposures Mentioned by the Owner Potentially Affecting the Animal's Overall State of Health Prior to the Event . (check all that apply)

- Indoor Outdoor Indoor & Outdoor Carrion Rodents Grapes or Raisins Nuts
- Plants Trash Hunt Pet Shows Sporting Events Pet Recreation Facilities
- Livestock Poultry Reptiles Pet Birds Small Mammals Untreated Surface Water
- Anti-freeze Mushrooms Heavy Metals Ticks Urban Suburban Rural

Comments:

HOUSEHOLD-Signalment of Additional Animals Given the Product mentioned by the owner.

Animal 1 Reacted

Animal 2 Reacted

Animal 3 Reacted

Comments

Report Details - EON-345831		
ICSR:	2040528	
Type Of Submission:	Initial	
Report Version:	FPSR.FDA.PETF.V.V1	
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)	
Reporting Type:	Voluntary	
Report Submission Date:	2018-01-22 18:04:59 EST	
Reported Problem:	Problem Description: One week prior to presentation at a local emergency and specialty clinic, owners noted (b) (6) to be tiring more quickly when playing catch. On the day of her initial presentation to the ER clinic, she had been coughing more than usual and sank to the ground when attempting to chase a ball, but recovered quickly. At the ER clinic she was diagnosed with atrial fibrillation and early congestive heart failure. Treatment was initiated there (b) (6), she was transferred to our clinic on (b) (6) for further evaluation and care. Her arrhythmia converted back to a normal sinus rhythm on (b) (6) prior to transfer and her congestive heart failure resolved with treatment. She had a second collapse episode prior to referral. Echocardiogram showed evidence of dilated cardiomyopathy with concurrent chronic degenerative valve disease.	
	Date Problem Started: (b) (6)	
	Concurrent Medical Problem: Yes	
	Pre Existing Conditions: On supplements - Missing Link and Sea Jerky History of focal facial seizures - owner occasionally uses an herbal supplement for calming when this occurs, unknown	
	Outcome to Date: Better/Improved/Recovering	
Product Information:	Product Name: California Natural Grain-Free Kangaroo and Red Lentils Recipe	
	Product Type: Pet Food	
	Lot Number:	
	Package Type: BAG	
	Purchase Date: 08/14/2017	
	Possess Unopened Product: No	
	Possess Opened Product: No	
	Product Use Information:	Description: (b) (6) had eaten this diet for years alongside her housemate who was being fed it for a food allergy.
		Last Exposure Date: 09/01/2017
		Product Use Stopped After the Onset of the Adverse Event: Yes
		Adverse Event Abate After Product Stop: No
Product Use Started Again: No		
Perceived Relatedness to Adverse Event: Probably related		
Manufacturer /Distributor Information:		
Purchase Location Information:		
Animal Information:	Name: (b) (6)	
	Type Of Species: Dog	

	Type Of Breed:	Retriever - Labrador	
	Gender:	Female	
	Reproductive Status:	Neutered	
	Weight:	32.1 Kilogram	
	Age:	8 Years	
	Assessment of Prior Health:	Excellent	
	Number of Animals Given the Product:	4	
	Number of Animals Reacted:	4	
	Owner Information:	Owner Information provided:	Yes
		Contact:	Name: (b) (6)
Phone: (b) (6)			
Other Phone: (b) (6)			
Email: (b) (6)			
Address:	(b) (6) United States		
Healthcare Professional Information:	Practice Name:	(b) (6)	
	Contact:	Name: (b) (6)	
		Phone: (b) (6)	
		Other Phone: (b) (6)	
Email: (b) (6)			
Address:	(b) (6) United States		
Sender Information:	Name:	(b) (6)	
	Address:	(b) (6) United States	
	Contact:	Phone:	(b) (6)
		Other Phone:	(b) (6)
		Email:	(b) (6)
	Permission To Contact Sender:	Yes	
	Preferred Method Of Contact:	Phone	
Reported to Other Parties:	Other		
Additional Documents:	Attachment:	cardio0009.pdf	
	Description:	Taurine level	
	Type:	Laboratory Report	
	Attachment:	cardio0008.pdf	

FDA-CVM-FOIA-2019-1704-000471

Description: Echo reports from initial presentation, 5/2017, and 12/2017

Type: Medical Records

16051

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/aal/aal.html>

Vet/Tech Contact: Account # _____ (b) (6) / Contact: (b) (6) Date: 1-23-17

Company Name: _____ (b) (6)

Address: _____ (b) (6)

Email: _____ (b) (6)

Tel: _____ (b) (6) Fax: _____ (b) (6)

Billing Contact: _____ (b) (6) (C) TAX ID: _____

Email: _____ (b) (6) Tel: _____ (b) (6)

Patient Name: (b) (6)

Species: hg

Owner's Name: (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____

Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: **236** Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

Patient Information

Patient:	(b) (6)	Age: 5	years	Referring Veterinarian:	(b) (6)
Patient Number:	(b) (6)	Weight:(kg)	25.60	Cardiologist:	(b) (6)
Breed:	Labrador Retriever	Sex:	FS	Client Number:	(b) (6)
Exam Date:	(b) (6) 08:19	BSA:	0.88		

History: (b) (6) was presented to (b) (6) today for evaluation of a new heart murmur and evaluation after being diagnosed with congestive heart failure on 8/18/17. (b) (6) was evaluated by her regular vet on 8/18/17 for heavy breathing and coughing. Radiographs and blood work were done at the time and (b) (6) was diagnosed with an enlarged heart and congestive heart failure at that time. (b) (6) was started on Furosemide, Vetmedin and Enalapril by her regular vet. (b) (6) was seen through (b) (6) on (b) (6) for reevaluation of congestive heart failure. The clients report that (b) (6) had improved some but had not improved a lot. Medications were adjusted based on recommendations from (b) (6) on (b) (6) until (b) (6) could get an appointment to be see by (b) (6). The clients report that since the medications were increased (b) (6) has improved, however they do still feel that she is breathing faster than normal at home and she is still panting a lot at home. (b) (6) is still eating very well at home and is currently on a low sodium kangaroo and lentil diet. The clients also report that there other dog, (b) (6), who we also see is (b) (6) aunt (b) (6) mother was a littermate of (b) (6). (b) (6) is also urinary incontinent which started prior to developing congestive heart failure and initiation of treatment. The clients report that (b) (6) was not well controlled on Incurin 1 mg tablets: 1 tablet by mouth once daily so Proin 50 mg tablets: Give 1/2 tablet by mouth every 12 hours was added in. They are unsure if (b) (6) incontinence is now controlled because there other dog has developed incontinence as well. The client feel that (b) (6) is less social and less active at home. (b) (6) is currently receiving furosemide 40 mg tablets: give 1 and 1/2 tablets by mouth every 12 hours, spironolactone 25 mg tablets: give 1 tablet by mouth every 12 hours, enalapril 10 mg tablets: give 1 tablet by mouth every 12 hours, Vetmedin 5 mg tablets: give 1 and 1/2 tablets by mouth every 12 hours, Incurin 1 mg tablets: give 1 tablet by mouth once every 24 hours, Proin 50 mg tablets: give 1/2 tablet by mouth every 12 hours and Apoquel 16 mg tablets: give 3/4 tablet by mouth every 24 hours.

Physical Examination: Temp: 101.5 HR: 150 RR: panting
Grade 4/6 left apical systolic murmur with radiation to the right. Adequate femoral pulses. Regular rhythm. Normal lung sounds. Eupneic. Normal abdominal palpation. PLNs WNL. MM pink/moist. CRT < 2 sec.

Diagnostic Tests: Blood Pressure: 110 mmHg (#4 cuff, left forelimb)
Thoracic Radiographs: Persistent cardiomegaly with mild decrease in severity. No evidence of cardiac decompensation.
Renal Panel: Mildly elevated SDMA (19 ug/dL). Otherwise clinically unremarkable.
Echo: See Below. Sinus rhythm on ECG.

Echocardiographic Report

2D ECHO

LA Systolic Diameter LX	6.3 cm	Aortic Root Diameter	2.1 cm
-------------------------	--------	----------------------	--------

DOPPLER

AV Peak Velocity	101 cm/s	PV Peak Velocity	76.1 cm/s
AV Peak Gradient	4.1 mmHg	PV Peak Gradient	2.3 mmHg
Mitral E Point Velocity	103 cm/s	TR Peak Velocity	251 cm/s
Mitral E to A Ratio	1.9	TR Peak Gradient	25.3 mmHg
MR Peak Velocity	435 cm/s		

M-MODE

LV Diastolic Diameter MM	7.5 cm	IVS Percent Thickening MM	0.22
LV Systolic Diameter MM	6.6 cm	LVPW Diastolic Thickness MM	0.67 cm
LV Fractional Shortening MM	12.1 %	LVPW Systolic Thickness MM	0.84 cm
LV Diastolic Volume Cube	427 cm ³	LVPW Percent Thickening MM	0.26
LV Systolic Volume Cube	290 cm ³	IVS to PW Ratio MM	1.1
LV Ejection Fraction Cube	0.32	LV Mass MM	238 g
IVS Diastolic Thickness MM	0.74 cm	LV Mass Normalized MM	272 g/m ²
IVS Systolic Thickness MM	0.9 cm	MV E Point Septal Separation	3.4 cm

Left Ventricle: Severe dilation with marked global myocardial dysfunction. Normalized LVIDd 2.9, normalized LVIDs 2.38.

Left Atrium: Severe dilation with septum bowing to the right.

Right Ventricle: Mild to moderate dilation with reduced myocardial function.

Right Atrium: Mild to moderate dilation.

Mitral Valve: Thickened valve leaflets. 3-4+ mitral regurgitation.

Aortic Valve: Mildly thickened valve leaflets. No aortic insufficiency.

Tricuspid Valve: Thickened valve leaflets. Two jets of 2-3+ tricuspid regurgitation. Normal regurgitant velocities.

Pulmonic Valve: Mildly thickened valve leaflets. Mild pulmonic insufficiency.

Aorta: Normal

Pericardium: Normal

Diagnosis

Endocardiosis (chronic degenerative valve disease) - Degenerative changes in one or more heart valves have caused leaking across these valves. This is the source of the heart murmur. As this disease progresses, the heart enlarges. Eventually this can lead to symptoms of cough and shortness of breath (airway compression and/or congestive heart failure). This is usually a slowly progressive disease.

Dilated cardiomyopathy - This is a disease characterized by weakening of the heart muscle and dilation of the heart chambers. As the disease progresses, it can lead to congestive heart failure (fluid in the lungs causing shortness of breath and cough). Abnormal heart rhythms are common and can result in sudden death. Most commonly this is an inherited disease, though it can occur secondary to a deficiency in an amino acid called taurine.

Recommendations

Please continue the following medications as previously directed:

Furosemide 40 mg tablets: Give 1 and 1/2 tablets by mouth every 12 hours.

This is a diuretic (water pill), that prevents the body from retaining excessive sodium and water. It will cause your pet to drink and urinate more frequently. It is important that fresh water is always available.

Enalapril 10 mg tablets: Give 1 tablet by mouth every 12 hours.

This medication is a strong drug that dilates blood vessels, permitting the heart to pump blood more efficiently. It can lower blood pressure (hypotension) and cause changes in kidney function and electrolyte values. If your pet develops weakness or depression, decrease the drug dose by 1/2 and call. A kidney panel and blood pressure should be reevaluated 7-10 days after beginning this medication.

Vetmedin 5 mg tablets: Give 1 and 1/2 tablets by mouth every 12 hours.

Pimobendan (Vetmedin) in Congestive Heart Failure (CHF) - This is a drug that is approved for the treatment of congestive heart failure secondary to dilated cardiomyopathy or chronic valve disease (endocardiosis). Studies have shown improved quality of life and increase survival time when this drug is added to other standard cardiac medications. In our experience, side effects are uncommon, but it is important that you advise us if you feel your pet is having any potential adverse effects from this medication. The reported potential side effects listed for this medication are increased heart rate, vomiting, diarrhea, inappetence, uneasiness, incoordination, convulsions, increased drinking and increase urinating.

Spironolactone 25 mg tablets: Give 1 tablet by mouth every 12 hours

This is a diuretic (water pill) that also blocks a hormone that can injure the heart muscle. It works well in combination with the furosemide and enalapril.

Proin 50 mg tablets: Give 1/2 tablet by mouth every 12 hours

Apoquel 16 mg tablets: Give 3/4 tablet by mouth once every 24 hours

INCREASE:

Incurin 1 mg tablets: INCREASE to 2 tablets by mouth once every 24 hours.

As we discussed, (b) (6) and (b) (6) unfortunately have very similar structural heart disease. Since they are related, this raises concern for a genetic component. You have expressed that there is no history of heart disease in their lineage. It is possible that the disease has remained silent in other related dogs or is inherited in a way that it is only expressed in certain individuals. The other common denominator that (b) (6) and (b) (6) have is the kangaroo diet. Even though we have not specifically associated this protein source with taurine/carnitine deficiency, it may be warranted to consider a diet with a different protein source since it is a novel protein and both dogs have very similar disease manifestations. Lamb should be avoided as it has been associated with taurine deficiency in dogs.

We did not check (b) (6) blood taurine level today- since (b) (6) was normal it is highly unlikely that (b) (6) will be deficient as they are related and eat the same food.

One thing that can be very helpful for home monitoring is checking sleeping or resting respiratory rates. A recent study showed that even pets with severe heart disease rarely have resting respiratory rates greater than 30 breaths per minute unless they are starting to decompensate for that disease. Elevated respiratory rates at home may be even more sensitive than chest radiographs at picking up early decompensation. Count your pet's respiratory rate when he/she is at rest or sleeping (not within 20 minutes of being active). If his/her respiratory rate is greater than 30 breaths per minute, recheck again in a couple of hours. If persistently elevated above this level, call.

With advanced heart disease, our biggest dietary concerns are adequate caloric content and low sodium content. We aim for less than 80mg sodium per 100 kilocalories (kcal) in patients that have developed congestive heart failure. We do not advise protein restriction unless there is concurrent kidney disease (i.e. kidney diets are not advised unless there is concurrent kidney disease). Please refer to our diet handouts with a list of currently adequate diets and treats, though this list is not exclusive. If you wish to feed a diet that is not on these lists, you will need to call the manufacturer of the diet to obtain a sodium content.

Exercise is also a concern in advanced heart disease. While cage rest is ideal with active heart failure, some exercise is permissible in asymptomatic disease. However, vigorous or extended exercise should be avoided.

***As long as (b) (6) does well at home we would like to re-evaluate her in 4-6 weeks. At this time we will recheck her kidney values/electrolytes and blood pressure as well as repeat chest x-rays.

(b) (6) (Cardiology)

(Electronically Signed)

Final Date:

Like us on Facebook!

www.facebook.com/ (b) (6)

Notes to our clients

- Please bring all medications to your pet's scheduled appointments.
- We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).
- Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.
- If an emergency arises with your pet, (b) (6) is a 24 hour facility.

JLR placed in cooler
(b) (6) e??



PATIENT CASE REPORT

Date: (b) (6) Time: 22:45

Client: (b) (6)
Address: (b) (6)

Patient: (b) (6)
Breed: Retriever, Labrador
Age: 5 Yrs. 2 Mos.

History:

(b) (6) returner to (b) (6) for increased respiratory rate. The owner reports after the visit yesterday and the lasix bolus, (b) (6) did well until evening. Throughout the evening and night her respiratory rate increased to over 40/min. This afternoon she began to cough. The owner reports she coughed up pink tinged fluid. She also had an episode where she was excited and collapsed. She has been taking all of her medications as previously directed. She had her midday dose of Vetmedin and lasix. She is currently on Proin 50 mg 1/2 BID, Apoquel 16 mg 3/4 PO SID PM, Incurin 1 mg 2 PO SID, Vetmedin 5 mg 1 and 1/2 PO TID (for the past 2 days), Enalapril 10 mg ! PO BID, Furosemide 40 mg 1 and 1/2 PO BID, and Spironalactone 25 mg po BID. She has been dry heaving on the way here this morning. She has a history of allergies and is on a Venison and Lentil diet.

Physical Exam:

Vitals:

	(b) (6)
	7:05 PM
Vital Sign	211
Weight	27.4 kilograms
Attitude	0 - BAR
Temp	101.4
HR	180
RQ	Panting
Muc	Pale Pink
Memb	
CRT	<2 sec

Mucous Membranes: Pale pink, moist
Heart and lungs: 4/6 murmur, Fine crackles right dorsal lung fields/no dyspnea, regular rhythm, strong and synchronous femoral pulses
Abdominal Exam: Soft and not painful to palpation,
Musculoskeletal and Integument: Ambulatory with no lameness
Eyes, Ears, Nose, and Throat: No ocular or nasal discharge
Peripheral lymph nodes: No peripheral lymphadenopathy
Urogenital Exam: Unremarkable
Neuro Exam: Intact CN, normal mentation, no ataxia seen
Rectal Examination: Formed stool with no blood

(b) (6) (b) (6) respiratory rate continued to increase throughout the night despite being on a lasix CRI. Called owner and discussed poor prognosis. Owner elect humane euthanasia. (b) (6) also spoke to owner for euthanasia consent per phone consultation. 10 ml fatal plus IV.

Diagnostics:

Radiographs-

The cardiac silhouette is again noted to be generally enlarged. There is an unstructured interstitial pulmonary pattern within the right

middle and right caudal lung lobes. There is mild enlargement of the cranial lobar pulmonary veins. There are no abnormalities of the pleural space.

Conclusion

1. Persistent generalized cardiomegaly with evidence of left-sided congestive heart failure characterized by cardiogenic pulmonary edema and pulmonary venous congestion.

(b) (6), DVM, Diplomate ACVR

The study includes 3 projections of the thorax dated (b) (6). The study is compared with a prior exam from yesterday

(b) (6)

The cardiac silhouette is again noted to be generally enlarged. There is a persistent unstructured interstitial pulmonary pattern within the right middle and right caudal lung lobes. This is relatively unchanged since the prior study. There is persistent enlargement of the cranial lobar pulmonary veins. There are no abnormalities of the pleural space.

Conclusion

1. Persistent generalized cardiomegaly with persistent left-sided congestive heart failure characterized by cardiogenic pulmonary edema and pulmonary venous congestion.

(b) (6), DVM, Diplomate ACVR

Diagnosis:

Endocardiosis

Dilated cardiomyopathy

Treatment:

Lasix bolus 80 mg IV

Placed IV catheter

Monitored respiratory rate

Lasix CRI at 0.5 mg/kg/hr throughout the night

Due to poor response to treatment and declining condition owners elected humane euthanasia

Releasing DVM:

Client Signature

(b) (6)

Client Name (Print)

Patient Information

Patient:	(b) (6)	Age: 4 years	Referring Veterinarian: (b) (6)
Patient Number:	(b) (6)	Weight:(kg) 14.80	Cardiologist: (b) (6) DVM, DACVIM (Cardiology)
Breed:	Cocker Spaniel	Sex:	Client Number: (b) (6)
Exam Date:	09/05/2017 08:21	BSA: 0.61	

History: (b) (6) was presented to (b) (6) for evaluation of an enlarged heart and congestive heart failure diagnosed on radiographs on 8/22/17. (b) (6) has no history of a heart murmur. (b) (6) was taken to his regular vet on 8/22/17 for evaluation of a week long progressive cough. The clients report that (b) (6) was coughing about two times per day and the cough became more severe over the course of the week. (b) (6)'s RDVM ran blood work and took thoracic radiographs at that time. The clients report that initially his RDVM was concerned with pneumonia and prescribed antibiotics, but once the radiographs were reviewed heart failure was diagnosed and furosemide and enalapril were recommended. The clients report that they had not started the antibiotics prior to starting the Lasix and Enalapril. The cough went away after starting the furosemide and enalapril, but (b) (6) regular vet wanted (b) (6) on the antibiotics as well. The cough is greatly improved, however (b) (6) does still cough every now and then. (b) (6) is currently receiving furosemide 12.5 mgs once daily, enalapril 5 mgs in the morning and 2.5 mgs in the evenings, amoxicillin 500mgs twice daily and doxycycline 100 mgs twice daily.

Physical Examination: Temp: 104.4 HR: 150 RR: panting. Quiet/distant heart sounds with gallop, no audible murmur on left, grade 1/6 systolic murmur left. Regular rhythm. Fine crackles bilaterally. Normal abdominal palpation. Femoral pulses difficult to assess due to shivering, suspect decreased. Palpable jugular pulsation. Good hydration, normal refill, pink mm. Suspect epulis on gingiva associated with left upper canine. Fundic exam WNL.

Diagnostic Tests:

- Blood Pressure: 160 mmHg with a 4 cm cuff on the left forelimb
- Profile: albumin 2.5 (increased from 2.1), otherwise unremarkable
- U/A: USG 1.015, pH 8.0, trace protein
- Echo: See Below. ECG during echo showed a sinus tachycardia, heart rate averaging about 200bpm.

Echocardiographic Report

2D ECHO

LA Systolic Diameter LX	4.3 cm	Aortic Root Diameter	1.5 cm
-------------------------	--------	----------------------	--------

DOPPLER

AV Peak Velocity	101 cm/s	PV Peak Velocity	64.6 cm/s
AV Peak Gradient	4.1 mmHg	PV Peak Gradient	1.7 mmHg
Mitral E Point Velocity	166 cm/s	TR Peak Velocity	337 cm/s
MR Peak Velocity	467 cm/s	TR Peak Gradient	45.4 mmHg

M-MODE

LV Diastolic Diameter MM	4.7 cm	LVPW Diastolic Thickness MM	0.71 cm
LV Systolic Diameter MM	4.2 cm	LVPW Systolic Thickness MM	0.82 cm
LV Fractional Shortening MM	11.4 %	LVPW Percent Thickening MM	0.15
LV Diastolic Volume Cube	104 cm ³	IVS to PW Ratio MM	0.99
LV Systolic Volume Cube	72.3 cm ³	LV Mass MM	105 g
LV Ejection Fraction Cube	0.3	LV Mass Normalized MM	173 g/m ²
IVS Diastolic Thickness MM	0.71 cm	LA Systolic Diameter MM	3.1 cm
IVS Systolic Thickness MM	0.71 cm	Aortic Root Diameter MM	1.5 cm
IVS Percent Thickening MM	0.011	MV E Point Septal Separation	1.8 cm

Left Ventricle: Moderate dilation with increased sphericity and severe global decrease in contractility.

Left Atrium: Moderate dilation.

Right Ventricle: Mild dilation with decreased contractility.

Right Atrium: Mild dilation with decreased contractility.

Mitral Valve: 3+ central regurgitation, fused inflow.

Aortic Valve: Normal.

Tricuspid Valve: Multiple 2-3+ jets of regurgitation. TR velocity is increased consistent with moderate pulmonary hypertension.

Pulmonic Valve: Normal.

Aorta: Normal.

Pericardium: Normal. No free fluid in the abdomen, distended hepatic vessels.

Diagnosis

Dilated cardiomyopathy - This is a disease characterized by weakening of the heart muscle and dilation of the heart chambers. As the disease progresses, it can lead to congestive heart failure (fluid in the lungs causing shortness of breath and cough). Abnormal heart rhythms are common and can result in sudden death. Most commonly this is an inherited disease, though it can occur secondary to a deficiency in an amino acid called taurine.

Congestive heart failure - We did not repeat radiographs today, but based on the finding of crackles on physical exam, I suspect that there is still some mild fluid in (b) (6) lungs today.

Mild decrease in blood albumin (protein) - This value is increased from the initial bloodwork, though still just mildly low today. We will keep an eye on this, and if it is persistent or progressive we can evaluate further. It is possible that this could be due to heart failure if there had previously been free fluid in (b) (6) abdomen as well as in his lungs.

Recommendations

Please INCREASE:

Furosemide (Lasix, Salix) 12.5mg tablets - INCREASE to 1 tablet by mouth every 12 hours. This is a diuretic (water pill), that prevents the body from retaining excessive sodium and water. It will cause your pet to drink and urinate more frequently. It is important that fresh water is always available.

Enalapril (Enacard, Vasotec) 5mg tablets - INCREASE to 1 and 1/2 tablets by mouth every 12 hours. This medication is a strong drug that dilates blood vessels, permitting the heart to pump blood more efficiently. It can lower blood pressure (hypotension) and cause changes in kidney function and electrolyte values. If your pet develops weakness or depression, decrease the drug dose by 1/2 and call. A kidney panel and blood pressure should be reevaluated 7-10 days after beginning this medication.

Please ADD:

Spironolactone (Aldactone) 25mg tablets - Give one tablet by mouth once daily. This is a diuretic (water pill) that also blocks a hormone that can injure the heart muscle. It works well in combination with the furosemide and enalapril.

Pimobendan (Vetmedin) 5mg tablets - Give one tablet by mouth in the morning and 1/2 tablet by mouth in the evening. This is a drug that is approved for the treatment of congestive heart failure secondary to dilated cardiomyopathy or chronic valve disease (endocardiosis). Studies have shown improved quality of life and increase survival time when this drug is added to other standard cardiac medications. In our experience, side effects are uncommon, but it is important that you advise us if you feel your pet is having any potential adverse effects from this medication. The reported potential side effects listed for this medication are increased heart rate, vomiting, diarrhea, inappetance, uneasiness, incoordination, convulsions, increased drinking and increase urinating.

While we are waiting for taurine results, consider supplementing:

Taurine 500mg tablets - Give one tablet by mouth every 12 hours.

L-carnitine 1g - Give 1g by mouth every 8 hours.

One thing that can be very helpful for home monitoring is checking sleeping or resting respiratory rates. A recent study showed that even pets with severe heart disease rarely have resting respiratory rates greater than 30 breaths per minute unless they are starting to decompensate for that disease. Elevated respiratory rates at home may be even more sensitive than chest radiographs at picking up early decompensation. Count your pet's respiratory rate when he/she is at rest or sleeping (not within 20 minutes of being active). If his/her respiratory rate is greater than 30 breaths per minute, recheck again in a couple of hours. If persistently elevated above this level, call.

I also recommend considering a new diet with a different protein source. While I do not know of any documented amino acid deficiencies associated with a kangaroo diet, I also have two littermates that I diagnosed with severe dilated cardiomyopathy that were both fed a kangaroo diet for a long time. In that case, an inherited form of disease is possible, and in (b) (6), either an inherited or taurine-associated form of disease is possible, but the connection does bother me. With advanced heart disease, our biggest dietary concerns are adequate calorie content and low sodium content. We aim for less than 80mg sodium per 100 kilocalories (kcal) in patients that have developed congestive heart failure. We do not advise protein restriction unless there is concurrent kidney disease (i.e. kidney diets are not advised unless there is concurrent kidney disease). Please refer to our diet handouts with a list of currently adequate diets and treats, though this list is not exclusive. If you wish to feed a diet that is not on these lists, you will need to call the manufacturer of the diet to obtain a sodium content.

Exercise is also a concern in advanced heart disease. While cage rest is ideal with active heart failure, some exercise is permissible in asymptomatic disease. However, vigorous or extended exercise should be avoided.

I would like to recheck (b) (6) again in another 7-10 days for chest radiographs, kidney panel, and bloodwork on the new medications. Please call if you have any questions or concerns in the meantime. We will call when we receive taurine level results (this can take a couple of weeks sometimes).

(b) (6) (Cardiology)

(Electronically Signed)

Final Date: 05 September 2017 12:17

Amended: 05 September 2017 12:28

Like us on Facebook!

www.facebook.com/ (b) (6)

Notes to our clients

- Please bring all medications to your pet's scheduled appointments.
- We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).
- Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.
- If an emergency arises with your pet, (b) (6) is a 24 hour facility.

(b) (6)

(b) (6)

Client ID:	(b) (6)	Patient ID:	(b) (6)
Client Name:	(b) (6)	Name:	(b) (6)
Spouse/Other:	(b) (6)	Breed:	Spaniel, Cocker
Address:	(b) (6)	Sex:	Neutered Male
		Color:	Black/ White/ Brown
Telephone:	(b) (6)	Age:	4 Yrs. 3 Mos.
		DOB:	(b) (6)

Referring Veterinarian: (b) (6)
Practice: (b) (6)
Phone: (b) (6)
FAX: (b) (6)

Cardiology Reevaluation

Reevaluation of:

Congestive heart failure, left sided, Dilated cardiomyopathy, Hypoalbuminemia.

(b) (6) continues to do well at home, without any weakness or collapse. The owners report that (b) (6) has great energy levels and loves to play. The owners have reported resting respiratory rates at 14-16 bpm, without any coughing. (b) (6) has a normal appetite with normal eliminations, though did vomit clear liquid once 1.5 weeks ago. The owner reports that the mass or (b) (6) gums does not seem to affect his chewing anymore.

Physical Exam:

	9/19/2017	9/19/2017	11/2/2017
	3:06 PM	3:12 PM	1:03 PM
Vital Sign	CAR	038	399
Weight	15.5 kilograms	15.5 kilograms	16.5 kilograms
Attitude	0 - BAR		0 - BAR
Temp	102.4	102.4	103.2
HR	178	180	168
RR			110
RQ	Panting	Panting	Panting
Muc			Pink
Memb			
CRT	<2 sec		<2 sec
BP	152	152	
	#4/LF	4/LF	

Quiet heart sounds. Gallop present. Fair femoral pulses. Regular rhythm. Normal lung sounds. Normal jugular veins. Palpable hepatomegaly. Epidermal collarettes with exudative crusting on ventral abdomen. PLNs WNL. Unchanged appearance to growth on gingiva. MM pink/moist. CRT < 2 sec.

Diagnostics:

Thoracic radiographs: Decrease in heart size as compared to previous films. No evidence of cardiac decompensation. Renal panel: BUN 32 mg/dL, otherwise unremarkable. Taurine level: pending, with call with results

Diagnosis:

Congestive heart failure, left sided
Dilated cardiomyopathy
Hypoalbuminemia

Superficial dermatitis

Recommendations:

Please give the following medications as directed:

ITEM DESCRIPTION	DIRECTIONS
Vetmedin 5 mg tab	Give 1 tablet by mouth in the mornings and 1/2 tablet by mouth in the evenings.
L-Carnitine 500mg tablets	Give 1 tablet by mouth every 12 hours.
Taurine 500mg tablets	Give 1 tablet by mouth every 8 hours.
Spirolactone 25mg tablets	Give 1 tablet by mouth once every 24 hours.
Enalapril 5mg tablets	Give 1 and 1/2 tablets by mouth every 12 hours.
Furosemide 12.5mg tablet	Give 1 tablet by mouth every 12 hours.

ADD:

Simplicef 100mg tablets - Give 1 tablet by mouth once every 24 hours for 10 days.

(b) (6) has some lesions on his abdomen that are characteristic of a superficial skin infection. Simplicef is a good antibiotic for uncomplicated skin infections. (b) (6) should be re-evaluated by your primary veterinarian or a veterinary dermatologist if he does not improve.

We will call you with (b) (6) bloodwork results when they are available.

Please continue to monitor (b) (6) for cough, lethargy and/or changes in respiratory rate/effort.

*** As long as (b) (6) continues to do well at home we would like to re-evaluate him in 3-4 months. At this time we will recheck his kidney values/electrolytes, repeat chest x-rays and repeat an echocardiogram.

Like us on Facebook!!

www.facebook.com, (b) (6)

*****Notes to our clients*****

-Please bring all medications to your pet's scheduled appointments.

-We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS OUTSIDE OF (b) (6) (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays, and weekends) MAY BE ASSOCIATED WITH AN AFTER HOURS FILLING FEE.

-Check out www.goodrx.com and enter your local zip code to search for the best prices on your medications at your local pharmacies.

-If an emergency arises with your pet, (b) (6) is only a phone call away. (b) (6) is a 24 hour facility and the emergency veterinarians can always reach the cardiologist on-call.

-Please schedule your recommended recheck as soon as possible. Our schedule tends to book up quite quickly and we want to make sure that we see your pet in a timely manner.

Report Details - EON-345835		
ICSR:	2040532	
Type Of Submission:	Initial	
Report Version:	FPSR.FDA.PETF.V.V1	
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)	
Reporting Type:	Voluntary	
Report Submission Date:	2018-01-22 18:42:17 EST	
Reported Problem:	Problem Description: (b) had been presented to his regular veterinarian for a week long history of cough on August 22, 2017. Radiographs were taken and (b) was initially placed on antibiotics for presumed pneumonia (amoxicillin and doxycycline). Radiographs were reviewed by a radiologist who revised the diagnosis to congestive heart failure and furosemide and enalapril were added. I evaluated (b) on September 5, 2017 and diagnosed him with dilated cardiomyopathy with congestive heart failure.	
	Date Problem Started: 08/15/2017	
	Concurrent Medical Problem: No	
	Outcome to Date: Better/Improved/Recovering	
Product Information:	Product Name: Zignature Kangaroo Limited Ingredient Formula Dry Dog Food	
	Product Type: Pet Food	
	Lot Number:	
	Package Type: BAG	
	Possess Unopened Product: No	
	Possess Opened Product: No	
	Product Use Information:	Description: (b) had been fed this diet since he was one year of age because it was recommended by Pet People. He was also given grain free treats, carrots and apples.
		Time Interval between Product Use and Adverse Event: 3 Years
		Product Use Stopped After the Onset of the Adverse Event: Yes
		Product Use Started Again: No
Perceived Relatedness to Adverse Event: Probably related		
Manufacturer /Distributor Information:		
Purchase Location Information:		
Animal Information:	Name: (b) (6)	
	Type Of Species: Dog	
	Type Of Breed: Spaniel - Cocker American	
	Gender: Male	
	Reproductive Status: Neutered	
	Weight: 15.5 Kilogram	
	Age: 4 Years	
Assessment of Prior Health: Excellent		

FDA-CVM-FOIA-2019-1704-000486

	Number of Animals Given the Product:	4										
	Number of Animals Reacted:	4										
	Owner Information:	<table border="1"> <tr> <td>Owner Information provided:</td> <td>Yes</td> </tr> <tr> <td rowspan="4">Contact:</td> <td>Name: (b) (6)</td> </tr> <tr> <td>Phone: (b) (6)</td> </tr> <tr> <td>Other Phone: (b) (6)</td> </tr> <tr> <td>Email: (b) (6)</td> </tr> <tr> <td rowspan="2">Address:</td> <td>(b) (6)</td> </tr> <tr> <td>United States</td> </tr> </table>	Owner Information provided:	Yes	Contact:	Name: (b) (6)	Phone: (b) (6)	Other Phone: (b) (6)	Email: (b) (6)	Address:	(b) (6)	United States
Owner Information provided:	Yes											
Contact:	Name: (b) (6)											
	Phone: (b) (6)											
	Other Phone: (b) (6)											
	Email: (b) (6)											
Address:	(b) (6)											
	United States											
	Healthcare Professional Information:	<table border="1"> <tr> <td>Practice Name:</td> <td>(b) (6)</td> </tr> <tr> <td rowspan="4">Contact:</td> <td>Name: (b) (6)</td> </tr> <tr> <td>Phone: (b) (6)</td> </tr> <tr> <td>Other Phone: (b) (6)</td> </tr> <tr> <td>Email: (b) (6)</td> </tr> <tr> <td rowspan="2">Address:</td> <td>(b) (6)</td> </tr> <tr> <td>United States</td> </tr> </table>	Practice Name:	(b) (6)	Contact:	Name: (b) (6)	Phone: (b) (6)	Other Phone: (b) (6)	Email: (b) (6)	Address:	(b) (6)	United States
Practice Name:	(b) (6)											
Contact:	Name: (b) (6)											
	Phone: (b) (6)											
	Other Phone: (b) (6)											
	Email: (b) (6)											
Address:	(b) (6)											
	United States											
Sender Information:	Name:	(b) (6)										
	Address:	(b) (6) United States										
	Contact:	<table border="1"> <tr> <td>Phone:</td> <td>(b) (6)</td> </tr> <tr> <td>Other Phone:</td> <td>(b) (6)</td> </tr> <tr> <td>Email:</td> <td>(b) (6)</td> </tr> </table>	Phone:	(b) (6)	Other Phone:	(b) (6)	Email:	(b) (6)				
Phone:	(b) (6)											
Other Phone:	(b) (6)											
Email:	(b) (6)											
	Permission To Contact Sender:	Yes										
	Preferred Method Of Contact:	Phone										
	Reported to Other Parties:	Other										
Additional Documents:	Attachment:	cardio0011.pdf										
	Description:	Initial visit with echo report and most recent recheck medical record										
	Type:	Medical Records										
	Attachment:	cardio0012.pdf										
	Description:	Taurine level										
	Type:	Laboratory Report										

Report Details - EON-345965							
ICSR:	2040808						
Type Of Submission:	Initial						
Report Version:	FPSR.FDA.PETF.V.V1						
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)						
Reporting Type:	Voluntary						
Report Submission Date:	2018-01-25 12:18:44 EST						
Reported Problem:	Problem Description: At his scheduled visit to my clinic, thoracic radiographs showed generalized cardiomegaly which had been progressive compared to prior chest radiographs from his regular veterinarian but there was no evidence of cardiogenic edema. Echocardiogram was performed which showed dilated cardiomyopathy. Fundic exam was abnormal with a suspected partial retinal detachment OS. Diet history revealed that (b) was eating a kangaroo based diet. At this time the patient was continued on the Cough-tabs, Lasix was discontinued, and Vetmedin (2.5mg a.m., 1.25mg p.m.), enalapril (1.25mg BID), and taurine (500mg BID) were started. Taurine was discontinued after a normal taurine level was received. Cough persisted despite these changes and a course of doxycycline was prescribed (50mg BID x 10days). The cough improved significantly but did not completely resolve so the doxycycline was continued an additional 14 days. The dog has since been lost to follow-up. I have attempted to contact the owner and am waiting for a response. I did contact the referring veterinarian and to their knowledge the dog is still alive.						
	Date Problem Started: 04/24/2017						
	Concurrent Medical Problem: Yes						
	Pre Existing Conditions: (b) was presented to me for evaluation of lethargy and progressive cough of 6 months duration. He had been treated with a cough suppressant (Cough-tabs 1/2 tab PO BID) and furosemide (5mg once daily) prior to presentation with no response.						
	Outcome to Date: Unknown						
Product Information:	Product Name: limited ingredient diet with kangaroo as protein source - manufacturer not specified in written history (we have attempted to contact the owner but they do not return phone calls)						
	Product Type: Pet Food						
	Lot Number:						
	Possess Unopened Product: Unknown						
	Possess Opened Product: Unknown						
	Storage Conditions: Unknown						
	Product Use Information: <table border="1"> <tr> <td>Description:</td> <td>History in medical record describes diet but does not indicate duration of administration.</td> </tr> <tr> <td>Product Use Stopped After the Onset of the Adverse Event:</td> <td>No</td> </tr> <tr> <td>Perceived Relatedness to Adverse Event:</td> <td>Probably related</td> </tr> </table>	Description:	History in medical record describes diet but does not indicate duration of administration.	Product Use Stopped After the Onset of the Adverse Event:	No	Perceived Relatedness to Adverse Event:	Probably related
	Description:	History in medical record describes diet but does not indicate duration of administration.					
	Product Use Stopped After the Onset of the Adverse Event:	No					
	Perceived Relatedness to Adverse Event:	Probably related					
Manufacturer /Distributor Information:							
Purchase Location Information:							
Name: (b)							
Animal Information:	Type Of Species: Dog						
	Type Of Breed: Shih Tzu						
	Gender: Male						

	Reproductive Status:	Neutered	
	Weight:	6.08 Kilogram	
	Age:	8 Years	
	Assessment of Prior Health:	Good	
	Number of Animals Given the Product:	5	
	Number of Animals Reacted:	5	
	Owner Information:	Owner Information provided:	Yes
		Contact:	Name: (b) (6) Phone: (b) (6) Other Phone: (b) (6) Email: (b) (6)
		Address:	(b) (6) United States
	Healthcare Professional Information:	Practice Name:	(b) (6)
	Contact:	Name: (b) (6) Phone: (b) (6) Other Phone: (b) (6) Email: (b) (6)	
	Address:	(b) (6) United States	
Sender Information:	Name:	(b) (6)	
	Address:	(b) (6) United States	
	Contact:	Phone: (b) (6) Other Phone: (b) (6) Email: (b) (6)	
	Permission To Contact Sender:	Yes	
	Preferred Method Of Contact:	Email	
	Reported to Other Parties:	Other	
Additional Documents:	Attachment:	cardio0030.pdf	
	Description:	Labwork including CBC, profile, taurine level and radiology report	
	Type:	Laboratory Report	
	Attachment:	cardio0029.pdf	
	Description:	Medical records from initial and follow-up visit	
	Type:	Medical Records	

FDA-CVM-FOIA-2019-1704-000489

Report Details - EON-350158			
ICSR:	2044632		
Type Of Submission:	Initial		
Report Version:	FPSR.FDA.PETF.V.V1		
Type Of Report:	Adverse Event (a symptom, reaction or disease associated with the product)		
Reporting Type:	Voluntary		
Report Submission Date:	2018-03-27 15:12:36 EDT		
Reported Problem:	Problem Description:	At the time of diagnosis (10/31/17), Lucy was a 13 year old female spayed Labrador retriever who had been maintained on a Zignature Kangaroo formula. She presented with a history of a progressive cough which, prior to presentation, became productive and she coughed up a small volume of pink foam (possible pulmonary edema). On examination she had a 2/6 left apical systolic heart murmur and on echo diagnosed with advanced dilated cardiomyopathy with severe left ventricular dilation, moderate to severe left ventricular systolic dysfunction, and moderate to severe left atrial dilation. Thoracic radiographs were suspicious for early congestive heart failure. A whole blood taurine level was submitted and was low at 168. She was treatment with furosemide, benazepril, pimobendan, spironolactone, taurine and l-carnitine and her diet was changed to Royal Canin Early Cardiac. At her recheck in 2/26/18, (b) (6) heart had improved significantly with now mild dilated cardiomyopathy with normalized left atrial dimensions, mild left ventricular dilation and low normal left ventricular systolic function. The furosemide was able to be discontinued at this time.	
	Date Problem Started:	10/31/2017	
	Concurrent Medical Problem:	No	
	Outcome to Date:	Better/Improved/Recovering	
Product Information:	Product Name:	Zignature Kangaroo Formula	
	Product Type:	Pet Food	
	Lot Number:		
	Package Type:	BAG	
	Possess Unopened Product:	Unknown	
	Possess Opened Product:	Unknown	
	Product Use Information:	Product Use Stopped After the Onset of the Adverse Event:	Yes
		Adverse Event Abate After Product Stop:	Yes
		Product Use Started Again:	No
		Perceived Relatedness to Adverse Event:	Probably related
		Other Foods or Products Given to the Animal During This Time Period:	Unknown
Manufacturer /Distributor Information:	Name:	Pets Global - Zignature	
	Type(s):	Manufacturer	
	Address:	28334 Industry Dr Valencia California 91355 United States	
FDA-CVM-FOIA-2019-1704-000491			

		Contact:	Phone: (661) 309-1235	
			Web Address: www.zignature.com	
		Possess One or More Labels from This Product:	Yes	
	Purchase Location Information:			
Animal Information:	Name:	(b)		
	Type Of Species:	Dog		
	Type Of Breed:	Retriever - Labrador		
	Gender:	Female		
	Reproductive Status:	Neutered		
	Weight:	33.18 Kilogram		
	Age:	13 Years		
	Assessment of Prior Health:	Good		
	Number of Animals Given the Product:	1		
	Number of Animals Reacted:	1		
	Owner Information:	Owner Information provided:	Yes	
		Contact:	Name:	(b) (6)
			Phone:	(b) (6)
			Other Phone:	(b) (6)
			Email:	(b) (6)
	Address:	(b) (6)		
		United States		
Healthcare Professional Information:	Practice Name:	(b) (6)		
	Contact:	Name:	(b) (6)	
		Phone:	(b) (6)	
		Email:	(b) (6)@cvcavets.com	
	Address:	(b) (6)		
		United States		
	Practice Name:	(b) (6)		
	Contact:	Name:	(b) (6)	
		Phone:	(b) (6)	
		Email:	(b) (6)@cvcavets.com	
	Address:	(b) (6)		
		United States		
	Type of Veterinarian:	Referred veterinarian		
	Permission to Release Records	Yes		

FDA-CVM-FOIA-2019-1704-000492

		to FDA:
Sender Information:	Name:	(b) (6)
	Address:	(b) (6) United States
	Contact:	Phone: (b) (6) Email: (b) (6)@cvcavets.com
	Permission To Contact Sender:	Yes
	Preferred Method Of Contact:	Email
	Reported to Other Parties:	Other
Additional Documents:	Attachment:	(b) (6) Echo Report 2017-10-31.pdf
	Description:	Echocardiogram 10-31-2017
	Type:	Echocardiogram
	Attachment:	(b) (6) Echo Report 2018-02-26.pdf
	Description:	Echocardiogram 2-26-2018
	Type:	Echocardiogram
	Attachment:	(b) (6) Taurine Level 2017-11-03.pdf
	Description:	BW Taurine Level 11-3-2017
	Type:	Laboratory Report



Selenium bioavailability: current knowledge and future research requirements^{1 5}

Susan J Fairweather-Tait, Rachel Collings, and Rachel Hurst

ABSTRACT

Information on selenium bioavailability is required to derive dietary recommendations and to evaluate and improve the quality of food products. The need for robust data is particularly important in light of recent suggestions of potential health benefits associated with different intakes of selenium. The issue is not straightforward, however, because of large variations in the selenium content of foods (determined by a combination of geologic/environmental factors and selenium supplementation of fertilizers and animal feedstuffs) and the chemical forms of the element, which are absorbed and metabolized differently. Although most dietary selenium is absorbed efficiently, the retention of organic forms is higher than that of inorganic forms. There are also complications in the assessment and quantification of selenium species within foodstuffs. Often, extraction is only partial, and the process can alter the form or forms present in the food. Efforts to improve, standardize, and make more widely available techniques for species quantification are required. Similarly, reliable and sensitive functional biomarkers of selenium status are required, together with improvements in current biomarker methods. This requirement is particularly important for the assessment of bioavailability, because some functional biomarkers respond differently to the various selenium species. The effect of genotype adds a potential further dimension to the process of deriving bioavailability estimates and underlines the need for further research to facilitate the process of deriving dietary recommendations in the future. *Am J Clin Nutr* 2010;91(suppl):1484S-91S.

INTRODUCTION

To derive selenium requirements and establish dietary recommendations for optimal health, estimates of selenium bioavailability are needed. A literature review on the bioavailability of selenium from foods was published in 2006 (1), and it highlights the dependence of bioavailability on food sources associated with different forms of selenium and emphasizes the importance of the assessment of bioavailability with the use of functional assays. Data on chemical speciation and metabolic transformations (in conjunction with information on the relation between selenium intake and status and health outcomes) are required to assess selenium bioavailability and the longer term health consequences that result from different intakes.

DIETARY REQUIREMENTS

The 1991 UK Dietary Reference Values (2) used data from older literature and estimated that between 55% and 65% of

dietary selenium is absorbed. The 1993 Population Reference Intakes published by the European Scientific Committee for Food (3) concluded that for selenium “all usual dietary forms are absorbed quite efficiently.” The 2000 report of the US Food and Nutrition Board (4) suggested that most dietary selenium is highly bioavailable: >90% of selenomethionine is absorbed; selenocysteine appears to be absorbed very well; ≈100% of selenate is absorbed, but a significant fraction is lost in the urine; and >50% of selenite is absorbed (depending on luminal interactions) and is better retained than selenate. There is clearly a need to review dietary recommendations in light of more recent data, in particular, information on dietary forms of selenium and the relationships between intake and health outcomes.

SELENIUM SPECIATION

A recent review (5) provides information on the forms of selenium in food and associated health effects; technical approaches used for speciation have also been reviewed recently (6, 7). The analysis of forms of selenium in food is a challenging task; there are currently no methods that can reliably extract 100% of the selenium from foods without potentially affecting the species, and the techniques are established in only a few laboratories worldwide. Therefore, care has to be taken to extract as much selenium as possible while still retaining the form that is present in the food as consumed; conditions that are devised to maximize the extraction of selenium from a food matrix may cause changes in chemical form. Ideally, the measurements should be made in food that has gone through processing (eg, cooking) followed by simulated gastrointestinal digestion, be

¹ From the School of Medicine, Health Policy and Practice, University of East Anglia, Norwich, United Kingdom.

² Presented at the workshop “Micronutrient Bioavailability: Priorities and Challenges for Setting Dietary Reference Values,” held in Barcelona, Spain, 11–12 June 2009.

³ This article does not necessarily reflect the views of the Commission of the European Communities and in no way anticipates future policy in this area.

⁴ Supported by the Commission of the European Communities, specific RTD Programme “Quality of Life and Management of Living Resources,” within the 6th Framework Programme (contract no. FP6 036196 2 EURRECA: EUROpean micronutrient RECommendations Aligned).

⁵ Address correspondence to SJ Fairweather Tait, School of Medicine, Health Policy & Practice, University of East Anglia, Norwich, NR4 7TJ, United Kingdom. E mail: s.fairweather.tait@uea.ac.uk.

First published online March 3, 2010; doi: 10.3945/ajcn.2010.28674J.

cause this is the form present in the lumen of the gut that is of interest. Although it has not been possible to produce comprehensive data that describe forms of selenium in food, there are limited data on the percentage distribution of different species (expressed as percentage of extractible or total selenium); examples are given in **Table 1**.

The selenium content and species of both plant and animal foodstuffs depend on environmental conditions, in particular, the quantity and species of selenium to which the animal/plant is exposed (6, 24). Selenomethionine is predominant in cereals, and selenium concentrations vary from 0.01 to 0.55 $\mu\text{g/g}$ fresh weight (5), whereas in other plant foods the content is generally lower, with the exception of Brazil nuts and vegetables, which are selenium-accumulating plants, namely those in the allium and brassica families. The selenium content of Brazil nuts varies depending on soil content and other environmental factors, and nuts from trees in the central part of Brazil contain ≤ 10 times more selenium than those from West Brazil (6). The reason for the high content of selenium in Brazil nuts is that the proteins are high in sulfur-containing amino acids, and selenomethionine can nonspecifically replace methionine. The major species in non-selenium-accumulating plant foods are selenate and selenomethionine, plus smaller amounts of selenocysteine. In contrast, the predominant form of selenium in selenium-accumulating plants is γ -glutamyl methylselenocysteine (13, 14). There are limited data on the forms of selenium in animal foodstuffs, but it appears that the major forms are selenomethionine and selenocysteine, which are incorporated nonspecifically into muscle protein (19). In addition, selenate and selenite have been detected in fish (18, 20) and there appear to be large differences between fish species in relation to selenoproteins (25). In foods of animal origin, supplementation with organic compared with inorganic selenium results in meat of higher selenium concentration. For example, when a comparison is made between the effect of selenium yeast and sodium selenite supplements, skeletal muscle from lambs contained 0.12 and 0.08 μg selenium/g fresh weight, respectively (26), and beef contained 0.41 and 0.30 mg/kg dry weight, respectively (27).

ABSORPTION, RETENTION, AND METABOLISM

Data on selenium metabolism from different foods and selenium supplements indicate differences in the absorption and use of selenium between inorganic and organic forms in humans (28, 29) and rats (30). The absorptive pathways have not yet been fully characterized, but selenium as selenate or selenite appears to be very well absorbed but less well retained in the body than organic forms of selenium, such as selenomethionine and selenocysteine (31–33). The proposed metabolic pathways for different forms of selenium are shown in **Figure 1** (5). Most forms of selenium are efficiently absorbed, but subsequent metabolism depends on the form in which they are present in plasma. Selenomethionine, selenocysteine, selenate, and selenite enter the selenide pool and from here the selenium is either used for selenoprotein synthesis or excreted in the urine as a selenosugar. Selenomethionine can, however, also be incorporated directly (and nonspecifically) into proteins through the replacement of methionine. A separate pathway is followed by the organic compound, γ -glutamyl methylselenocysteine, found in brassica and allium vegetables, whereby it is first converted to Se-methylselenocysteine and

then transformed by β -lyase into methylselenol, which is primarily excreted in breath and urine but may also enter the selenide pool.

Several approaches have been used to measure the bioavailability of selenium in various foods, as summarized in **Table 2**. These include the measurement of changes in plasma selenium concentration, measurement of glutathione peroxidase (GPx) enzyme activity, and absorption/retention studies. For the last, intrinsic techniques with the use of stable isotopes of selenium have been developed to label the endogenous forms of selenium in foods (40). In general, selenium is absorbed efficiently, but it is not possible to assign specific figures for retention and use (bioavailability) to individual forms of selenium because of the complexity of many foods (Table 1). However, a study by Bügel et al (39), on the assumption that selenomethionine is the major form in meat, showed that most of the selenium was absorbed and just over half retained in the body (ie, not excreted in the urine). Selenium in Brazil nuts appeared to be better used than selenomethionine, in terms of the response of plasma selenium concentration and red blood cell GPx activities: the plasma selenium increase was similar despite the fact that the daily intake from Brazil nuts was half that from selenomethionine (35). Changes in selenium status that reflect changes in intake occur over a period of several weeks or months, although the feeding trial of Hawkes et al (38) showed a significant difference between a beef, rice, and powdered milk diet with low selenium content and one with high selenium content after only 14 d. In a study by Kirby et al (11), the plasma selenium response in a feeding trial appeared to be related to the form of selenium in wheat flour biscuits: intake of selenomethionine in biofortified wheat-biscuits resulted in a greater increase in plasma selenium after 6 mo than the oxidized selenomethionine (selenomethionine selenoxide) in fortified biscuits (Table 2).

FUNCTIONAL MEASURES

There are 25 known selenoprotein genes in humans (41, 42), which encode selenoproteins with a variety of functions, as summarized in **Table 3**. Several of the selenoproteins, which include selenoproteins P and W and the GPx 1, 3, and 4, have been used widely as biomarkers of selenium status. Functional biomarkers are only useful if they can be measured in readily accessible tissues, such as blood. At present, the most promising biomarker appears to be selenoprotein P, which appears to reach a plateau after 2–4 wk of supplementation (88, 89) and is well correlated with plasma selenium across a wide range of selenium status (90), up to a plasma selenium concentration of ≈ 125 ng/mL (33). Selenoprotein P typically accounts for approximately half of the selenium in plasma (46). It is generally more sensitive than other selenoproteins, such as GPx, in both deficiency (90) and after supplementation (89–91), and, in addition, the response of selenoprotein P to different forms of selenium appears to be similar (92).

Biomarkers of selenium status have recently been the subject of a systematic review (93), in which the response of each biomarker to either depletion or supplementation (only studies that intervened with selenomethionine or selenium-enriched yeast were included) was assessed and evaluated for different population groups. However, for most biomarkers there was



TABLE 1

Examples of forms of selenium (percentage of total or extractable selenium) in foods

Food (reference)	Typical selenium content ¹	Forms of selenium
	<i>µg/g fresh weight</i>	
Selenium-enriched yeast (5, 8)	1200–2200	60–84% Selenomethionine, usual percentage in high-quality commercial preparation of selenium-enriched yeast but values for selenomethionine content can vary: 23–83% Selenomethionine 3–21% Selenocysteine 1–20% γ -Glutamyl-Se-methylselenocysteine 4% Selenate 13–51% Other forms
Brazil nuts (<i>Bertholletia excelsa</i>) (9)	2.54 (0.85–6.86)	≈25% Selenomethionine
Wheat (8, 10)	0.1–30	12–19% Selenate/ite
	0.08–44	56–83% Selenomethionine 4–12% Selenocysteine 1–4% Se-methylselenocysteine ≈55% Selenomethionine
Wheat (biofortified) (11)	8.3	76–85% Selenomethionine
Wheat-flour (biofortified) biscuits (11)	4.4	76–85% Selenomethionine
Wheat flour (unfortified) soaked in aqueous solution of selenomethionine and baked into biscuits (11)	8.5	55% Selenomethionine selenoxide 5% Selenomethionine
Broccoli (selenium enriched) (12)	62.3 ²	45% Se-methylselenocysteine 20% Selenate 20% Selenate 12% Selenomethionine
Onions (<i>Allium cepa</i>) (13)	<0.5	100% Selenate (extractable selenium)
Onions (selenium enriched) (13)	140	63% γ -Glutamyl-Se-methylselenocysteine 10% Selenate 5% Selenomethionine
Garlic (<i>Allium sativum</i>) (13)	<0.5	53% Selenomethionine 31% γ -Glutamyl-Se-methylselenocysteine 12% Se-methylselenocysteine 4% Selenate
Garlic (selenium enriched) (14)	296	73% γ -Glutamyl-Se-methylselenocysteine (total eluted selenium) 13% Selenomethionine 4% γ -Glutamyl-selenomethionine 3% Se-methylselenocysteine 2% Selenate
Lentils (<i>Lens culinaris</i> L.) (15)	0.24–0.36	90% Organic selenium 10% Selenate
Carrots (16)	<0.05	Undetectable
Carrots (selenium enriched) (16)	0.4–2.2	Selenium-enriched with the use of selenate (% extractable): ≈54% Selenomethionine 32% Selenate ≈14% γ -Glutamyl-selenomethionine Selenium-enriched with the use of selenite: ≈71% Selenomethionine 17% Selenite ≈12% γ -Glutamyl-selenomethionine
Potatoes (17)	0.12	50% Selenomethionine (extractable) 50% Selenate (extractable)
Shellfish (18)	0.36–1.33	7.6–44.8% Selenate
Cod (19, 20)	1.5	70% Selenomethionine 12% Selenite
Tuna (canned in water) (21)	5.6	29% Selenomethionine (extractable)
Shark (21)	2.0	56% Selenomethionine (extractable)
Swordfish (22)	Not quantified	Selenomethionine, selenenyl sulfide, selenite
Chicken (23)	0.5	56–66% Selenomethionine (extractable) 20–31% Selenocysteine (extractable)
Lamb (23)	0.4	56–60% Selenomethionine (extractable) 50% Selenocysteine (extractable)

¹ Values are means and/or ranges² $\mu\text{g/g}$ dry weight




FIGURE 1. Metabolic pathway of dietary selenium in humans. Se, selenium; SeMet, selenomethionine; SeCys, selenocysteine; GSSeSG, selenodiglutathione; γ glutamyl CH_3SeCys , γ glutamyl Se methylselenocysteine; H_2Se , hydrogen selenide; HSePO_3^{2-} , selenophosphate; CH_3SeCys , Se methylselenocysteine; CH_3SeH , methylselenol; $(\text{CH}_3)_2\text{Se}$, dimethyl selenide; SeO_2 , selenium dioxide; $(\text{CH}_3)_3\text{Se}^+$, trimethyl selenonium ion. Reproduced with permission from reference 5.

a paucity of data for meaningful subgroup or dose response analysis. In the included studies plasma selenium was the most commonly measured biomarker, and it responded positively to intervention, as did whole blood and erythrocyte selenium, plasma selenoprotein P, and platelet, plasma, erythrocyte and whole blood GPx activity, albeit with significant heterogeneity in each case. The review concluded that further large scale in interventions are required to assess the usefulness of selenium

responsive biomarkers, and these could conceivably include aspects of speciation. Plasma selenium concentration reflects dietary exposure to most forms of selenium, but in the absence of a well described homeostatic regulation there is no absolute plateau, although the concentration will reach a steady state at any constant level of intake after ≈ 10 – 12 wk (33, 91, 92, 94–97). In addition to dose, the plasma response to dietary selenium is species dependent, so consumption of 2 different forms may result in different plasma selenium concentrations (33, 92, 95, 96, 98, 99).

EFFECT OF GENOTYPE

The response by individuals to 6 wk of selenium supplementation with $100 \mu\text{g}$ sodium selenite/d has been shown to be influenced by genetic polymorphisms in the selenoprotein P gene (*SEPP*) (100) and *GPX4* gene (101). Biomarkers that are commonly used to assess selenium bioavailability (plasma selenium, selenoprotein P, and GPx3) were associated with 2 common single nucleotide polymorphisms in *SEPP* in both baseline and postsupplementation samples (100). The *GPX4* polymorphism was shown to influence lymphocyte GPx4 concentration and other selenoproteins in vivo (101). A single nucleotide polymorphism in GPx1 (Pro198Leu) was associated with selenium deficiency and impaired GPx1 activity (102) and also may be associated with a different response of GPx1 activity to selenium (103). This observation raises the issue of whether common polymorphisms in selenoprotein genes, such as *SEPP*, *GPX1*, *GPX4*, and selenoprotein S (*SELS*) (92), will

TABLE 2

Bioavailability of selenium from various foods¹

Food (reference)	Technique used	Results
Selenium (Se) yeast, 300 $\mu\text{g}/\text{d}$ for 10 wk, then single dose of ⁷⁷ Se yeast (34)	Absorption from stable isotopically labeled material (327 μg selenium)	89%
	Retention (absorption minus urinary excretion)	74%
Brazil nuts, 53 $\mu\text{g}/\text{d}$ for 3 mo (35)	Plasma selenium increase	64.2%
	Plasma GPx increase	8.2%
	Whole blood GPx increase	13.2%
Selenomethionine, 100 $\mu\text{g}/\text{d}$ for 3 mo (35)	Plasma selenium increase	61%
	Plasma GPx increase	3.4%
	Whole blood GPx increase	5.3%
Biofortified wheat flour biscuits, mean intake 172 $\mu\text{g}/\text{d}$ for 6 mo (11)	Plasma selenium increase after 6 mo feeding trial	72 $\mu\text{g}/\text{L}$ increase
Fortified wheat flour biscuits, mean intake 208 $\mu\text{g}/\text{d}$ for 6 mo (11)	Plasma selenium increase after 6 mo feeding trial	16 $\mu\text{g}/\text{L}$ increase
Basal diet, 52 μg selenium + cow milk, 15 μg selenium (36)	Fractional absorption in ileostomists	65.5%
		73.3%
Shrimp, 88 $\mu\text{g}/\text{d}$ for 6 wk (37)	Plasma selenium increase	6.3 $\mu\text{g}/\text{L}$ increase
	Apparent absorption	83%
Beef, rice, and powdered milk, 14 $\mu\text{g}/\text{d}$ (low) compared with 297 $\mu\text{g}/\text{d}$ (high) for 14 d (38)	Plasma selenium change	–40 $\mu\text{g}/\text{L}$ (low); 97 $\mu\text{g}/\text{L}$ (high)
	Muscle selenium	–0.37 $\mu\text{g}/\text{g}$ protein (low); 0.57 $\mu\text{g}/\text{g}$ protein (high)
	Platelet GPx	–120 nkat/g protein (low); 100 nkat/g protein (high)
	Red blood cell selenium	–42 $\mu\text{g}/\text{L}$ (low); 106 $\mu\text{g}/\text{L}$ (high)
	Red blood cell GPx	–15 nkat/g protein (low); 13 nkat/g protein (high)
Pork, 106 $\mu\text{g}/\text{d}$ for 3 wk; 7 d metabolic balance in final week (39)	Apparent absorption	94%
	Retention	58%

¹ GPx, glutathione peroxidase; nkat, nanokatal.



TABLE 3Summary of human selenoproteins and their functions¹

Group/name (reference)	Abbreviation(s)	Location	Main functions
Glutathione peroxidases			
Glutathione peroxidase-1 (43)	GPx1, cGPx, GSH-Px	Widely expressed throughout the body, intracellular enzyme	Cystolic enzyme, antioxidant activity
Glutathione peroxidase-2 (44)	GPx2, GI-GPx	Mainly in gastrointestinal tissue, also in liver	Protection of GI tract from oxidative damage
Glutathione peroxidase-3 (45)	GPx3, eGPx	Plasma [accounts for 10–30% selenium in plasma (46)] and extracellular fluid, expressed in liver, kidney, heart, lung, thyroid, GI tract, and breast (47)	Plasma antioxidant, can decrease lipid hydroperoxides (48)
Glutathione peroxidase-4	GPx4, PHGPx	Widely expressed, high expression in the testes (49, 50); cytosolic and membrane-bound forms (51, 52)	Antioxidant activity, protects membranes from peroxidative degradation (51); can decrease phospholipid, cholesterol and cholesterol ester hydroperoxides to less toxic derivatives (52); protection against oxidatively damaged DNA (53); regulation of 15-lipoxygenase pathway (54) and 5-lipoxygenase (55); important for male fertility and sperm maturation/function/motility (56–59)
Glutathione peroxidase-6 (41)	GPx6	Embryo and olfactory epithelium	Unknown
Thioredoxin reductases			
Thioredoxin reductase-1 (60, 61)	TrxR-1, TR1, Txnrd1	Intracellular (cytosolic/nuclear), widely distributed	Regulation of intracellular redox state, cell signaling; decreases thioredoxin
Thioredoxin reductase-2 (60, 61)	TrxR-2, TR2	Mitochondrial, widely distributed	Regulation of intracellular redox state; decreases thioredoxin
Thioredoxin reductase-3 (60, 61)	TrxR-3, TR3	Testis-specific	Regulation of intracellular redox state
Iodothyronine deiodinases			
Iodothyronine 5' deiodinase-1, type 1	DIO-1, DI1, 5'IDI	Kidney, liver, thyroid, and brown adipose tissue (62–64)	Thyroid hormone metabolism, converts inactive thyroxine to active 3,3'-5'triiodothyronine; activation of thyroid hormones (65)
Iodothyronine 5' deiodinase-2, type 2 (63, 66, 67)	DIO-2, DI2	Thyroid, CNS, pituitary, brown adipose tissue, skeletal muscle	Activation of thyroid hormones
Iodothyronine 5 deiodinase-3, type 3 (68)	DIO-3, DI3	Placenta, CNS, fetus	Inactivation of thyroid hormones
Selenoproteins and other			
Selenoprotein-P (69, 70)	SeIP, Sepp1	Plasma [accounts for 30–50% of selenium in plasma (46, 71)] and also ubiquitously expressed in most tissues; high expression in brain, liver, and testes	Selenium homeostasis (72) and transport of selenium to tissues; antioxidant activity and decrease of lipid hydroperoxides (73)
Selenoprotein-W (74, 75)	SeIW	Most tissues, abundant in brain, colon, heart, skeletal muscle, and prostate	Involved in skeletal and cardiac muscle metabolism/function, antioxidant function
Selenoprotein-N	SeIN	Most tissues, ubiquitous expression, transmembrane glycoprotein associated with endoplasmic reticulum (76, 77)	Unknown, may be important in muscle and development (76)
Selenoprotein-S	SeIS	Membrane protein, located in the endoplasmic reticulum, widely expressed	Inflammatory response, regulation of inflammatory cytokines (interleukin 1 β and 6 and tumor necrosis factor alpha) (78), removal of misfolded proteins from the endoplasmic reticulum (79)
Selenoprotein-K (80)	SeIK	Membrane protein, localized to endoplasmic reticulum	Possible antioxidant activity
Selenoprotein-R (81)	SeIR / MsrB1	Cytosol and nucleus; widely expressed	Antioxidant, protein repair and methionine metabolism (82)
Selenoprotein-H (83)	SeIH	Widely expressed in tissues, localized to the nucleus	DNA binding protein, regulation of glutathione synthesis genes, and phase II detoxification
Selenoprotein-I (41)	SeII	Unknown	Unknown

(Continued)



TABLE 3 (Continued)

Group/name (reference)	Abbreviation(s)	Location	Main functions
Selenoprotein-M (84)	SelM	Localized in the endoplasmic reticulum	Protein folding in the endoplasmic reticulum, antioxidant activity
Selenoprotein-O (41)	SelO	Unknown	Unknown
Selenoprotein-T (41)	SelT	Unknown	Unknown
Selenoprotein-V (41)	SelV	Testes	Unknown
15 kD selenoprotein (85)	Sel15	Localized in the endoplasmic reticulum	Thioredoxin-like, role in unfolded protein response (86)
Selenophosphate synthetase-2	SPS-2	Unknown	Selenoprotein biosynthesis (87)

¹ The list of selenoproteins that contain selenocysteine was generated from information on the selenoprotein database SelenoDB (42). Glutathione peroxidases 5, 7, and 8 (GPx5, GPx7, GPx8); selenoproteins R2, R3, and W2 (SelR2, SelR3, SelW2); selenium-binding protein 2 (SBP2); selenophosphate synthetase1 (SPS1); and eukaryotic elongation factor (eEFSec) are not listed in the table of selenoproteins because they are homologs that contain cysteine or other amino acids that do not contain selenocysteine (SelenoDB). CNS, central nervous system; GI, gastrointestinal

have a significant effect on the metabolism of dietary selenium and will generate different figures for bioavailability. However, it is most likely that the effect of genotype on the biomarkers used to predict bioavailability is subtle and only becomes relevant when longer-term health outcomes are considered (104–106).

RESEARCH REQUIREMENTS

The bioavailability of different selenium species requires further investigation with the use of stable isotope labels, and the mechanism of absorption of the different forms of selenium needs to be elucidated. Further data on selenium species in food are required, but with the well-known extraction constraints it will not be possible to generate comprehensive information for all foods; therefore, dietary intervention studies may be required to study foods that make a major contribution to selenium intake. The native forms of selenium need to be labeled intrinsically with stable isotopes of selenium to measure uptake and retention of food selenium in acute studies, and longer-term studies need to be undertaken to measure changes in functional biomarkers; the most promising at present is selenoprotein P, but other novel biomarkers should be sought. Interactions between selenium and other micronutrients, such as vitamin E, should be taken into consideration [possibly with the use of a network biology approach (107)], particularly in relation to health outcomes that are associated with antioxidant nutrients, such as inflammation. Finally, the effect of common selenoprotein gene polymorphisms on metabolism (and hence requirements) remains to be clarified.

The authors' responsibilities were as follows—SJF-T: first draft of the manuscript; and RC and RH: contribution of sections to the manuscript draft. All authors approved the final manuscript. The authors had no personal or financial conflicts of interest.

REFERENCES

- 1 Finley JW Bioavailability of selenium from foods. *Nutr Rev* 2006;64:146–51
- 2 Department of Health. Dietary Reference Values for food energy and nutrients for the United Kingdom. London, United Kingdom: HMSO, 1991
- 3 Reports of the Scientific Committee for Food. Nutrient and energy intakes for the European Community. Luxembourg: Commission of the European Communities, 1993
- 4 Institute of Medicine. Dietary Reference Intakes: vitamin C, vitamin E, selenium, and carotenoids. Washington, DC: National Academies Press, 2000
- 5 Rayman MP, Infante HG, Sargent M. Food-chain selenium and human health: spotlight on speciation. *Br J Nutr* 2008;100:238–53
- 6 Dumont E, Vanhaecke F, Cornelis R. Selenium speciation from food source to metabolites: a critical review. *Anal Bioanal Chem* 2006;385:1304–23
- 7 Pedrero Z, Madrid Y. Novel approaches for selenium speciation in foodstuffs and biological specimens: a review. *Anal Chim Acta* 2009;634:135–52
- 8 Whanger PD. Selenocompounds in plants and animals and their biological significance. *J Am Coll Nutr* 2002;21:223–32
- 9 Barclay MNI, MacPherson A, Dixon J. Selenium concentration of a range of UK foods. *J Food Compos Anal* 1995;8:307–18
- 10 Wolf WR, Goldschmidt RJ. Updated estimates of the selenomethionine content of NIST wheat reference materials by GC-IDMS. *Anal Bioanal Chem* 2007;387:2449–52
- 11 Kirby JK, Lyons GH, Karkkainen MP. Selenium speciation and bioavailability in biofortified products using species-unspecific isotope dilution and reverse phase ion pairing-inductively coupled plasma-mass spectrometry. *J Agric Food Chem* 2008;56:1772–9
- 12 Finley JW, Ip C, Lisk DJ, Davis CD, Hintze KJ, Whanger PD. Cancer-protective properties of high-selenium broccoli. *J Agric Food Chem* 2001;49:2679–83
- 13 Kotrbai M, Tyson JF, Uden PC, Birringer M, Block E. Selenium speciation in enriched and natural samples by HPLC-ICP-MS and HPLC-ESI-MS with perfluorinated carboxylic acid ion-pairing agents. *Analyst* 2000;125:71–8
- 14 Ip C, Birringer M, Block E, et al. Chemical speciation influences comparative activity of selenium-enriched garlic and yeast in mammary cancer prevention. *J Agric Food Chem* 2000;48:2062–70
- 15 Thavarajah D, Vandenberg A, George GN, Pickering IJ. Chemical form of selenium in naturally selenium-rich lentils (*Lens culinaris* L.) from Saskatchewan. *J Agric Food Chem* 2007;55:7337–41
- 16 Kopolná E, Hillestrøm PR, Laursen KH, Husted S, Larsen EH. Effect of foliar application of selenium on its uptake and speciation in carrot. *Food Chem* 2009;115:1357–63
- 17 Cuderman P, Kreft I, Germ M, Kovačević M, Stibilj V. Selenium species in selenium-enriched and drought-exposed potatoes. *J Agric Food Chem* 2008;56:9114–20
- 18 Cappon CJ, Smith JC. Chemical form and distribution of mercury and selenium in edible seafood. *J Anal Toxicol* 1982;6:10–21
- 19 Díiz Huerta V, Fernández Sánchez ML, Sanz-Medel A. Quantitative selenium speciation in cod muscle by isotope dilution ICP-MS with a reaction cell: comparison of different reported extraction procedures. *J Anal At Spectrom* 2004;19:644–8
- 20 Crews HM, Clarke PA, Lewis DL, Owen LM, Strutt PR, Izquierdo A. Investigation of selenium speciation in *in vitro* gastrointestinal extracts of cooked cod by high-performance liquid chromatography-inductively coupled plasma mass spectrometry and electrospray mass spectrometry. *J Anal At Spectrom* 1996;11:1177–82
- 21 Reyes LH, Mar JL, Rahman GM, Seybert B, Fahrenholz T, Kingston HM. Simultaneous determination of arsenic and selenium species in



- fish tissues using microwave-assisted enzymatic extraction and ion chromatography-inductively coupled plasma mass spectrometry *Talanta* 2009;78:983–90
- 22 George GN, Singh SP, Prince RC, Pickering IJ Chemical forms of mercury and selenium in fish following digestion with simulated gastric fluid *Chem Res Toxicol* 2008;21:2106–10
 - 23 Bierla K, Dernovics M, Vaccina V, Szpunar J, Bertin G, Lobinski R Determination of selenocysteine and selenomethionine in edible animal tissues by 2D size-exclusion reversed-phase HPLC-ICP MS following carbamidomethylation and proteolytic extraction *Anal Bioanal Chem* 2008;390:1789–98
 - 24 Li H-F, McGrath SP, Zhao F-J Selenium uptake, translocation and speciation in wheat supplied with selenate or selenite *New Phytol* 2008;178:92–102
 - 25 Bergdahl IA Fractionation of soluble selenium compounds from fish using size-exclusion chromatography with on-line detection by inductively coupled plasma mass spectrometry *Analyst (Lond)* 1999;124:1435–8
 - 26 Steen A, Strom T, Bernhoft A Organic selenium supplementation increased selenium concentrations in ewe and newborn lamb blood and in slaughter lamb meat compared to inorganic selenium supplementation *Acta Vet Scand* 2008;50:7
 - 27 Juniper DT, Phipps RH, Ramos-Morales E, Bertin G Effect of dietary supplementation with selenium-enriched yeast or sodium selenite on selenium distribution and meat quality in beef cattle *J Anim Sci* 2008;86:3100–9
 - 28 Butler JA, Thomson CD, Whanger PD, Robinson MF Selenium distribution in blood fractions of New Zealand women taking organic or inorganic selenium *Am J Clin Nutr* 1991;53:748–54
 - 29 Brown KM, Pickard K, Nicol F, Beckett GJ, Duthie GG, Arthur JR Effects of organic and inorganic selenium supplementation on selenoenzyme activity in blood lymphocytes, granulocytes, platelets and erythrocytes *Clin Sci* 2000;98:593–9
 - 30 Finley JW, Davis CD Selenium (Se) from high-selenium broccoli is utilized differently than selenite, selenate and selenomethionine, but is more effective in inhibiting colon carcinogenesis *Biofactors* 2001;14:191–6
 - 31 Thomson CD, Burton CE, Robinson MF On supplementing the selenium intake of New Zealanders 1 Short experiments with large doses of selenite or selenomethionine *Br J Nutr* 1978;39:579–87
 - 32 Schrauzer GN Selenomethionine: a review of its nutritional significance, metabolism and toxicity *J Nutr* 2000;130:1653–6
 - 33 Burk RF, Norsworthy BK, Hill KE, Motley AK, Byrne DW Effects of chemical form of selenium on plasma biomarkers in a high-dose human supplementation trial *Cancer Epidemiol Biomarkers Prev* 2006;15:804–10
 - 34 Bügel S, Larsen EH, Sloth JJ, et al Absorption, excretion, and retention of selenium from a high selenium yeast in men with a high intake of selenium *Food Nutr Res*, 2008:52
 - 35 Thomson CD, Chisholm A, McLachlan SK, Campbell JM Brazil nuts: an effective way to improve selenium status *Am J Clin Nutr* 2008;87:379–84
 - 36 Chen J, Lindmark-Månsson H, Drevelius M, et al Bioavailability of selenium from bovine milk as assessed in subjects with ileostomy *Eur J Clin Nutr* 2004;58:350–5
 - 37 Bügel SH, Sandström B, Larsen EH Absorption and retention of selenium from shrimps in man *J Trace Elem Med Biol* 2001;14:198–204
 - 38 Hawkes WC, Alkan FZ, Oehler L Absorption, distribution and excretion of selenium from beef and rice in healthy North American men *J Nutr* 2003;133:3434–42
 - 39 Bügel S, Sandström B, Skibsted LH Pork meat: a good source of selenium? *J Trace Elem Med Biol* 2004;17:307–11
 - 40 Fairweather-Tait SJ Bioavailability of selenium *Eur J Clin Nutr* 1997;51:S20–3
 - 41 Kryukov GV, Castellano S, Novoselov SV, et al Characterization of mammalian selenoproteomes *Science* 2003;300:1439–43
 - 42 Castellano S, Gladyshev VN, Guigo R, Berry MJ SelenoDB 1.0: a database of selenoprotein genes, proteins and SECIS elements *Nucleic Acids Res* 2008;36:D332–8
 - 43 Flohe L, Gunzler WA, Schock HH Glutathione peroxidase: a selenoenzyme *FEBS Lett* 1973;32:132–4
 - 44 Chu FF, Doroshow JH, Esworthy RS Expression, characterization, and tissue distribution of a new cellular selenium-dependent glutathione peroxidase, GSHPx-GI *J Biol Chem* 1993;268:2571–6
 - 45 Takahashi K, Avissar N, Whitin J, Cohen H Purification and characterization of human plasma glutathione peroxidase: a selenoglycoprotein distinct from the known cellular enzyme *Arch Biochem Biophys* 1987;256:677–86
 - 46 Deagen JT, Butler JA, Zachara BA, Whanger PD Determination of the distribution of selenium between glutathione peroxidase, selenoprotein P, and albumin in plasma *Anal Biochem* 1993;208:176–81
 - 47 Chu FF, Esworthy RS, Doroshow JH, Doan K, Liu XF Expression of plasma glutathione peroxidase in human liver in addition to kidney, heart, lung, and breast in humans and rodents *Blood* 1992;79:3233–8
 - 48 Yamamoto Y, Takahashi K Glutathione peroxidase isolated from plasma reduces phospholipid hydroperoxides *Arch Biochem Biophys* 1993;305:541–5
 - 49 Roveri A, Casasco A, Maiorino M, Dalan P, Calligaro A, Ursini F Phospholipid hydroperoxide glutathione peroxidase of rat testis Gonadotropin dependence and immunocytochemical identification *J Biol Chem* 1992;267:6142–6
 - 50 Roveri A, Maiorino M, Nisii C, Ursini F Purification and characterization of phospholipid hydroperoxide glutathione peroxidase from rat testis mitochondrial membranes *Biochim Biophys Acta* 1994;1208:211–21
 - 51 Ursini F, Maiorino M, Valente M, Ferri L, Gregolin C Purification from pig liver of a protein which protects liposomes and biomembranes from peroxidative degradation and exhibits glutathione peroxidase activity on phosphatidylcholine hydroperoxides *Biochim Biophys Acta* 1982;710:197–211
 - 52 Maiorino M, Gregolin C, Ursini F Phospholipid hydroperoxide glutathione peroxidase *Methods Enzymol* 1990;186:448–57
 - 53 Bao Y, Jemth P, Mannervik B, Williamson G Reduction of thymine hydroperoxide by phospholipid hydroperoxide glutathione peroxidase and glutathione transferases *FEBS Lett* 1997;410:210–2
 - 54 Schnurr K, Belkner J, Ursini F, Schewe T, Kuhn H The selenoenzyme phospholipid hydroperoxide glutathione peroxidase controls the activity of the 15-lipoxygenase with complex substrates and preserves the specificity of the oxygenation products *J Biol Chem* 1996;271:4653–8
 - 55 Imai H, Narashima K, Arai M, Sakamoto H, Chiba N, Nakagawa Y Suppression of leukotriene formation in RBL-2H3 cells that overexpressed phospholipid hydroperoxide glutathione peroxidase *J Biol Chem* 1998;273:1990–7
 - 56 Nam SY, Fujisawa M, Kim JS, Kurohmaru M, Hayashi Y Expression pattern of phospholipid hydroperoxide glutathione peroxidase messenger ribonucleic acid in mouse testis *Biol Reprod* 1998;58:1272–6
 - 57 Maiorino M, Wissing JB, Brigelius-Flohe R, et al Testosterone mediates expression of the selenoprotein PHGPx by induction of spermatogenesis and not by direct transcriptional gene activation *FASEB J* 1998;12:1359–70
 - 58 Ursini F, Heim S, Kiess M, et al Dual function of the selenoprotein PHGPx during sperm maturation *Science* 1999;285:1393–6
 - 59 Foresta C, Flohe L, Garolla A, Roveri A, Ursini F, Maiorino M Male fertility is linked to the selenoprotein phospholipid hydroperoxide glutathione peroxidase *Biol Reprod* 2002;67:967–71
 - 60 Tamura T, Stadtman TC A new selenoprotein from human lung adenocarcinoma cells: purification, properties, and thioredoxin reductase activity *Proc Natl Acad Sci USA* 1996;93:1006–11
 - 61 Sun QA, Wu Y, Zappacosta F, et al Redox regulation of cell signaling by selenocysteine in mammalian thioredoxin reductases *J Biol Chem* 1999;274:24522–30
 - 62 Arthur JR, Nicol F, Hutchinson AR, Beckett GJ The effects of selenium depletion and repletion on the metabolism of thyroid hormones in the rat *J Inorg Biochem* 1990;39:101–8
 - 63 Berry MJ, Banu L, Larsen PR Type I iodothyronine deiodinase is a selenocysteine-containing enzyme *Nature* 1991;349:438–40
 - 64 Berry MJ, Larsen PR The role of selenium in thyroid hormone action *Endocr Rev* 1992;13:207–19
 - 65 Arthur JR, Nicol F, Beckett GJ Selenium deficiency, thyroid hormone metabolism, and thyroid hormone deiodinases *Am J Clin Nutr* 1993;57:236S–9S
 - 66 Davey JC, Becker KB, Schneider MJ, St Germain DL, Galton VA Cloning of a cDNA for the type II iodothyronine deiodinase *J Biol Chem* 1995;270:26786–9



- 67 Arthur JR, Beckett GJ New metabolic roles for selenium Proc Nutr Soc 1994;53:615-24
- 68 Arthur JR, Bermanno G, Mitchell JH, Hesketh JE Regulation of selenoprotein gene expression and thyroid hormone metabolism Biochem Soc Trans 1996;24:384-8
- 69 Burk RF, Hill KE Selenoprotein P: a selenium-rich extracellular glycoprotein J Nutr 1994;124:1891-7
- 70 Saito Y, Takahashi K Characterization of selenoprotein P as a selenium supply protein Eur J Biochem 2002;269:5746-51
- 71 Harrison I, Littlejohn D, Fell GS Distribution of selenium in human blood plasma and serum Analyst 1996;121:189-94
- 72 Burk RF, Hill KE Selenoprotein P-Expression, functions, and roles in mammals Biochim Biophys Acta (Epub ahead of print 1 April 2009)
- 73 Saito Y, Hayashi T, Tanaka A, et al Selenoprotein P in human plasma as an extracellular phospholipid hydroperoxide glutathione peroxidase Isolation and enzymatic characterization of human selenoprotein P J Biol Chem 1999;274:2866-71
- 74 Vendeland SC, Beilstein MA, Chen CL, Jensen ON, Barofsky E, Whanger PD Purification and properties of selenoprotein W from rat muscle J Biol Chem 1993;268:17103-7
- 75 Whanger PD Selenoprotein W: a review Cell Mol Life Sci 2000;57:1846-52
- 76 Lescure A, Deniziak M, Rederstorff M, Krol A Molecular basis for the role of selenium in muscle development and function Chem Biodivers 2008;5:408-13
- 77 Petit N, Lescure A, Rederstorff M, et al Selenoprotein N: an endoplasmic reticulum glycoprotein with an early developmental expression pattern Hum Mol Genet 2003;12:1045-53
- 78 Curran JE, Jowett JB, Elliott KS, et al Genetic variation in selenoprotein S influences inflammatory response Nat Genet 2005;37:1234-41
- 79 Gao Y, Feng HC, Walder K, et al Regulation of the selenoprotein SeS by glucose deprivation and endoplasmic reticulum stress - SeS is a novel glucose-regulated protein FEBS Lett 2004;563:185-90
- 80 Lu C, Qiu F, Zhou H, et al Identification and characterization of selenoprotein K: an antioxidant in cardiomyocytes FEBS Lett 2006;580:5189-97
- 81 Kryukov GV, Kumar RA, Koc A, Sun Z, Gladyshev VN Selenoprotein R is a zinc-containing stereo-specific methionine sulfoxide reductase Proc Natl Acad Sci USA 2002;99:4245-50
- 82 Lee BC, Dikiy A, Kim HY, Gladyshev VN Functions and evolution of selenoprotein methionine sulfoxide reductases Biochim Biophys Acta 2009;1790:1471-7
- 83 Panee J, Stoytcheva ZR, Liu W, Berry MJ Selenoprotein H is a redox-sensing high mobility group family DNA-binding protein that up-regulates genes involved in glutathione synthesis and phase II detoxification J Biol Chem 2007;282:23759-65
- 84 Hwang DY, Sin JS, Kim MS, et al Overexpression of human selenoprotein M differentially regulates the concentrations of antioxidants and H₂O₂, the activity of antioxidant enzymes, and the composition of white blood cells in a transgenic rat Int J Mol Med 2008;21:169-79
- 85 Gladyshev VN, Jeang KT, Wootton JC, Hatfield DL A new human selenium-containing protein Purification, characterization, and cDNA sequence J Biol Chem 1998;273:8910-5
- 86 Labunskyy VM, Yoo MH, Hatfield DL, Gladyshev VN Sep15, a thioredoxin-like selenoprotein, is involved in the unfolded protein response and differentially regulated by adaptive and acute ER stresses Biochemistry 2009;48:8458-65
- 87 Xu XM, Carlson BA, Irons R, et al Selenophosphate synthetase 2 is essential for selenoprotein biosynthesis Biochem J 2007;404:115-20
- 88 Persson-Moschos M, Alfthan G, Akesson B Plasma selenoprotein P levels of healthy males in different selenium status after oral supplementation with different forms of selenium Eur J Clin Nutr 1998;52:363-7
- 89 Moschos MP Selenoprotein P Cell Mol Life Sci 2000;57:1836-45
- 90 Hill KE, Xia Y, Akesson B, Boelin ME, Burk RF Selenoprotein-P concentration in plasma is an index of selenium status in selenium-deficient and selenium-supplemented Chinese subjects J Nutr 1996;126:138-45
- 91 Duffield AJ, Thomson CD, Hill KE, Williams S An estimation of selenium requirements for New Zealanders Am J Clin Nutr 1999;70:896-903
- 92 Xia Y, Hill KE, Byrne DW, Xu J, Burk RF Effectiveness of selenium supplements in a low-selenium area of China Am J Clin Nutr 2005;81:829-34
- 93 Ashton K, Hooper L, Harvey LJ, Hurst R, Casgrain A, Fairweather-Tait SJ Methods of assessment of selenium status in humans: a systematic review Am J Clin Nutr 2009;89:2025S-39S
- 94 Combs GF Jr, Midthune DN, Patterson KY, et al Effects of selenomethionine supplementation on selenium status and thyroid hormone concentrations in healthy adults Am J Clin Nutr 2009;89:1808-14
- 95 Alfthan G, Aro A, Arvilommi H, Huttunen JK Selenium metabolism and platelet glutathione peroxidase activity in healthy Finnish men: effects of selenium yeast, selenite, and selenate Am J Clin Nutr 1991;53:120-5
- 96 Thomson CD, Robinson MF, Campbell DR, Rea HM Effect of prolonged supplementation with daily supplements of selenomethionine and sodium selenite on glutathione peroxidase activity in blood of New Zealand residents Am J Clin Nutr 1982;36:24-31
- 97 Neve J, Vertongen F, Capel P Selenium supplementation in healthy Belgian adults: response in platelet glutathione peroxidase activity and other blood indices Am J Clin Nutr 1988;48:139-43
- 98 Thomson CD, Robinson MF, Butler JA, Whanger PD Long-term supplementation with selenate and selenomethionine: selenium and glutathione peroxidase (EC 1.11.1.9) in blood components of New Zealand women Br J Nutr 1993;69:577-88
- 99 Levander OA, Alfthan G, Arvilommi H, et al Bioavailability of selenium to Finnish men as assessed by platelet glutathione peroxidase activity and other blood parameters Am J Clin Nutr 1983;37:887-97
- 100 Méplan C, Crosley LK, Nicol F, et al Genetic polymorphisms in the human selenoprotein P gene determine the response of selenoprotein markers to selenium supplementation in a gender-specific manner (the SELGEN study) FASEB J 2007;21:3063-74
- 101 Méplan C, Crosley LK, Nicol F, et al Functional effects of a common single-nucleotide polymorphism (GPX4c718t) in the glutathione peroxidase 4 gene: interaction with sex Am J Clin Nutr 2008;87:1019-27
- 102 Lei C, Niu X, Wei J, Zhu J, Zhu Y Interaction of glutathione peroxidase-1 and selenium in endemic dilated cardiomyopathy Clin Chim Acta 2009;399:102-8
- 103 Jablonska E, Gromadzinska J, Reszka E, et al Association between GPx1 Pro198Leu polymorphism, GPx1 activity and plasma selenium concentration in humans Eur J Nutr (Epub ahead of print 5 May 2009)
- 104 Seiderer J, Dambacher J, Kühnlein B, et al The role of the selenoprotein S (SELS) gene -105G>A promoter polymorphism in inflammatory bowel disease and regulation of SELS gene expression in intestinal inflammation Tissue Antigens 2007;70:238-46
- 105 Shibata T, Arisawa T, Tahara T, et al Selenoprotein S (SEPS1) gene -105G>A promoter polymorphism influences the susceptibility to gastric cancer in the Japanese population BMC Gastroenterol 2009;9:2
- 106 Bermanno G, Pagmantidis V, Holloway N, et al Evidence that a polymorphism within the 3' UTR of glutathione peroxidase 4 is functional and is associated with susceptibility to colorectal cancer Genes Nutr 2007;2:225-32
- 107 van Ommen B, Fairweather-Tait S, Freidig A, Kardinaal A, Scalbert A, Wopereis S A network biology model of micronutrient related health Br J Nutr 2008;99(suppl 3):S72-80



Erratum

Fairweather Tait SJ, Collings R, Hurst R. Selenium bioavailability: current knowledge and future research. *Am J Clin Nutr* 2010;91(suppl):1484S-91S.

In the print version of this article, the term “selenium methylselenocysteine” appears in error. The correct term, as provided by the authors, is “Se methylselenocysteine.” This error occurs in Table 1 on page 1486S, in the legend to Figure 1 on page 1487S, and in text on page 1485S. The correct term appears in the online version.

doi: 10.3945/ajcn.2010.30169.

Erratum

Morris MS, Jacques PF, Rosenberg IH, Selhub J. Circulating unmetabolized folic acid and 5-methyltetrahydrofolate in relation to anemia, macrocytosis, and cognitive test performance in American seniors. *Am J Clin Nutr* 2010;91:1733-44.

On page 1735, the last sentence of the first paragraph of the section entitled “Classification of subjects according to vitamin B12 status and folate fractions” contains a detection limit. The published detection limit for the assay is 0.18 nmol/L, not 0.027 nmol/L. The incorrect limit was also given in the footnotes to Table 1 and Table 4. There were 7 eligible subjects with measured serum folic acid concentrations between 0 and 0.18 nmol/L who were included in the group with detectable unmetabolized folic acid. In addition, in Table 4, the multivariate model was not specified for the row labeled “Mean cell volume.” The results in that row were controlled for sex, age, race/ethnicity, current smoking, current alcohol intake, BMI, self-reported cancer history, and serum concentrations of ferritin, cystatin C, and C-reactive protein.

doi: 10.3945/ajcn.2010.30170.

Erratum

Yatabe MS, Yatabe J, Yoneda M, et al. Salt sensitivity is associated with insulin resistance, sympathetic overactivity, and decreased suppression of circulating renin activity in lean patients with essential hypertension. *Am J Clin Nutr* 2010;92:77-82.

In the print version of this article, 2 lines appear for each group represented in Figure 7; however, only one line should appear for each group. The correct version of Figure 7 appears below. The online version of this figure is correct.




FIGURE 7. Relation between glucose infusion rate and change in urinary norepinephrine (Δu -NE) from the low- to high-salt diet in salt-resistant (SR) subjects ($n = 14$, open circles) and salt-sensitive (SS) subjects ($n = 10$, filled circles) with essential hypertension. Statistical significance was estimated with Pearson's correlation coefficient.

doi: 10.3945/ajcn.2010.30173.

Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997–2001)

Andrea J. Fascetti, VMD, PhD, DACVN, DACVIM; John R. Reed, DVM, MS, DACVIM; Quinton R. Rogers, PhD, DACVN; Robert C. Backus, DVM, PhD

Objective—To determine signalment, history, clinical signs, blood and plasma taurine concentrations, electrocardiographic and echocardiographic findings, treatment, and outcome of dogs with low blood or plasma taurine concentrations and dilated cardiomyopathy (DCM).

Design—Retrospective study.

Animals—12 client-owned dogs with low blood or plasma taurine concentrations and DCM.

Procedure—Medical records were reviewed, and clinical data were obtained.

Results—All 12 dogs were being fed a commercial dry diet containing lamb meal, rice, or both as primary ingredients. Cardiac function and plasma taurine concentration improved with treatment and taurine supplementation. Seven of the 12 dogs that were still alive at the time of the study were receiving no cardiac medications except taurine.

Conclusions and Clinical Relevance—Results suggest that consumption of certain commercial diets may be associated with low blood or plasma taurine concentrations and DCM in dogs. Taurine supplementation may result in prolonged survival times in these dogs, which is not typical for dogs with DCM. Samples should be submitted for measurement of blood and plasma taurine concentrations in dogs with DCM, and taurine supplementation is recommended while results of these analyses are pending. (*J Am Vet Med Assoc* 2003;223:1137–1141)

Large-breed dogs, especially males, are predisposed to developing dilated cardiomyopathy (DCM).¹ Because the long-term prognosis for dogs with this disease is poor, methods for preventing the disease would be beneficial. However, in most affected dogs, the underlying cause is unknown.

In 1987, Pion et al² reported an association between low plasma taurine concentrations and DCM in cats. Oral supplementation of affected cats with taurine sig-

From the Departments of Molecular Biosciences (Fascetti, Rogers, Backus) and Surgery and Radiological Sciences (Reed), School of Veterinary Medicine, University of California, Davis, CA 95616-8741; and the Highlands Veterinary Hospital, 3451 Elkhorn Blvd, North Highlands, CA 95660 (Reed).

Supported in part by a grant from the Center of Companion Animal Health, School of Veterinary Medicine, University of California, Davis. Presented in part at the 18th Annual Veterinary Medical Forum of the American College of Veterinary Internal Medicine, Seattle, May 2000.

The authors thank Drs. Sean Delaney, Melanie Morgan, and Lorie Siemens for their assistance.

Address correspondence to Dr. Fascetti.

nificantly improved clinical signs, restored myocardial function, and improved survival times.³ Since then, the addition of taurine to commercial diets for cats has resulted in a marked decrease in the number of cats developing this disease.

Traditionally, dogs have not been recognized as having a dietary need for taurine, because they are able to synthesize taurine from the dietary sulfur amino acids methionine and cysteine.⁴ Recently, however, a cardiologist in private practice (JRR) brought to the attention of the authors 4 unrelated, large-breed dogs with DCM. At the time of initial examination, all 4 dogs were found to have low blood taurine concentrations. One common factor among the dogs was consumption of the same lamb meal and rice commercial dry diet. Later, a Border Collie with DCM and low blood taurine concentrations was brought to our attention by a second local cardiologist in private practice. This dog was also consuming a lamb meal and rice diet, but one produced by another manufacturer. The common diet history for these 5 dogs suggested that diet may have had a role in the development of low blood taurine concentrations and DCM in these dogs. The purpose of the study reported here was to determine the signalment, history (including diet history), clinical signs, blood and plasma taurine concentrations, electrocardiographic and echocardiographic findings, treatment, and outcome of dogs with low blood or plasma taurine concentrations and DCM. In addition, we wanted to determine whether diet may have had any role in the development of DCM.

Criteria for Selection of Cases

The cardiology database at the Veterinary Medical Teaching Hospital of the University of California, Davis, was searched for dogs examined between October 1997 and August 2001 in which a diagnosis of DCM had been made. Dogs were included in the study only if DCM had been diagnosed by a veterinary cardiologist; the diagnosis had been confirmed by means of echocardiography; samples had been submitted to the Amino Acid Laboratory at the University of California, Davis, and blood or plasma taurine concentration had been found to be low; and a complete diet history was available. In addition, the 5 dogs brought to the authors' attention by local cardiologists were included in the study.

Procedures

Information collected for all dogs included signalment, history, diet history, initial clinical signs, electrocardiographic and echocardiographic findings, blood and

plasma taurine concentrations, and treatment. Follow-up information was obtained by reviewing medical records or through telephone conversations with the attending cardiologist or primary care veterinarian.

Measurement of blood and plasma taurine concentrations—In all dogs, blood and plasma taurine concentrations were determined by the Amino Acid Laboratory at the University of California, Davis. At least 1 mL of heparinized blood, plasma, or both was submitted for analysis. Blood and plasma taurine concentrations were determined as described⁷ with an automated amino acid analyzer.⁸ Plasma taurine concentrations < 40 nmol/mL and blood taurine concentrations < 150 nmol/mL were considered indicative of a taurine deficiency.

Statistical analyses—Descriptive statistics (mean, SD, median, and range) were calculated for taurine concentrations and echocardiographic findings. Paired *t* tests were used to compare taurine concentrations and echocardiographic findings obtained at the time of diagnosis of DCM with values obtained after treatment. All analyses were performed with commercial software^b; values of *P* ≤ 0.05 were considered significant.

Results

From October 1997 through August 2001, 64 dogs with DCM were evaluated at the University of California veterinary teaching hospital. Blood samples from 24 of these dogs were submitted for analysis of taurine concentration, and 14 of the dogs were classified as having a taurine deficiency on the basis of a low blood or plasma taurine concentration. Six of the 14 dogs were excluded from the study, because they were American Cocker Spaniels, a breed well documented to have taurine- and carnitine-responsive DCM.⁶ A seventh dog was excluded because of an inadequate diet history. Thus, 12 dogs were included in the study, including the 5 dogs brought to the authors' attention by local cardiologists and the 7 dogs in which DCM was diagnosed at the veterinary teaching hospital during the study period.

The 12 dogs included in the study consisted of 8 males (4 sexually intact) and 4 females (2 sexually intact). There were 3 English Setters, an Alaskan Malamute, a Border Collie, a German Shepherd Dog, a Golden Retriever, a Gordon Setter, a Great Pyrenees, a Labrador Retriever, a Newfoundland, and a Scottish Terrier. Dogs ranged from 4.5 to 11 years old at the time DCM was diagnosed (mean and median, 8.3 years).

Lethargy and anorexia were the 2 most common clinical abnormalities (7 dogs each). Other clinical signs included cough (*n* = 5), dyspnea (5), weight loss (3), trembling (1), and collapse (1). Three of the dogs were clinically normal at the time of initial examination. In 2 of these dogs, an arrhythmia was detected during a routine examination. The third dog was evaluated at the request of the owner, because 2 of her other dogs had recently been found to have low taurine concentrations and DCM.

Ten of the 12 dogs had ECG abnormalities. Left ventricular enlargement (LVE) was the most common abnormality (*n* = 7), followed by left atrial enlargement

(6), atrial fibrillation (3), left bundle branch block (2), and ventricular premature contractions (1). All of the dogs underwent echocardiography. Mean ± SD E-point to septal separation at the time of initial examination was 14.5 ± 5.6 mm (*n* = 10; median, 14.5 mm; range, 7.5 to 26 mm). Mean fractional shortening at the time of initial examination was 15.7 ± 7.6% (*n* = 12; median, 15.9%; range, 7.0 to 28.9%). Mean plasma taurine concentration at the time DCM was diagnosed was 16 ± 20 nmol/mL (*n* = 12; median, 7 nmol/mL; range, 2 to 64 nmol/mL). Blood taurine concentrations had also been measured in 8 dogs. Mean blood taurine concentration at the time DCM was diagnosed was 121 ± 76 nmol/mL (median, 135 nmol/mL; range, 8 to 229 nmol/mL).

All 12 dogs were consuming a commercial dry diet with lamb meal, rice, or both as primary ingredients. Eight of the 12 dogs were consuming the same commercial lamb meal and rice dry diet^c (diet A). The remaining 4 dogs were each consuming 1 of 4 other commercial diets^{d–g} (diets B through E). Nutrient composition of the 5 diets was determined from the manufacturers' reported data (Table 1).

All 12 dogs were treated with taurine (1,000 to 3,000 mg, PO, q 24 h) beginning at the time DCM was diagnosed or when the low blood or plasma taurine concentration was discovered. In addition, 8 dogs were treated with inotropic agents, 7 were treated with diuretics, 7 were treated with angiotensin-converting enzyme inhibitors, and 1 was treated with a calcium channel blocker to manage cardiac abnormalities. Eleven of the dogs were changed to a different commercial dry diet at the time DCM was diagnosed or when the low blood or plasma taurine concentration was discovered.

Dogs were reevaluated between 1 and 12 months after the time of initial diagnosis (mean ± SD, 4.3 ± 3.5 months; median, 3 months). Electrocardiography was repeated on 4 dogs. Results were normal for 1 of the 4 dogs; abnormalities identified in the other 3 included left ventricular enlargement (*n* = 1), atrial fibrillation (2), and ventricular premature contractions (1). Echocardiography was repeated on 9 dogs. Mean E-point to septal separation was 8.9 ± 4.1 mm (*n* = 8; median, 8.3 mm; range, 2.9 to 17 mm). This was significantly (*P* = 0.002) lower than the mean value at the time of initial examination of these 8 dogs. Mean fractional shortening was 22.5 ± 6.3% (*n* = 9; median, 20%;

Table 1—Proximate dry matter composition of diets fed to dogs with dilated cardiomyopathy and low blood or plasma taurine concentrations

Variable	Diet				
	A	B	C	D	E
Crude protein (%)	24	23	17	22	28
Crude fat (%)	14	13	7	11	13
NFE* (%)	48	57	63	62	46
Crude fiber (%)	2	1	7	5	4
Ash (%)	10	6	6	NR	9
ME (kcal/g)	3.8	3.9	3.4	3.5	3.8

All values were obtained from manufacturers' reported data.

*Calculated or reported nitrogen-free extract.

NR = Not reported. ME = Metabolizable energy.

range, 16.6 to 36%). This was significantly ($P = 0.002$) greater than the mean value at the time of initial examination of these 9 dogs. Blood and plasma taurine concentrations increased in the 3 dogs in which analyses were repeated. Mean plasma taurine concentration was 226 ± 54 nmol/mL ($n = 3$; median, 208 nmol/mL; range, 183 to 286 nmol/mL), and mean blood taurine concentration was 455 ± 88 nmol/mL ($n = 3$; median, 443 nmol/mL; range, 373 to 548 nmol/mL). Plasma taurine concentrations were significantly ($P = 0.02$) increased in these 3 dogs, compared with concentrations prior to institution of taurine supplementation. Blood taurine concentrations were not significantly increased, but the P value was close to the cutoff for significance ($P = 0.06$).

Seven of the 12 dogs were alive at the time of the study. Mean \pm SD survival time for all 12 dogs was 585 ± 499 days (median, 456 days; range, 1 to 1,460 days). One dog died within 1 day, 3 dogs died within 180 days, and 1 dog died within 365 days of initial examination. Survival times for the 7 dogs still alive at the time of the study ranged from 365 to 1,460 days (913 ± 380 days; median, 1,095 days). None of the 7 surviving dogs were receiving any cardiac medications at the time of the study other than taurine.

Discussion

With the exception of the single Newfoundland, none of the dogs in this study were of breeds predisposed to developing DCM. Breeds recognized to have a high prevalence of DCM include the Scottish Deerhound, Doberman Pinscher, Irish Wolfhound, Great Dane, Boxer, Saint Bernard, Afghan Hound, Newfoundland, and Old English Sheepdog.⁷ The dogs in the present study also generally had longer survival times than are typically reported in the literature for dogs with DCM. For instance, 2 recent multibreed retrospective studies^{8,9} reported overall probabilities of survival 1 year after diagnosis of DCM of 17.5 and 37.5% and median survival times of 27 and 65 days. In contrast, the median survival time for the 12 dogs in the present study was 456 days, and many of these dogs were still alive at the time of the study. Finally, several of the dogs in the present study regained substantial cardiac function and were weaned off all medications except taurine. This is unusual for most dogs with DCM, in which the disease is typically progressive and fatal. Information on follow-up cardiac evaluations and measurements of taurine concentrations was not available for all dogs in the present study. However, in those dogs in which this information was available, cardiac function improved and taurine concentrations increased to concentrations greater than those considered evidence of a deficiency. Furthermore, in all of the dogs that lived > 1 year, all cardiac medications other than taurine were discontinued. These characteristics suggest that the dogs in the present study did not have the typical form of DCM.

Consumption of commercial dry diets containing lamb meal or rice as a primary ingredient was a common finding among dogs in the present study. Three of the 5 diets that these dogs consumed contained both lamb meal and rice (diets A, C, and E); 1 contained

chicken meal and rice (diet B); and 1 contained ground whole wheat, lamb meal, and rice (diet D). Most of the dogs were changed to another diet at the time DCM was diagnosed. However, the owner of the Border Collie decided to keep the dog on the same commercial lamb meal and rice diet (diet D) but supplemented the dog with taurine. Increases in taurine concentration and cardiac function in this dog indicated a response to taurine supplementation and supported the hypothesis that diet played a contributing role in the development of DCM by causing taurine depletion.

Taurine is the most abundant of the free amino acids in tissues. High concentrations of taurine are found in animal tissues, especially muscle, viscera, and brain. In cats, 3 manifestations of taurine deficiency have been identified: central retinal degeneration, reproductive failure and impaired fetal development, and DCM.¹⁰ Taurine deficiency can also cause hearing loss, platelet hyperaggregation, and impaired immune function, although specific clinical disorders have not been recognized.¹⁰ The mechanism of heart failure in taurine-deficient cats and dogs is poorly understood, but in the myocardium, taurine appears to participate in many functions, including cellular osmoregulation, free-radical scavenging, and modulation of contraction strength through regulation of calcium concentration.³

In cats, inadequate provision of dietary taurine results in DCM.³ Cats have a limited ability to synthesize taurine because of low concentrations of the enzymes cysteine sulfonic acid decarboxylase and cysteine dioxygenase.³ In contrast, dogs have not been generally recognized to have a need for dietary taurine, because they have the metabolic capacity to synthesize taurine from cysteine and methionine.⁴ The concentration of taurine necessary to prevent clinical signs of a taurine deficiency in cats varies with diet composition and processing, but ranges from 400 mg of taurine/kg of diet to $> 2,000$ mg of taurine/kg of diet.^{11,12}

Possible causes of the taurine deficiency in the dogs in the present report include insufficient synthesis of taurine, extraordinary loss of taurine or its precursors in urine, extraordinary gastrointestinal tract loss of taurine in bile acid conjugates as found in cats, and a reduction in protein digestibility.¹³ On the basis of our clinical findings, in conjunction with the blood and plasma taurine concentrations and common diet histories in these dogs, we hypothesize that the consumption of diets with inadequate or unavailable sulfur amino acid precursors of taurine resulted in taurine deficiency and low blood taurine concentrations that, in turn, led to the development of abnormal cardiac function and DCM.

Recent experimental and clinical observations in dogs are supportive of the possibility that insufficient synthesis of taurine from sulfur amino acid precursors results in the development of DCM. Sanderson et al¹⁴ found low plasma taurine concentrations in Beagles fed an energy-dense, protein-restricted (10% protein on a dry-matter basis) diet for 48 months. One dog developed DCM, and taurine supplementation (500 mg, PO, q 12 h) reversed the cardiac changes in this dog. Prolonged feeding of a commercial prescription diet with a protein-to-calorie ratio similar to one used by

Sanderson et al¹⁴ may have induced development of DCM in Dalmatians,¹⁵ but whether taurine deficiency caused DCM in these dogs is unclear. Blood and plasma taurine concentrations in the Dalmatians were within reference limits at the time of clinical evaluation. However, taurine concentrations in blood and tissues at the time DCM developed were not known. It is possible that a period of taurine deficiency could produce myocardial damage in dogs that cannot be reversed. Following a change in diet, blood taurine concentrations may indicate normal body taurine stores and not reveal that a period of deficiency had occurred.

Alternatively, the rice bran and whole rice products in the commercial diets consumed by most of the dogs in the present study may have been a factor in the development of low blood and plasma taurine concentrations. Rice bran and whole-rice products are sources of moderately soluble fiber and contain relatively high amounts of fat. The fiber, fat, or protein content of the rice bran may accelerate excretion of bile acids, predisposing dogs to taurine deficiency. Stratton-Phelps et al¹⁶ recently demonstrated that cats fed a purified diet with 26% (dry-matter basis) full-fat rice bran had critically low plasma and blood taurine concentrations after 6 and 12 weeks, respectively. Extraordinary intestinal loss of taurine secondary to increases in bacterial populations appears to be contributing to taurine deficiency in these cats.¹⁶ Preliminary results from a dose-response study conducted by the authors of the present study indicate that cats can develop critically low blood taurine concentrations when consuming full-fat rice bran at concentrations as low as 4% (dry-matter basis).

Dogs in the present study were all fed a commercial dry diet containing high quantities of lamb meal or rice products, but why this should be associated with taurine deficiency was not readily apparent, as the diets appeared sufficient in protein and sulfur amino acid contents and had passed testing in accordance with the Association of American Feed Control Officials' feeding trials for all life stages. However, because lamb meals may be of low quality, limited bioavailability of sulfur amino acids required for taurine synthesis in the diet was suspected. Relative to other meat meal sources, lamb meal has been shown to have poor ileal nitrogen and cystine digestibility in dogs.¹⁷ Results of a study involving feeding a lamb meal and rice diet to young Beagles for 8 months indicate that plasma taurine concentrations decreased substantially during the first month after dogs were switched to the diet but did not change thereafter.¹⁸ Thus, the lamb meal and rice diet appeared to have an effect on taurine status, but not to the point of a depletion sufficient to cause DCM.

Recently, low blood taurine concentrations have been identified in Newfoundland¹⁹. Reduced reproductive performance, small litter sizes, poor litter growth, and small puppies were reported, and similar findings have been reported for cats with taurine deficiency.²⁰ Diet appeared to be the cause of the taurine deficiency in these dogs, in that 7 of the 12 dogs with plasma taurine concentrations < 40 nmol/mL were

consuming diets containing lamb meal and rice. Methionine supplementation in dogs consuming lamb meal and rice diets resulted in substantial improvement in taurine concentrations, and plasma taurine concentrations increased when a dietary change was instituted but were unchanged when the diet was not changed.

The authors have also examined 2 dogs with taurine deficiency that were being fed a home-prepared, low-protein, tofu-based diet that meet the National Research Council's requirements for adult maintenance.²¹ Taurine deficiency in these dogs was attributed to inadequate synthesis, and it was assumed that the low concentrations of protein in general and of sulfur amino acids in particular provided an inadequate supply of precursors for taurine synthesis. An additional contributing factor may have been an increase in taurine loss, as soybean protein augments bile acid loss through microbial degradation and accelerates cholecystokinin-mediated bile acid turnover.^{5,22} Similarly, we are aware of 3 Golden Retrievers with taurine deficiency that lived in the same household and were consuming a vegetarian diet formulated by the owner.

Taken together, these findings suggest that taurine deficiency may result in DCM in dogs other than American Cocker Spaniels fed a diet that contains insufficient amino acid precursors for adequate taurine synthesis or that accelerates taurine loss. We recommend that a complete diet history be obtained for every dog each time it is examined by a veterinarian, including the name and amount of food fed, the name and amount of any snacks and treats, a description of the manner in which the dog is fed, whether the dog has access to other food sources, and whether any dietary supplements are given. We also recommend that all home-cooked diets be evaluated by a veterinary nutritionist.

Blood and plasma taurine concentrations should be measured in all dogs with DCM, just as measurement of blood and plasma taurine concentrations is recommended for all cats with DCM.²³ Although blood taurine concentration is only a fraction of the concentration in the tissues, blood and plasma taurine concentrations do change in proportion with tissue concentrations.^{4,24} Blood taurine concentrations may be used to substantiate a diagnosis of taurine deficiency when plasma concentrations are equivocal. In addition, blood taurine concentrations are only slightly altered after eating, whereas plasma taurine concentration may change substantially in taurine-depleted animals.²³ A substantial increase in plasma or serum taurine concentration can occur secondary to taurine leakage from granulocytes and platelets, as occurs with clotting or hemolysis, but analysis of blood taurine concentration is not confounded by these effects. Serum taurine concentrations are of questionable clinical value because of the variations in time of clotting and the method of serum separation, and in our experience, the variability in serum taurine concentrations is greater than the variability in plasma taurine concentrations.

Finally, we recommend that taurine be administered to all dogs with DCM, pending results of taurine

analysis. Taurine is inexpensive and readily available and has no reported adverse effects when administered orally. Follow-up measurement of blood and plasma taurine concentrations should be performed after 1 to 2 months of taurine supplementation to confirm that taurine status has improved and verify owner compliance with regard to administration.

One manufacturer produced 3 of the diets (diets A, B, and C) associated with taurine deficiency in the present study. Since identification of these dogs, the manufacturer has added taurine to 1 of the 3 diets (diet A). Diets D and E, each produced by other manufacturers at the time the dogs consuming them developed low taurine concentrations and DCM, have recently been acquired by other companies. The formulation of diet E has been changed, but neither diet D nor diet E includes additional taurine. Nevertheless, we suggest that veterinarians not focus on particular diets but on the issue of taurine deficiency as a whole. Further research is needed to identify dietary factors inducing taurine deficiency and determine the mechanisms of their effects.

^aSystem 7300 and Model 121-M automated amino acid analyzers, Beckman Instruments Inc, Palo Alto, Calif.

^bSYSTAT, version 9.0, SPSS Inc, Chicago, Ill.

^cNutro Natural Choice lamb meal and rice formula dog food, Nutro Products Inc, City of Industry, Calif.

^dNutro Natural Choice senior dog food, Nutro Products Inc, City of Industry, Calif.

^eNutro Natural Choice lite rice and lamb meal formula for overweight dogs, Nutro Products Inc, City of Industry, Calif.

^fNature's Recipe adult maintenance lamb meal and rice, Nature's Recipe Pet Foods, Heinz Pet Products, Pittsburgh, Pa.

^gSensible Choice adult lamb meal and rice formula, Pet Products Plus Inc, St Charles, Mo.

^hStratton-Phelps M, Rogers QR, Backus RC, et al. Oral antibiotics do not replete taurine in cats consuming 26% dietary rice bran (abstr), in *Proceedings. Fed Am Soc Exp Biol* 2002;5055.

References

- Buchanan JW. Causes and prevalence of cardiovascular disease. In: Kirk RW, Bonagura JD, eds. *Current veterinary therapy XI*. Philadelphia: WB Saunders Co, 1992;647-655.
- Pion PD, Kittleson MD, Rogers QR, et al. Myocardial failure in cats is associated with low plasma taurine: a reversible cardiomyopathy. *Science* 1987;237:764-768.
- Pion PD, Sanderson SL, Kittleson MD. The effectiveness of taurine and levocarnitine in dogs with heart disease. *Vet Clin North Am Small Anim Pract* 1998;28:1495-1514.
- Hayes KC. Taurine nutrition. *Nutr Res Rev* 1988;1:99-113.
- Kim SW, Morris JG, Rogers QR. Dietary soybean protein decreases plasma taurine in cats. *J Nutr* 1995;125:2831-2837.
- Kittleson MD, Keene B, Pion PD, et al. Results of the Multicenter Spaniel Trial (MUST): taurine- and carnitine-responsive dilated cardiomyopathy in American Cocker Spaniels with decreased plasma taurine concentration. *J Vet Intern Med* 1997;11:204-211.
- Sisson DD, Thomas WP, Keene BW. Primary myocardial disease in the dog. In: Ettinger SJ, Feldman EC, eds. *Textbook of veterinary internal medicine*. 5th ed. Philadelphia: WB Saunders Co, 2000; 874-895.
- Tidholm A, Svensson H, Sylven C. Survival and prognostic factors in 189 dogs with dilated cardiomyopathy. *J Am Anim Hosp Assoc* 1997;33:364-368.
- Monnet E, Orton EC, Salam M, et al. Idiopathic dilated cardiomyopathy in dogs: survival and prognostic indicators. *J Vet Intern Med* 1995;9:12-17.
- Kirk CA, Debraekeleer J, Armstrong PJ. Normal cats. In: Hand MS, Thatcher CD, Remillard RL, et al, eds. *Small animal clinical nutrition*. 4th ed. Marcline, Mo: Walsworth Publishing Co, 2000; 291-347.
- National Research Council. *Nutrient requirements of cats*. Washington, DC: National Academy Press, 1986;13-15.
- 2002 Official publication. Oxford, England. In: Association of American Feed Control Officials Inc, 2002;140-141.
- Morris JG, Rogers QR, Kim SW, et al. Dietary taurine requirement of cats is determined by microbial degradation of taurine in the gut. In: Huxtable RJ, Michalk D, eds. *Taurine in health and disease*. New York: Plenum Press, 1994;59-70.
- Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L-carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am J Vet Res* 2001;62:1616-1623.
- Freeman LM, Michel KE, Brown DJ, et al. Idiopathic dilated cardiomyopathy in Dalmatians: nine cases (1990-1995). *J Am Vet Med Assoc* 1996;209:1592-1596.
- Stratton-Phelps M, Backus RC, Rogers QR, et al. Dietary rice bran decreases plasma and whole-blood taurine in cats. *J Nutr* 2002;132(6 suppl 2):1745S-1747S.
- Johnson ML, Parsons CM, Fahey GC, et al. Effects of species raw material source, ash content and processing temperature on amino acid digestibility of animal by-product meals by cecectomized and ileally cannulated dogs. *J Anim Sci* 1998;76:1112-1122.
- Torres CL, Backus RC, Fascetti AJ, et al. Taurine status in normal dogs fed a commercial diet associated with taurine deficiency and dilated cardiomyopathy. *J Anim Physiol Anim Nutr* 2003;in press.
- Backus RC, Cohen GL, Pion PD, et al. Taurine deficiency in Newfoundland dogs fed commercially available complete and balanced diets. *J Am Vet Med Assoc* 2003;223:1130-1136.
- Sturman JA, Gargano AD, Messing JM, et al. Feline maternal taurine deficiency: effect on mother and offspring. *J Nutr* 1985; 116:655-667.
- National Research Council. *Nutrient requirements of dogs*. Washington, DC: National Academy Press, 1985;11-12.
- Backus RC, Rogers QR, Rosenquist GL, et al. Diets causing taurine depletion in cats substantially elevate postprandial plasma cholecystokinin (CCK) concentration. *J Nutr* 1995;125:2650-2657.
- Pion PD, Lewis J, Greene K, et al. Effect of meal-feeding and food deprivation on plasma and whole blood taurine concentrations in cats. *J Nutr* 1991;121(suppl 11):S177-S178.
- Pacioretty L, Hickman MA, Morris JG, et al. Kinetics of taurine depletion and repletion in plasma, serum, whole blood and skeletal muscle in cats. *Amino Acids* 2001;21:417-427.

Iron Laboratory Studies in Pediatric Patients With Heart Failure from Dilated Cardiomyopathy



David Higgins, MD^a, Jessica Otero, PharmD^b, Christa Jefferis Kirk, PharmD^b, Jennifer Pak, PharmD^b, Neal Jorgensen, MS^c, Mariska Kemna, MD^a, Erin Albers, MD^a, Borah Hong, MD^a, Joshua Friedland-Little, MD^a, and Yuk Law, MD^{a,*}

Iron deficiency (FeD), with or without anemia, in adults with heart failure (HF) is associated with poor outcomes, which can be improved with replacement therapy. A similar therapeutic opportunity may exist for children; however, iron laboratory measurements and FeD have not been described in pediatric patients with HF. A single-center, retrospective study was conducted on 28 patients <21 years old with a diagnosis of dilated cardiomyopathy and HF who had iron laboratories (serum iron, iron saturation, and ferritin) performed. The mean (standard deviation) age at time of laboratory collection was 10.3 (5.5) years. Twenty-seven patients (96.4%) met the criteria for FeD. Serum iron and iron saturation were significantly associated with inpatient hospitalization, being on inotropic medications, or having stage D HF. Low-serum iron was associated with a higher left ventricular end-diastolic dimension and left ventricular end-systolic dimension z-score by echocardiography (β -2.58, 95% confidence interval [CI] -4.76, -0.40, p = 0.02) and (β -2.43, 95% CI -4.70, -0.17, p = 0.04), respectively. Low ferritin was associated with higher mortality (relative risk 0.29, 95% CI 0.12, 0.70, p = 0.006). In conclusion, FeD was common in this pediatric cohort with more advanced HF. Iron profile abnormalities were associated with worse HF severity and outcomes including mortality. © 2017 Elsevier Inc. All rights reserved. (Am J Cardiol 2017;120:2049–2055)

Iron deficiency (FeD) is the most common nutrient deficiency worldwide, and is particularly prevalent in children and females of childbearing age.^{1,2} Iron is an essential nutrient and a cofactor that is crucial for multiple cellular processes and is likely even more important in metabolically active cells such as cardiac myocytes.³ FeD in adult patients with heart failure (HF) has been studied extensively.^{4–9} Several studies have estimated the prevalence of FeD in adult patients with HF to be as high as 73%.^{10–13} Recent trials of iron supplementation in adult HF patients demonstrated significant improvements in New York Heart Association functional class, quality of life scores, exercise capacity, anemia, and a reduction in the risk of hospitalization.^{4–9,14} No studies have investigated the coexistence of FeD with HF in pediatric patients where growth may place a further demand on proper iron metabolism. Therefore, children with HF may be at an especially high risk for FeD and the associated comorbidities. Timely identification of FeD may also allow early intervention. In this study we evaluated the iron profiles in a cohort of children with HF from dilated cardiomyopathy to examine the prevalence and outcomes of FeD in this disease population.

Methods

This was a single-center, retrospective observational study. The study was approved by the local Institutional Review Board. Inclusion criteria were age of <21 years old, the diagnosis of dilated cardiomyopathy, and at least one set of iron laboratories that included ferritin, serum iron, and iron saturation [Tsat] plus a same-day complete blood count. Patients were identified through a systematic search of the institution's electronic medical record for cardiomyopathy or HF based on the International Classification of Diseases-Ninth Revision-Clinical Modification (ICD-9-CM) codes 425.1, 425.11, 425.18, 425.4, 425.8, 425.9, 428.0, 428.1, 428.20, 428.21, 428.22, 428.23, 428.32, 428.41, 428.42, 428.43, 428.9, 428.0, 429.3, 429.89, and 429.9 for the period from 2005 to 2014. This list of identified patients was adjudicated by the investigators to ensure inclusion criteria were met, including the presence of a complete set of iron laboratories and complete blood count. Patients were excluded if they were already receiving iron supplementation, post-cardiac transplantation, receiving chemotherapy or erythropoietin, received a transfusion within 2 weeks before obtaining iron laboratories, exposed to cardiopulmonary bypass 2 weeks before iron laboratories, or have renal dysfunction on renal replacement therapy as these conditions are likely to affect the iron profile. Patients with congenital heart defects leading to HF were excluded as this group is likely to have a significantly different etiology to their HF.

A comprehensive chart review was performed on the 28 patients included in the study to collect their demographics including age, gender, race/ethnicity, location at time of the iron laboratories (outpatient, inpatient, or cardiac intensive

^aDivision of Cardiology, Seattle Children's Hospital, University of Washington, Seattle, Washington; ^bDivision of Pharmacy, Seattle Children's Hospital, University of Washington, Seattle, Washington; and ^cDepartment of Biostatistics, University of Washington, Seattle, Washington. Manuscript received May 15, 2017; revised manuscript received and accepted August 14, 2017.

See page 2054 for disclosure information.

*Corresponding author: Tel: (206) 987-1417; fax: (206) 987-3839.

E-mail address: Yuk.Law@seattlechildrens.org (Y. Law).

care unit [ICU]), echocardiogram, presence of anemia and microcytic anemia (based on hemoglobin or MCV less than the 5th percentile for age),¹⁵ and the use of HF medications including beta blockers, angiotensin-converting enzyme inhibitors/angiotensin receptor blockers, diuretics, anticoagulants, or inotropes within 48 hours of laboratory collection. Outcome measures included mortality and use of mechanical circulatory support (MCS) following collection of iron profile laboratories. Patients were followed up until the end of 2014 with a mean (standard deviation [SD]) time from laboratory collection to end of follow-up of 2.05 (2.11) years. FeD was assessed by examining serum iron level (mcg/dL), total iron binding capacity (mcg/dL), Tsat, and ferritin (ng/mL). Because the definition of FeD in healthy adults and children is nearly identical, we used the definition of FeD in adults with HF which has been used consistently in major studies and trials: ferritin <100 ng/mL alone or a ferritin 100 to 300 ng/mL and Tsat <20%.¹⁴

Echocardiography data to assess the severity of dilated cardiomyopathy were abstracted from the echocardiogram reports at the time of iron laboratories, including the left ventricular shortening fraction (%), left ventricular end-diastolic dimension z-score, and left ventricular end-systolic dimension z-score. The reviewed echocardiograms were performed on average within 14 days of iron laboratory collection. To grade the severity of HF at the time of iron laboratory collection, HF stage,^{16,17} plasma B-type natriuretic peptide (BNP), location at time of laboratory collection (outpatient vs inpatient), and use of an inotrope were collected.

Descriptive data were presented as counts with percentages or mean values and SDs where appropriate. Multivariable linear regression models were used to examine the relation between iron laboratory levels with BNP and echocardiographic parameters. Multivariable relative risk regression was used to assess the relation between iron laboratory levels with location of laboratories (inpatient/outpatient), inotrope use, and HF stage D. Binary outcomes for mortality, need for MCS, and a “composite” outcome defined as either mortality or need for MCS were analyzed using univariate relative risk regression as the number of events was not adequate for a multivariable analysis. All linear and relative risk regression models used robust standard errors. Natural log transformations were used for BNP, serum iron, Tsat, and ferritin in the linear regression models. For the relative risk regression models, iron values and BNP were transformed by taking the base-2 logarithm. Multivariable general additive models adjusted for age and race were used to illustrate the relation of left ventricular end-systolic dimension (LVESD) and left ventricular end-diastolic dimension (LVESD) (z-scores) with iron laboratory values on the nontransformed scale using 3 degrees of freedom. Data analysis was performed using Stata 14.0 (StataCorp LP, College Station, Texas).

Results

A total of 28 patients were included in this study. The baseline characteristics of the study population are listed in [Table 1](#). A majority of patients were hospitalized when laboratories were obtained with 53.6% of patients in the cardiac ICU, 21.4% in the acute care ward (non-ICU), and 25.0% in outpatient clinics. A majority of patients had HF stage D (71.4%),

Table 1
Summary of patient characteristics

Demographics	Value, % or mean (SD) (n = 28)
Age (years)	10.3 (5.5)
Male	13 (46%)
Race/ethnicity	
White	11 (39%)
Black	4 (14%)
Asian	1 (4%)
Other or “Did not indicate”	12 (43%)
Weight (kg)	41.0 (27)
Weight z-score	-0.4 (2.0)
Height (cm)	134 (36)
Height z-score	-0.7 (1)
BMI	20 (7)
BMI z-score	0.01 (1)
Location at Lab	
Outpatient	7 (25%)
Inpatient (non-ICU)	6 (21%)
CICU	15 (54%)
Heart failure stage D*	20 (71%)
Anemia (Hgb)	11 (39%)
Microcytic anemia	2 (7%)
Heart Failure Medications	
Beta-Blocker	9 (32%)
ACE-I/ARB	25 (89%)
Diuretic	22 (79%)
Anticoagulation	15 (54%)
Inotrope	18 (64%)
BNP (pg/mL)	745 (912)
Clinical Outcomes	
Mortality	4 (14%)
Listed for transplant	12 (43%)
Mechanical Circulatory Support	4 (14%)

* Based on Heart Failure Staging for Infants and Children modifications made to the ACC/AHA staging guidelines.^{14,15}

Table 2
Summary of iron profile variables

Iron Status	
Deficiency*	27 (96%)
Ferritin <100 (ng/mL)	26 (93%)
Tsat <20%	22 (79%)
Iron Lab Profile	mean (SD)
Serum Iron (µg/dL)	60 (35)
Tsat %	17 (10)
Ferritin (ng/mL)	45 (30)

* Iron deficiency defined as ferritin < 100 ng/mL or ferritin = 100–300 ng/mL and iron saturation (Tsat) < 20%.

and the mean (SD) plasma BNP for the entire cohort was 745 (912) pg/mL.

The cohort’s iron profile is listed in [Table 2](#). Based on the previously published definition of FeD in adult HF patients, 27 patients (96.4%) met the criteria for FeD. Twenty-six patients (92.9%) met criteria for FeD based on ferritin <100 ng/mL alone. Of the 26 patients with low ferritin, 21 (80.8%) also had a Tsat <20%. Anemia for age was found in 11 (39.3%) patients and only 2 of those patients had a microcytic anemia.

Table 3

(A) Multivariable relative risk model for clinical heart failure severity with iron profile variables. (B) Multivariable linear regression modeling B-type natriuretic peptide (BNP) and echocardiographic measures of dilated cardiomyopathy with iron profile variables

(A)						
Variable [†]	Inpatient (n = 21/28)		Inotrope (n = 18/28)		Heart failure stage D* (n = 20/28)	
	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p
Log2(Serum Iron)	0.71 [0.56,0.90]	0.006	0.67 [0.51,0.87]	0.003	0.82 [0.66,1.03]	0.095
Log2 (Tsat)	0.69 [0.53,0.92]	0.011	0.63 [0.48,0.81]	0.001	0.80 [0.61,1.04]	0.092
Log2 (Ferritin)	1.00 [0.82,1.22]	0.982	0.94 [0.73,1.20]	0.607	0.99 [0.80,1.21]	0.887
(B)						
Variable [†]	Ln(BNP) (n = 28)		LVEDD (z-score) (n = 28)		LVESD (z-score) (n = 28)	
	β (95% CI)	p	β (95% CI)	p	β (95% CI)	p
Ln(Serum Iron)	-1.16 [-2.02,-0.29]	0.011	-2.58 [-4.76,-0.40]	0.023	-2.43 [-4.70,-0.17]	0.036
Ln(Tsat)	-1.04 [-1.99,-0.09]	0.033	-2.50 [-4.96,-0.05]	0.046	-2.50 [-4.91,-0.09]	0.043
Ln(Ferritin)	-0.02 [-0.54,0.50]	0.928	-0.93 [-2.38,0.53]	0.201	-1.33 [-2.95,0.28]	0.101

* Based on Heart Failure Staging for Infants and Children modifications made to the ACC/AHA staging guidelines.^{14,15}

[†] Models adjusted for Age and Race with each Iron Profile variable added one at a time.

Additionally, there was no association between patient weight or body mass index (BMI) z-scores and iron laboratory values.

We compared the degree of iron profile abnormalities with the severity of the cardiomyopathy and HF. Patients were grouped into HF severity based on location (outpatient vs inpatient), the use of inotropes, and HF stage at the time of laboratory collection. Table 3A lists the multivariable analyses between these markers of HF severity and iron laboratory values. While ferritin did not show an independent association with the severity of HF using these measures, both serum iron and Tsat were associated with being inpatient and being on inotropic medications (Table 3).

Echocardiography parameters relevant to dilated cardiomyopathy were also compared with the iron profile. Table 3B lists the independent relation between iron laboratory values and these echocardiographic parameters. Lower levels of serum iron and Tsat were significantly associated with a higher LVEDD z-score and a higher LVESD z-score, but ferritin was not associated with remodeling on echocardiography. Figure 1 provides an illustration of the relation between LVEDD, LVESD, and iron levels. BNP was also examined in relation to iron values. Lower serum iron and Tsat were significantly associated with higher BNP levels, a marker of increased wall stress.

We compared the degree of iron profile abnormalities with clinical outcome measures including mortality and MCS. A total of 8 patients had a poor clinical outcome (death or need for MCS). Table 4 lists the associations between iron laboratory values and these clinical outcome measures. Higher serum iron trended toward not requiring MCS. High ferritin level was significantly associated with lower mortality and lower composite outcome. It is important to note that among the other clinical and laboratory indices including age, weight, BMI, patient location, HF stage, presence of anemia, echocardiographic indices of remodeling, and inotrope

use, none were independently associated with clinical outcomes.

Discussion

In this study, we observed a high prevalence of FeD in a pediatric cohort with HF. We also examined individual components of the conventional iron laboratory profile and their relation to left ventricular remodeling by echocardiography, severity of HF, and HF related cardiovascular outcomes. Lower levels of iron storage and availability were in general associated with higher severity of remodeling, HF, and worst clinical outcomes, whereas BNP and clinical or echocardiographic features were not.

In our cohort, the frequency of FeD, 91%, is quite alarming. Although not reported before in the pediatric literature, the prevalence in our study is higher than reports from adult HF studies, and this is likely due to several different factors. First, the cohort studied has higher severity of illness as demonstrated by the proportion admitted to the hospital and on inotropes as well as by the overall mortality and need for MCS, whereas the adult studies focused on ambulatory patients with HF. Additionally, defining FeD in HF patients is challenging, and this difficulty emanates from the fact that FeD can be understood as total body depletion or a functional metabolic deficiency.¹⁸ Because patients with HF are susceptible to both types of deficiency, diagnosing FeD in HF patients is complex.¹⁹ In other medical conditions, the diagnosis of FeD is a ferritin <30 μg/L and a Tsat <16%. However, these definitions have serious limitations as well because in long-term illnesses and inflammatory states, ferritin can be increased independently of iron status.¹⁸ For example, similar to the accepted adult definition for iron deficiency, in pediatric chronic kidney disease, a ferritin <100 μg/L is also the accepted cutoff for iron deficiency.²⁰ Furthermore, a study in adult patients with HF demonstrated that 73% had confirmed FeD on bone




Figure 1. Relation between (A) LVESD and (B) LVEDD (z-scores) and iron profile values. LVEDD = left ventricular end-diastolic dimension; LVESD = left ventricular end-systolic dimension.

Table 4
Univariate relative risk regression modeling clinical outcomes

Variable	Mortality (n = 4/28)		MCS (n = 4/28)		Composite* (n = 8/28)	
	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p
Age (years)	1.11 [0.95,1.30]	0.197	1.00 [0.86,1.17]	0.954	1.05 [0.95,1.16]	0.357
Male	1.15 [0.18,7.32]	0.879	0.38 [0.04,3.39]	0.390	0.69 [0.20,2.41]	0.563
White	3.00 [0.34,26.48]	0.323	0.33 [0.04,2.94]	0.323	1.00 [0.30,3.30]	1.000
Weight (kg)	1.01 [0.99,1.04]	0.287	1.01 [0.98,1.03]	0.633	1.01 [0.99,1.02]	0.249
Weight z-score	1.09 [0.66,1.78]	0.742	0.97 [0.48,1.95]	0.934	1.03 [0.68,1.56]	0.880
Height (cm)	1.02 [0.99,1.04]	0.1900	1.01 [0.97,1.05]	0.687	1.01 [0.99,1.03]	0.291
Height z-score	1.05 [0.69,1.61]	.809	1.41 [0.46,4.33]	0.549	1.20 [0.73,1.96]	0.471
BMI	1.03 [0.92,1.14]	0.634	1.01 [0.91,1.11]	0.915	1.01 [0.95,1.08]	0.649
BMI z-score	1.01 [0.30,3.43]	0.985	0.93 [0.34,2.52]	0.881	0.96 [0.40,2.31]	0.922
Location at Lab		0.231		—		1.000
Outpatient	ref				ref	
Inpatient	0.33 [0.06,2.01]		All MCS		1.00 [0.25,3.96]	
Heart failure stage D [†]	0.40 [0.07,2.45]	0.322	All MCS	—	1.20 [0.30,4.86]	0.798
Anemia (Hgb)	1.55 [0.25,9.74]	0.643	0.52 [0.06,4.52]	0.549	0.93 [0.27,3.19]	0.905
Beta-Blocker	6.33 [0.73,54.84]	0.094	No MCS	—	1.27 [0.38,4.26]	0.703
Diuretic	0.82 [0.10,6.77]	0.852	All MCS	—	1.91 [0.28,13.09]	0.510
Anticoagulation	All Mortality	—	All MCS	—	All Composite	—
Inotrope	0.56 [0.09,3.48]	0.530	All MCS	—	1.67 [0.40,6.94]	0.483
Cardiac Function						
Log2(BNP)	0.86 [0.72,1.04]	0.122	1.71 [0.97,3.02]	0.064	1.09 [0.84,1.41]	0.507
LVEDD (z-score)	1.01 [0.85,1.20]	0.939	1.51 [0.57,4.00]	0.403	1.27 [0.84,1.93]	0.256
LVESD (z-score)	0.99 [0.83,1.17]	0.906	1.35 [0.92,1.99]	0.130	1.15 [0.94,1.40]	0.164
Log2(Serum Iron)	1.66 [0.82,3.35]	0.161	0.19 [0.03,1.07]	0.060	0.71 [0.34,1.45]	0.340
Log2(TSAT)	1.33 [0.67,2.65]	0.414	0.19 [0.03,1.31]	0.091	0.53 [0.20,1.38]	0.193
Log2(Ferritin)	0.29 [0.12,0.70]	0.006	0.74 [0.48,1.15]	0.177	0.53 [0.35,0.81]	0.003

* Composite of mortality or need for MCS.

[†] Based on Heart Failure Staging for Infants and Children modifications made to the ACC/AHA staging guidelines.^{16,17}

marrow aspiration despite a high ferritin.²¹ Therefore, traditional definitions in the general population may not be applicable to patients with HF, particularly those who are more advanced, have active symptoms, or are in a decompensated state. For this reason and the fact that most of our cohort has higher severity of HF, we also examined individual iron profile components as continuous variables.

Multiple studies in the adult HF literature have described the definition of FeD in HF as a serum ferritin level <100 µg/L (a measure of total body FeD), or serum ferritin level between 100 and 300 µg/L combined with T_{sat} <20% (a measure of functional FeD).^{22,23} This definition of FeD based on therapeutic responses has since been included in guidelines on the treatment of adults with HF.²⁴ Although there is no validated definition of FeD in children with HF, the observed association of lower available iron stores with higher severity of HF suggests that FeD should be further examined, including its repletion as a therapeutic target in HF in children as few exist in children with HF.

Iron has merit as a therapeutic target because it is essential for many cellular functions. For example, the importance of iron homeostasis goes beyond the commonly acknowledged effect on erythropoiesis. In fact, a majority of patients with FeD in this study did not have concurrent anemia, which is a similar observation in adult studies.¹⁸ Iron is a critical part of cellular processes such as oxygen transport, oxygen storage, mitochondrial respiration, and cellular immunity.³ Hence, it is theoretically plausible that FeD can lead to disease

progression and poor outcomes as suggested by the limited preliminary outcome analysis in this study. Even without a well-elucidated mechanism to explain these associations, multiple adult studies have demonstrated FeD as an independent predictor of all-cause mortality,^{10,13,25} whereas parenteral iron replacement improved markers of cardiac remodeling and HF.^{5,26,27} It is worth noting that weight and BMI were not associated with the iron profile, suggesting FeD may not simply emanate from poor intake in which case parenteral replacement therapy is required.

In our study, there were differences between the various components of the traditional iron laboratory profile that warrant discussion. First, lower serum iron and T_{sat} were associated with higher BNP and a higher severity of remodeling, whereas the level of ferritin was not. However, ferritin was associated with mortality and composite outcome. These observed differences between markers of iron availability (serum iron and T_{sat}) and iron storage (ferritin) may be due, in part, to the independent elevation of ferritin with inflammation, which may be heightened and more prevalent during acute decompensated HF.^{21,28} However, it is also possible that iron availability is more important than depleted total body stores in acute HF where the cardiomyocyte iron requirements may theoretically increase. Conversely, ferritin as a storage marker may be more related to the duration and hence underlying reserve and reversibility of the diseased heart, which may contribute more to the association with major adverse outcomes other than admission for acute decompensated HF. Future large

studies encompassing a broader group of patients at all stages of HF will be important to determine the significance of these findings.

There are several limitations to this study. In addition to the potential selection bias of chronically ill and hospitalized patients enrolled in the study, the sample size is small. Therefore, prevalence is not likely to be accurate from this study. Pediatric HF encompasses heterogeneity of diagnoses. By keeping the cohort to dilated cardiomyopathy and excluding patients whose iron profile could be affected by transfusions and other disease states that can affect iron metabolism, the sample size was further diminished. We graded severity of HF by how patients were treated instead of symptoms because it is difficult to assign a symptom-based functional class to children, especially in a retrospective study. This is a limitation in the sense that it would be more difficult to compare our findings with the existing literature on FeD in adults with HF, as severity in adults is symptom-based and graded by a well-established and validated functional class system. Because of the potential patient selection bias, and because a numerical cutoff does not adequately define “deficiency” in disease states, we purposely examined the association of iron laboratory values as a continuous variable with severity of HF and cardiomyopathy to demonstrate the presence of robust relations in children.

Despite recognition of the importance of FeD in adults with HF, this study describes FeD as a new, potentially serious comorbidity in children. FeD appears to be common, particularly in sicker, more advanced HF patients. Iron profiling may also serve as biomarkers for clinical outcome. The observations in this study should help guide future studies investigating the effects of iron metabolism on pediatric HF outcomes and whether FeD can be a therapeutic target for HF therapy in this orphan disease population.

Disclosures

The authors have no conflicts of interest to disclose.

- (CDC) CfDcCaP. Iron deficiency—United States, 1999–2000. *MMWR Morb Mortal Wkly Rep* 2002;51:897–899.
- Pfeiffer CM, Sternberg MR, Schleicher RL, Haynes BM, Rybak ME, Pirkle JL. The CDC’s Second National Report on Biochemical Indicators of Diet and Nutrition in the U.S. Population is a valuable tool for researchers and policy makers. *J Nutr* 2013;143:938S–947S.
- Beard JL. Iron biology in immune function, muscle metabolism and neuronal functioning. *J Nutr* 2001;131:568S–579S.
- Bolger AP, Bartlett FR, Penston HS, O’Leary J, Pollock N, Kaprielian R, Chapman CM. Intravenous iron alone for the treatment of anemia in patients with chronic heart failure. *J Am Coll Cardiol* 2006;48:1225–1227.
- Toblli JE, Lombraña A, Duarte P, Di Gennaro F. Intravenous iron reduces NT-pro-brain natriuretic peptide in anemic patients with chronic heart failure and renal insufficiency. *J Am Coll Cardiol* 2007;50:1657–1665.
- Okonko DO, Grzeslo A, Witkowski T, Mandal AK, Slater RM, Roughton M, Foldes G, Thum T, Majda J, Banasiak W, Missouri CG, Poole-Wilson PA, Anker SD, Ponikowski P. Effect of intravenous iron sucrose on exercise tolerance in anemic and nonanemic patients with symptomatic chronic heart failure and iron deficiency FERRIC-HF: a randomized, controlled, observer-blinded trial. *J Am Coll Cardiol* 2008;51:103–112.
- Anker SD, Comin Colet J, Filippatos G, Willenheimer R, Dickstein K, Drexler H, Lüscher TF, Bart B, Banasiak W, Niegowska J, Kirwan BA, Mori C, von Eisenhart Rothe B, Pocock SJ, Poole-Wilson PA, Ponikowski P, Investigators F-HT. Ferric carboxymaltose in patients with heart failure and iron deficiency. *N Engl J Med* 2009;361:2436–2448.
- Ponikowski P, van Veldhuisen DJ, Comin-Colet J, Ertl G, Komajda M, Mareev V, McDonagh T, Parkhomenko A, Tavazzi L, Levesque V, Mori C, Roubert B, Filippatos G, Ruschitzka F, Anker SD, Investigators C-H. Beneficial effects of long-term intravenous iron therapy with ferric carboxymaltose in patients with symptomatic heart failure and iron deficiency. *Eur Heart J* 2015;36:657–668.
- Ponikowski P, Veldhuisen DJ, Comin-Colet J, Ertl G, Komajda M, Mareev V, McDonagh TA, Parkhomenko A, Tavazzi L, Levesque V, Mori C, Roubert B, Filippatos G, Ruschitzka F, Anker SD. Rationale and design of the CONFIRM-HF study: a double-blind, randomized, placebo-controlled study to assess the effects of intravenous ferric carboxymaltose on functional capacity in patients with chronic heart failure and iron deficiency. *ESC Heart Fail* 2014;1:52–58.
- Jankowska EA, Rozentryt P, Witkowska A, Nowak J, Hartmann O, Ponikowska B, Borodulin-Nadzieja L, Banasiak W, Polonski L, Filippatos G, McMurray JJ, Anker SD, Ponikowski P. Iron deficiency: an ominous sign in patients with systolic chronic heart failure. *Eur Heart J* 2010;31:1872–1880.
- Parikh A, Natarajan S, Lipsitz SR, Katz SD. Iron deficiency in community-dwelling US adults with self-reported heart failure in the National Health and Nutrition Examination Survey III: prevalence and associations with anemia and inflammation. *Circ Heart Fail* 2011;4:599–606.
- Kasner M, Aleksandrov AS, Westermann D, Lassner D, Gross M, von Haehling S, Anker SD, Schultheiss HP, Tschöpe C. Functional iron deficiency and diastolic function in heart failure with preserved ejection fraction. *Int J Cardiol* 2013;168:4652–4657.
- Klip IT, Comin-Colet J, Voors AA, Ponikowski P, Enjuanes C, Banasiak W, Lok DJ, Rosentryt P, Torrens A, Polonski L, van Veldhuisen DJ, van der Meer P, Jankowska EA. Iron deficiency in chronic heart failure: an international pooled analysis. *Am Heart J* 2013;165:575–582.
- von Haehling S, Jankowska EA, van Veldhuisen DJ, Ponikowski P, Anker SD. Iron deficiency and cardiovascular disease. *Nat Rev Cardiol* 2015;11:659–669.
- Lanzkowsky P. Iron-deficiency anemia. In: Lanzkowsky’s Manual of Pediatric Hematology and Oncology. Sixth ed. San Diego, CA: Academic Press; 2016:69–83 [Chapter 6].
- Rosenthal D, Chrisant MR, Edens E, Mahony L, Canter C, Colan S, Dubin A, Lamour J, Ross R, Shaddy R, Addonizio L, Beerman L, Berger S, Bernstein D, Blume E, Boucek M, Checchia P, Dipchand A, Drummond-Webb J, Fricker J, Friedman R, Hallowell S, Jaquiss R, Mital S, Pahl E, Pearce FB, Pearce B, Rhodes L, Rotondo K, Rusconi P, Scheel J, Pal Singh T, Towbin J. International Society for Heart and Lung Transplantation: practice guidelines for management of heart failure in children. *J Heart Lung Transplant* 2004;23:1313–1333.
- Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013;128:1810–1852.
- Wong CC, Ng AC, Kritharides L, Sindone AP. Iron deficiency in heart failure: looking beyond anaemia. *Heart Lung Circ* 2016;25:209–216.
- Zimmermann MB, Hurrell RF. Nutritional iron deficiency. *Lancet* 2007;370:511–520.
- KDIGO Clinical practice guideline for anemia in chronic kidney disease. *Kidney Int Suppl* 2012;2:279–335.
- Nanas JN, Matsouka C, Karageorgopoulos D, Leonti A, Tsolakis E, Drakos SG, Tsalagou EP, Maroulidis GD, Alexopoulos GP, Kanakakis JE, Anastasiou-Nana MI. Etiology of anemia in patients with advanced heart failure. *J Am Coll Cardiol* 2006;48:2485–2489.
- Ebner N, von Haehling S. Iron deficiency in heart failure: a practical guide. *Nutrients* 2013;5:3730–3739.
- Jankowska EA, von Haehling S, Anker SD, Macdougall IC, Ponikowski P. Iron deficiency and heart failure: diagnostic dilemmas and therapeutic perspectives. *Eur Heart J* 2013;34:816–829.

24. McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, Falk V, Filippatos G, Fonseca C, Gomez-Sanchez MA, Jaarsma T, Køber L, Lip GY, Maggioni AP, Parkhomenko A, Pieske BM, Popescu BA, Rønnevik PK, Rutten FH, Schwitzer J, Seferovic P, Stepinska J, Trindade PT, Voors AA, Zannad F, Zeiher A, Bax JJ, Baumgartner H, Ceconi C, Dean V, Deaton C, Fagard R, Funck-Brentano C, Hasdai D, Hoes A, Kirchhof P, Knuuti J, Kolh P, McDonagh T, Moulin C, Reiner Z, Sechtem U, Sirnes PA, Tendera M, Torbicki A, Vahanian A, Windecker S, Bonnet LA, Avraamides P, Ben Lamin HA, Brignole M, Coca A, Cowburn P, Dargie H, Elliott P, Flachskampf FA, Guida GF, Hardman S, Jung B, Merkely B, Mueller C, Nanas JN, Nielsen OW, Orn S, Parissis JT, Ponikowski P, ESC Committee for Practice Guidelines. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2012;14:803–869.
25. Okonko DO, Mandal AK, Missouriis CG, Poole-Wilson PA. Disordered iron homeostasis in chronic heart failure: prevalence, predictors, and relation to anemia, exercise capacity, and survival. *J Am Coll Cardiol* 2011;58:1241–1251.
26. Usmanov RI, Zueva EB, Silverberg DS, Shaked M. Intravenous iron without erythropoietin for the treatment of iron deficiency anemia in patients with moderate to severe congestive heart failure and chronic kidney insufficiency. *J Nephrol* 2008;21:236–242.
27. Toblli JE, Di Gennaro F, Rivas C. Changes in echocardiographic parameters in iron deficiency patients with heart failure and chronic kidney disease treated with intravenous iron. *Heart Lung Circ* 2015;24:686–695.
28. Dick SA, Epelman S. Chronic heart failure and inflammation: what do we really know? *Circ Res* 2016;119:159–176.

Reproduced with permission of copyright owner. Further reproduction prohibited without permission.

Chronic oral exposure to the aldehyde pollutant acrolein induces dilated cardiomyopathy

Mohamed Ameen Ismahil,¹ Tariq Hamid,¹ Petra Haberzettl,¹ Yan Gu,^{1,3} Bysani Chandrasekar,² Sanjay Srivastava,¹ Aruni Bhatnagar,¹ and Sumanth D. Prabhu^{1,3}

¹Department of Medicine, Institute of Molecular Cardiology, University of Louisville, Louisville, ²Heart and Vascular Institute, Tulane University School of Medicine and Southeast Louisiana Veterans Health Care System, New Orleans; and ³Medical Service, Louisville Veterans Affairs Medical Center, Louisville, Kentucky

Submitted 3 February 2011; accepted in final form 24 August 2011

Ismahil MA, Hamid T, Haberzettl P, Gu Y, Chandrasekar B, Srivastava S, Bhatnagar A, Prabhu SD. Chronic oral exposure to the aldehyde pollutant acrolein induces dilated cardiomyopathy. *Am J Physiol Heart Circ Physiol* 301: H2050–H2060, 2011. First published September 9, 2011; doi:10.1152/ajpheart.00120.2011.—Environmental triggers of dilated cardiomyopathy are poorly understood. Acute exposure to acrolein, a ubiquitous aldehyde pollutant, impairs cardiac function and cardioprotective responses in mice. Here, we tested the hypothesis that chronic oral exposure to acrolein induces inflammation and cardiomyopathy. C57BL/6 mice were gavage-fed acrolein (1 mg/kg) or water (vehicle) daily for 48 days. The dose was chosen based on estimates of human daily unsaturated aldehyde consumption. Compared with vehicle-fed mice, acrolein-fed mice exhibited significant ($P < 0.05$) left ventricular (LV) dilatation (LV end-diastolic volume 36 ± 8 vs. 17 ± 5 μ l), contractile dysfunction (dP/dt_{max} $4,697 \pm 1,498$ vs. $7,016 \pm 1,757$ mmHg/s), and impaired relaxation (τ 15.4 ± 4.3 vs. 10.4 ± 2.2 ms). Histological and biochemical evaluation revealed myocardial oxidative stress (membrane-localized protein-4-hydroxy-*trans*-2-nonenal adducts) and nitritative stress (increased protein-nitrotyrosine) and varying degrees of plasma and myocardial protein-acrolein adduct formation indicative of physical translocation of ingested acrolein to the heart. Acrolein also induced myocyte hypertrophy (~ 2.2 -fold increased myocyte area, $P < 0.05$), increased apoptosis (~ 7.5 -fold), and disrupted endothelial nitric oxide synthase in the heart. DNA binding studies, immunohistochemistry, and PCR revealed significant ($P < 0.05$) activation of nuclear factor- κ B in acrolein-exposed hearts, along with upregulated gene expression of proinflammatory cytokines tumor necrosis factor- α and interleukin- 1β . Long-term oral exposure to acrolein, at an amount within the range of human unsaturated aldehyde intake, induces a phenotype of dilated cardiomyopathy in the mouse. Human exposure to acrolein may have analogous effects and raise consideration of an environmental, aldehyde-mediated basis for heart failure.

acrolein; oxidative stress; cardiomyopathy; environmental pollution

IDIOPATHIC DILATED CARDIOMYOPATHY (DCM) is the underlying diagnosis in approximately one-third of cases of heart failure (HF) (15). While often attributed to remote infectious, metabolic, or toxic injury to the heart, in most circumstances the etiological factors responsible for DCM are difficult to identify. Epidemiological studies have established that pollution exposure is associated with increased mortality from several cardiovascular diseases, including HF (3, 5, 28). The biological mechanisms proposed to explain these adverse effects have included pollutant-induced alterations in autonomic tone, the

elaboration of proinflammatory and prooxidant mediators, and the physical translocation of soluble constituents of pollutants into the circulation that have direct effects on the heart and vasculature. Theoretically, all of these broad mechanisms can unfavorably impact pathogenetic alterations and/or modifiers of DCM and HF (16). Nonetheless, little is known about the potential environmental triggers of DCM and the specific effects induced by individual constituents of the pollutant mix.

Aldehydes are ubiquitous pollutants in air and water generated by burning fossil fuels (10). They are also readily found in food and are natural products of lipid peroxidation and glucose oxidation (10). More than 300 different aldehydes have been identified in various foods, and at least 36 are present in water, often at levels exceeding maximal recommended concentrations (2, 10). Unsaturated aldehydes are highly reactive; form adducts with cell thiols and amine groups in sugars, phospholipids, proteins, and DNA bases (9, 25); and provoke oxidative stress and proinflammatory responses in tissue (30, 38). Nonetheless, the *in vivo* cardiovascular effects of exposure to aldehyde pollutants are not well defined.

Because toxicological profiles of environmental aldehyde mixtures are difficult to determine, we have previously focused on the cardiac effects of acrolein, a prototypical reactive α,β -unsaturated aldehyde classified by the Environmental Protection Agency (EPA) as a high-priority air and water toxic (7). These studies demonstrated that acute exposure to acrolein at concentrations documented in human disease, or doses approximating human oral total aldehyde intake, impaired cardiac function and intrinsic cardioprotective responses in mice (19, 42). However, the cardiac effects of long-term acrolein exposure, an issue with greater implications for public health, remain unknown. Notably, the abundance of acrolein and other aldehydes derived endogenously from lipid peroxidation (and their protein-aldehyde adducts) are known to be elevated in the failing heart (14, 33, 40, 41). In the current study, we evaluated whether long-term oral exposure to acrolein would engender inflammation, oxidant stress, and cardiomyopathy.

METHODS

Eight-week-old male C57BL/6 mice weighing ~ 20 g were used. All animal studies were performed in compliance with the National Institutes of Health (NIH) *Guide for the Care and Use of Laboratory Animals* [Department of Health and Human Services Publication No. (NIH) 85-23, revised 1996] and were approved by the University of Louisville Institutional Animal Care and Use Committee.

Address for reprint requests and other correspondence: S. D. Prabhu, Medicine/Cardiology, Univ. of Louisville, ACB, 3rd Floor, 550 South Jackson St., Louisville, KY 40202 (e-mail: sprabhu@louisville.edu).

Acrolein dosage and administration. Acrolein was prepared daily from the acid hydrolysis of diethyl acetal acrolein as previously described (19, 42) and used within 4 h. In our previous study, we estimated the maximal human daily unsaturated aldehyde consumption to be $5 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ and maximal acrolein exposure to be $0.1\text{--}0.2 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ (42). Based on these estimates, and with the intent of using acrolein as a representative unsaturated aldehyde, we tested the chronic effects of $1 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ acrolein, representing a 5- to 10-fold greater dose than the expected human acrolein intake but only 20% of the expected overall unsaturated aldehyde intake. Animals were gavage-fed acrolein (in $200 \mu\text{l}$ water, $n = 15$) or the same volume of water (vehicle, $n = 18$) daily for 48 days.

Echocardiography. M-mode, two-dimensional, and Doppler echocardiography in mice were performed under tribromoethanol sedation (0.25 mg/g ip) using a Philips Sonos 5500 machine and 15-MHz linear array transducer as previously described (14, 41). The two echocardiographers performing the study were blinded as to the assigned experimental group of each mouse. Measured parameters included end-diastolic (ED) and end-systolic (ES) diameter (D), end-diastolic anterior and posterior wall thickness (AWT and PWT, respectively), and the ejection time (ET) and heart rate as determined from the aortic Doppler trace. Left ventricular (LV) systolic function was indexed by the fractional shortening [$\text{FS} = (\text{EDD} - \text{ESD})/\text{EDD}$] and the mean velocity of circumferential fiber shortening ($V_{\text{cf}} = \text{FS}/\text{ET}$) (34, 35). LV hypertrophy and/or wall thinning was assessed by the relative wall thickness [$\text{RWT} = (\text{AWT} + \text{PWT})/\text{LVEDD}$]. Echocardiographic imaging was performed at baseline and after 48 days of acrolein feeding.

LV pressure-volume studies. Closed-chest LV pressure-volume (P-V) studies were performed in adult C57/BL6 mice ($n = 8/\text{group}$) anesthetized with $80 \mu\text{g/g}$ ip pentobarbital and mechanically ventilated (155–160 breaths/min, tidal volume $15 \mu\text{l/g}$) as previously described (19). Body temperature was maintained at 37°C using a heating pad and lamps. A Millar 1.4-Fr conductance catheter (SPR-839) was inserted in the LV via the carotid artery, and pressure and conductance signals were visualized on-line using the ARIA-1 system (Millar). A small ($<1\text{-cm}$) abdominal incision was made to gain access to the subdiaphragmatic inferior vena cava (IVC). After hemodynamic stabilization for 15 min, recordings of pressure and conductance were performed under steady-state conditions and during transient mechanical IVC occlusion [to vary load and allow determination of the end-systolic pressure-volume relation (ESPVR)]. Intravenous hypertonic saline ($0.5\text{--}1 \mu\text{l/g}$) was then given to determine parallel conductance, and LV volume (μl) was derived from the parallel conductance and ex vivo cuvette calibration with heparinized, warm blood. LV systolic function was indexed by dP/dt_{max} , stroke work (area bounded by the P-V loop), maximal power (peak value of the product of LV pressure and flow), and end-systolic elastance (E_{es} , the slope of the ESPVR) (19, 41). LV diastolic function was assessed by the LVEDP, dP/dt_{min} , and tau, the time constant of LV relaxation (ms) (19, 33, 41).

Immunohistological studies. Formalin-fixed, paraffin-embedded short-axis LV sections ($5 \mu\text{m}$) were deparaffinized and rehydrated for histological and immunohistochemical staining using standard techniques as previously described (14, 34, 41). Hematoxylin and eosin-stained sections were used to evaluate cardiomyocyte cross-sectional area. In separate studies, immunostaining was performed for the activated p65 subunit of nuclear factor (NF)- κB using anti-p65 antibody (Chemicon) as described previously (31). Nuclear staining intensity was quantified with a MetaMorph 4.5 imaging system and software (Universal Imaging). Digital images were acquired from six fields at standard intervals in each of five short-axis sections from each group. The threshold for p65 staining was predetermined and held constant for all sections analyzed.

Immunohistochemical staining for protein-nitrotyrosine was performed to index peroxynitrite generation in the heart. Deparaffinized and rehydrated tissue sections were incubated for 20 min

with 10 mmol/l citric acid (pH 6.0) and then treated with enzymatic antigen retrieval to recover antigenicity. Nonspecific binding was blocked with 5% normal goat serum and 0.05% saponin (Sigma) in PBS (pH 7.4) for 30 min, followed by incubation with monoclonal anti-nitrotyrosine antibody (1:200; Santa Cruz Biotechnology) in PBS with 1% BSA and 0.05% saponin for 1 h at 37°C . Tissue sections were then incubated for 30 min at room temperature with Alexa fluor-555 anti-mouse IgG (1:500) secondary antibody (Invitrogen), which labeled nitrotyrosinated protein residues red, and counterstained with 4',6-diamidino-2-phenylindole (DAPI) (Invitrogen), which labels nuclei blue. Images were made with a $\times 40$ objective lens at 12 different locations in each tissue section. Mean fluorescence intensity was evaluated using MetaMorph software in 12 images/heart. Sections treated with peroxynitrite (1 mmol/l) were used as positive controls.

Western blotting. Total protein extraction, SDS-PAGE Western blotting, and immunodetection using electrochemiluminescence protocols (Amersham Biosciences) were performed as previously described (19). IgG-purified polyclonal 1:2,000 anti-KLH-acrolein primary antibody and horseradish peroxidase-linked secondary antibody were used to evaluate protein-acrolein adducts (19). Protein adducts with 4-hydroxy-*trans*-2-nonenal (HNE) in the membrane fraction (isolated using differential centrifugation) were probed using both dot blots and Western blotting. Polyclonal anti-KLH-HNE primary antibody was used as previously described (34). For dot blots, protein ($1.0 \mu\text{g}$) was loaded in the wells of a Bio-Dot apparatus (Bio-Rad) and microfiltered through nitrocellulose membranes under vacuum. Primary antibodies for the detection of endothelial nitric oxide (NO) synthase (eNOS), phospho-eNOS-Ser¹¹⁷⁷, inhibitor of κB ($\text{I}\kappa\text{B}\alpha$), and α -tubulin were obtained from Santa Cruz Biotechnology.

For immunoblot analysis of the monomeric and dimeric forms of eNOS, equal amounts of total protein lysates were subjected to low-temperature SDS-PAGE (LT-PAGE) (43). Briefly, the gel running buffer, 6% SDS-containing polyacrylamide gels, and the gel assembly were equilibrated to 4°C before running the samples. The samples were mixed with SDS containing gel-loading buffer and were not heated. The temperature of the gels was maintained below 10°C during electrophoresis by immersing the gel tanks in ice. Following LT-PAGE, the gels were transferred, and the blots were probed with anti-eNOS antibody and the corresponding secondary antibody. The intensity of the immunoreactive bands was quantified by ImageQuant TL software.

Electrophoretic mobility shift assay. NF- κB DNA binding activity was quantified by electrophoretic mobility shift assay (EMSA). Nuclear protein extraction from frozen myocardium, the EMSA protocol, autoradiography, and densitometry were all performed as previously described (14). ³²P-labeled consensus double-stranded oligonucleotides (sense, 5'-AGTTGAGGGGACTTCCAGGC-3') containing the NF- κB binding site were used as probes. Specificity of NF- κB DNA binding activity was confirmed in competition studies using cold consensus or mutant oligonucleotides.

Real-time PCR and mRNA quantitation. Total RNA isolation from LV tissue, cDNA synthesis, and quantitative real-time PCR were performed as previously described (14). mRNA transcripts for atrial natriuretic factor (ANF), tumor necrosis factor- α (TNF- α), and interleukin (IL)-1 β were determined and normalized to glyceraldehyde-3-phosphate dehydrogenase expression using primer pairs previously described (14).

Apoptosis quantitation. Myocardial apoptosis was assessed by using the DeadEnd Fluorometric terminal deoxytransferase-mediated dUTP nick-end labeling (TUNEL) assay kit from Promega, which catalytically incorporates fluorescein-12-dUTP at the 3'-ends of fragmented DNA in apoptotic cells using recombinant terminal deoxynucleotidyl transferase (rTdT). Deparaffinized and rehydrated tissue sections were treated with Proteinase K ($20 \mu\text{g/ml}$) for 15 min at 37°C and then fixed with 4% methanol-free formaldehyde solution in PBS.




Fig. 1. Chronic acrolein exposure depresses left ventricular (LV) function. **A**: M-mode echocardiograms from two mice, one acrolein-fed and the other vehicle-fed (control). AW and PW, anterior and posterior wall, respectively; ESD and EDD, end-systolic and end-diastolic diameter, respectively. **B**: LV pressure-volume loops and the corresponding end-systolic pressure-volume relations in representative control and acrolein-fed mice. E_{es} , end-systolic elastance.

All subsequent steps were performed following the manufacturer's instructions. All sections were counterstained with DAPI to label nuclei. Cardiomyocytes were identified by staining with anti-troponin I antibody (Santa Cruz Biotechnology) followed by Alexa Fluor 555-conjugated secondary antibody (Invitrogen). TUNEL-positive nuclei (cyan staining) were visualized directly by confocal microscopy (Zeiss LSM510) with nuclear staining confirmed by z -axis sections. Images were taken with a $\times 63$ objective lens at six different locations in each tissue section, and nine sections per heart were evaluated to determine the overall apoptotic rate (total 54 fields/heart). DNase (10 U/ml)-treated sections were used as positive controls. Sections without rTdT treatment were considered as negative controls.

Statistical analysis. Continuous variables are presented as means \pm SD. Two-group comparisons were performed using an unpaired t -test. A P value <0.05 was considered significant.

RESULTS

Chronic acrolein consumption induces LV remodeling and dysfunction. Mice gavage-fed acrolein at $1 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ for 48 days displayed no overt abnormalities or distress and no significant mortality. Body weight was similar between vehicle-fed and acrolein-fed animals after 48 days (control 26.1 ± 2.3 g; acrolein 26.0 ± 2.0 g). Baseline echocardiographic variables (before the start of feeding) were similar between the two groups. M-mode echocardiographic images obtained after the 48-day feeding period are shown in Fig. 1A. The acrolein-exposed mouse exhibited increased LV size and decreased FS compared with control. Group echocardiographic data (Table 1) indicate that acrolein exposure induced LV dilatation (increased EDD and ESD), LV systolic dysfunction (reduced FS and V_{cf}), and wall thinning (decreased RWT) consistent with a phenotype of DCM. While these changes were not severe (generally between ~ 8 and 20% change), they were highly consistent and statistically significant. To evaluate LV function more precisely, P-V analysis was performed. Figure

1B shows representative P-V loops from control and acrolein-exposed mice during IVC occlusion, together with the corresponding ESPVRs. Consistent with the echocardiographic results, acrolein exposure induced LV dilatation with increased end-diastolic volume and end-systolic volume and depressed LV systolic function as indicated by the smaller E_{es} . Group data (Table 2) demonstrated consistent LV enlargement and more profound reductions in systolic function with diminished dP/dt_{max} , maximal power, E_{es} , and stroke work. Also evident was impairment of LV relaxation with decreased dP/dt_{min} and increased tau. Hence, chronic acrolein exposure induced pathological remodeling and LV dysfunction.

Chronic acrolein exposure generates myocardial oxidative stress and protein-acrolein adducts. α,β -Unsaturated aldehydes are products of lipid-peroxidation and as such are sensitive markers of oxidative stress (8, 37, 40). Moreover, reac-

Table 1. Echocardiography in control and acrolein-exposed mice

	Control ($n = 16$)	Acrolein ($n = 14$)	P Value
HR, beats/min	469 ± 60	481 ± 62	0.585
LVEDD, mm	3.7 ± 0.1	$4.0 \pm 0.2^*$	<0.001
LVESD, mm	2.1 ± 0.2	$2.5 \pm 0.2^*$	<0.001
FS, %	43 ± 4	$36 \pm 3^*$	<0.001
ET, ms	51 ± 4	52 ± 7	0.771
V_{cf} , circ/s	8.5 ± 1.1	$7.1 \pm 1.0^*$	0.0022
AWT, mm	0.78 ± 0.04	0.74 ± 0.06	0.060
PWT, mm	0.79 ± 0.03	$0.76 \pm 0.03^*$	0.0089
RWT	0.42 ± 0.02	$0.38 \pm 0.02^*$	<0.001

Values are means \pm SD; n , no. of mice. HR, heart rate; LV, left ventricular; EDD, end-diastolic diameter; ESD, end-systolic diameter; FS, fractional shortening; ET, ejection time; V_{cf} , velocity of circumferential fiber shortening; AWT and PWT, anterior and posterior wall thickness at end-diastole, respectively; RWT, relative wall thickness. *Statistical significance.

Table 2. Pressure-volume parameters in control and acrolein-exposed mice

	Control (n = 8)	Acrolein (n = 8)	P Value
HR, beats/min	501 ± 63	451 ± 45	0.073
LVEDV, μ l	17 ± 5	36 ± 8*	<0.001
LVESV, μ l	8 ± 2	29 ± 7*	<0.001
LVSP, mmHg	94 ± 9	80 ± 14*	0.028
LVEDP, mmHg	7 ± 3	11 ± 6	0.134
SW, mmHg· μ l	601 ± 214	378 ± 153*	0.025
dP/dt _{max} , mmHg/s	7,016 ± 1,757	4,697 ± 1,498*	0.010
Maximal power, mW	3.45 ± 1.61	1.98 ± 0.81*	0.029
E _{es} , mmHg/ μ l	4.93 ± 1.16	3.35 ± 0.98*	0.049
dP/dt _{min} , mmHg/s	-8,002 ± 1,995	-5,291 ± 1,957*	0.013
Tau, ms	10.4 ± 2.2	15.4 ± 4.3*	0.0094

Values are means ± SD; n, no. of mice. EDV, end-diastolic volume; ESV, end-systolic volume; PSP, peak systolic pressure; EDP, end-diastolic pressure; SW, stroke work; dP/dt_{max} and dP/dt_{min}, maximal and minimal rate of change in LV pressure, respectively; E_{es}, end-systolic elastance; tau, time constant of LV relaxation. *Statistical significance.

tive aldehydes can induce cellular toxicity by adducting with cysteine, histidine, and lysine residues on proteins (9, 37). To index oxidative stress in the hearts of control and acrolein-exposed mice, we measured protein-HNE adducts. The abundance of protein-HNE adducts in total heart homogenates did not change in acrolein-exposed mice (data not shown). However, examination of the membrane fraction of the cardiac homogenates revealed robust augmentation of protein-HNE adducts as assessed by dot blot and Western blotting (Fig. 2A), indicating membrane-localized oxidative stress. We next determined whether acrolein-exposed mice exhibited greater formation of acrolein-protein adducts in serum and heart tissue. Hearts harvested from mice chronically fed acrolein did not exhibit appreciable increases in the abundance of protein-acrolein adducts over control (data not shown). Because these results were not striking, we further examined the abundance of plasma and myocardial acrolein adducts 1 and 24 h after a single oral dose. Plasma protein-acrolein adducts (~150 kDa) increased markedly at both time points with the highest levels seen at 1 h (Fig. 2B), suggesting that ingested acrolein reaches the blood. Myocardial protein-acrolein adducts, involving proteins of varying molecular weight, were more modestly increased at 1 h but returned to baseline by 24 h (Fig. 2C), approximating the adduct levels observed in the hearts from chronically fed mice. These results suggest that, following oral exposure, sufficient acrolein translocates via the circulation to the heart to modify proteins. However, these adducts accumulate transiently and are then metabolically removed or degraded. Presumably, adduct formation is less pronounced after chronic exposure because of the metabolic disposition of extant tissue adducts.

Chronic acrolein exposure disrupts myocardial eNOS function and induces nitrate stress. We next examined whether acrolein disrupts eNOS function and promotes nitrate stress in the heart. As shown in Fig. 3A, a single oral dose of acrolein (5 mg/kg) profoundly suppressed eNOS phosphorylation at Ser¹¹⁷⁷, an indicator of eNOS activation (4), without affecting overall eNOS abundance in the heart. In contrast, chronic exposure to acrolein significantly diminished eNOS dimers and increased relative levels of eNOS monomers (Fig. 3B), suggestive of eNOS uncoupling (36). Uncoupling of eNOS would

be expected to promote the generation of reactive oxygen species (ROS) and peroxynitrite (36, 39). Indeed, hearts from mice chronically fed acrolein exhibited significantly greater staining for protein nitrotyrosine, an index of peroxynitrite generation (Fig. 3C). These results indicate that chronic acrolein exposure disrupted and uncoupled eNOS and induced nitrate stress in the heart.

Chronic acrolein exposure induces myocyte hypertrophy and apoptosis. As shown in Fig. 4A, histological evaluation of acrolein-exposed hearts revealed myocyte hypertrophy, with a twofold increase in myocyte cross-sectional area compared with control hearts. There was no substantial difference in interstitial fibrosis (data not shown). Gene expression of the hypertrophic marker ANF was similarly augmented over twofold in acrolein-exposed hearts compared with control (Fig. 4B). Despite these observations, gravimetric analysis of the LV and whole heart did not reveal differences in LV or whole heart weight (normalized to body wt) between the groups. This suggested that the increase in myocyte size was offset by myocyte loss. Indeed, as shown in Fig. 5, we observed a greater frequency of TUNEL-positive nuclei in the hearts of acrolein-exposed mice; these were primarily in cardiomyocytes. Quantitation of the apoptotic rate revealed a more than sixfold increase in TUNEL-positive nuclei compared with control. Hence, chronic oral acrolein exposure induced prohypertrophic and proapoptotic effects in the heart.

Chronic acrolein exposure promotes myocardial inflammation. Reactive aldehydes are known to promote inflammation (30, 38), which is a hallmark of chronic HF (14, 22). NF- κ B is a central transcriptional regulator of proinflammatory mediators such as TNF- α and IL-1 β . To evaluate NF- κ B activation, we performed EMSA using pooled cardiac tissue from animals with either acute (24 h after single dose of 1 mg/kg) or chronic oral acrolein exposure, along with appropriate controls. As seen in Fig. 6A, heart tissue from chronically exposed (but not acutely exposed) mice demonstrated robust activation of NF- κ B. Figure 6B depicts activated NF- κ B p65 subunit immunostaining and quantitation of nuclear immunoreactivity from control and acrolein-exposed hearts. Consistent with the DNA binding studies, the hearts from acrolein-exposed mice exhibited a robust (~5-fold) increase in the nuclear localization of p65. Additionally, protein levels of I κ B α (which binds cytoplasmic NF- κ B thereby preventing its nuclear translocation) were decreased in hearts from acrolein-exposed mice (Fig. 6C). Moreover, in parallel with NF- κ B activation, hearts from acrolein-exposed mice also exhibited significant (~2-fold) upregulation of TNF- α and IL-1 β mRNA expression compared with controls (Fig. 6D), which is indicative of sustained inflammation.

DISCUSSION

In this study, we demonstrate for the first time that oral exposure to acrolein, a prototypical α,β -unsaturated aldehyde pollutant, at concentrations within the estimated range of human total unsaturated aldehyde exposure, induces a phenotype of DCM in the mouse. Specifically, 48 days of acrolein exposure induced: 1) LV dilatation, wall thinning, impairment of LV relaxation, and depressed contractility; 2) chronic membrane-localized oxidative stress associated with varying degrees of systemic and myocardial protein-acrolein adduct for-




Fig. 2. Chronic acrolein exposure induces cardiac oxidative stress and protein modification. *A*: representative dot blot (DB) and Western blot (WB) performed on the membrane fractions of cardiac homogenates derived from acrolein (ACR)-fed and vehicle-fed [control (C)] mice and corresponding WB densitometry. *B* and *C*: WB and densitometry for protein-acrolein adducts in plasma (*B*) and myocardium (*C*) from mice fed a single dose of acrolein (1 mg/kg) or water 1 and 24 h after exposure. Augmented protein bands at different molecular weights are indicated by the arrows. M, molecular weight markers. **P* < 0.05 vs. control; *n* = 4 mice/group.

mation; 3) diminished levels and uncoupling of eNOS with associated myocardial nitrate stress; 4) myocyte hypertrophy and apoptosis without fibrosis; and 5) myocardial inflammation with activation of NF-κB and upregulation of TNF-α and IL-1β. The features of oxidant stress, hypertrophy, apoptosis,

and inflammation are pathological hallmarks of the failing heart. Taken together, the results suggest that analogous environmental exposure to acrolein in humans can contribute to the development of DCM and/or exacerbate pathological remodeling in humans with preexisting disease. Our results further




Fig. 3. Acrolein increases myocardial nitrosative stress. *A*: WB and quantitation for phospho (P)-endothelial nitric oxide synthase (eNOS)-Ser¹¹⁷⁷ and total eNOS performed on total cardiac homogenates from mice 24 h after a single oral dose of acrolein (5 mg/kg) or vehicle ($n = 5/\text{group}$). Hela, Hela cell lysate. *B*: WB and densitometry for eNOS dimer and monomer performed on cardiac homogenates derived from mice chronically fed acrolein or water for 48 days ($n = 4-5/\text{group}$). *C*: immunofluorescent stains for protein-nitrotyrosine (NT, red) with 4',6-diamidino-2-phenylindole (DAPI) costain for nuclei (blue) in hearts harvested from acrolein-fed and control-fed mice as in *B*, along with fluorescence quantitation (control, $n = 3$; acrolein, $n = 5$). Peroxynitrite-treated sections were used as a positive control. * $P < 0.05$ vs. control.

suggest the possibility that acrolein (and potentially other unsaturated aldehydes) can serve as a dietary xenobiotic mediator and/or modulator of cardiomyopathy.

Epidemiological data indicate that pollution exposure increases cardiovascular morbidity and mortality (3, 5, 28), with the most robust associations related to ischemic heart disease, dysrhythmias, HF, and cardiac arrest (28). A recent study of elderly survivors of acute myocardial infarction revealed that air pollution exposure increased both the risk of mortality and the risk for new-onset HF within four to five years (44). Because the development of new-onset HF following infarction is related to the progression of underlying LV remodeling over time (16), this suggests that exposure to one or a variety of constituent pollutants can exacerbate underlying structural remodeling. One proposed mechanism of pollution-related cardiovascular risk is the physical translocation of soluble pollutant constituents into the heart and vasculature via the circula-

tion (5). However, little is known about the specific pathophysiological responses to individual constituents of source mixtures of environmental pollutants.

Acrolein is a ubiquitous aldehyde pollutant of considerable importance to public health (7). High levels of acrolein have been detected in several foods (ranging from 10 to 600 $\mu\text{g}/\text{kg}$), cigarette smoke (10–140 $\mu\text{g}/\text{cigarette}$), water samples, heated oils, automobile exhaust, coal, and industrial waste (10, 11, 42). Volatile aldehydes such as acrolein are important constituents of the vapor phase of urban air pollution and diesel exhaust and are considered hazardous air pollutants by the EPA (7, 29). Given the large number of environmental sources of acrolein and its potential for long-term toxicity, we sought to determine the effects of chronic acrolein exposure on the heart. In this study, we chose to examine the effects of ingested (as opposed to inhaled) acrolein because, in humans, even in smokers, the highest level of acrolein exposure is through food




Fig. 4. Chronic acrolein exposure induces myocyte hypertrophy. *A*: representative histomicrographs of heart tissue from control and acrolein-fed mice demonstrating myocytes in cross section and corresponding quantitation of myocyte cross-sectional area. Also shown is the expression of the atrial natriuretic factor (ANF) gene in the heart by quantitative real-time PCR (*B*) and tissue gravimetric data (*C*) from the same experimental groups. GAPDH, glyceraldehyde-3-phosphate dehydrogenase; NS, not significant. * $P < 0.05$ vs. control.

substances (42). Nevertheless, our findings that acrolein translocates to plasma and heart tissue following exposure (evidenced by the formation of adducts) and induces chronic changes in cardiac gene expression suggest the possibility that analogous exposure to acrolein in ambient air may, via physical transport in blood, produce similar responses. This is consistent with the high cardiovascular toxicity associated with the aldehyde-containing components of air pollution, diesel exhaust, and cigarette smoke (3, 18).

We have previously estimated the maximal human acrolein exposure from food and water to be $0.1 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ (with an additional $0.1 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ from cigarette smoking) and the maximal human unsaturated aldehyde consumption to be $5 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ (42). In the current study, we evaluated the chronic effects of $1 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ acrolein, a dose fivefold lower than in our acute studies (42), representing a level 5- to 10-fold greater than maximal human acrolein consumption but only ~20% of total estimated unsaturated aldehyde intake. We chose this intermediate dose given that the sensitivity to acrolein varies among experimental animals; compared with rabbits (LD_{50} 7 mg/kg), mice are relatively less sensitive (LD_{50} 40 mg/kg acrolein) (10). Human sensitivity to acrolein, however, has not been assessed. Whether different acrolein dosing regimens (e.g., lower but more frequent doses) would influence the results differently should be explored in future investigations.

Our results establish that environmental exposure to acrolein, via the oral route, induces a state of inflammation and oxidant stress in the heart, along with LV systolic dysfunction, myocyte hypertrophy, and apoptosis, all consistent with xenobiotic-mediated DCM. These effects are consistent with the known prooxidant and proinflammatory effects of α,β -unsaturated

aldehydes, which have been shown to activate inflammatory genes and signaling (including NF- κ B) (27, 30, 38) and promote monocyte adhesion to endothelial cells (13). Similarly, in our study, acrolein-exposed hearts exhibited NF- κ B activation, proinflammatory cytokine (TNF- α , IL-1 β) gene expression, and oxidative and nitritative stress. Furthermore, in our prior study (19), we have shown that oxidant stress is required for acrolein-induced contractile dysfunction, since such effects were prevented by the antioxidant *N*-acetylcysteine. These findings are of significance, since chronic inflammation and oxidant stress are hallmarks of HF and considered to be important mediators of pathological LV remodeling (12, 16, 22). Plasma TNF- α is an independent predictor of patient mortality in HF (6), and, in experimental models, TNF- α induces many aspects of HF, including contractile depression, hypertrophy, apoptosis, matrix metalloproteinase activation, and oxidative stress (14, 22). Similarly, systemic oxidant stress in human HF correlates with the degree of ventricular dysfunction (21). Signaling related to ROS has been strongly implicated in the induction of pathological cardiac hypertrophy, and ROS can also mediate apoptosis, alter calcium channels and calcium flux, and reduce myofilament calcium sensitivity (12, 20, 32). Moreover, *in vivo* treatment with ROS scavengers improves pathological LV remodeling (17).

The stimulus for inflammatory cytokines and oxidant stress in HF is generally thought to reflect a response to injury, hemodynamic abnormalities, neurohormonal activation, and alterations in tissue perfusion. Our data suggest that environmental triggers may also contribute to this process and thereby exacerbate the course and progression of HF, and that those with preexisting LV dysfunction may be especially sensitive to environmental acrolein exposure. Interestingly, epidemiologi-




Fig. 5. Confocal microscopic images of terminal deoxytransferase-mediated dUTP nick-end labeling (TUNEL) staining in hearts from mice chronically fed acrolein or vehicle (control) and quantitation of TUNEL-positive nuclei. Myocytes were stained with anti-troponin I (red), and nuclei were stained with DAPI (blue). TUNEL-positive nuclei appear green-cyan on the overlaid images from the acrolein-exposed heart as shown in the magnified (zoom) image. Arrows denote a TUNEL-positive nucleus. Scale bar: standard magnification 20 μm ; zoomed magnification 5 μm . $n = 5/\text{group}$.

cal studies have established that human subjects with HF are more vulnerable to the adverse cardiovascular effects of pollution exposure (5). Moreover, similar to our results obtained with acrolein exposure, environmental carbon monoxide also induces pathological remodeling in hearts of normal rats (1), supporting the idea that pollutant exposure could also lead to adverse changes in the heart in the absence of underlying cardiomyopathy.

One underlying mechanism for acrolein-mediated cardiac remodeling may be related to the induced abnormalities in eNOS function. Alterations in eNOS coupling and NO synthesis can contribute substantially to pathological cardiac remodeling (24, 36, 39). When electron transfer from its reductase to oxidase domains is normally coupled, eNOS is generally cardioprotective and antihypertrophic (39). However, during pathological hypertrophy and HF, both eNOS downregulation and uncoupling can occur, thereby augmenting superoxide generation, diminishing NO bioavailability, and increasing peroxynitrite formation (12, 24, 36, 39). In our study, acute acrolein exposure suppressed eNOS activation, whereas chronic acrolein exposure decreased overall eNOS abundance and reduced the eNOS dimer-to-monomer ratio, consistent with eNOS uncoupling. The biological relevance of these changes was demonstrated by the approximately twofold increase in protein-nitrotyrosine levels in the heart, indicative of increased peroxynitrite generation. Hence, disruption of eNOS function may be in part responsible for increased free radicals and oxidant stress induced by acrolein.

The observed cardiomyopathic phenotype may have resulted from both direct and indirect effects of acrolein. We have demonstrated that oral acrolein exposure induces protein-acrolein adducts in both plasma and myocardium with adduct abundance decreasing in a time-dependent manner following exposure. This suggests that consumed acrolein physically circulates to remote sites such as the heart to directly disrupt protein function, thereby secondarily inducing cardiac injury and inflammation. In our previous studies, we demonstrated that acrolein primarily modifies sarcomeric, cytoskeletal, and mitochondrial proteins in the context of acute exposure (19, 42). The time dependence of adduct levels in the current study suggests ongoing metabolic disposition and turnover of protein-acrolein adducts both systemically and in the heart. This is consistent with prior studies that have demonstrated lability of aldehyde-adducted proteins and degradation by the proteasome and lysosomes in minutes to hours (23, 26). Long-term exposure and/or reduced metabolic capacity for aldehyde detoxification may therefore enhance the adverse effects of acrolein. In this regard, we have previously shown that aldose reductase, the main aldehyde-reducing enzyme in the heart, is significantly downregulated in HF (33). Hence, the cardiotoxic effects of environmental acrolein may be heightened in subjects with preexisting HF.

In summary, we have shown that long-term environmental exposure to acrolein, at an amount within the range of human unsaturated aldehyde intake, induces DCM in the mouse. Primary features included the induction of myocardial inflam-




Fig. 6. Chronic acrolein exposure induces inflammation in the heart. **A**: EMSA to determine nuclear factor (NF)-κB DNA binding activity of pooled myocardial nuclear protein extracts from control and acrolein-fed mice. Acute control and acrolein-fed mice were given a single dose of 1 mg/kg vehicle or acrolein, and tissue was harvested at 24 h. Chronic control and acrolein-fed mice were administered daily vehicle or acrolein (1 mg/kg) for 48 days, and tissue was harvested 24 h after the final dose. NF-κB DNA binding is indicated by the arrowhead. The circle indicates unbound oligonucleotide probe. **B**: representative immunohistochemical stains for the activated p65 subunit of NF-κB in hearts from control mice and mice chronically fed acrolein, together with quantitation of staining intensity by image analysis. Note the nuclear localization of p65 in the acrolein-exposed mouse heart. **C**: WB and densitometry for inhibitor of κBα (IκBα) in hearts from control mice and mice chronically fed acrolein as in **A**. **D**: myocardial gene expression of tumor necrosis factor (TNF) and interleukin (IL)-1β by real-time PCR in the same hearts as in **B**. **P* < 0.05 vs. control.

mation and oxidative/nitrative stress, which may represent responses to the formation of detrimental acrolein-protein adducts in the heart, together with myocyte hypertrophy and apoptosis. These results suggest that human exposure to acro-

lein can have analogous deleterious effects, especially in those with preexisting structural heart disease and/or reduced capacity for aldehyde detoxification. Moreover, our findings raise consideration of an underrecognized environmental basis for

idiopathic DCM related to aldehyde constituents of natural food and the pollutant mix.

GRANTS

This work was supported by a Veterans Affairs Merit Award (S. D. Prabhu); National Institutes of Health Grants ES-11860 (A. Bhatnagar, S. D. Prabhu), HL-78825 (S. D. Prabhu, A. Bhatnagar), HL-99014 (S. D. Prabhu), HL-95593 (S. Srivastava), and RR-024489 (A. Bhatnagar, S. Srivastava, S. D. Prabhu); and an American Heart Association Scientist Development Grant (T. Hamid).

DISCLOSURES

There are no conflicts of interest to disclose.

REFERENCES

- Andre L, Boissiere J, Reboul C, Perrier R, Zalvidea S, Meyer G, Thireau J, Tanguy S, Bideaux P, Hayot M, Boucher F, Obert P, Cazorla O, Richard S. Carbon monoxide pollution promotes cardiac remodeling and ventricular arrhythmia in healthy rats. *Am J Respir Crit Care Med* 181: 587–595, 2010.
- Assembly of Life Sciences (U.S.) Committee on Aldehydes. *Formaldehyde and Other Aldehydes*. Washington, DC: Natl Acad, 1981, p. ix, 340 p.
- Bhatnagar A. Environmental cardiology: studying mechanistic links between pollution and heart disease. *Circ Res* 99: 692–705, 2006.
- Boo YC, Kim HJ, Song H, Fulton D, Sessa W, Jo H. Coordinated regulation of endothelial nitric oxide synthase activity by phosphorylation and subcellular localization. *Free Radic Biol Med* 41: 144–153, 2006.
- Brook RD, Rajagopalan S, Pope CA, 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC Jr, Whitsel L, Kaufman JD. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 121: 2331–2378, 2010.
- Deswal A, Petersen NJ, Feldman AM, Young JB, White BG, Mann DL. Cytokines and cytokine receptors in advanced heart failure: an analysis of the cytokine database from the Vesnarinone trial (VEST). *Circulation* 103: 2055–2059, 2001.
- DeWoskin RS and United States Environmental Protection Agency. *Toxicological Review of Acrolein (CAS No. 107-02-8) in Support of Summary Information on the Integrated Risk Information System (IRIS)*. Washington, DC: U.S. Environmental Protection Agency, 2003.
- Eaton P, Li JM, Hearse DJ, Shattock MJ. Formation of 4-hydroxy-2-nonenal-modified proteins in ischemic rat heart. *Am J Physiol Heart Circ Physiol* 276: H935–H943, 1999.
- Esterbauer H, Schaur RJ, Zollner H. Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free Radic Biol Med* 11: 81–128, 1991.
- Feron VJ, Til HP, de Vrijer F, Woutersen RA, Cassee FR, van Bladeren PJ. Aldehydes: occurrence, carcinogenic potential, mechanism of action and risk assessment. *Mutat Res* 259: 363–385, 1991.
- Ghilarducci DP, Tjeerdema RS. Fate and effects of acrolein. *Rev Environ Contam Toxicol* 144: 95–146, 1995.
- Giordano FJ. Oxygen, oxidative stress, hypoxia, and heart failure. *J Clin Invest* 115: 500–508, 2005.
- Go YM, Halvey PJ, Hansen JM, Reed M, Pohl J, Jones DP. Reactive aldehyde modification of thioredoxin-1 activates early steps of inflammation and cell adhesion. *Am J Pathol* 171: 1670–1681, 2007.
- Hamid T, Gu Y, Ortines RV, Bhattacharya C, Wang G, Xuan YT, Prabhu SD. Divergent tumor necrosis factor receptor-related remodeling responses in heart failure: role of nuclear factor-kappaB and inflammatory activation. *Circulation* 119: 1386–1397, 2009.
- He J, Ogden LG, Bazzano LA, Vupputuri S, Loria C, Whelton PK. Risk factors for congestive heart failure in US men and women: NHANES I epidemiologic follow-up study. *Arch Intern Med* 161: 996–1002, 2001.
- Jessup M, Brozena S. Heart failure. *N Engl J Med* 348: 2007–2018, 2003.
- Kinugawa S, Tsutsui H, Hayashidani S, Ide T, Suematsu N, Satoh S, Utsumi H, Takeshita A. Treatment with dimethylthiourea prevents left ventricular remodeling and failure after experimental myocardial infarction in mice: role of oxidative stress. *Circ Res* 87: 392–398, 2000.
- Logue JM, Small MJ, Stern D, Maranche J, Robinson AL. Spatial variation in ambient air toxics concentrations and health risks between industrial-influenced, urban, and rural sites. *J Air Waste Manag Assoc* 60: 271–286, 2010.
- Luo J, Hill BG, Gu Y, Cai J, Srivastava S, Bhatnagar A, Prabhu SD. Mechanisms of acrolein-induced myocardial dysfunction: implications for environmental and endogenous aldehyde exposure. *Am J Physiol Heart Circ Physiol* 293: H3673–H3684, 2007.
- Luo J, Xuan YT, Gu Y, Prabhu SD. Prolonged oxidative stress inverts the cardiac force-frequency relation: role of altered calcium handling and myofilament calcium responsiveness. *J Mol Cell Cardiol* 40: 64–75, 2006.
- Mak S, Lehotay DC, Yazdanpanah M, Azevedo ER, Liu PP, Newton GE. Unsaturated aldehydes including 4-OH-nonenal are elevated in patients with congestive heart failure. *J Card Fail* 6: 108–114, 2000.
- Mann DL. Inflammatory mediators and the failing heart: past, present, and the foreseeable future. *Circ Res* 91: 988–998, 2002.
- Marques C, Pereira P, Taylor A, Liang JN, Reddy VN, Szweda LI, Shang F. Ubiquitin-dependent lysosomal degradation of the HNE-modified proteins in lens epithelial cells. *FASEB J* 18: 1424–1426, 2004.
- Moens AL, Takimoto E, Tocchetti CG, Chakir K, Bedja D, Cormaci G, Ketner EA, Majmudar M, Gabrielson K, Halushka MK, Mitchell JB, Biswal S, Channon KM, Wolin MS, Alp NJ, Paolucci N, Champion HC, Kass DA. Reversal of cardiac hypertrophy and fibrosis from pressure overload by tetrahydrobiopterin: efficacy of recoupling nitric oxide synthase as a therapeutic strategy. *Circulation* 117: 2626–2636, 2008.
- Nath RG, Chung FL. Detection of exocyclic 1,N2-propanodeoxyguanosine adducts as common DNA lesions in rodents and humans. *Proc Natl Acad Sci USA* 91: 7491–7495, 1994.
- Okada K, Wangpoengtrakul C, Osawa T, Toyokuni S, Tanaka K, Uchida K. 4-Hydroxy-2-nonenal-mediated impairment of intracellular proteolysis during oxidative stress. Identification of proteasomes as target molecules. *J Biol Chem* 274: 23787–23793, 1999.
- Parola M, Bellomo G, Robino G, Barrera G, Dianzani MU. 4-Hydroxynonenal as a biological signal: molecular basis and pathophysiological implications. *Antioxid Redox Signal* 1: 255–284, 1999.
- Pope CA 3rd, Burnett RT, Thurston GD, Thun MJ, Calle EE, Krewski D, Godleski JJ. Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease. *Circulation* 109: 71–77, 2004.
- Pratt GC, Palmer K, Wu CY, Ollaei F, Hollerbach C, Fenske MJ. An assessment of air toxics in Minnesota. *Environ Health Perspect* 108: 815–825, 2000.
- Rahman I, Marwick J, Kirkham P. Redox modulation of chromatin remodeling: impact on histone acetylation and deacetylation, NF-kappaB and pro-inflammatory gene expression. *Biochem Pharmacol* 68: 1255–1267, 2004.
- Ramana KV, Chandra D, Srivastava S, Bhatnagar A, Aggarwal BB, Srivastava SK. Aldose reductase mediates mitogenic signaling in vascular smooth muscle cells. *J Biol Chem* 277: 32063–32070, 2002.
- Sawyer DB, Siwik DA, Xiao L, Pimentel DR, Singh K, Colucci WS. Role of oxidative stress in myocardial hypertrophy and failure. *J Mol Cell Cardiol* 34: 379–388, 2002.
- Srivastava S, Chandrasekar B, Bhatnagar A, Prabhu SD. Lipid peroxidation-derived aldehydes and oxidative stress in the failing heart: role of aldose reductase. *Am J Physiol Heart Circ Physiol* 283: H2612–H2619, 2002.
- Srivastava S, Chandrasekar B, Gu Y, Luo J, Hamid T, Hill BG, Prabhu SD. Downregulation of CuZn-superoxide dismutase contributes to beta-adrenergic receptor-mediated oxidative stress in the heart. *Cardiovasc Res* 74: 445–455, 2007.
- Syed F, Diwan A, Hahn HS. Murine echocardiography: a practical approach for phenotyping genetically manipulated and surgically modeled mice. *J Am Soc Echocardiogr* 18: 982–990, 2005.
- Takimoto E, Champion HC, Li M, Ren S, Rodriguez ER, Tavazzi B, Lazzarino G, Paolucci N, Gabrielson KL, Wang Y, Kass DA. Oxidant stress from nitric oxide synthase-3 uncoupling stimulates cardiac pathologic remodeling from chronic pressure load. *J Clin Invest* 115: 1221–1231, 2005.
- Uchida K, Kanematsu M, Sakai K, Matsuda T, Hattori N, Mizuno Y, Suzuki D, Miyata T, Noguchi N, Niki E, Osawa T. Protein-bound acrolein: potential markers for oxidative stress. *Proc Natl Acad Sci USA* 95: 4882–4887, 1998.
- Uchida K, Shiraiishi M, Naito Y, Torii Y, Nakamura Y, Osawa T. Activation of stress signaling pathways by the end product of lipid

- peroxidation 4-hydroxy-2-nonenal is a potential inducer of intracellular peroxide production. *J Biol Chem* 274: 2234–2242, 1999.
39. **Umar S, van der Laarse A.** Nitric oxide and nitric oxide synthase isoforms in the normal, hypertrophic, and failing heart. *Mol Cell Biochem* 333: 191–201, 2010.
 40. **Vasilyev N, Williams T, Brennan ML, Unzek S, Zhou X, Heinecke JW, Spitz DR, Topol EJ, Hazen SL, Penn MS.** Myeloperoxidase-generated oxidants modulate left ventricular remodeling but not infarct size after myocardial infarction. *Circulation* 112: 2812–2820, 2005.
 41. **Wang G, Hamid T, Keith RJ, Zhou G, Partridge CR, Xiang X, Kingery JR, Lewis RK, Li Q, Rokosh DG, Ford R, Spinale FG, Riggs DW, Srivastava S, Bhatnagar A, Bolli R, Prabhu SD.** Cardioprotective and antiapoptotic effects of heme oxygenase-1 in the failing heart. *Circulation* 121: 1912–1925, 2010.
 42. **Wang GW, Guo Y, Vondriska TM, Zhang J, Zhang S, Tsai LL, Zong NC, Bolli R, Bhatnagar A, Prabhu SD.** Acrolein consumption exacerbates myocardial ischemic injury and blocks nitric oxide-induced PKCepsilon signaling and cardioprotection. *J Mol Cell Cardiol* 44: 1016–1022, 2008.
 43. **Yang YM, Huang A, Kaley G, Sun D.** eNOS uncoupling and endothelial dysfunction in aged vessels. *Am J Physiol Heart Circ Physiol* 297: H1829–H1836, 2009.
 44. **Zanobetti A, Schwartz J.** Particulate air pollution, progression, and survival after myocardial infarction. *Environ Health Perspect* 115: 769–775, 2007.



Assessment of protein and amino acid concentrations and labeling adequacy of commercial vegetarian diets formulated for dogs and cats

Kayo Kanakubo, BVSc; Andrea J. Fascetti, VMD, PhD; Jennifer A. Larsen, DVM, PhD

Objective—To determine measured crude protein (CP) and amino acid (AA) concentrations and assess labeling adequacy of vegetarian diets formulated for dogs and cats.

Design—Cross-sectional study.

Sample—13 dry and 11 canned vegetarian diets for dogs and cats.

Procedures—Concentrations of CP and AAs were determined for each diet. Values were compared with the Association of American Feed Control Officials (AAFCO) Dog and Cat Food Nutrient Profiles. Product labels were assessed for compliance with AAFCO regulations.

Results—CP concentration (dry-matter basis) ranged from 19.2% to 40.3% (median, 29.8%). Minimum CP concentrations for the specified species and life stage were met by 23 diets; the remaining diet passed appropriate AAFCO feeding trials. Six diets did not meet all AA minimums, compared with the AAFCO nutrient profiles. Of these 6 diets, 1 was below AAFCO minimum requirements in 4 AAs (leucine, methionine, methionine-cystine, and taurine), 2 were below in 3 AAs (methionine, methionine-cystine, and taurine), 2 were below in 2 AAs (lysine and tryptophan), and 1 was below in 1 AA (tryptophan). Only 3 and 8 diets (with and without a statement of calorie content as a requirement, respectively) were compliant with all pet food label regulations established by the AAFCO.

Conclusion and Clinical Relevance—Most diets assessed in this study were not compliant with AAFCO labeling regulations, and there were concerns regarding adequacy of AA content. Manufacturers should ensure regulatory compliance and nutritional adequacy of all diets, and pets fed commercially available vegetarian diets should be monitored and assessed routinely. (*J Am Vet Med Assoc* 2015;247:385–392)

Popularity of vegetarian and vegan diets for humans has increased for ethical, ecological, and health reasons, and this influences pet food choices for some families.^{1,2} In addition, vegetarian diets are often used for veterinary patients with conditions such as hepatic encephalopathy, food allergies, and urate and cystine urolithiasis. However, for several reasons, vegetarian pet foods have been linked to concerns related to nutritional adequacy. Vegetarian protein sources are often poor sources of specific essential vitamins (vitamin D, vitamin A, niacin, and cobalamin), fatty acids (arachidonic acid, docosahexaenoic acid, and eicosapentaenoic acid), and minerals (calcium and potassium).³ In addition, plants are highly variable in protein concentration and provide incomplete AA profiles for meeting the needs of pets. Therefore, vegetarian diets must be appropriately formulated and balanced, including the use of proper supplementation with purified sources of essential AAs when indicated.

Adequate protein and AA intake is an important consideration for both dogs and cats. Cats are more lim-

From the Veterinary Medical Teaching Hospital (Kanakubo) and the Department of Molecular Biosciences (Fascetti, Larsen), School of Veterinary Medicine, University of California-Davis, Davis, CA 95616.

Supported by the Center for Companion Animal Health, School of Veterinary Medicine, University of California-Davis.

Address correspondence to Dr. Larsen (jalarsen@vmth.ucdavis.edu).

ABBREVIATIONS

AA	Amino acid
AAFCO	Association of American Feed Control Officials
CP	Crude protein
DM	Dry matter
ME	Metabolizable energy

ited than dogs in their ability to conserve nitrogen and AAs in the face of inadequate dietary intake.⁴ In addition, sulfur-containing AAs (methionine, cystine, and taurine) are found primarily in animal protein. Although it is not used for protein synthesis, taurine is a required dietary nutrient for cats and is important for several physiologic processes, including retinal function, cardiac function, reproduction, and growth.⁵ Taurine is considered conditionally essential for dogs because they have the metabolic capacity to synthesize it when adequate concentrations of sulfur-containing AA precursors (methionine and cysteine) are available, except for specific breeds⁶ and diseases associated with decreased taurine synthesis.⁷ Taurine deficiency has also been identified in dogs fed low-protein diets for extended periods or fed diets limited in sulfur-containing AAs.^{8,9}

Pet foods sold in the United States are regulated by both federal and state laws. Manufacturers are re-

sponsible for proper formulation and labeling of products to meet requirements set by the US FDA as well as those mandated by each state, many of which have adopted AAFCO model regulations for pet food.¹⁰ Information from the label is often used by pet owners and veterinarians to assess pet foods; therefore, accuracy and compliance with regulations are expected.¹¹ However, to our knowledge, there have been no studies conducted on the incidence of noncompliance of any category of pet foods with AAFCO model regulations, although there is evidence that some diets may provide CP at concentrations below the minimum guaranteed analysis value.¹²

The primary objective of the study reported here was to measure CP and AA concentrations in commercial vegetarian foods formulated for dogs and cats and to compare those values with minimum required concentrations for the intended species and life stage as established by the AAFCO. A secondary objective was to compare the information on pet food labels with required components as established by the AAFCO. We hypothesized that all diets would meet all nutritional and labeling requirements.

Materials and Methods

Sample—Commercially available over-the-counter diets (foods distributed directly to the public without veterinary oversight) consisting of dry and canned products for dogs and cats that were labeled or marketed as vegan or vegetarian and available during June and July 2014 were obtained from local pet stores and online sources. Similarly labeled or marketed veterinary therapeutic dry and canned diets for dogs were obtained from a local veterinary clinic,^a and 1 diet was donated by an employee of the University of California-Davis Veterinary Medical Teaching Hospital.

Procedures—Information from the labels was compared with 9 AAFCO labeling requirements¹³ (product and brand name, species specification, quantity statement, guaranteed analysis, ingredient statement, nutritional adequacy statement, feeding directions, name and address of manufacturer or distributor, and calorie content). The new labeling requirement for inclusion of the calorie content statement on all pet food labels was included in the AAFCO 2014 official publication.¹³ However, the AAFCO recommended in that publication that enforcement be delayed 18 months for new products in development and 3 years for existing products.¹⁴ Therefore, labels were assessed both including and excluding the statement of calorie content as a requirement. Information that was not provided on the label but was required for assessment was obtained from the product website or by contacting the manufacturer.

A sample of each diet was placed in a plastic bag, labeled with a number corresponding to the product, and submitted for analysis; all analytic laboratories were not aware of the commercial source for each sample submitted for analysis.

A sample of each of the canned diets was manually crushed within the plastic bag until a paste consistency was achieved, whereas dry diets were analyzed without any processing. Dry matter values were obtained

by drying representative samples of each diet (20 g of canned diets and 5 g of dry diets) to a constant weight in a vacuum oven at 95° to 100°C.

In addition, 100 g of each canned diet and 50 g of each dry diet were stored in individual containers and frozen at -80°C. These samples were placed into a freeze-drier for 7 days prior to analysis, and canned diets then were manually crushed into a powder to ensure homogeneity. Approximately 5 g of each freeze-dried diet was submitted to a reference laboratory^b for measurement of total nitrogen concentration via a combustion method.¹⁵ This method was not included in the methods cited by the AAFCO¹⁶; however, results of a comparison study¹⁷ with Kjeldahl analysis revealed that the combustion method had improved repeatability and reproducibility for SD ranges. Twenty diets were measured as single samples, and 4 diets were measured as duplicate samples in accordance with the laboratory's standard procedures. The laboratory's acceptable variance was 6.7%, and analytic variation for the 4 duplicate samples was 0.3%. Crude protein content was determined by use of the following equation: CP percentage = nitrogen percentage \times 6.25.

For AA analysis, all freeze-dried samples were ground until they could pass through a 2-mm screen (80 mesh). Approximately 10 mg of each ground sample was hydrolyzed in a vacuumed-sealed glass ampule with 2 mL of 6M HCl at 115°C for 24 hours. The hydrolysate was then dried with nitrogen gas, and the resulting residue was reconditioned with lithium hydroxide loading buffer. This solution was filtered by use of a 0.45- μ m polytetrafluorethylene syringe filter. The AA composition was determined in the filtrate by use of a norleucine internal standard with an automated high-performance liquid chromatography AA analyzer^c at the Amino Acid Laboratory at the University of California-Davis, with methods described elsewhere.¹⁸ Cystine and methionine concentrations were determined by use of performic acid oxidation with acid hydrolysis (hydrobromic acid method¹⁹), and tryptophan concentration was determined by use of a method described elsewhere.²⁰ All diets were measured as single samples. In addition to the internal standard used by the laboratory, a reference sample of purified casein was analyzed concurrently with each batch of sample diets; analytic variation was within 5%.

Measured CP and AA concentrations were compared with the minimum requirement in the AAFCO Dog and Cat Food Nutrient Profiles for the intended species and life stage.¹³ Diets formulated for both dogs and cats were compared with the AAFCO food nutrient profiles for cats. Concentrations of CP and AA were corrected for energy density if the diet contained > 4,000 or > 4,500 kcal/kg of DM for canine or feline diets, respectively.²¹ When assessing whether measured concentrations met the minimum values of the AAFCO food nutrient profiles, consideration was given to the allowed analytic variation for CP and lysine (AAFCO does not specify allowable variations for other AAs).¹⁶

Calorie content was obtained from the label or manufacturer; if calorie content was not provided or could not be obtained, it was calculated from the guaranteed analysis. For calculation of calorie content, measured CP and moisture concentrations were

used; modified Atwater values of 3.5 kcal/g for CP and nitrogen-free extract and 8.5 kcal/g for crude fat were used.¹³ Ash concentration was obtained from the label or manufacturer or were estimated by use of the mean value of the ash concentrations measured for diets.

Statistical analysis—A Shapiro-Wilk test was used to confirm data were nonparametric. Spreadsheet software^d was used to calculate descriptive statistics (median and range).

Results

Twenty-four diets were assessed, consisting of 13 dry diets (9 for dogs,^{e-m} 3 for cats,^{n-p} and 1 for both dogs and cats^q) and 11 canned diets (8 for dogs,^{r-y} 2 for cats,^{z,aa} and 1 for both dogs and cats^{bb}). One dry diet for dogs was donated; the other 23 diets were purchased. There were 21 over-the-counter diets for dogs or cats (or both) and 3 veterinary therapeutic diets for dogs. Dry diets represented 9 manufacturers, and canned diets represented 8 manufacturers. There were 2 dry diets that were manufactured at facilities outside the United States.

Only 3 diets (1 dry and 2 canned), including the statement of calorie content as a requirement, and 8 diets (4 dry and 4 canned), excluding the statement of calorie content as a requirement, were compliant with all pet food label regulations as established by the AAFCO. As indicated by the nutritional adequacy statement (or, when not provided, other label information), 14 diets (7 dry and 7 canned) were intended for adult maintenance, 9 diets (5 dry and 4 canned) were intended for all life stages, and 1 diet (dry) was intended for both growth and adult maintenance. Nutritional adequacy for the designated life stage or stages was substantiated by the formulation method to meet the AAFCO Dog and Cat Food Nutrient Profiles for all but 1 diet (1 dry diet for dogs); that diet successfully completed appropriate AAFCO-recognized animal feeding trials.

When label information was compared with the 9 AAFCO requirements, 20 diets (9 dry and 11 canned) met the requirement for product and brand name, 23 diets (13 dry and 10 canned) met the requirement for species specification, 18 diets (7 dry and 11 canned) met the requirement for quantity statement, 17 diets (6 dry and 11 canned) met the requirement for guaranteed analysis, 17 diets (6 dry and 11 canned) met the requirement for ingredient statement, 18 diets (7 dry and 11 canned) met the requirement for nutritional adequacy statement, 12 diets (7 dry and 5 canned) met the requirement for feeding directions, 20 diets (9 dry and 11 canned) met the requirement for name and address of the manufacturer or distributor, and 8 diets (2 dry and 6 canned) met the requirement for statement of calorie content. Of the diets that failed to meet the AAFCO labeling requirements, 4 had the product name outside of the principal display panel, 1 did not have a species-specification statement on the principal display panel, 6 did not have a quantity statement, 4 did not have a guaranteed analysis and 3 did not have an appropriate guaranteed analysis format (terms used and order of items), 5 had misspelled or duplicated words in the ingredient statement and 2 did not have an appropriate

ingredient statement format (ingredients listed under 2 separate headings [ie, composition and additives]), 6 did not have a nutritional adequacy statement, 4 did not have feeding directions, 2 had misspelled words in the feeding directions, 6 did not have frequency or species specifications in the feeding directions, 4 did not have the name and address of the manufacturer, 14 did not have a statement of calorie content, and 2 did not have an appropriate calorie content format (not listed under a heading of calorie content or no information on method of determination). Both diets manufactured outside the United States did not meet 6 of the AAFCO labeling requirements (including not having a statement of calorie content), whereas some of the 22 diets manufactured within the United States did not meet up to 8 of the 9 requirements (including not having a statement of calorie content). Overall, 9 diets (4 dry and 5 canned) had labels with misspelled words.

None of the diets exceeded the maximum moisture percentage as reported on guaranteed analysis. Median measured moisture concentration of the diets was 4.8% (range, 3.3% to 7.8%) for dry diets and 69.9% (range, 61.4% to 74.3%) for canned diets.

Dried eggs were listed as an ingredient in 1 canned diet, whereas the other 23 diets listed only plant-sourced ingredients. Nineteen diets (11 dry and 8 canned) were supplemented with 1 or more AAs: methionine (7 dry and 4 canned), taurine (10 dry and 7 canned), lysine (7 dry and 2 canned), and tryptophan (5 dry and 0 canned); 1 dry diet was supplemented with both cystine and glycine. All 7 diets formulated for cats were supplemented with taurine. Two dry diets included a minimum taurine concentration claim in the guaranteed analysis (which is optional); both of these diets contained taurine in concentrations that exceeded the AAFCO minimum value. However, 1 of the 7 taurine-supplemented diets contained a measured taurine concentration that was 85% of the minimum listed in the guaranteed analysis.

Median measured CP concentration (DM basis) was 29.8% (range, 19.2% to 40.3%) for all diets. Measured CP concentrations were above the minimum requirement for the AAFCO Dog and Cat Food Nutrient Profiles (DM basis or corrected for energy density when necessary) for the intended species and life stage for 23 diets (12 dry and 11 canned). The dry diet for dogs that did not meet the minimum requirement contained 94% of the minimum required value but had completed an AAFCO-recognized animal feeding trial. One additional canned diet for dogs that exceeded 4,000 kcal/kg of DM contained only 91% of the reported minimum CP for the guaranteed analysis on an as-fed basis but met the AAFCO minimum CP on a DM basis when corrected for energy density. All other diets met the reported minimum CP for the guaranteed analysis.

Eighteen diets (10 dry and 8 canned) contained all AAs in concentrations that met or exceeded the minimum values for the AAFCO Dog and Cat Food Nutrient Profiles (DM basis or corrected for energy density when necessary) for the designated life stage (Table 1). Five diets (all for cats; 3 dry and 2 canned) provided 1 or more AAs at concentrations below the AAFCO minimum value. Of these 5 diets, 1 was below the AAFCO

Table 1—The AA concentrations of vegetarian dry and canned diets formulated for dogs and cats and values for the AAFCO Dog and Cat Food Nutrient Profiles.

AA	Median	Range	AAFCO	
			Growth and reproduction (minimum)	Adult maintenance (minimum)
Canine (n = 17)				
Arginine	1.66	1.08–2.83	0.62	0.51
Histidine	0.59	0.40–0.96	0.22	0.18
Isoleucine	1.05	0.84–1.81	0.45	0.37
Leucine	1.88	1.45–4.74	0.72	0.59
Lysine	1.40	0.99–2.47	0.77	0.63
Methionine-cystine	0.85	0.46–3.62	0.53	0.43
Phenylalanine-tyrosine	2.39	1.92–3.90	0.89	0.73
Threonine	1.13	0.90–1.53	0.58	0.48
Tryptophan	0.25	0.18–0.40	0.20	0.16
Valine	1.29	1.01–2.00	0.48	0.39
Taurine	0.19	0.11–0.30	—	—
Feline (n = 7)*				
Arginine	1.85	1.49–2.50	1.25	1.04
Histidine	0.77	0.68–0.88	0.31	0.31
Isoleucine	1.44	1.28–1.58	0.52	0.52
Leucine	3.41	0.43–4.81	1.25	1.25
Lysine	1.46	1.12–2.18	1.20	0.83
Methionine-cystine	1.63	0.59–3.14	1.10	1.10
Methionine†	0.62	0.51–1.32	0.62	0.62
Phenylalanine-tyrosine	3.20	3.00–3.88	0.88	0.88
Phenylalanine	1.89	1.80–2.22	0.42	0.42
Threonine	1.42	1.10–1.60	0.73	0.73
Tryptophan	0.36	0.16–0.41	0.25	0.16
Valine	1.72	1.51–1.80	0.62	0.62
Taurine (extruded)‡	0.18	0.15–0.18	0.10	0.10
Taurine (canned)§	0.12	0.11–0.15	0.20	0.20

Values reported are percentage DM.
 *Includes results for 2 diets formulated for both dogs and cats. †Methionine is the only AA with a maximum allowed value, and only for feline diets (1.5% DM). ‡Values are for 4 extruded diets. §Values are for 3 canned diets.
 — = Not applicable.

minimum requirements in 4 AAs (leucine, methionine, methionine-cystine, and taurine), 1 was below in 3 AAs (methionine, methionine-cystine, and taurine), 2 were below in 2 AAs (lysine and tryptophan), and 1 was below in 1 AA (tryptophan). An additional canned diet intended for both dogs and cats exceeded the AA minimum values for dogs but was below the minimum values for cats for 3 AAs (methionine, methionine-cystine, and taurine), despite inclusion of dried eggs as an ingredient. All of the canned diets formulated for cats (2 for cats and 1 for both dogs and cats) were below the AAFCO minimum value for taurine; dry diets for cats exceeded this value. Overall, of the diets that contained 1 or more AAs at concentrations below AAFCO minimum values, the AA concentrations ranged from 34% to 98% (median, 82%) of the minimum requirement stated in the AAFCO Dog and Cat Food Nutrient Profile. The 2 diets below the minimum value for lysine (98% and 93% of the minimum requirement) were within the analytic variation (20%) allowed by the AAFCO regulations; lysine was the only AA for which the AAFCO provided an allowance for analytic variation. All other AAs that did not meet the AAFCO minimum requirement exceeded the range of analytic variation provided by the laboratory.

Calorie content was provided on the label for 10 diets (4 dry and 6 canned). Calorie content was obtained from the product website for 2 diets (1 dry and 1

canned) and from the manufacturer (on a volume basis only [can or cup]) for 10 diets (6 dry and 4 canned). Calorie content information could not be obtained for 2 diets (both dry). Calorie content was calculated for 4 canned diets by use of the per-unit calorie content provided by the manufacturer, 4 dry diets by use of the modified Atwater factor and ash content provided by the manufacturer, and 4 dry diets by use of the modified Atwater factors and mean ash content calculated for dry diets (n = 8) for which the ash concentration was measured (5.76% on an as-fed basis). Median calorie content (DM basis) for all 24 diets was 3,758 kcal of ME/kg of diet (range, 2,915 to 4,316 kcal of ME/kg of diet). Median calorie content (DM basis) of the 17 diets for dogs was 3,725 kcal of ME/kg of diet (range, 3,233 to 4,316 kcal of ME/kg of diet) and of the 7 diets for cats or for both cats and dogs was 3,843 kcal of ME/kg of diet (range, 2,915 to 4,050 kcal of ME/kg of diet). One diet (canned maintenance diet for dogs) required adjustments in nutrient concentrations on the basis of the correction for calorie content.

Discussion

One objective for the present study was to assess product labeling by comparing diet labels with the AAFCO model regulations.¹³ Although all pet foods must comply with federal labeling requirements,²²

many states also mandate specific aspects of the label, often by adopting the AAFCO labeling and formulation requirements in full or in part.¹⁰ Despite the fact all 24 diets were sold in most or all states, and even with exclusion of calorie content as a requirement, only 8 diets (including all 3 veterinary therapeutic diets) were compliant with all label regulations as established by the AAFCO.

There are 3 means of substantiating claims that pet foods are complete and balanced, and the label's nutritional adequacy statement must specify which method is used.²³ The first method is to formulate the diet to meet the AAFCO Dog and Cat Food Nutrient Profiles. The second method is to conduct a feeding trial by use of AAFCO-recognized protocols for the specified life stage; in the case of successful completion of an appropriate feeding trial, the pet food is exempt from meeting nutrient profiles. Third, if a food is a member of a nutritionally similar product family for which the designated lead product has successfully completed an AAFCO-recognized feeding trial, the label of the products for that food family may state that AAFCO feeding trials substantiate the claim of complete and balanced and the nutritional adequacy statements are indistinguishable. In both cases, the label will state that the product has passed animal feeding tests. When a product fails to meet 1 of the aforementioned 3 methods and is not clearly labeled on the principal display as a snack, treat, or dietary supplement, the product must contain a statement that indicates "intended for intermittent or supplemental feeding only." One diet in the present study had a nutritional adequacy statement that indicated it had successfully completed AAFCO-recognized animal feeding trials (which we confirmed by contacting the manufacturer) and was assessed as adequately formulated, although the CP concentration was 94% of the AAFCO nutrient profile minimum value; all AA concentrations exceeded the AAFCO minimum values. Of the 6 diets that did not have nutritional adequacy statements, none were labeled snack or treat, and they did not have a statement to indicate that the product was intended for intermittent or supplemental feeding only. Rather, the labels of those 6 diets included wording that indicated that they were intended to be complete and balanced (phrasing such as "100% complete" and "ideal maintenance"), which was inadequate.

The AAFCO Dog and Cat Food Nutrient Profiles provide minimum values for CP and essential AA concentrations (as well as a maximum value for methionine concentration in foods formulated for cats) for pet foods made with complex, nonpurified ingredients and to account for effects of processing and impacts on digestibility. Most (23/24) diets assessed in the present study met guaranteed analysis claims for minimum CP concentration, and most (23/24) diets exceeded CP minimum values for the AAFCO nutrient profiles; however, CP concentration was assessed with *in vitro* methods that provided an estimate of protein content calculated by use of the nitrogen concentration. As such, the calculated CP value provided no information related to protein quality, which is defined by the digestibility of the protein and the pattern and bioavailability of the AAs. It is generally recognized that plant protein

sources have lower digestibility than do animal protein sources²⁴; however, studies^{25,26} of dogs have found equal total digestibility for soy-based protein when the soy product is adequately processed. Both animal and plant protein sources can vary in quality. Although protein digestibility was not assessed in the present study, short-term studies^{27,28} revealed that animal-protein meals differ in their ability to support nitrogen retention in cats, with chicken and fish meals not differing from corn gluten meal, whereas meat meal is superior to corn gluten meal. Because digestibility, AA pattern, and AA bioavailability are not provided on product labels, protein quality cannot be assessed from a pet food's ingredient list or guaranteed analysis regardless of the fact that nutrients may be present in concentrations that satisfy the corresponding AAFCO nutrient profile. Investigators of 1 study²⁹ reported limitations of measured CP concentrations for the assessment of protein quality of pet food as evaluated with feeding trials on growing rats. They reported that the biological variables for assessment of protein quality (including weight gain, feed efficiency, protein efficiency ratio, net protein ratio, and net protein utilization) had poor correlation with measured CP concentrations.²⁹ Furthermore, the sum of essential AA concentrations was not correlated with measured CP concentration or biological variables (protein efficiency ratio and net protein ratio).²⁹

Concentrations and proportions of AAs are arguably more important than is CP concentration per se, and AA bioavailability should also be considered. Dogs and cats differ from many other species in that they have obligatory bile acid conjugation with taurine rather than glycine, which is associated with variable losses of taurine through feces. Effects of intestinal bacteria on taurine loss appear to be substantial^{30,31} and may be exacerbated by dietary factors. Studies^{32,33} have revealed that cats fed canned versus frozen-preserved diets, or diets with soybean versus casein protein, had lower plasma taurine concentrations, even though the diets were equal in taurine content. The negative effect on taurine status appears to be secondary to augmented loss of bile acids through microbial degradation and accelerated cholecystokinin-mediated turnover of bile acids.³⁴ In addition, fiber likely increases taurine losses in the feces by influencing intestinal bacterial populations as well as through other effects on bile acid metabolism.³⁵ In the present study, the 3 diets for dogs that provided methionine-cystine concentrations closest to the AAFCO minimum value (8%, 25%, and 35% above the minimum value) were all canned diets that did not provide additional purified sulfur-containing AAs. In addition, all 3 canned diets for cats were too low in taurine concentration despite supplementation. Because plant-based diets are typically lower in sulfur-containing AAs and higher in fiber, these factors may contribute to an increased risk of taurine deficiency in both dogs and cats fed vegetarian diets, especially canned products and products that do not provide supplemental taurine or its precursors.

Processing of pet foods impacts protein digestibility as well as AA bioavailability. Conditions for ingredient rendering, extrusion cooking, and can retorting include application of heat, moisture, pressure, or

mechanical shear to inactivate food-borne pathogens, increase shelf-life, increase digestibility of certain nutrients (denaturation of protein, gelatinization of starch, and inactivation of trypsin inhibitors in vegetable protein), and promote desirable flavor and texture.³⁶ However, despite these beneficial effects of processing, some nutrients are lost during processing. Nonenzymatic browning of foods during processing as a result of Maillard reactions is considered a major factor that negatively affects the quality of protein. Depending on the exact conditions and nutrients present, variable AA losses occur (especially losses of lysine, methionine, cystine, and tryptophan).³⁷ Concentrations of 3 of the 4 AAs (all but lysine) were too low in some of the diets assessed in the present study.

Two diets for cats, including 1 diet with purified L-lysine in the ingredient list, did not meet the minimum concentration for lysine as per the AAFCO food nutrient profiles, but the values for these 2 diets were within the analytic variation allowed by the AAFCO. A third diet provided lysine at only 1% above the minimum AAFCO value. However, bioavailability is an important consideration. Acid hydrolysis of protein, which is required for the measurement of AAs in food, results in reversion of damaged (unavailable) lysine and falsely increases the estimate of bioavailable lysine. In 1 study,³⁸ measurement of total lysine overestimated by 87% the bioavailable lysine concentration of 20 diets formulated for cats (10 dry and 10 canned). Lysine is commonly the limiting AA in cereals, and the impact of processing on lysine availability together with a limited ability to accurately assess available lysine concentrations with routine methods is of particular concern for commercially available vegetarian pet foods.

Notably, 6 of the 24 diets assessed in the study reported here were inadequate in 1 or more AAs; 3 of these diets were too low in sulfur-containing AAs (methionine, methionine-cystine, and taurine). However, on the basis of the ingredient lists, all 3 of those diets were supplemented with taurine, and 2 of those 3 diets were also supplemented with methionine. This finding is similar to that in a study³⁹ conducted to investigate nutritional adequacy of 2 commercially available vegan diets for cats. The authors of that study³⁹ found that both diets had inadequate concentrations of taurine, methionine, methionine-cystine, arachidonic acid, and pyridoxine. One of the diets had additional deficiencies of CP, arginine, lysine, calcium, phosphorus, vitamin A, niacin, and vitamin B₁₂, despite label claims of nutritional adequacy and the fact that limiting AAs were listed in the ingredient list as additive supplements. Dietary deficiencies in sulfur-containing AAs and lysine could result in decreased food intake, low growth rate, and negative nitrogen balance in both dogs and cats.⁴⁰⁻⁴⁶ Furthermore, dermatitis has been reported in dogs⁴⁷ and cats^{48,49} with methionine and lysine deficiency, and retinal and cardiac dysfunction has been reported in dogs^{8,9} and cats⁵⁰ with taurine deficiency.

Analysis of results of the study reported here indicated problems with compliance with labeling regulations in addition to concerns regarding adequacy of AA concentrations in commercially available vegetarian pet foods. Overall, only 5 of 21 over-the-counter diets, but

all 3 of the veterinary therapeutic diets, met all requirements for labeling and nutritional adequacy (excluding the recently published regulation for a calorie content statement); however, the sample size was small. Another important limitation of this study was that samples were collected at 1 time point and from 1 batch of each product. The samples that were assessed for CP and AA concentrations may not have been representative because of variations in composition for each batch. In addition, although assay variability for both AA and CP analysis was low, substantial variations in results attributable to laboratory methods were possible. Regardless, all nutritional and labeling requirements should be met consistently, and manufacturers are responsible for quality assurance. It may be informative to measure the CP and AA concentrations across numerous batches to assess variation and more accurately determine the deviation from nutritional adequacy and regulatory compliance.

In the present study, we assessed only a limited number of essential nutrients in commercially available vegetarian pet foods. A more thorough evaluation of other essential nutrients is warranted, especially because important inadequacies of other nutrients in vegan pet foods have been reported.³⁹ In addition, there was no assessment of the animals while consuming the diets; evaluation of blood AA concentrations would provide valuable information for assessing the AA adequacy of pet foods.⁵¹ Only 1 diet had a nutritional adequacy statement indicating that it had passed AAFCO feeding trials to substantiate a claim of complete and balanced for the specified life stages. Given that both the present study and a previous report³⁹ documented deficiencies of nutrients that were declared to have been included in purified form, this may be evidence that manufacturing errors occur or that diets are not formulated properly. Veterinary therapeutic diets may be more appropriate options for vegetarian pet foods because all 3 veterinary diets assessed in the study reported here met current nutritional adequacy and labeling requirements, compared with only 5 of 21 over-the-counter diets that met the nutritional adequacy and labeling requirements. In addition, the US FDA provides allowance for the marketing of veterinary therapeutic diets under the presumption that they are used only under the direction of a licensed veterinarian who is providing recommendations for appropriate use of the product and for monitoring of individual patients.⁵² It may be prudent that such monitoring includes measurement of plasma AA and whole blood taurine concentrations as well as routine assessment of general health to more fully evaluate the status of pets eating vegetarian diets. Given the findings of the present study, this may be of even greater importance for dogs and cats eating canned vegetarian diets, in which case regular monitoring of taurine status in particular is strongly recommended. For all animals and regardless of diet, general routine monitoring and assessment are necessary for adequate nutritional evaluation and to enable clinicians to provide recommendations for individual animals.⁵³

a. Sacramento Animal Hospital, Sacramento, Calif.

b. UC Davis Analytical Lab, University of California-Davis, Davis, Calif.

- c. Biochrom 30, Biochrom Ltd, Holliston, Mass.
- d. Microsoft Office Excel 2008, Microsoft Corp, Redmond, Wash.
- e. Ami Dog, Ami, Padova, Italy.
- f. Gourmet Fondue Veggie Cheese Burger Flavor, Evolution Diet Pet Food, Saint Paul, Minn.
- g. Incredibly Delicious Gourmet Pasta, Evolution Diet Pet Food, Saint Paul, Minn.
- h. Vegetarian Formula for Dogs, Dick Van Patten's Natural Balance Pet Foods, Pacoima, Calif.
- i. Vegan Garden Medley Adult, Halo Purely for Pets, Tampa, Fla.
- j. Nature's Recipe Healthy Skin Vegetarian Recipe, Big Heart Pet Brands, San Francisco, Calif.
- k. Veterinary Diets HA Hypoallergenic Canine Formula, Néstle Purina, St Louis, Mo.
- l. Veterinary Diet Canine Vegetarian, Royal Canin, Charles, Mo.
- m. V-dog, V-dog Food, Sacramento, Calif.
- n. Ami Cat, Ami, Padova, Italy.
- o. Gourmet Fondue Veggie Cheese Burger Flavor, Evolution Diet Pet Food, Saint Paul, Minn.
- p. Incredibly Delicious Gourmet Pasta, Evolution Diet Pet Food, Saint Paul, Minn.
- q. Vegan, Wysong Corp, Midland, Mich.
- r. AvoDerm Natural Vegetarian Formula, Central Garden and Pet Co, Walnut Creek, Calif.
- s. Vegetable Stew Entrée, Evolution Diet Pet Food, Saint Paul, Minn.
- t. Vegetarian Formula, Dick Van Patten's Natural Balance Pet Foods, Pacoima, Calif.
- u. Vegan Garden Medley for Dogs, Halo, Purely for Pets, Tampa, Fla.
- v. Nature's Recipe Stew Healthy Skin Vegetarian Recipe Cuts in Gray, Big Heart Pet Brands, San Francisco, Calif.
- w. Organic Vegan Formula, PetGuard, Green Cove Springs, Fla.
- x. Vegetarian Feast Dinner, PetGuard, Green Cove Springs, Fla.
- y. Veterinary Diet Canine Vegetarian, Royal Canin, Charles, Mo.
- z. Gourmet Entrée, Evolution Diet Pet Food, Saint Paul, Minn.
- aa. Vegetable Stew Entrée, Evolution Diet Pet Food, Saint Paul, Minn.
- bb. Vegetarian Dinner, Evanger's Dog and Cat Food Co, Wheeling, Ill.

References

1. Wakefield LA, Shofer FS, Michel KE. Evaluation of cats fed vegetarian diets and attitudes of their caregivers. *J Am Vet Med Assoc* 2006;229:70–73.
2. Joshi M, Mehta MK, Sharma SK. Feeding practices and common nutritional deficiency disorders in dogs. *Vet Pract* 2007;8:83–84.
3. Craig WJ, Mangels AR, American Dietetic Association. Position of the American Dietetic Association: vegetarian diets. *J Am Diet Assoc* 2009;109:1266–1282.
4. Rogers QR, Morris JG. Do cats really need more protein? *J Small Anim Pract* 1982;23:521–532.
5. Sturman JA. Taurine in development. *Physiol Rev* 1993;73:119–147.
6. Kittleson MD, Keene B, Pion PD, et al. Results of the multicenter Spaniel trial (MUST): taurine- and carnitine-responsive dilated cardiomyopathy in American Cocker Spaniels with decreased plasma taurine concentration. *J Vet Intern Med* 1997;11:204–211.
7. Sanderson SL, Osborne CA, Lulich JP, et al. Evaluation of urinary carnitine and taurine excretion in 5 cystinuric dogs with carnitine and taurine deficiency. *J Vet Intern Med* 2001;15:94–100.
8. Sanderson SL, Gross KL, Ogburn PN, et al. Effects of dietary fat and L-carnitine on plasma and whole blood taurine concentrations and cardiac function in healthy dogs fed protein-restricted diets. *Am J Vet Res* 2001;62:1616–1623.
9. Fascetti AJ, Reed JR, Rogers QR, et al. Taurine deficiency in dogs with dilated cardiomyopathy: 12 cases (1997–2001). *J Am Vet Med Assoc* 2003;223:1137–1141.
10. Dzanis DA. Understanding regulations affecting pet foods. *Top Companion Anim Med* 2008;23:117–120.
11. WSAVA Global Nutrition Committee. WSAVA Global Nutrition Committee: recommendations on selecting pet foods. Available at: www.wsava.org/sites/default/files/Recommendations%20on%20Selecting%20Pet%20Foods.pdf. Accessed Apr 18, 2015.
12. Hill RC, Choate CJ, Scott KC, et al. Comparison of the guaranteed analysis with the measured nutrient composition of commercial pet foods. *J Am Vet Med Assoc* 2009;234:347–351.
13. Association of American Feed Control Officials. Model regulations for pet food and specialty pet food under the model bill. In: *2014 official publication*. Oxford, Ind: Association of American Feed Control Officials, 2014;136–164.
14. Association of American Feed Control Officials. Recommendation of enforcement dates. In: *2014 official publication*. Oxford, Ind: Association of American Feed Control Officials, 2014;iii.
15. AOAC official method 990.03. Protein (crude) in animal feed, combustion method, chapter 4. In: Horowitz W, Latimer GW Jr, eds. *Official methods of analysis of AOAC International*. 18th ed. Revision 1. Gaithersburg, Md: AOAC International, 2006;30–31.
16. Association of American Feed Control Officials. Analytical variations (AV) based on AAFCO Check Sample Program. In: *2014 official publication*. Oxford, Ind: Association of American Feed Control Officials, 2014;296–298.
17. Sweeney RA. Generic combustion method for determination of crude protein in feeds: collaborative study. *J Assoc Off Anal Chem* 1989;72:770–774.
18. Spitze AR, Wong DL, Rogers QR, et al. Taurine concentrations in animal feed ingredients; cooking influences taurine content. *J Anim Physiol Anim Nutr (Berl)* 2003;87:251–262.
19. AOAC official method 994.12. Chapter 4: amino acids in feeds. In: Horowitz W, Latimer GW Jr, eds. *Official methods of analysis of AOAC International*. 18th ed. Gaithersburg, Md: AOAC International, 2006;9–19.
20. AOAC official method 988.15. Chapter 45: tryptophan in foods and food and feed ingredients. In: Horowitz W, Latimer GW Jr, eds. *Official methods of analysis of AOAC International*. 18th ed. Gaithersburg, Md: AOAC International, 2006;88–89.
21. Association of American Feed Control Officials. Correcting for energy density. In: *2014 official publication*. Oxford, Ind: Association of American Feed Control Officials, 2014;160–161.
22. US FDA. Pet food labels—general. Available at: www.fda.gov/AnimalVeterinary/ResourcesforYou/UCM047113. Accessed Sep 19, 2014.
23. Association of American Feed Control Officials. Regulation PF7. Nutritional adequacy. In: *2014 official publication*. Oxford, Ind: Association of American Feed Control Officials, 2014;142–143.
24. Neirinck K, Istasse L, Gabriel A, et al. Amino acid composition and digestibility of four protein sources for dogs. *J Nutr* 1991;121:S64–S65.
25. Clapper GM, Grieshop CM, Merchen NR, et al. Ileal and total tract nutrient digestibilities and fecal characteristics of dogs as affected by soybean protein inclusion in dry, extruded diets. *J Anim Sci* 2001;79:1523–1532.
26. Bednar GE, Murray SM, Patil AR, et al. Selected animal and plant protein sources affect nutrient digestibility and fecal characteristics of ileally cannulated dogs. *Arch Tierernahr* 2000;53:127–140.
27. Funaba M, Oka Y, Kobayashi S, et al. Evaluation of meat meal, chicken meal, and corn gluten meal as dietary sources of protein in dry cat food. *Can J Vet Res* 2005;69:299–304.
28. Funaba M, Tanak T, Kaneko M, et al. Fish meal vs. corn gluten meal as a protein source for dry cat food. *J Vet Med Sci* 2001;63:1355–1357.
29. Hegedüs M, Fekete S, Solti L, et al. Assessment of nutritional adequacy of the protein in dog foods by trials on growing rats. *Acta Vet Hung* 1998;46:61–70.
30. Kim SW, Rogers QR, Morris JG. Dietary antibiotics decrease taurine loss in cats fed a canned heat-processed diet. *J Nutr* 1996;126:509–515.
31. Hickman MA, Rogers QR, Morris JG. Effect of processing on fate of dietary [¹⁴C]taurine in cats. *J Nutr* 1990;120:995–1000.
32. Hickman MA, Bruss ML, Morris JG, et al. Dietary protein source (soybean vs. casein) and taurine status affect kinetics of the enterohepatic circulation of taurocholic acid in cats. *J Nutr* 1992;122:1019–1028.
33. Kim SW, Morris JG, Rogers QR. Dietary soybean protein decreases plasma taurine in cats. *J Nutr* 1995;125:2831–2837.
34. Backus RC, Rogers QR, Rosenquist GL, et al. Diets causing taurine depletion in cats substantially elevate postprandial plasma cholecystokinin concentration. *J Nutr* 1995;125:2650–2657.
35. Stratton-Phelps M, Backus RC, Rogers QR, et al. Dietary rice bran decreases plasma and whole-blood taurine in cats. *J Nutr* 2002;132:1745S–1747S.

36. van Boekel M, Fogliano V, Pellegrini N, et al. A review on the beneficial aspects of food processing. *Mol Nutr Food Res* 2010;54:1215–1247.
37. Hendriks WH, Emmens MM, Trass B, et al. Heat processing changes the protein quality of canned cat foods as measured with a rat bioassay. *J Anim Sci* 1999;77:669–676.
38. Rutherford SM, Rutherford-Markwick KJ, Moughan PJ. Available (ileal digestible reactive) lysine in selected pet foods. *J Agric Food Chem* 2007;55:3517–3522.
39. Gray CM, Sellon RK, Freeman LM. Nutritional adequacy of two vegan diets for cats. *J Am Vet Med Assoc* 2004;225:1670–1675.
40. Burns RA, Milner JA. Sulfur amino acid requirements of immature Beagle dogs. *J Nutr* 1981;111:2117–2124.
41. Blaza SE, Burger IH, Holme DW, et al. Sulfur-containing amino acid requirements of growing dogs. *J Nutr* 1982;112:2033–2042.
42. Hirakawa DA, Baker DH. Lysine requirement of growing puppies fed practical and purified diets. *Nutr Res* 1986;6:527–538.
43. Milner JA. Lysine requirements of the immature dog. *J Nutr* 1981;111:40–45.
44. Rogers QR, Morris JG. Essentiality of amino acids for the growing kitten. *J Nutr* 1979;109:718–723.
45. Morris JG, Rogers QR, O'Donnell JA. Lysine requirement of kittens given purified diets for maximal growth. *J Anim Physiol Anim Nutr (Berl)* 2004;88:113–116.
46. Teeter RG, Baker DH, Corbin JE. Methionine and cystine requirements of the cat. *J Nutr* 1978;108:291–295.
47. Hirakawa DA, Baker DH. Sulfur amino acid nutrition of the growing puppy: determination of dietary requirements for methionine and cystine. *Nutr Res* 1985;5:631–642.
48. Strieker MJ, Werner A, Morris JG, et al. Excess dietary cystine intensifies the adverse effect of a methionine deficiency in the cat. *J Anim Physiol Anim Nutr (Berl)* 2006;90:440–445.
49. Larsen JA, Outerbridge CA, Fascetti AJ, et al. Skin lesions associated with lysine deficiency in kittens are characterized by inflammation. *Int J Appl Res Vet Med* 2014;12:61–66.
50. Burger IH, Barnett KC. The taurine requirement of the adult cat. *J Small Anim Pract* 1982;23:533–537.
51. Zicker S, Rogers QR. Use of plasma amino acid concentrations in the diagnosis of nutritional and metabolic diseases in veterinary medicine, in *Proceedings. IVth Cong Int Soc Anim Clin Biochem* 1990;1–16.
52. US FDA. Draft compliance policy guide: labeling and marketing of nutritional products intended for use to diagnose, cure, mitigate, treat, or prevent diseases in dogs and cats. www.fda.gov/downloads/ICECI/ComplianceManuals/CompliancePolicyGuidanceManual/UCM318761.pdf. Accessed Oct 4, 2014.
53. Freeman L, Becvarova I, Cave N, et al. WSAVA Nutritional Assessment Guidelines. *J Small Anim Pract* 2011;52:385–396.



From this month's *AJVR*

Electrocardiogram reference intervals for clinically normal wild-born chimpanzees (*Pan troglodytes*)

Rebeca Atencia et al


Objective—To generate reference intervals for ECG variables in clinically normal chimpanzees (*Pan troglodytes*).

Animals—100 clinically normal (51 young [< 10 years old] and 49 adult [≥ 10 years old]) wild-born chimpanzees.

Procedures—Electrocardiograms collected between 2009 and 2013 at the Tchimpounga Chimpanzee Rehabilitation Centre were assessed to determine heart rate, PR interval, QRS duration, QT interval, QRS axis, P axis, and T axis. Electrocardiographic characteristics for left ventricular hypertrophy (LVH) and morphology of the ST segment, T wave, and QRS complex were identified. Reference intervals for young and old animals were calculated as mean \pm 1.96•SD for normally distributed data and as 5th to 95th percentiles for data not normally distributed. Differences between age groups were assessed by use of unpaired Student *t* tests.

Results—Reference intervals were generated for young and adult wild-born chimpanzees. Most animals had sinus rhythm with small or normal P wave morphology; 24 of 51 (47%) young chimpanzees and 30 of 49 (61%) adult chimpanzees had evidence of LVH as determined on the basis of criteria for humans.

Conclusions and Clinical Relevance—Cardiac disease has been implicated as the major cause of death in captive chimpanzees. Species-specific ECG reference intervals for chimpanzees may aid in the diagnosis and treatment of animals with, or at risk of developing, heart disease. Chimpanzees with ECG characteristics outside of these intervals should be considered for follow-up assessment and regular cardiac monitoring. (*Am J Vet Res* 2015;76:688–693)



See the midmonth issues
of *JAVMA*
for the expanded
table of contents
for the *AJVR*
or log on to
avmajournals.avma.org
for access
to all the abstracts.



Influence of mercury and selenium chemistries on the progression of cardiomyopathy in pygmy sperm whales, *Kogia breviceps*

Colleen E. Bryan^{a,b,*}, W. Clay Davis^a, Wayne E. McFee^c, Carola A. Neumann^d, Jennifer Schulte^d, Gregory D. Bossart^e, Steven J. Christopher^a

^a Analytical Chemistry Division, National Institute of Standards and Technology, Hollings Marine Laboratory, 331 Fort Johnson Road, Charleston, SC 29412, USA

^b Marine Biomedicine and Environmental Science Center, Medical University of South Carolina, 221 Fort Johnson Road, Charleston, SC 29412, USA

^c Center for Coastal Environmental Health and Biomolecular Research, National Ocean Service, National Oceanic and Atmospheric Administration, 219 Fort Johnson Road, Charleston, SC 29412, USA

^d Department of Cell and Molecular Pharmacology and Experimental Therapeutics, Medical University of South Carolina, 173 Ashley Avenue, MSC-505, Charleston, SC 29425, USA

^e Georgia Aquarium, 225 Baker Street, Atlanta, GA 30313, USA

HIGHLIGHTS

- ▶ More than half of stranded pygmy sperm whales exhibit signs of cardiomyopathy.
- ▶ Hg and Se balance and oxidative stress may influence progression of cardiomyopathy.
- ▶ Adults have significantly greater Hg:Se liver molar ratios than younger age classes.
- ▶ Hg:Se molar ratios were greater in males and increased with heart disease progression.
- ▶ Protein oxidation was greater in males and increased with heart disease progression.

ARTICLE INFO

Article history:

Received 16 September 2011
Received in revised form 20 January 2012
Accepted 16 May 2012
Available online 15 June 2012

Keywords:

Mercury
Selenium
Protein oxidation
Cardiomyopathy
Pygmy sperm whale

ABSTRACT

More than half of pygmy sperm whales (*Kogia breviceps*) that strand exhibit signs of cardiomyopathy (CMP). Many factors may contribute to the development of idiopathic CMP in *K. breviceps*, including genetics, infectious agents, contaminants, biotoxins, and dietary intake (e.g. selenium, mercury, and pro oxidants). This study assessed trace elements in *K. breviceps* at various stages of CMP progression using fresh frozen liver and heart samples collected from individuals that stranded along US Atlantic and Gulf coasts between 1993 and 2007. Standard addition calibration and collision cell inductively coupled plasma mass spectrometry (ICP MS) were employed for total Se analysis and pyrolysis atomic absorption (AA) was utilized for total Hg analysis to examine if the Se/Hg detoxification pathway inhibits the bioavailability of Se. Double spike speciated isotope dilution gas chromatography ICP MS was utilized to measure methyl Hg and inorganic Hg. Immunoblot detection and colorimetric assays were used to assess protein oxidation status. Data collected on trace elements, selenoproteins, and oxidative status were evaluated in the context of animal life history and other complementary histological information to gain insight into the biochemical pathways contributing to the development of CMP in *K. breviceps*. Cardiomyopathy was only observed in adult pygmy sperm whales, predominantly in male animals. Both Hg:Se molar ratios and overall protein oxidation were greater in males than females and increased with progression of CMP.

Published by Elsevier Ltd.

1. Introduction

Greater than fifty percent of pygmy sperm whales (*Kogia breviceps*) that strand show signs of cardiac degradation and cardiomy

opathy (CMP) (Bossart et al., 1985, 2007). Natural occurrence of CMP in *K. breviceps* is greater than all other mammal species studied to date. The idiopathic nature of this disease requires studying many of the possible factors that may contribute to the development of CMP in *K. breviceps*, including genetics, infectious agents, chemical toxins, biotoxins, contaminants, and nutritional abnormalities (e.g. trace elements, vitamins, and pro oxidants). This study focuses contaminant uptake and tissue distribution involving selenium (Se) and mercury (Hg) chemistries, along with a

* Corresponding author at: Analytical Chemistry Division, National Institute of Standards and Technology, Hollings Marine Laboratory, 331 Fort Johnson Road, Charleston, SC 29412, USA. Tel.: +1 843 762 8832; fax: +1 843 762 8742.

E-mail address: colleen.bryan@nist.gov (C.E. Bryan).

marker for protein oxidative stress, which may contribute to CMP onset and progression. Nutritional deficiencies of Se, accumulation of Hg, and imbalance of oxidants have been shown in humans, mice, hamsters, cattle, and dogs to play roles in the development and progression of CMP (Kennedy et al., 1987; Bartfay et al., 1998; Freeman et al., 1998; Kannan et al., 2004; Ichihara et al., 2006).

Marine mammals exhibit an Hg detoxification metabolism wherein Se is specifically involved in Hg detoxification pathways. Methyl Hg is demethylated to inorganic Hg (iHg) that can bind with elemental Se to form Hg Se crystals, which are stored in the liver and cannot be mobilized. Selenium binds strongly to Hg and has a greater affinity for Hg than sulfur due to a low solubility product constant ($K_{sp} = 1 \times 10^{-59}$ at 298 K). This mechanism appears to be uniquely self limiting in marine mammals; measured Hg:Se molar ratios near or less than 1 are typical. In other species (e.g. birds) that are more sensitive to Hg contamination, Hg:Se molar ratios can exceed 10:1 (Scheuhammer et al., 2008). Relating the Hg and Se chemistries and oxidation measurements to CMP in pygmy sperm whales allows for a first look into whether the Hg detoxification pathway could play a role in CMP progression by sequestering bioavailable Se that may otherwise be channeled into chemoprotective pathways, for example in selenoprotein/selenoenzyme synthesis, and Se antioxidant biochemistry.

Marine mammals can serve as key sentinel species for environmental health since they are long lived, top predators that utilize many of the same ocean resources as humans (Wells et al., 2004; Bossart, 2006). Increased human activity in recent decades has accelerated inputs of Hg into the marine environment resulting in larger amounts entering marine food chains. Mercury concentrations have been well documented in marine mammal species that have strong interactions with humans through sharing coastal water habitats, subsistence hunting, or being caught as fisheries by catch, while little is known about pygmy sperm whales that rarely have interactions with humans until stranding (Wells et al., 2004; Booth and Zeller, 2005). During 2003, an increased number of pygmy sperm whales stranded along the southeastern coasts of the United States and renewed marine mammal community interest in understanding health problems associated with the species (Berini, 2009). Pygmy sperm whales are pelagic odontocetes that feed at higher trophic levels than mysticetes and other marine organisms, translating to relatively higher accumulation of toxic metals such as Hg. Pygmy sperm whales primarily consume a cephalopod diet that is high in polyunsaturated fatty acids (PUFAs) and free radicals. Lacking sufficient antioxidant defenses, PUFAs can turn rancid very rapidly and can impact cellular membrane structure and function (Kennedy et al., 1987). Stomach contents of pygmy sperm whales have revealed their main prey items are squid in the family Histioteuthidae, which are predatory squid that concentrate Hg and other contaminants in the food chain (Bustamante et al., 2006; Santos et al., 2006). It is uncertain if the restrictive diet of *K. breviceps* is playing a role in the development and progression of CMP by requiring an excess in antioxidants, while simultaneously adding large enough doses of Hg that can contribute to oxidative damage and affect Se biochemistry response. This working hypothesis was studied by analyzing Hg and Se chemistries in conjunction with protein oxidation in the context of cardiac disease state.

2. Materials and methods

2.1. Study population and sample collection

Pygmy sperm whale samples were collected from animals that stranded from 1993 to 2007 along the United States Atlantic and

Gulf of Mexico coasts. All animals that are part of this study were either euthanized or freshly dead in order to minimize postmortem degradation of samples. Liver tissue from individuals ($n = 30$) were collected according to NISTIR 6279 for the National Marine Mammal Tissue Bank (NMMTB) housed at the National Institute of Standards and Technology (NIST), Charleston, SC and stored under cryogenic conditions (approximately -150°C) (Becker et al., 1999). Additional heart ($n = 11$) and liver ($n = 32$) samples donated for use in this project were collected by the Center for Coastal Environmental Health and Biomolecular Research (CCEHBR) NOS/NOAA, Charleston, SC and stored at -80°C until analysis. Fresh heart tissue was available for collection at necropsies from some individual animals ($n = 36$) for histology preparation and histopathological evaluation. Individual animal life history and gross pathology data for this project was generated by the collaborating institutions and communicated in the form of necropsy reports. Animals were divided into three age classes based on the following criteria: calf, dependent animal that is observed with its presumed mother or <200 cm; subadult, independent sexually immature animals $200 - 270$ cm; adult, independent animals with observed indication of sexual maturity or >270 cm (Bossart et al., 1985).

2.2. Sample preparation

Fresh frozen tissue was cryogenically homogenized to produce a uniform sample composition of fresh frozen powder for analysis. Liver samples from the NMMTB were homogenized in class 100 clean room conditions at NIST, Charleston, SC using cryogenic procedures developed by Zeisler et al. (1983) and Pugh et al. (2007). Left ventricle heart and liver tissue samples donated by CCEHBR/NOS/NOS were cryogenically homogenized using a bench top 6850 Freezer/Mill (SPEX SamplePrep, Metuchen, NJ). Samples were placed in vials with a stainless steel impactor, capped, placed in the mill, submerged in LN_2 , and shaken at 10 Hz for 3 min. Homogenized powder was transferred into pre cleaned polypropylene jars and stored at -80°C .

Acid assisted microwave digestion using PTFE pressurized vessels was utilized to decompose liver and heart tissue samples prior to performing inductively coupled plasma mass spectrometry (ICP-MS) analyses of total Se. Sample digestion methods are described in detail in the Supplementary Material. Tissue samples were aliquotted directly in nickel weigh boats and weighed for total Hg (THg) measurements. Sample preparation methods for MeHg and iHg were described in detail by Davis et al. (2007).

Hematoxylin and eosin (H&E) stained histology slides for each animal case were evaluated independently without knowledge of life history or chemical analyses data by a veterinary pathologist and assigned a heart score according to criteria put forth by Bossart et al. (2007). Heart histology and gross pathology was used to categorize heart disease in the following three stages: no pathological findings (NPF), myocardial degeneration (MCD), and cardiomyopathy (CMP).

2.3. Analytical techniques

2.3.1. Instrumentation

Total Se mass fraction measurements were collected using a Thermo Electron X Series II ICP-MS (Bremen, Germany) with a standard low volume glass impact bead spray chamber (Peltier cooled at $+3^\circ\text{C}$), concentric glass nebulizer, and operating in collision cell mode utilizing 8% H_2 in 92% He as the collision cell gas. The collision cell ICP-MS working conditions were optimized with a 10 ng/g 68 element tuning solution and a Se calibrant prior to sample analysis. The mass fraction of THg was determined with a direct mercury analyzer DMA 80 (Milestone Scientific, Shelton, CT) by pyrolytic sample decomposition, catalytic reduction to

FDA-CVM-FOIA-2019-1704-000536

Hg⁰, and trapping on a gold amalgamation trap. The Hg was then thermally desorbed and the Hg atomic absorbance was measured at 254 nm. Methyl Hg and iHg measurements were made using a Thermo Trace GC Ultra gas chromatograph (ThermoFinnigan, Austin, TX) equipped with a 30 m DB 5MS + DG 250 μ m i.d. capillary column coated with a 0.25 μ m thick film of (5% Phenyl) methylpolysiloxane (J & W Scientific, Folsom, CA). The GC was coupled to a Thermo Elemental X7 quadrupole ICP MS (Winsford, UK) by a Thermo GC/ICP MS commercial interface.

2.3.2. Calibration methods and sample measurements

An analytical quantification and validation scheme using the method of standard additions was employed for Se mass fraction measurements in liver and heart samples (Christopher et al., 2005). A NIST interlaboratory comparison exercise control material, QC03LH3 pygmy sperm whale liver homogenate was used to build matrix matched standard addition calibration curves for liver tissue analysis by spiking at different concentration levels. Calibration curve slopes were used to assign Se concentrations in unknown samples that were unspiked. Single point standard addition methods were used for heart tissue analysis since a whale heart CRM is not available and calibration curves from one tissue (e.g. liver) are not transferable to another tissue type (e.g. heart) to produce accurate data. Single point methods avoid matrix interferences by splitting a single sample and spiking one of the sample splits. The spike was prepared from SRM 3149 Selenium Standard Solution (NIST, Gaithersburg, MD).

Total Hg concentrations were determined by external calibration utilizing SRM 1946 Lake Superior fish tissue (NIST, Gaithersburg, MD) and QC03LH3 pygmy sperm whale liver homogenate by aliquotting different masses of the certified reference materials (CRMs) in nickel sample boats. The slope and intercept from the established calibration curves for the CRMs were used to calculate the concentration in the heart, liver, and control material samples. MeHg and iHg were measured using double spike speciated isotope dilution methods as described in detail by Davis et al. (2007). Details of quality assurance are provided in the Supplementary Material. Reported concentrations for Se, THg, and MeHg are presented as mass fraction values, expressed in μ g/g on a wet mass fraction basis.

2.4. Protein oxidation

Protein oxidation status was examined in liver samples from 30 individual pygmy sperm whales that were banked in the NMMTB, have been cryogenically homogenized, and were prepared for analysis under anoxic conditions. Only NMMTB livers were measured for protein oxidation since sample collection and storage integrity minimized outside oxidation exposure to the samples. Oxyblot™ Protein Oxidation Detection Kit (Chemicon International, Temecula, CA) was used for immunoblot detection and quantification of proteins that have been modified by free radicals. Methods for protein oxidation and calculating the oxidation ratio are described in detail in the Supplementary Material. Normalizing for varying protein concentrations between animal samples was achieved by performing a Bio Rad protein assay (Bio Rad Laboratories, New York), which is based on the Bradford method using bovine serum albumin as a standard protein (Bradford, 1976).

Cayman's GSH assay kit (Cayman Chemical Company, Ann Arbor, MI) was used to measure total free glutathione by measuring both total GSH and GSSG. The kit utilized glutathione reductase for enzymatic recycling to quantify GSH (Tietze, 1969; Eyer and Podhradský, 1986; Baker et al., 1990). Variation between 96 well plates was normalized by running QC03LH3 on each plate and dividing the sample concentration by the QC03LH3 concentration on the plate to produce a relative sample concentration that

accounted for inter plate variability to facilitate among sample, between plate comparisons.

2.5. Statistical analyses

Statistics were performed using JMP 7 (SAS Institute Inc., Cary, North Carolina) and Microsoft Excel (Redmond, Washington). Data was first tested for normality using a Shapiro Wilk goodness of fit test and equal variance was tested with the Levene median test. Pearson's correlation analyses were carried out to determine if molar concentrations between Hg and Se were linearly associated within each tissue; and to determine whether Hg and Se correlated with protein oxidation levels within liver tissue. Analysis of variance (ANOVA) was used to analyze the relationship of trace element concentrations, molar ratios, and protein oxidation with the factors of age class, gender, and heart disease stage. Tukey honestly significant difference (HSD) tests and least squares (LS) mean plots were performed on statistically significant ($p < 0.05$) data to determine how the means varied within a factor.

3. Results and discussion

3.1. Concentrations of trace elements in liver and heart tissue

Summary data for Se, THg, and MeHg concentrations, % MeHg, and Hg:Se molar ratios determined in pygmy sperm whale liver and heart tissue are presented in Table 1. The US Environmental Protection Agency (USEPA) reference doses for THg and MeHg in edible fish tissue are 0.300 μ g/g, wet mass fraction and 0.100 μ g/g, wet mass, respectively (USEPA, 1999). All pygmy sperm whale liver and heart samples analyzed reflect THg and MeHg concentrations that were over USEPA action limits, which may reflect levels that are potentially hazardous to pygmy sperm whale health. Odontocetes are exposed to high levels of Hg, primarily in the MeHg form, through squid, crustacean, and fish consumption. Oceanic squid in the family Histioteuthidae comprise 80% of the pygmy sperm whale diet (Santos et al., 2006). In *Histioteuthis reversa*, MeHg represents 83% of the THg (0.015 ± 0.005 μ g/g, wet mass fraction) in whole squids (Bustamante et al., 2006). While Hg concentrations in squid may appear relatively low, dietary Hg exposure allows biomagnification up the food chain along with bioaccumulation by individuals yielding high Hg concentrations in pygmy sperm whales. Similar to pygmy sperm whales, squid is the major food item for long finned pilot whales (*Globicephala melas*) and short finned pilot whales (*Globicephala macrorhynchus*). Methyl Hg concentrations in *K. breviceps* liver tissues were comparable to both pilot whale species (Caurant et al., 1996; Bustamante et al., 2003). Many humans consume squid and the diet of Faroe islanders, indigenous people in the Arctic, and some people in Japan consists partly of marine mammal tissue, which poses possible health risks from the dietary intake of Hg mainly in the more toxic MeHg form (Myers and Davidson, 1998; Wagemann et al., 1998; Booth and Zeller, 2005; Endo et al., 2005; Bustamante et al., 2006).

Previous studies on trace elements in for *K. breviceps* include small data sets which primarily examined total element concentrations (Supplementary material, Table S2). The data set in this study is the largest and most comprehensive to date for *K. breviceps* assessing element speciation, and it is the first to present data for heart tissue. Pearson's correlation analysis indicated there were no correlations between heart and liver tissue concentrations of Se ($p = 0.422$, $r = 0.270$) or Hg ($p = 0.931$, $r = 0.030$), which was unexpected since internal organ correlations have been found for these elements in other odontocete species (Meador et al., 1999; Yang et al., 2002; Bryan et al., 2007). Selenium concentrations were

Table 1
Trace element concentrations ($\mu\text{g/g}$, wet mass fraction), % MeHg, and Hg:Se molar ratios in heart and liver from pygmy sperm whales.

	THg	MeHg	% MeHg	Se	Hg:Se
<i>Heart</i>					
<i>n</i>	11	11	11	11	11
Mean	1.409	1.786	92.761	2.389	0.242
SD	0.523	0.819	3.861	0.704	0.115
Range	0.301–2.482	0.248–3.360	81.950–95.840	1.405–3.792	0.084–0.522
<i>Liver</i>					
<i>n</i>	62	7	7	62	62
Mean	11.537	1.102	33.306	9.444	0.416
SD	10.627	0.507	14.019	4.399	0.266
Range	0.385–56.888	0.156–1.650	17.030–51.540	2.005–21.551	0.009–1.039

approximately 4 times greater in the liver than the heart, and THg concentrations were approximately 8 times greater in the liver than the heart. Methyl Hg reflected $92.761 \pm 3.861\%$ of the THg in heart tissue and showed no trend with increasing THg concentrations, while $33.306 \pm 14.019\%$ of THg in the liver is MeHg and percent MeHg exponentially decreased with increasing THg concentrations (Supplementary material, Fig. S3). At low THg concentrations the metal is mainly in the MeHg form and at greater THg concentrations demethylation is occurring resulting in iHg as the predominant form. Increasing THg concentrations in conjunction with decreasing percent MeHg verified that the liver is the site of Hg deposition. Similar exponential decrease trends in percent MeHg relative to increasing THg were additionally observed in pilot whale and striped dolphin (*Stenella coeruleoalba*) livers and the authors of these studies also concluded that the liver is the site of Hg demethylation and iHg storage (Palmisano et al., 1995; Caurant et al., 1996). Heart tissue was comparable to blood and skeletal muscle tissue in other cetacean species regarding THg concentrations, percent MeHg, and no relationship between percent MeHg and increasing THg. Since the heart is muscle tissue and pumps blood, it was anticipated that Hg concentrations and elemental species would behave similar to skeletal muscle and blood. Comparable to percent MeHg in *K. breviceps* heart tissue, bottlenose dolphins (*Tursiops truncatus*) were found to have $0.512 \pm 0.363 \mu\text{g/g}$ THg and 91% MeHg in blood (Bryan et al., 2005, 2007). In other studies that examined percentage MeHg to THg in liver and muscle tissue, pilot whale and beluga (*Delphin apterus leucas*) percent MeHg ranged from approximately 3–33% in liver and 78–97% in muscle (Caurant et al., 1996; Wagemann et al., 1998).

A strong positive correlation existed between Se and THg concentrations in liver ($p < 0.001$, $r = 0.773$) and no correlation was present between Se and THg concentrations in heart ($p = 0.498$, $r = 0.229$). Strong positive correlations between Se and Hg concentrations in pygmy sperm whale tissues were expected since this has been reported commonly in internal organs, such as liver and kidney, of other marine mammal species (Becker, 2000). The mean molar ratio of Hg:Se in liver was 0.416 ± 0.266 (0.009–1.039). Law et al. (2001) found a Hg:Se molar ratio of 0.32 in a one pygmy sperm whale liver, emphasizing the need for greater sampling effort. Many individuals fell close to having 1:1 molar ratios between Hg and Se in liver. The 1:1 molar ratio relationship between Hg and Se in liver was first noted in marine mammals in 1973 by Koeman et al. and has since been observed in several marine mammal species trace element studies (Koeman et al., 1973; Becker, 2000; Scheuhammer et al., 2008). In *K. breviceps* liver, Hg:Se molar ratios increased in individual animals as a function of increasing THg molar masses as illustrated in Fig. 1. Studies on fish eating bird species have shown onset of neurological and birth defects once the Hg:Se molar ratio exceeded 1 (Scheuhammer et al., 2008). To date, deleterious health impacts have not been reported in marine mammals




Fig. 1. Relationship between THg molar mass and Hg:Se molar ratio in livers of pygmy sperm whales, partitioned as a function of gender and age class.

in conjunction with high Hg concentrations suggesting that marine mammals have developed a mechanism to deal with high Hg in take. Selenium aids in detoxifying Hg by forming mercury selenide crystals which could limit Se bioavailability for seleno amino acid formation, protein synthesis, and metalloprotein binding. Seleno proteins, selenoenzymes, and Se binding proteins play important roles in antioxidant biochemistry. Unlike many enzymes that can be recycled once used, each time a selenoenzyme is used it must be broken down and synthesized all over again making the Se accessible to bind to free Hg. Selenium bioavailability may be restricted in the presence of high Hg concentrations measured in pygmy sperm whales. Some studies suggest that all Hg binds to Se and the quantity of non Hg bound Se can be calculated by the molar difference of the two elements ($\text{Se}_{\text{nmol/g}} - \text{Hg}_{\text{nmol/g}}$) (Drasch et al., 2000; Falnoga et al., 2006). Pygmy sperm whales with an Hg:Se molar ratio greater than 1 had negative values for the quantity of non Hg bound Se signifying that all bioavailable Se may be bound to Hg. Total trace element concentrations do not give much insight into the actual mechanisms or pathways by which Se and Hg interact and how the Hg sequestering mechanism influences Se bioavailability and biochemistry, but allow associations to be made between concentration data, animal life history information, and heart disease state.

3.2. Protein oxidation in pygmy sperm whale liver

Proteins are a key target of pro oxidants and free radicals. Immunoblot was used to assess overall protein oxidation since this method allowed immunodetection and quantification of carbonyl groups that were introduced to proteins which were modified by oxidation. Oxidation ratio differences were not detected in correlation to Se or Hg concentrations. Total free GSH and GSSG assays measured antioxidants in a specific pathway that aid in protecting

cells from pro oxidants and free radicals at the level of interception (Arteel and Sies, 2001). Glutathione can function to detoxify xenobiotics and this correlation was examined with Hg and Se concentrations in pygmy sperm whale liver. Correlations were not observed between Hg concentration and GSH states, which may be a limitation of the assay only measuring free GSH and not other pools of GSH such as that bound to proteins. While this study could not examine the GPx activity in *K. breviceps* liver tissue, GPx produces GSSG during the reduction of hydroperoxides and GSSG is reduced to two GSH molecules. Activity of GPx is thought to be directly proportional to the rate of GSSG produced (Reed, 1990). Glutathione peroxidase contains the seleno amino acid selenocysteine in its polypeptide chain. Although not statistically significant, there is a positive correlation between Se concentration and GSSG ($p = 0.064$, $r = 0.349$) in pygmy sperm whale liver (Supplementary material, Fig. S4). Humans suffering from chronic liver disease have been found to have positive and highly significant correlations between Se concentrations and GPx activities and the study concluded that these correlations may be well correlated with GSH and Se accessibility (Czuczajko et al., 2003).

The liver is secondarily affected in pygmy sperm whales with CMP and hepatic congestion was observed in these animals. Examining protein oxidation status in the liver relative to CMP was of key interest for this study since the liver is the primary organ for selenoprotein synthesis and xenobiotic uptake, detoxification, and storage or excretion. To date, there is no literature available that examines oxidative stress markers in relationship to diseases affecting marine mammals. Effects of dietary fat intake on anti oxidative state have been studied in captive bottlenose dolphins. Animals fed a high fat fish diet resulted in blood serum lipid peroxidase levels that were significantly higher than animals fed a low fat fish diet and the study concluded that decreased antioxidant states may be strongly influenced by high amounts of PU FAs and fat in diet (Kasamatsu et al., 2001). The mainly squid diet of pygmy sperm whales is relatively low fat (approximately 2% total lipid) when compared to a fish diet, however squid have PUFA levels greater than most fish species (Kirsch et al., 1998; Iversen et al., 2002; Recks and Seaborn, 2008). Pygmy sperm whales are potentially exposed to oxidative stress by ingesting large amounts of PUFAs and Hg. PUFAs are prone to oxidation because of multiple double bonds that rapidly transform into epoxides. Mercury promotes free radical formation through univalent redox reactions and has a very high affinity for thiol groups, which can lead to oxidative stress and lipid peroxidation (Ganther, 1980). Antioxidants, such as selenoproteins, glutathione, and GPx, defend against oxidants by prevention, interception, and repair of oxidative damage. Oxidative stress can impair protein function, damage DNA, and damage membrane lipids which can lead onset set of disease (Arteel and Sies, 2001).

3.3. Trace element concentrations and protein oxidation in relation to age class, gender, and heart disease stage

3.3.1. Age class

Animals were grouped into age classes based on total body length and observations of sexual maturity. Pygmy sperm whale adults can range in age from 4 to 22 years old, which may mask some statistical patterns due to the broad range of ages in the adult age class. Adult animals have THg and Se concentrations in liver that are significantly greater ($p < 0.0001$) than younger age classes. Selenium concentrations in liver are greater in calves and subadults than THg concentrations and this pattern reverses in adult animals (Fig. 2). Mercury increases little with age in fast growing calf and subadult pygmy sperm whales because of growth dilution, while in adults as growth slows Hg continuously bioaccumulates, prey are larger in size, and quantities of food ingested increase.




Fig. 2. Mean (+SE) THg and Se concentrations in pygmy sperm whale liver, partitioned as a function of age class. Significantly different mean Hg and Se concentrations between age classes are denoted with an asterisk ($p < 0.0001$).

Mercury has been shown to accumulate in liver in relationship to age in bottlenose dolphin (Meador et al., 1999). In heart tissue, Se concentrations are greater in adults than calves ($p = 0.151$) and THg concentrations are significantly greater in adults than calves ($p = 0.016$). When heart tissue concentrations were compared within an age class, Se is greater than THg in calves and adult pygmy sperm whales, which does not reflect the pattern that was observed in adult liver tissue. As a comparison, significant differences in Se concentrations between age classes were not observed in bottlenose dolphins (Bryan et al., 2007). Age plays a role in the molar relationship between Hg and Se in liver. Adult animals bioaccumulate higher concentrations of Hg, their Hg:Se molar ratios are significantly greater ($p < 0.0001$) than younger age classes, and their Hg:Se molar ratio approaches one (Fig. 1). Age class differences were not observed for protein oxidation.

3.3.2. Gender

Full factorial ANOVA showed no significant differences in Se ($p = 0.590$) and Hg ($p = 0.097$) concentrations in liver between adult males and females. Statistically significant differences between genders have also not been observed for Se concentrations in bottlenose dolphins and humans; and Hg concentrations in humans (Drasch et al., 2000; Bryan et al., 2007). In adult *K. breviceps*, Hg:Se molar ratios were greater in males than females ($p = 0.072$) (Fig. 3A). Bottlenose dolphin and pilot whale females, more specifically lactating pilot whale females, exhibit greater Hg concentrations and Hg:Se molar ratios in liver tissue than males (Caurant




Fig. 3. Mean (±SE) (A) Hg:Se molar ratios in livers of adult females ($n = 22$) and males ($n = 22$) and (B) immunoblot oxidation ratios in livers of females ($n = 13$) and males ($n = 17$).

et al., 1996; Meador et al., 1999; Bryan et al., 2007). Adult females are thought to have greater bioaccumulation of Hg than adult males due to increased dietary consumption to keep up with the energy demands of gestation and lactation (Bryan et al., 2007). The same gender pattern was expected to prevail in pygmy sperm whale females, especially since they reach sexual maturity at a younger age of approximately 3.5 years old than bottlenose dolphin females, which become sexually mature at 5–10 years old. Since age class differences were not observed for protein oxidation, all age classes were included in statistical analyses of gender and protein oxidation. Immunoblot oxidation ratio gender patterns were similar to Hg:Se molar ratio gender patterns (Fig. 3). While not statistically significant ($p = 0.160$), males have a greater mean oxidation ratio than females. Total GSH means were not different between male and females.

3.3.3. Heart disease stage

Only a subset of 5 of the 11 heart samples measured for Hg and Se content possessed the complementary histological information needed to assign a heart disease score, preventing statistical comparisons of trace element concentrations in heart tissue relative to heart disease stage. A heart score was assigned for 36 of 62 liver samples. Animals that have NPF have significantly lower ($p = 0.004$) Se concentrations and lower ($p = 0.094$) Hg concentrations than animals affected with MCD and CMP. Selenium concentrations increase in conjunction with increasing Hg concentrations and both relate to MCD and CMP in pygmy sperm whales. While not statistically significant ($p = 0.236$), animals with MCD and CMP have greater liver Hg:Se molar ratios than animals with NPF (Fig. 4A). Mercury concentrations drive the overall Hg:Se ratio when Hg is present in greater concentrations than Se. Selenium may be bound to Hg to detoxify Hg and is no longer bioavailable in heart disease affected animals.

Of the 30 pygmy sperm whale livers analyzed for protein oxidation, heart scores were able to be assigned to 21 individual animals. Although not statistically significant ($p = 0.453$), animals affected with CMP have greater mean immunoblot oxidation ratios than animals with NPF or MCD (Fig. 4B, and supplemental Fig. S1). Gender may be interrelated with heart disease stage in adult animals, since prevalence of CMP was greater in males and males had both greater Hg:Se and oxidation ratios (Fig. 3), which were both elevated in animals with CMP (Fig. 4). Overall, stage of CMP progression did not relate to reduced or oxidized glutathione concentrations (data not shown); these measurements provide complementary information about selenium specific oxidation pathways, and not an overall protein oxidation status. These findings point out that it is difficult to determine whether oxidative stress directly or indirectly plays a role in CMP progression. Other studies have pointed out that redox equilibrium is essential to bio-




Fig. 4. Mean (\pm SE) (A) Hg:Se molar ratios (NPF ($n = 15$), MCD ($n = 13$), CMP ($n = 8$)) and (B) immunoblot oxidation ratios (NPF ($n = 8$), MCD ($n = 9$), CMP ($n = 4$)) in pygmy sperm whale livers as a function of heart disease stage.

logical systems and that imbalance due to oxidative stress or swinging the pendulum too far in the direction of reductive stress can result in similar deleterious health effects (Rajasekaran et al., 2007).

Pygmy sperm whales could possibly be a species that are more susceptible to developing CMP and environmental factors put them at greater risk for disease onset. Pygmy sperm whales may perhaps serve as an indicator species to tie human health concerns back with how trace elements and oxidative stress can play a role in CMP. Myocardial degeneration was observed in subadult and adult pygmy sperm whales, while CMP was only observed in adult animals. In human epidemiological studies, cardiac disease usually presented in adulthood (Virmani, 2004). Cardiomyopathy was more prevalent in adult male pygmy sperm whales and males were found to have greater Hg:Se molar ratios and immunoblot oxidation ratios, which may relate to prevalence of CMP. Human men are 2–3 times more likely than women to develop dilated CMP, and men are affected 1–1.5 times more frequently than women with hypertrophic CMP (Virmani, 2004). In human population studies examining benefits of fish consumption with heart disease risk, findings showed that greater Hg exposure reduces benefits of fish consumption relative to heart disease risks (Rissanen et al., 2000). Many marine mammal studies have put forth the idea that Se status may be impacted by sequestration chemistry wherein Se binds Hg in the process of detoxifying Hg. This study represents an initial attempt to study Hg speciation and detoxification chemistry in marine mammals and its impact on development of CMP, but a range of complementary studies needs to be performed to ascertain the ultimate biochemical impact of this detoxification mechanism and its role in CMP progression and more generally, in antioxidant and Se biochemistry.

Disclaimer

Certain commercial products and instruments are identified in this paper to adequately specify the experimental procedures. Such identification does not imply recommendation or endorsement by the National Institute of Standards and Technology. Nor does it imply that the items mentioned are the best for the intended purpose.

Acknowledgments

Collectors from the Southeast Regional Stranding Program are appreciated for their time and effort in responding to strandings and collecting pygmy sperm whale samples. Michelle Fleetwood (AFIP), Kenny Kroell (HBOI), and David Rotstein (U of TN) were extremely helpful in gathering histology slides and reports. Lynn Thorsell and Ed Wirth (HML/NOS/NOAA) are thanked for use of instrumentation for total mercury measurements. Members of the Woodley Laboratory (HML/NOS/NOAA) are thanked for use of equipment and assistance with cryogenic homogenization. Rebecca Pugh, Amanda Moors, and Michael Ellisor were key to sample banking and cryogenic homogenization. Teresa Rowles of NOAA is thanked for providing support for these studies under the Marine Mammal Health and Stranding Response Program.


Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.chemosphere.2012.05.051>.

References

Arteel, G.E., Sies, H., 2001. The biochemistry of selenium and the glutathione system. *Environmental Toxicology and Pharmacology* 10, 153–158.

- Baker, M.A., Cerniglia, G.J., Zaman, A., 1990. Microtiter plate assay for the measurement of glutathione and glutathione disulfide in large numbers of biological samples. *Analytical Biochemistry* 190, 360–365.
- Bartfay, W.J., Hou, D., Brittenham, G.M., Bartfay, E., Sole, M.J., Lehotay, D., Liu, P.P., 1998. The synergistic effects of vitamin E and selenium in iron-overloaded mouse hearts. *Canadian Journal of Cardiology* 14, 937–941.
- Becker, P.R., 2000. Concentration of chlorinated hydrocarbons and heavy metals in Alaska Arctic marine mammals. *Marine Pollution Bulletin* 40, 819–829.
- Becker, P.R., Porter, B.J., Mackey, E.A., Schantz, M.M., Demiralp, R., Wise, S.A., 1999. National Marine Mammal Tissue Bank and Quality Assurance Program: protocols, inventory, and analytical results. NISTIR6279. USDOC, National Institute of Standards and Technology, Gaithersburg, MD.
- Berini, C., 2009. Pygmy sperm whale (*Kogia breviceps*, De Blainville 1838) strandings along the Atlantic coast of the southeastern United States: analysis of correlation with environmental factors. Grice Marine Biology Program. College of Charleston, Charleston, pp. 97.
- Booth, S., Zeller, D., 2005. Mercury, food webs, and marine mammals: implications of diet and climate change for human health. *Environmental Health Perspectives* 113, 521–526.
- Bossart, G.D., 2006. Marine mammals as sentinel species for oceans and human health. *Oceanography* 19, 134–137.
- Bossart, G.D., Hensley, G., Goldstein, J.D., Kroell, K., Manire, C.A., Defran, R.H., Reif, J.S., 2007. Cardiomyopathy and Myocardial Degeneration in Stranded Pygmy (*Kogia breviceps*) and Dwarf (*Kogia sima*) Sperm Whales. *Aquatic Mammals* 33, 214–222.
- Bossart, G.D., Odell, D.K., Altman, N.H., 1985. Cardiomyopathy in Stranded Pygmy and Dwarf Sperm Whales. *Journal of the American Veterinary Medical Association* 187, 1137–1140.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantification of microgram quantities of protein using the principle of protein dye binding. *Annals of Biochemistry* 72, 248–254.
- Bryan, C.E., Christopher, S.J., Balmer, B.C., Wells, R.S., 2007. Establishing baseline levels of trace elements in blood and skin of bottlenose dolphins in Sarasota Bay, Florida: implications for non-invasive monitoring. *Science of the Total Environment* 388, 325–342.
- Bryan, C.E., Christopher, S.J., Davis, W.C., Day, R.D., Hohn, A.A., Wells, R.S., 2005. Establishing baseline trace element and methylmercury concentrations for bottlenose dolphins in Sarasota Bay, Florida as an indicator of health status. In: 16th Biennial Conference on the Biology of Marine Mammals, San Diego, CA.
- Bustamante, P., Garrigue, C., Breaud, L., Caurant, F., Dabin, W., Greaves, J., Dodemont, R., 2003. Trace elements in two odontocete species (*Kogia breviceps* and *Globicephala macrorhynchus*) stranded in New Caledonia (South Pacific). *Environmental Pollution* 124, 263–271.
- Bustamante, P., Lahaye, V., Durnez, C., Churlaud, C., Caurant, F., 2006. Total and organic Hg concentrations in cephalopods from the North Eastern Atlantic waters: influence of geographical origin and feeding ecology. *Science of the Total Environment* 368, 585–596.
- Caurant, F., Navarro, M., Amiard, J.-C., 1996. Mercury in pilot whales: possible limits to the detoxification process. *Science of the Total Environment* 186, 95–104.
- Christopher, S.J., Day, R.D., Bryan, C.E., Turk, G.C., 2005. Improved calibration strategy for measurement of trace elements in biological and clinical whole blood reference materials via collision-cell inductively coupled plasma mass spectrometry. *Journal of Analytical Atomic Spectrometry* 20, 1035–1043.
- Czuczajko, J., Zachara, B.A., Staubach-Topczewska, E., Halota, W., Kedziora, J., 2003. Selenium, glutathione and glutathione peroxidases in blood of patients with chronic liver diseases. *Acta Biochimica Polonica* 50, 1147–1154.
- Davis, W.C., Christopher, S.J., Pugh, R.S., Donard, O.F.X., Krupp, E.A., Point, D., Horvat, M., Gibicar, D., Kljakovic-Gaspic, Z., Porter, B.J., Schantz, M.M., 2007. Certification of methylmercury content in two fresh-frozen reference materials: SRM 1947 Lake Michigan fish tissue and SRM 1974b organics in mussel tissue (*Mytilus edulis*). *Analytical and Bioanalytical Chemistry* 387, 2335–2341.
- Drasch, G., Mailander, S., Schlosser, C., 2000. Content of non-mercury-associated selenium in human tissues. *Biological Trace Element Research* 77, 219–230.
- Endo, T., Haraguchi, K., Hotta, Y., Hisamichi, Y., Lavery, S., Dalebout, M.L., Baker, C.S., 2005. Total mercury, methyl mercury, and selenium levels in the red meat of small cetaceans sold for human consumption in Japan. *Environmental Science and Technology* 39, 5703–5708.
- Eyer, P., Podhradský, D., 1986. Evaluation of the micromethod for determination of glutathione using enzymatic cycling and Ellman's reagent. *Analytical Biochemistry* 153, 57–66.
- Falnoga, I., Tusek-Znidaric, M., Stegnar, P., 2006. The influence of long-term mercury exposure on selenium availability in tissues: an evaluation of data. *Biometals* 19, 283–294.
- Freeman, L.M., Brown, D.J., Rush, J.E., 1998. Antioxidant status in dogs with idiopathic dilated cardiomyopathy. *Journal of Nutrition* 128, 2768S–2770S.
- Ganther, H.E., 1980. Interactions of vitamin E and selenium with mercury and silver. *Annals of the New York Academy of Sciences* 355, 212–226.
- Ichihara, S., Yamada, Y., Ichihara, G., Kanazawa, H., Hashimoto, K., Kato, Y., Matsushita, A., Oikawa, S., Yokota, M., Iwase, M., 2006. Attenuation of oxidative stress and cardiac dysfunction by bisoprolol in an animal model of dilated cardiomyopathy. *Biochemical and Biophysical Research Communications* 350, 105–113.
- Iverson, S.J., Frost, K.J., Lang, S.L.C., 2002. Fat content and fatty acid composition of forage fish and invertebrates in Prince William Sound, Alaska: factors contributing to among and within species variability. *Marine Ecology-Progress Series* 241, 161–181.
- Kannan, M., Wang, L., Kang, Y.J., 2004. Myocardial oxidative stress and toxicity induced by acute ethanol exposure in mice. *Experimental Biology and Medicine* 229, 553–559.
- Kasamatsu, M., Tsunokawa, M., Taki, M., Higuchi, H., Nagahata, H., 2001. Serum lipid peroxide and alpha-tocopherol concentrations and superoxide dismutase activity in captive bottle-nosed dolphins. *American Journal of Veterinary Research* 62, 1952–1956.
- Kennedy, S., Rice, D., Davidson, W., 1987. Experimental myopathy in vitamin E- and selenium-depleted calves with and without added dietary polyunsaturated fatty acids as a model for nutritional degenerative myopathy in ruminant cattle. *Research Veterinary Science* 43, 384–394.
- Kirsch, P.E., Iverson, S.J., Bowen, W.D., Kerr, S.R., Ackman, R.G., 1998. Dietary effects on the fatty acid signature of whole Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences* 55, 1378–1386.
- Koeman, J.H., Peeters, W.H.M., Koudstaa, Ch, Tjioe, P.S., Goeij, J., 1973. Mercury–Selenium correlations in marine mammals. *Nature* 245, 385–386.
- Law, R.J., Bennett, M.E., Blake, S.J., Allchin, C.R., Jones, B.R., Spurrier, C.J.H., 2001. Metals and organochlorines in pelagic cetaceans stranded on the coasts of England and Wales. *Marine Pollution Bulletin* 42, 522–526.
- Meador, J.P., Ernest, D., Hohn, A.A., Tilbury, K., Gorzelany, J., Worthy, G., Stein, J.E., 1999. Comparison of elements in bottlenose dolphins stranded on the beaches of Texas and Florida in the Gulf of Mexico over a one-year period. *Archives of Environmental Contamination and Toxicology* 36, 87–98.
- Myers, G.J., Davidson, P.W., 1998. Prenatal methylmercury exposure and children: neurologic, developmental, and behavioral research. *Environmental Health Perspectives* 106, 841–847.
- Palmisano, F., Cardellicchio, N., Zamboni, P.G., 1995. Speciation of mercury in dolphin liver: a two-stage mechanism for the demethylation accumulation process and role of selenium. *Marine Environmental Research* 40, 109–121.
- Pugh, R.S., Ellisor, M.B., Moors, A.J., Porter, B.J., Becker, P.R., 2007. Marine environmental specimen bank: clean room and specimen bank protocols. NISTIR7389. USDOC, National Institute of Standards and Technology, Gaithersburg, MD.
- Rajasekaran, N.S., Connell, P., Christians, E.S., Yan, L.J., Taylor, R.P., Orosz, A., Zhang, X.Q., Stevenson, T.J., Peshock, R.M., Leopold, J.A., Barry, W.H., Loscalzo, J., Odelberg, S.J., Benjamin, I.J., 2007. Human alpha B-crystallin mutation causes oxido-reductive stress and protein aggregation cardiomyopathy in mice. *Cell* 130, 427–439.
- Recks, M.A., Seaborn, G.T., 2008. Variation in fatty acid composition among nine forage species from a southeastern US estuarine and nearshore coastal ecosystem. *Fish Physiology and Biochemistry* 34, 275–287.
- Reed, D.J., 1990. Glutathione – toxicological implications. *Annual Review of Pharmacology and Toxicology* 30, 603–631.
- Rissanen, T., Voutilainen, S., Nyyssonen, K., Lakka, T.A., Salonen, J.T., 2000. Fish oil-derived fatty acids, docosahexaenoic acid and docosapentaenoic acid, and the risk of acute coronary events: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Circulation* 102, 2677–2679.
- Santos, M.B., Pierce, G.J., Lopez, A., Reid, R.J., Ridoux, V., Mente, E., 2006. Pygmy sperm whales *Kogia breviceps* in the Northeast Atlantic: New Information on Stomach Contents and Strandings. *Marine Mammal Science* 22, 600–616.
- Scheuhammer, A.M., Basu, N., Burgess, N.M., Elliott, J.E., Campbell, G.D., Wayland, M., Champoux, L., Rodrigue, J., 2008. Relationships among mercury, selenium, and neurochemical parameters in common loons (*Gavia immer*) and bald eagles (*Haliaeetus leucocephalus*). *Ecotoxicology* 17, 93–101.
- Tietze, F., 1969. Enzymic method for quantitative determination of nanogram amounts of total and oxidized glutathione: Applications to mammalian blood and other tissues. *Analytical Biochemistry* 27, 502–522.
- USEPA, 1999. Integrated risk information system (IRIS) on elemental mercury. Washington, DC: National Center for Environment Assessment, Office of Research and Development.
- Virmani, R., 2004. Pathology of Cardiomyopathies in Man. In: ACVP, ASVCP (Eds.), 55th Annual Meeting of the American College of Veterinary Pathologists (ACVP) & 39th Annual Meeting of the American Society of Clinical Pathology (ASVCP). International Veterinary Information Service, Middleton, WI.
- Wagemann, R., Trebacz, E., Boila, G., Lockhart, W.L., 1998. Methylmercury and total mercury in tissues of arctic marine mammals. *The Science of the Total Environment* 218, 19–31.
- Wells, R.S., Rhinehart, H.L., Hansen, L.J., Sweeney, J.C., Townsend, F.I., Stone, R., Casper, D.R., Scott, M.D., Hohn, A.A., Rowles, T.K., 2004. Bottlenose dolphins as marine ecosystem sentinels: developing a health monitoring system. *EcoHealth* 1, 246–254.
- Yang, Y., Kunito, T., Tanabe, S., Amano, M., Miyazaki, N., 2002. Trace elements in skin of Dall's porpoises (*Phocoenoides dalli*) from the northern waters of Japan: an evaluation for utilization as non-lethal tracers. *Marine Pollution Bulletin* 45, 230–236.
- Zeisler, R., Langland, J.K., Harrison, S.H., 1983. Cryogenic homogenization of biological tissues. *Analytical Chemistry* 55, 2431–2434.



Pathobiochemistry

Selenium protein identification and profiling by mass spectrometry: A tool to assess progression of cardiomyopathy in a whale model



Colleen E. Bryan^{a,*}, Gregory D. Bossart^b, Steven J. Christopher^a, W. Clay Davis^a,
Lisa E. Kilpatrick^c, Wayne E. McFee^d, Terrence X. O'Brien^e

^a Chemical Sciences Division, National Institute of Standards and Technology, Hollings Marine Laboratory, 331 Fort Johnson Road, Charleston, SC 29412, USA

^b Georgia Aquarium, 225 Baker Street NW, Atlanta, GA 30313, USA

^c Biomolecular Measurement Division, National Institute of Standards and Technology, 100 Bureau Drive, Gaithersburg, MD 20899, USA

^d Center for Coastal Environmental Health and Biomolecular Research, National Ocean Service, National Oceanic and Atmospheric Administration, 219 Fort Johnson Road, Charleston, SC 29412, USA

^e Department of Medicine, Division of Cardiology, Medical University of South Carolina, Carolina 25 Courtenay Dr. ART 7063 and the Office of Research and Development, Ralph H. Johnson Veterans Administration Medical Center, Charleston, SC 29425, USA

ARTICLE INFO

Keywords:

Selenium
Selenoprotein
Cardiomyopathy
Kogia breviceps
ICP-MS
LC-ESI-MS/MS

ABSTRACT

Non-ischemic cardiomyopathy is a leading cause of congestive heart failure and sudden cardiac death in humans and in some cases the etiology of cardiomyopathy can include the downstream effects of an essential element deficiency. Of all mammal species, pygmy sperm whales (*Kogia breviceps*) present the greatest known prevalence of cardiomyopathy with more than half of examined individuals indicating the presence of cardiomyopathy from gross and histo-pathology. Several factors such as genetics, infectious agents, contaminants, biotoxins, and inappropriate dietary intake (vitamins, selenium, mercury, and pro-oxidants), may contribute to the development of idiopathic cardiomyopathy in *K. breviceps*. Due to the important role Se can play in antioxidant biochemistry and protein formation, Se protein presence and relative abundance were explored in cardiomyopathy related cases. Selenium proteins were separated and detected by multi-dimension liquid chromatography inductively coupled plasma mass spectrometry (LC-ICP-MS). Se protein identification was performed by liquid chromatography electrospray tandem mass spectrometry (LC-ESI-MS/MS), and Se protein profiles were examined in liver (n = 30) and heart tissue (n = 5) by SEC/UV/ICP-MS detection. Data collected on selenium proteins was evaluated in the context of individual animal trace element concentration, life history, and histological information. Selenium containing protein peak profiles varied in presence and intensity between animals with no pathological findings of cardiomyopathy and animals exhibiting evidence of cardiomyopathy. In particular, one class of proteins, metallothioneins, was found to be associated with Se and was in greater abundance in animals with cardiomyopathy than those with no pathological findings. Profiling Se species with SEC/ICP-MS proved to be a useful tool to identify Se protein pattern differences between heart disease stages in *K. breviceps* and an approach similar to this may be applied to other species to study Se protein associations with cardiomyopathy.

1. Introduction

This study focuses on how selenium (Se) is associated with a primary dilated cardiomyopathy in pygmy sperm whales, *Kogia breviceps*. Studying cardiomyopathy in pygmy sperm whales is of particular interest since greater than 50% of the whales that strand are affected by this disease. Determining the etiology may be relevant to humans since the clinically the cause(s) of non ischemic cardiomyopathy commonly cannot be determined, even in cases that lead to end stage heart failure or heart transplantation [1,2]. The natural occurrence of

cardiomyopathy in *K. breviceps* is greater than all other mammal species studied to date warranting further examination of factors that are associated with the onset of cardiomyopathy in this whale species. The presence of cardiomyopathy in pygmy sperm whales was first identified and described in 1985 by Bossart et al. [3]. Cardiomyopathy in *K. breviceps*, as in human, is a myocardial disease that results in deterioration of cardiomyocyte number, size and function, and diminished left and/or right ventricular function. Pygmy sperm whales present pathology of a mixed form comprising dilated and hypertrophic cardiomyopathy [1].

* Corresponding author at: NIST—Hollings Marine Laboratory, 331 Ft. Johnson Road, Charleston, SC, 29412 USA.
E-mail address: colleen.bryan@nist.gov (C.E. Bryan).

<http://dx.doi.org/10.1016/j.jtemb.2017.05.005>

Received 14 September 2016; Received in revised form 18 May 2017; Accepted 19 May 2017
0946-672X/ Published by Elsevier GmbH.

Selenium is an essential micronutrient for animals that is primarily acquired through diet. The major biological role of Se is in antioxidant defense, but it can be toxic at high concentrations [4]. Total Se concentration in blood reflects total Se status in an individual indicating whether Se concentrations are balanced, toxic, or deficient. However, total Se concentration provides little information about the element's bioactivity since the biological function of Se is primarily mediated by incorporation or association with proteins. The liver is the predominant site of seleno amino acid formation, Se protein synthesis, and excretion, therefore, making liver one of the tissues of choice for which to study Se speciation [5,6]. Additionally, the liver is affected secondarily in late stage cardiomyopathy through hepatic congestion.

Etiology of cardiomyopathy in pygmy sperm whales is currently idiopathic. Nutritional deficiencies of Se have been shown in humans, mice, dogs, and cattle to play roles in development and progression of cardiomyopathy [7–11]. Humans that suffer from Keshan disease have congestive cardiomyopathy caused by a combination of dietary deficiency of selenium and presence of Coxsackievirus or patients that are on a ketogenic diet to treat intractable epilepsy are often nutritionally deficient in Se and either experience cardiomyopathy or are at greater risk for heart disease development [9,12,13]. There have been relatively few studies on altered selenoprotein synthesis and research addressing whether Se roles in cardiomyopathy are correlative or causative. Many factors may contribute to development and progression of cardiomyopathy, such as genetics, infectious agents, biotoxins, and dietary intake (vitamins, Se, Hg, pro oxidants). These factors could act singly or additively and may be interconnected to one another. Other studies have shown that multiple dietary factors can interact, such as deficiency of vitamin E and Se, to further progression of myocardium degeneration [14–16].

Many prior marine mammal studies have put forth the idea that Se status may be impacted by sequestration chemistry wherein Se binds Hg in the process of detoxifying Hg [17–19]; however the ultimate biochemical impact of this detoxification mechanism has been less studied. To properly address Se bioavailability and determine whether Se activity is truly deficient due to Hg robbing *K. breviceps* of bioactive Se leading to onset of cardiomyopathy, presence of selenoenzymes, selenoproteins, and Se containing proteins must first be characterized in terms of abundance and form. Here, methods have been developed for the complex matrices of pygmy sperm whale liver and heart tissue to extract, purify, detect, and identify Se proteins. This study seeks to utilize mass spectrometry to identify changes in Se protein presence and abundance within animals exhibiting various states of cardiomyopathy.

2. Materials and methods

2.1. Sample collection

Since 1998, the National Institute of Standards and Technology (NIST) has cryogenically banked liver tissue from stranded individual *K. breviceps* in the National Marine Mammal Tissue Bank (NMMTB) housed at the National Institute of Standards and Technology (NIST), Charleston, SC. Liver samples obtained for the NMMTB were collected, processed, and frozen by trained field collectors during animal necropsies and stored in the specimen according to NISTIR 6279 [20]. Frozen heart samples were available from some of the same individual animals that had liver samples in the NMMTB and were donated for use in this project by the Center for Coastal Environmental Health and Biomolecular Research/National Ocean Service/National Oceanographic and Atmospheric Administration (CCEHBR/NOS/NOAA), Charleston, SC. Heart tissue samples for analytical analyses were placed in polypropylene centrifuge tubes and stored at -80°C . Fresh heart tissue that was collected at the necropsy for histology preparation and histopathological evaluation was fixed in 10% neutral buffered formalin and tissue was sectioned at $5\ \mu\text{m}$ for slide preparation. Individual animal life history and gross pathology data for this project was

generated by necropsy principal investigators at collaborating institutions.

The liver for QC03LH3 pygmy sperm whale liver homogenate was collected in 1994 from a female (MMES9469SC 8) that stranded and tissues were donated to NIST, Charleston, SC by CCEHBR/NOS/NOAA for use in making an interlaboratory comparison exercise control material. QC03LH3 was used throughout method development and for Se protein identification in *K. breviceps* due to sample abundance and integrity of sample collection, homogenization, and storage.

2.2. Heart disease stage assignment

Hematoxylin and eosin (H&E) stained histology slides for each animal case were evaluated independently and blind of protein and chemical analyses data by a veterinary pathologist and assigned a heart score according to criteria put forth by Bossart et al. [1] and the Dallas cardiomyopathy criteria in humans [21]. Heart histology and gross pathology was used to categorize cardiomyopathy progression in the following three stages: no pathological findings (NPF), myocardial degeneration (MCD), and cardiomyopathy (CMP). Descriptions of each stage are provided in the Supplementary materials along with histology images.

2.3. Sample preparation for trace element and protein analysis

Fresh frozen tissue was cryogenically homogenized to produce a uniform sample composition of fresh frozen powder for analysis. Liver samples were homogenized in class 100 clean room conditions at NIST, Charleston, SC using cryogenic procedures developed by Zeisler et al. and Pugh et al. [22,23]. Left ventricle heart tissue was cryogenically homogenized using a bench top 6850 Freezer/Mill (SPEX SamplePrep, Metuchen, NJ). Samples were placed in vials with a stainless steel impactor, capped, placed in the mill, submerged in LN_2 , and shaken at 10 Hz for 3 min. Homogenized powder was transferred into sterile polypropylene jars (Nalge Nunc International, Rochester, NY) and stored at -80°C until analysis.

RIPA lysis buffer (Pierce, Rockford, IL) with 1X concentration Halt™ protease inhibitor cocktail and 5 mM ethylenediamine tetracetic acid (EDTA) was placed in 6 mL aliquots into polypropylene centrifuge tubes. Approximately 0.5 g homogenized sample was weighed into each centrifuge tube with buffer solution and vortexed. The centrifuge tubes were laid on ice and rocked for 15 min. The tubes were then centrifuged for 3 min at 1500g. The supernatant was pulled off in 1.5 mL aliquots and placed into protein LoBind microcentrifuge tubes (Eppendorf, Hauppauge, NY). A Microfuge 22R (Beckman Coulter, Fullerton, CA) was used to centrifuge microcentrifuge tubes for 15 min at 14,000g and 4°C . The supernatant from each microcentrifuge tube was filtered with a $0.2\ \mu\text{m}$ PTFE membrane filter (SunSri, Rockwood, TN) into a glass auto sampler vial. All samples were kept on ice between steps to prevent protein degradation. Protein extraction was controlled and optimized by protein concentration measurements to ensure that tissue cells were thoroughly lysed. Total protein concentration was measured at 595 nm according to the Bradford method [24] using a Coomassie Plus assay kit (Pierce) and a DU800 spectrophotometer (Beckman Coulter). Bovine serum albumin (Pierce) was used as the calibration standard.

2.4. Analytical methods

2.4.1. Instrumentation

Total Se mass fraction measurements were made using a Thermo Electron X Series II ICP MS (Bremen, Germany) with a standard low volume glass impact bead spray chamber (Peltier cooled at $+3^{\circ}\text{C}$), concentric glass nebulizer, and operating in collision cell mode utilizing 8% H_2 in 92% He as the collision cell gas. Detailed total Se measurement methods and statistical analyses were described by Bryan et al. [25]. Fig. 1 outlines the multiple steps and instrumentation used for Se




Fig. 1. Flow chart of selenium species separation, detection, and identification methods. 1st dimension LC – size exclusion chromatography (SEC); 2nd dimension LC – strong anion exchange (SAX).

separation, detection, and identification. Liquid chromatography coupled to UV spectrophotometer and Thermo Elemental X7 quadrupole ICP MS (Winsford, UK) (LC/UV/ICP MS) were used for ^{80}Se separation, detection, and fraction collection. A DX 600 (Dionex, Sunnyvale, CA) ion chromatography system consisting of GP50 gradient pump, AS autosampler (cooled to 4 °C) (150 μL injection loop), and AD25 UV/Vis absorbance detector (set at 280 nm) was used for chromatographic separation. The chromatographic system was coupled to the ICP MS by 0.010 in. internal diameter (id) PEEK tubing. Once proteins were detected by the UV absorbance detector and presence of Se was identified by ICP MS, the chromatographic system was uncoupled from the ICP MS and the chromatographic system was set in line with a fraction collector (Foxy 200, ISCO, Lincoln, NE) to collect protein fractions. Liquid chromatography coupled to a Surveyor LC pump and autosampler and LTQ ESI MS/MS (ThermoFisher, Waltham, MA) were utilized for peptide sequencing.

2.4.2. Instrument calibration methods and quality assurance

Collision cell ICP MS working conditions were optimized with a 10 ng/g 68 element tuning solution (High Purity Standards, North Charleston, SC) and a Se calibrant prior to sample analysis and coupling to other instruments. LTQ ESI MS/MS signal intensity was optimized using ThermoFisher Xcalibur software tune methods with angiotensin (Sigma, Sigma Aldrich, St. Louis, MO). Commercial availability of selenoprotein standards is limited to GPx1 from bovine erythrocytes (84.5 kDa) and TrxR from rat liver (55 kDa 67 kDa) (Sigma Aldrich). The glutathione peroxidase standard along with procedural blanks was taken through the same analysis steps as unknown pygmy sperm whale samples. QC03LH3 and GPx standard were used throughout method development and sample analyses for method validation, reproducibility, and protein identification. The GPx standard was used to verify protein recovery along each chromatographic separation and verify that

the selenoprotein could be properly identified by peptide sequence.

2.4.3. Protein separation

Protein separation steps are outlined in Fig. 1. The 1st dimension of protein separation was carried out by LC, protein presence was detected by UV/Vis, and Se containing peaks were detected by ICP MS. Then Se containing fractions were collected from each of the 1st dimension Se containing protein peak elution times for 2nd dimension LC protein separation, ICP MS Se detection, and fraction collection when Se species began to elute. Mobile phase compositions, chromatographic programs, and sample injection volumes are outlined in the Supplementary materials (Table S1) for each LC separation. Liquid chromatography first dimension separation was carried out on a size exclusion column (SEC) that had an effective separation range of 1 kDa to 300 kDa since the column has notable tolerance of complex matrices. Strong anion exchange liquid chromatography (SAX/UV/ICP MS) was used for second dimension separation additional clean up of Se species. Replicate injections and LC separations were performed in order to collect multiple fractions from the same sample. Strong anion exchange liquid chromatography (SAX) Se containing protein peak were then combined and pre concentrated using Centrifugal Filter tubes (Millipore, Billerica, MA) which have a 3 kDa molecular weight cutoff. An Avanti J 20 XPI centrifuge (Beckman Coulter) was used to spin down samples for approximately 20 min at 7000g and 4 °C. Pre concentrated sample that did not pass through the filter and contained proteins greater than 3 kDa was removed from the top chamber of the tube and placed in protein LoBind microcentrifuge tubes for measurement of total protein concentration or tryptic digestion.

2.4.4. Protein identification

Selenium containing protein fractions collected required protein alkylation and tryptic digestion in order to prevent formation of disulfide bridges and to obtain peptides of a suitable length for identification by LC ESI MS/MS. Bradford assay was used to estimate total protein concentration in each protein fraction in order to decide how much sample was required for digestion. Protein was weighed out and 0.2% RapiGest (Waters, Milford, MA) in 50 mmol/L Tris buffer (pH 8.0) along with 200 mmol/L 1, 4 dithio DL threitol (DTT) (Fluka, Sigma Aldrich) solution was added to the protein. Samples were heated at 60 °C for 30 min followed by 30 min at 37 °C. Then 200 mmol/L iodoacetamide (Sigma, Sigma Aldrich) solution was added and samples were incubated at room temperature in the dark with vortex mixing for 1 h. The alkylation reaction was stopped with the addition of 200 mmol/L DTT solution and samples were again incubated at room temperature in the dark with vortex mixing for 30 min. Trypsin (Promega, Madison, WI) was added to alkylated protein to achieve a ratio of trypsin:protein (g/g) of 1:50 to 1:100 and incubated for 20 h at 37 °C. Digestion was stopped and RapiGest was cleaved by adding 25.5% trifluoroacetic acid (TFA) (Supelco, Bellefonte, PA) to bring sample pH < 2 prior to incubation at 37 °C for 60 min. After incubation, 0.1% FA (aq) formic acid (Fluka, Sigma Aldrich) was added and samples were centrifuged at 14,000g for 10 min at 4 °C to precipitate RapiGest. The supernatant was removed and transferred to protein LoBind microcentrifuge tubes which were stored frozen (–20 °C) until analysis.

Reverse Phase Liquid chromatography electrospray ionization tandem mass spectrometry (RPLC ESI MS/MS) was used for peptide sequencing and protein identification. Mobile phase compositions, chromatographic program, and sample injection volume are outlined in the Supplementary materials (Table S2) for chromatographic separation of peptides. SEQUEST software (ThermoFisher, Waltham, MA) was used to compare peptide fragmentation patterns to theoretical fragmentation of peptides. The FASTA database was created by downloading all currently known protein sequences that contain the amino acid seleno cysteine (U) from all species in the UniProtKB/Swiss Prot database (downloaded from www.expasy.org on 06/10/08) since there is no




Fig. 2. SEC/UV/ICP-MS chromatogram (^{80}Se intensity) of QC03LH3 pygmy sperm whale liver homogenate overlaid with UV absorption profile. Numbers indicate Se containing protein peaks.

protein database available for pygmy sperm whales. In addition, a custom made database was created that contained FASTA sequences from known selenoproteins and Se containing protein in humans and bovines. A list of proteins included in the custom database can be found in the Supplementary materials (Table S3). Scoring and cutoff criteria were set in SEQUEST for database searches to decide that a MS/MS data set indicated the presence of a protein in the sample. Monoisotopic precursor and fragment mass types were allowed along with two missed cleavage sites, cysteine alkylation, and methionine oxidation. The mass range of the peptides was 200 m/z to 6000 m/z . The XCORR versus charges state values were set at 1.50, 2.50, and 3.00; and the delta CORR cutoff value was 0.100. Maximum peptide probability was set at 0.05. This probability limit minimized the false positive rate of randomly matching a peptide in the database.

2.4.5. Selenium profiling

Liver samples from animals in the NMMTB, heart samples donated by CCEHBR/NOS/NOAA, and QC03LH3 were taken through sample preparation, separated by size exclusion chromatography, and Se species were detected by ICP MS in a single batch to eliminate variation in ICP MS response between batches and instrument drift was monitored over the course of the single sequence. Chromatograms were plotted for each sample to compare Se maximum peak intensities with total trace element concentrations and heart disease stage. Statistical analyses were performed using JMP 7 (SAS Institute Inc., Cary, NC) and Microsoft Excel (Redmond, Washington). Pearson's correlation analyses were carried out to determine if total trace element concentrations were linearly associated with individual Se peak maximum intensities. Analysis of variance (ANOVA) was used to analyze the relationship between individual Se peak maximum intensities and heart disease stage. Variations between heart disease stage mean maximum intensity within a peak were examined with least squares (LS) mean plots.

3. Results

3.1. Quality assurance

A single Se containing protein peak for the GPx1 standard from bovine erythrocytes had a retention time on the SEC column of approximately 9.5 min (Supplementary materials Fig. S7). The GPx1

standard from bovine erythrocytes had 97.56% protein sequence coverage with bovine GPx1 and a list of matched peptides with their probability scores can be found in Supplementary materials (Table S4). As a selenoprotein, glutathione peroxidase uniquely contains the selenoamino acid selenocysteine, which is coded for in a peptide sequence as a "U". The active site of GPx contains the selenocysteine residue that carries out the catalytic function of redox reactions [26]. The seleno cysteine containing selenopeptide GKVLLENVASLUGTTVR was identified in the GPx standard from bovine erythrocytes (Supplementary materials Fig. S8).

3.2. Selenium protein separation and identification

Sample separation has been identified as the limiting step in Se speciation of complex biological samples that contain a variety of Se proteins. Selenium in proteins containing selenoamino acids is covalently bound therefore the metal is less likely to separate from the protein of interest during protein separation and clean up steps. However, Se is not covalently bound in proteins that transport Se or Se small molecules and is more easily lost during some preparations, such as SDS PAGE, resulting in loss of detection of presence of these proteins by ICP MS. Liquid chromatography coupled to ICP MS was the most effective tool for separating intact Se proteins and retaining their native composition during Se separation and Se detection. Two dimensions of liquid chromatography were used for further separation and purification of Se proteins in the complex matrix of QC03LH3 pygmy sperm whale liver homogenate to improve protein identification with tandem mass spectrometry.

Fig. 2 shows a SEC/UV/ICP MS chromatogram of the pygmy sperm whale liver homogenate QC03LH3. Proteins and smaller molecules elute from the size exclusion column according to molecular weight resulting in larger proteins eluting first before smaller proteins. There were three prominent protein peaks containing Se from QC03LH3 that were separated by the size exclusion column. The UV peaks coinciding with Se peaks shows the absorbance for all the proteins that elute at the same time, but the selenium proteins were the only ones of interest in this study. The large Se peak with a retention time of approximately 16 min contained both organic and inorganic small molecule Se species and was not evaluated in this study.

Tandem mass spectrometry was used to identify proteins that were

Table 1
QC03LH3 Se protein peptide matches, species, % protein sequence coverage, and peptide probability (P) value.

Protein	Species	% protein sequence coverage	Peptides	P value
Glutathione peroxidase 1	Bovine	54.63%	CEVNGEK	8.32E-05*
			SAAALAAAAPR	1.38E-04
			NEEILNCLK	5.70E-05*
			FITWSPVCR	3.56E-04*
			FLVGPDPGVPVR	1.22E-05
			FLVGPDPGVPVRR	5.88E-05*
			GLVVLGFPCNQFGHQENAKNEEILNCLK	1.06E-11
			YVRPGGGFEPNFMLFEK	5.13E-07
			FLTIDIEPDIETLLSQGASA	8.23E-04
			AHPLFAFLR	3.24E-06
	Human	30.35%	GLVVLGFPCNQFGHQENAK	5.56E-11*
			VLLIENVASLUGTTVR	2.93E-09*
			YVRPGGGFEPNFMLFEK	2.62E-09*
			NDVAWNFEK	3.79E-07*
			NDVAWNFEK	9.04E-06
			DYTEMNDLQKR	5.34E-07
			EALPTPSDDATALMTDPK	1.09E-10
Common marmoset	4.48%	YVRPGGGFEPNFMLFQK	1.44E-05	
		Mouse	5.47%	
		Pig	8.74%	
		Rabbit	8.50%	
		Bovine	8.11%	
Glutathione S-transferase A1	Bovine	7.62%	AILNYIATKYNLGKDMK	4.83E-09
			FQDGDLTLYQSNAILR	4.85E-09
Glutathione S-transferase P	Long-tailed hamster	7.62%	FEDGDLTLYQSNAILR	5.63E-07
			Pig	5.31%
Glyceraldehyde-3-phosphate dehydrogenase	Bovine	37.24%	PPYTTYFPVVR	1.35E-04
			AENGLKLVINGK	2.47E-03*
			LEKPAKYDEIKK	3.50E-05*
			GAAQNIIPASTGAAK	1.09E-07*
			VVDLMVHMASKE	7.73E-05*
			IVSNASCTTNCLAPLAK	1.11E-09*
			ATIFQERDPANIK	3.51E-04
			VPTPNVSVVDLTCR	4.77E-04*
			LTGMAFRVPTPNVSVVDLTCR	3.09E-05*
			RVIISAPSADAPMFVMGVNHEK	8.69E-05
			WGDAAGAEYVVESTGVFTTMEK	5.00E-13*
			VKVGVNGFGR	4.73E-05*
			LISWYDNEFGYSNR	2.93E-06*
			RVIISAPSADAPMFVMGVNHEK	2.36E-05*
			WGDAAGAEYVVESTGVFTTMEK	9.63E-10
Glyceraldehyde-3-phosphate dehydrogenase	Human	20.00%	GASDKCSCCA	2.19E-06
			CAQGCVCCK	9.66E-04
			CAQGCVCCKGASDK	1.85E-06
			CAQGCVCCKGASDKCSCCA	2.65E-13
Metallothionein-1A	Greater Egyptian jerboa	5.79%	GASDKCSCCA	3.89E-07*
			Bovine	29.51%
Metallothionein-2	Human	16.39%	SCCSCCPAECEK	1.24E-05*
			Metallothionein-3	17.65%
Metallothionein-3	Wild yak	17.65%	SCCSCCPAECEK	4.34E-06
			Metallothionein-4	29.03%
Phospholipid hydroperoxide glutathione peroxidase	Dog	29.03%	CAQGCICKGGSDKCSCCA	2.87E-08
			Human	6.09%
Selenium-binding protein 1	Bovine	5.30%	YGPMEEPLVIEK	5.14E-06*
			GGPVQVLEDEQELK	1.26E-06
Selenium-binding protein 1	Human	7.84%	TKLLPLSLISSR	8.77E-07*
			GGPVQVLEDEELK	6.94E-06*
			LTGQLFLGGSIVK	3.66E-07
			EEIVYLPDIYR	4.62E-06*
			FLIATGERPR	2.79E-04*
			AGQPLQLLDASWYLPK	2.81E-08
Thioredoxin reductase 1	Human	1.54%	ALVSAQWVAEALR	7.53E-05*
			3-mercaptopyruvate sulfurtransferase	9.76%

* P value was calculated from the custom made human and bovine protein database since peptides associated with these values were not identified from the SwissProt database.

present in QC03LH3 pygmy sperm whale liver from selenium containing protein fractions separated by liquid chromatography and detected by ICP MS. Only QC03LH3 and GPx1 standard from bovine erythrocytes were taken through the complete sequence of steps for protein identification. Table 1 provides percent sequence coverage in each protein and a list of peptides matched with the respective probability scores. Peptide matches used for identification of Se proteins in pygmy sperm whales were made assuming that a high degree of protein homology exists between species because the proteome for *K. breviceps* has not been sequenced. Protein matches between pygmy sperm whales and species for which protein databases were available indicated that peptide sequences for many proteins are likely highly conserved between species. Some Se protein sequences in *Kogia* spp. could be greatly different from those sequenced in other animals therefore preventing identification of some Se proteins in pygmy sperm whales. Additionally,

the likelihood of false positive identifications is increased. Low abundance Se proteins that contribute to the ⁸⁰Se intensity may not have been identified since peptides for these proteins were not detectable at such low concentrations relative to greater abundance proteins. Several Se proteins were identified by a single peptide MS/MS spectrum match and should be verified by complementary means. Western blots with antibody against selenoproteins and Se containing proteins would be the next step in protein verification.

Selenium proteins were identified in the three prominent SEC/ICP MS protein peaks containing Se from QC03LH3 pygmy sperm whale liver homogenate (Fig. 2). Greater molecular weight selenoproteins, selenium containing, and selenium binding proteins were found in peaks 1 and 2; and proteins that fall into each of these Se protein classifications are identified in Table 2. Metallothioneins were identified in peak 3, which are small low molecular weight selenium binding

Table 2
Properties and functions of Se proteins identified in QC03LH3 pygmy sperm whale liver.

Protein	Protein Class	Species	Sequence Length	MW (kDa)	Known function
Glutathione peroxidase 1	Selenoprotein	Bovine	205 AA	22.7	protects hemoglobin in red blood cells from oxidative breakdown
		Human	201 AA	21.9	
		Common marmoset	201 AA	21.8	
		Mouse	201 AA	22.3	
		Pig	206 AA	22.6	
		Rabbit	200 AA	21.9	
Glutathione S-transferase A1	Se-binding	Bovine	222 AA	25.5	conjugation of reduced glutathione to hydrophobic electrophiles; conjugation of reduced glutathione to hydrophobic electrophiles; aids in detoxification with xenobiotic metabolism
Glutathione S-transferase P	Se-binding	Bovine	210 AA	23.6	
		Long-tailed hamster	210 AA	23.6	
Glyceraldehyde-3-phosphate dehydrogenase	Se-binding	Pig	207 AA	23.5	involved in metabolic switch under oxidative stress allowing cells to produce more NADPH
		Bovine	333 AA	35.9	
		Human	335 AA	36.1	
		Greater Egyptian jerboa	363 AA	39.4	
Metallothionein-1A	Se-binding	Bovine	61 AA	6.0	high cysteine residue content binds heavy metals
Metallothionein-2	Se-binding	Human	61 AA	6.0	high cysteine residue content binds heavy metals
Metallothionein-3	Se-binding	Human	68 AA	6.9	binds heavy metals
		Wild yak	68 AA	6.9	
Metallothionein-4	Se-binding	Dog	62 AA	6.2	binds heavy metals
Phospholipid hydroperoxide glutathione peroxidase	Selenoprotein	Human	197 AA	22.2	protects cells against membrane lipid peroxidation and oxidative damage
Selenium-binding protein 1	Se-binding	Bovine	472 AA	52.6	selenium-binding protein; involved in sensing reactive xenobiotics in the cytoplasm
		Human	472 AA	52.4	
Serum albumin	Se-containing	Bovine	607 AA	69.3	most abundant protein in plasma, regulates colloidal osmotic pressure of blood, acts as a plasma carrier by non-specific binding
		Human	609 AA	69.4	
Thioredoxin reductase 1	Selenoprotein	Human	649 AA	70.9	reduces thioredoxin using NADPH
3-mercaptopyruvate sulfurtransferase	Se-binding	Human	297 AA	33.2	transfers sulfur-containing groups (Se can substitute) to thiol compounds; participates in cysteine metabolism

proteins. Select LC ESI MS/MS peptide spectra of Se proteins identified from SEC peaks after SAX can be found in the Supplementary materials. Table 2 highlights the properties and functions of Se protein classes (selenoproteins, Se containing proteins, and Se binding proteins) identified in QC03LH3 pygmy sperm whale liver. Serum albumin was included in Table 2 since this protein can contain selenomethionine, transport small Se molecules, and was identified in several sample fractions. Serum albumin is a high abundance protein in many tissues, including liver, making it difficult to eliminate during sample preparation.

3.3. Selenium profiling in pygmy sperm whale liver and heart tissue

Selenium species profiling was performed along with total Se and Hg concentration measurements on 30 frozen liver samples from the NMMTB and 5 frozen heart samples donated by CCEHBR/NOS/NOAA. Heart disease stages were assigned from histologic heart tissue preparations to complement 21 animals with liver samples from the NMMTB and for all 5 animal heart samples. Total Se ($9.19 \pm 4.00 \mu\text{g/g}$, wet mass fraction) was positively correlated with the magnitude of certain Se protein species peaks in pygmy sperm whale livers. The maximum peak heights (cps) for peaks 2 and 3 increase significantly as total Se concentrations increase (Fig. 3B and C). In contrast, maximum peak heights for peak 1 do not change in relationship to increasing total Se concentrations (Fig. 3A). Fig. 4 shows a representative comparison between the SEC/ICP MS chromatogram for liver samples from individuals at different stages of heart disease progression. Selenium protein profiles differ in liver in relationship to stage of cardiomyopathy progression. To assess the differences in Se protein peak height in intensities between heart disease stages, mean individual peak intensities were calculated at each disease state as a percentage of the total sum of

the three peak height intensities for that stage (Fig. 5). Animals with no pathological findings had Se protein percentages among peaks that were equivalent. Whales with cardiomyopathy had the greatest peak 3 intensities when compared to other heart disease stages. Animals with myocardial degeneration and cardiomyopathy had greater peak 3 intensities than peak 1 intensities. While peak 2 intensities increased significantly (ANOVA, $p = 0.020$) with increasing total Se, the peaks mean percentage remained constant relative to heart disease stage. Pygmy sperm whales with NPF have significantly (ANOVA, $p = 0.020$) lower total Se concentrations ($(7.159 \pm 1.255) \mu\text{g/g}$, wet mass fraction) than animals with MCD ($(10.502 \pm 1.183) \mu\text{g/g}$, wet mass fraction) or CMP ($(11.759 \pm 1.775) \mu\text{g/g}$, wet mass fraction) [25]. Fig. 6 shows a comparison between the SEC/ICP MS chromatograms for all heart samples at different stages of heart disease progression. Peak patterns in heart tissue were similar to liver tissue while ^{80}Se count rates were an order of magnitude lower in heart tissue than liver tissue (Figs. 2 and 6). The heart tissue chromatograms have several small Se peaks and have inferior peak resolution compared to liver tissue possibly due to poorer sample collection and storage conditions that may have allowed protein degradation, relative to specimens preserved using strict NMMTB protocols. The number and quality of heart tissue samples prevented further statistical analyses.

4. Discussion and conclusion

Cardiomyopathy in pygmy sperm whales is a chronic, progressive disease in which varying degrees of cardiac degeneration occur leading to the terminal state of advanced cardiomyopathy [1]. This appears to be the first study to assess Se protein profiles with LC/UV/ICP MS in this mammalian species affected by a non ischemic cardiomyopathy. For whales with MCD and CMP, Se peak patterns showed that low




Fig. 3. Effect of the total selenium concentration on the intensity (cps) of selenium species in individual peaks in pygmy sperm whale livers ($n = 30$). Relationship between maximum peak intensity and total Se concentration for peaks 1 (A), 2 (B), and 3 (C). Data points are fitted with linear trend lines and data was collected in a single day.

molecular weight proteins, such as metallothionein, were in greater abundance than animals with NPF. Once more, the relative abundance of high molecular weight Se proteins increased parallel to increasing total Se concentration. These findings may be a model for Se related non ischemic cardiomyopathy in humans.

These Se associated proteins, glutathione peroxidase, selenium binding protein, and metallothioneins, have critical functions in the protection from oxidative damage, metal detoxification, detecting xenobiotics, and binding xenobiotics. These protective roles are important to pygmy sperm whales because these animals are continuously exposed to oxidative stress and contaminants in the marine environment [25,27].

4.1. Glutathione peroxidase 1

The protective effect of GPx is of particular importance when an organism is under oxidative stress [5]. Glutathione peroxidase 1 is a cellular or cytosolic enzyme that prevents lipid peroxidation of cell

membranes by reducing pro oxidants such as hydrogen peroxide and organic hydroperoxide consequently protecting cells from oxidative damage [28]. The selenoprotein glutathione peroxidase 1 protein was identified in QCO3LH3 by several separation schemes utilizing single (SEC) and two dimensional separations (SEC and SAX). Glutathione peroxidase is the most extensively studied selenoprotein for which protein functions and structure have been widely characterized and this protein has been identified in many animal species [4]. While GPx activity has been studied in tissues from bottlenose dolphins (*Tursiops truncatus*) [29,30] and ringed seals (*Pusa hispida*) [31], this is the first study to separate and identify GPx1 at the protein level in a marine mammal species.

4.2. Selenium binding protein 1

Selenium binding protein 1 (SBP1) was identified in QCO3LH3 pygmy sperm whale liver and selenium binding proteins may act in sensing reactive xenobiotics in the cytoplasm [32]. Pygmy sperm whales are exposed to PCBs and high concentrations of PCBs have been measured in many other marine mammal species [33]. Rats that have been exposed to toxic coplanar polychlorinated biphenyls (PCBs) have shown up regulation of selenium binding protein [32]. Exposure to chemicals known to be peroxisome proliferators, such as dibutyl phthalate, Wy 14 643, and ciprofibrate, have been shown in a mouse model to decrease abundance of selenium binding proteins [34]. Studies of the effects of different chemicals illustrate that regulation of selenium binding protein expression may be chemical dependent. Porat et al. [35] have suggested that selenium binding protein mediates the intracellular transport of Se. Selenium deficiency may limit SBP1 expression therefore reducing biological function of the protein.

4.3. Metallothioneins

Metallothioneins (MTs), which are Se binding proteins, were the only Se proteins identified in peak 3 of the size exclusion chromatography separation (Fig. 2). Additional low abundance Se proteins with similar molecular weights to MTs may have been present that minimally contributed to Se intensity in peak 3, but their peptides were undetectable for identification. Metallothioneins are low in molecular weight and rich in cysteine residues. Metals bind easily to MTs due to the thiol groups (SH) in the cysteine residues [36]. Selenium has a high binding affinity for cysteine. Metallothionein peptide fragments, identified in Se containing peak 3 and shown in Tables 1 and 2, contain several cysteine residues in each fragment. Metallothioneins are synthesized in a high capacity in tissues that uptake, store, and eliminate metals such as liver [37]. Both essential and toxic trace elements can induce MTs through chelation of cysteine residues. Metallothioneins act in maintaining homeostasis and detoxification by restricting availability of metal cations at harmful sites [36]. Metallothioneins have been proposed as biomarkers to assess marine organisms for exposure and impact of toxic metals in the marine ecosystem [37]. Kwohn et al. [38,39] were the first to isolate and identify MT1 and MT2 in striped dolphins (*Stenella coeruleoalba*). Metallothioneins have been detected in liver and associated with metals in several marine mammal species including sperm whales (*Physeter macrocephalus*), bottlenose dolphins, striped dolphins, pilot whales (*Globicephala melas*), narwhals (*Monodon monoceros*), Dall's porpoises (*Phocoenoides dalli*), and California sea lions (*Zalophus californianus*) [40–43].

Since the Se binding proteins metallothionein were the only Se proteins identified in peak 3, increased relative abundance of MTs suggest their potential utility as a biomarker for onset of early stages of heart disease leading to cardiomyopathy in pygmy sperm whales. Metallothioneins have been proposed in other mammal studies as biomarkers of exposure for metal pollution [41] and findings in this study could lead to another application of MTs as biomarkers for non ischemic cardiomyopathy. Metallothioneins have been suggested in





Fig. 4. Pygmy sperm whale liver SEC/ICP-MS ⁸⁰Se profiles for a representative individual at each heart disease stage; no pathological findings (NPF), myocardial degeneration (MCD), cardiomyopathy (CMP). Inset shows protein peak region of the SEC/ICP-MS chromatogram.



Heart Disease Stage	Peak 1	Peak 2	Peak 3
NPF (n = 8)	31.3 ± 6.2 %	34.4 ± 3.2 %	34.3 ± 4.0 %
MCD (n = 9)	24.0 ± 5.9 %	34.7 ± 3.1 %	41.3 ± 3.8 %
CMP (n = 4)	21.4 ± 8.8 %	35.4 ± 4.6 %	43.2 ± 5.6 %

Fig. 5. Pygmy sperm whale liver Se protein distribution in each protein peak as a percentage (mean ± SE) of the sum of the three Se containing protein peak heights relative to heart disease stage (n = number of whales at disease stage); no pathological findings (NPF), myocardial degeneration (MCD), cardiomyopathy (CMP).

playing a cardioprotective role by regulating metal homeostasis and anti oxidant response [44–46]. Recently, a human study found that individuals with the MT1A genetic polymorphism are predisposed to developing cardiovascular disease when there is an imbalance between oxidant production and antioxidant defenses [47]. In our study, MT1A was specifically identified in pygmy sperm whales liver. Variations in Se protein profiles in tissues of pygmy sperm whales at different heart disease stages may lend insight into how Se protein presence and relative abundance changes throughout this type of cardiomyopathy disease progression.

4.4. Selenium concentration

Since peaks 2 and 3 increased in intensity relative to total Se concentration increase in liver, there could be similar Se incorporation mechanisms between proteins eluting in these peaks. Total Se and Hg concentrations have been shown to be closely positively correlated in liver. However, total Hg (10.452 ± 8.744 µg/g, wet mass fraction) does not have a significant correlation with the concentration of Se protein species in individual peaks in pygmy sperm whale livers (total Hg measurements were discussed in Bryan et al. [25]). This may be due

to hydrogen selenide aiding in detoxifying methyl Hg and binding to inorganic Hg forming mercury selenide (HgSe) crystals, which are small inert inorganic molecules that are stored in the liver [48]. Thus, there appears to be sequestering of the bioavailable Se pool for Hg detoxification rather than for interactions involving protein formation.

4.5. Selenium species profiling in pygmy sperm whale liver and heart

Relative peak heights of Se protein species in Se profiles were related to cardiomyopathy progression in pygmy sperm whales. Differences in Se protein distribution among tissues and sub cellular fractions have previously been identified and suggested that these proteins are involved in several metabolic pathways [49,50]. Further biological importance of Se proteins was recognized when preferential routing of Se for formation of specific Se proteins was discovered with insufficient Se intake [49]. Selenium protein profile differences between animals with no pathological findings and animals affected by cardiac disease indicate a potential altered metabolic pathway of protein homeostasis. Peak 2 Se protein relative intensities do not change as a function of heart disease progression suggesting that Se proteins in this peak are not related to heart disease and that peak 2 Se protein abundance increases as a consequence of total Se concentration increasing. Peak 1 intensities are not affected by total Se concentration therefore the Se proteins found in this peak could actually remain stable throughout heart disease progression even though peak 1 appears to be of greater intensity in NPF animals (Fig. 4). Given that total Se concentrations are greater in whales with CMP, accumulation of total Se in conjunction with increases in peak 3 intensities may indicate that in the presence of heart disease correlated Se proteins found in peak 3 are up regulated and involved in the progression of the disease state.

4.6. Selenoproteins and human heart failure

Selenoproteins are required for normal cardiac health, in particular with regards to oxidative stress, which can be associated with progressive human chronic heart failure [51]. The role of micronutrients in heart failure is also being increasingly recognized. With wider




Fig. 6. SEC/ICP-MS ^{80}Se profiles for 5 individual pygmy sperm whale hearts with different heart disease stages; no pathological findings (NPF), myocardial degeneration (MCD), cardiomyopathy (CMP).

applications in higher cardiac risk populations of various forms of bariatric surgery and supplemental nutrition it becomes increasingly important to understand the pathophysiology of co factors such as Se [52]. This is in addition to the understudied role of environmental exposures such as Se and Hg in the development or progression of heart failure. For example, a reversible dilated cardiomyopathy secondary to Se deficiency has been long recognized (i.e., Keshan disease) [53].

Profiling Se species with SEC/ICP MS was a useful tool in identifying differences in Se protein containing peak patterns between stages of dilated cardiomyopathy disease progression. For whales with MCD and CMP, Se peak patterns showed that low molecular weight proteins were in greater abundance than animals with NPF. Relative abundance of high molecular weight Se proteins increased parallel to total Se concentration increasing. Protein identification and profiling was the first step in gaining insight to how selenium proteins are related to cardiomyopathy in pygmy sperm whales.

Many of the factors that can contribute to onset and progression of cardiomyopathy in pygmy sperm whales may not stand alone but rather act collectively and require further investigation. Methods developed and used in this study to identify and profile Se proteins in *K. breviceps* at various stages of cardiomyopathy progression could be applied to other species that are affected by cardiomyopathy to gain further insight into disease progression and the role of Se in non ischemic cardiomyopathy.

Disclaimer

Certain commercial products and instruments are identified in this paper to adequately specify the experimental procedures. Such identification does not imply recommendation or endorsement by the National Institute of Standards and Technology. Nor does it imply that the items mentioned are the best for the intended purpose.

Acknowledgements

Collectors from the Southeast Regional Stranding Program are appreciated for their time and effort in responding to strandings and collecting pygmy sperm whale samples. Michelle Fleetwood, Kenny

Kroell, and David Rotstein were extremely helpful in gathering histology slides and reports. Members of the Woodley Laboratory (HML/NOS/NOAA) are thanked for use of equipment. Rebecca Pugh, Amanda Moors, and Michael Ellisor were key to sample banking and cryogenic homogenization. Guillaume Ballihaut was instrumental in teaching some of the protein separation techniques. Teresa Rowles of NOAA is thanked for providing support for these studies under the Marine Mammal Health and Stranding Response Program (Permit No. 932 1905 00/MA 009526).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jtemb.2017.05.005>.

References

- [1] G.D. Bossart, G. Hensley, J.D. Goldstein, K. Kroell, C.A. Manire, R.H. Defran, J.S. Reif, Cardiomyopathy and myocardial degeneration in stranded pygmy (*Kogia breviceps*) and dwarf (*Kogia sima*) sperm whales, *Aquat. Mamm.* 33 (2) (2007) 214–222.
- [2] J.A. Towbin, N.E. Bowles, The failing heart, *Nature* 415 (6868) (2002) 227–233.
- [3] G.D. Bossart, D.K. Odell, N.H. Altman, Cardiomyopathy in stranded pygmy and dwarf sperm whales, *J. Am. Vet. Med. Assoc.* 187 (11) (1985) 1137–1140.
- [4] G.E. Arteel, H. Sies, The biochemistry of selenium and the glutathione system, *Environ. Toxicol. Pharmacol.* 10 (4) (2001) 153–158.
- [5] D. Behne, A. Kyriakopoulos, Mammalian selenium-containing proteins, *Annu. Rev. Nutr.* 21 (2001) 453–473.
- [6] P.D. Whanger, Metabolism of selenium in humans, *J. Trace Elem. Exp. Med.* 11 (2–3) (1998) 227–240.
- [7] L.M. Freeman, D.J. Brown, J.E. Rush, Assessment of degree of oxidative stress and antioxidant concentrations in dogs with idiopathic dilated cardiomyopathy, *J. Am. Vet. Med. Assoc.* 215 (5) (1999) 644–646.
- [8] W.J. Bartfay, D. Hou, G.M. Brittenham, E. Bartfay, M.J. Sole, D. Lehotay, P.P. Liu, The synergistic effects of vitamin E and selenium in iron-overloaded mouse hearts, *Can. J. Cardiol.* 14 (7) (1998) 937–941.
- [9] A.G.C. Bergqvist, C.M. Chee, L. Lutchka, J. Rychik, V.A. Stallings, Selenium deficiency associated with cardiomyopathy: a complication of the ketogenic diet, *Epilepsia* 44 (4) (2003) 618–620.
- [10] L.M. Freeman, D.J. Brown, J.E. Rush, Antioxidant status in dogs with idiopathic dilated cardiomyopathy, *J. Nutr.* 128 (12) (1998) 2768S–2770S.
- [11] S. Hara, Y. Shoji, A. Sakurai, K. Yuasa, S. Himeno, N. Imura, Effects of selenium deficiency on expression of selenoproteins in bovine arterial endothelial cells, *Biol. Pharmacol. Bull.* 24 (7) (2001) 754–759.

- [12] L.H. Foster, S. Sumar, Selenium in health and disease: a review, *Crit. Rev. Food Sci. Nutr.* 37 (3) (1997) 211–228.
- [13] Y. Xia, K.E. Hill, R.F. Burk, Biochemical studies of a selenium-deficient population in China: measurement of selenium, glutathione peroxidase and other oxidant defense indices in blood, *J. Nutr.* 119 (9) (1989) 1318–1326.
- [14] M.A. Baker, G.J. Cerniglia, A. Zaman, Microtiter plate assay for the measurement of glutathione and glutathione disulfide in large numbers of biological samples, *Anal. Biochem.* 190 (2) (1990) 360–365.
- [15] S. Kennedy, D. Rice, Selective morphologic alterations of the cardiac conduction system in calves deficient in vitamin E and selenium, *Am. J. Pathol.* 130 (2) (1988) 315–325.
- [16] S. Kennedy, D. Rice, W. Davidson, Experimental myopathy in vitamin E- and selenium-depleted calves with and without added dietary polyunsaturated fatty acids as a model for nutritional degenerative myopathy in ruminant cattle, *Res. Vet. Sci.* 43 (1987) 384–394.
- [17] F. Caurant, M. Navarro, J.-C. Amiard, Mercury in pilot whales: possible limits to the detoxification process, *Sci. Total Environ.* 186 (1–2) (1996) 95–104.
- [18] J.H. Koeman, W.H.M. Peeters, a.Ch Koudsta, P.S. Tjioe, J. Goeij, Mercury-selenium correlations in marine mammals, *Nature* 245 (5425) (1973) 385–386.
- [19] A. Wang, D. Barber, C.J. Pfeiffer, Protective effects of selenium against mercury toxicity in cultured Atlantic spotted dolphin (*Stenella plagiodon*) renal cells, *Arch. Environ. Contam. Toxicol.* 41 (4) (2001) 403–409.
- [20] P.R. Becker, B.J. Porter, E.A. Mackey, M.M. Schantz, R. Demiralp, S.A. Wise, National Marine Mammal Tissue Bank and Quality Assurance Program: Protocols, Inventory, and Analytical Results. NISTIR6279, USDOC, National Institute of Standards and Technology, Gaithersburg, MD, 1999.
- [21] H.T. Aretz, Myocarditis – the Dallas criteria, *Hum. Pathol.* 18 (6) (1987) 619–624.
- [22] R.S. Pugh, M.B. Ellis, A.J. Moors, B.J. Porter, P.R. Becker, Marine Environmental Specimen Bank: Clean Room and Specimen Bank Protocols. NISTIR7389, USDOC, National Institute of Standards and Technology, Gaithersburg, MD, 2007.
- [23] R. Zeisler, J.K. Langland, S.H. Harrison, Cryogenic homogenization of biological tissues, *Anal. Chem.* 55 (1983) 2431–2434.
- [24] M.M. Bradford, A rapid and sensitive method for the quantification of microgram quantities of protein using the principle of protein dye binding, *Ann. Biochem.* 72 (1976) 248–254.
- [25] C.E. Bryan, W.C. Davis, W.E. McFee, C.A. Neumann, J. Schulte, G.D. Bossart, S.J. Christopher, Influence of mercury and selenium chemistries on the progression of cardiomyopathy in pygmy sperm whales, *Kogia breviceps*, *Chemosphere* 89 (5) (2012) 556–562.
- [26] K.T. Suzuki, Metabolomics of selenium: Se metabolites based on speciation studies, *J. Health Sci.* 51 (2) (2005) 107–114.
- [27] S. Booth, D. Zeller, Mercury, food webs, and marine mammals: implications of diet and climate change for human health, *Environ. Health Perspect.* 113 (5) (2005) 521–526.
- [28] J. Czuczajko, B.A. Zachara, E. Staubach-Topczewska, W. Halota, J. Kedziora, Selenium, glutathione and glutathione peroxidases in blood of patients with chronic liver diseases, *Acta Biochim. Pol.* 50 (4) (2003) 1147–1154.
- [29] M.D. Pine, K. Greer, D. Busbee, Comparison of reactive oxygen scavenging systems between a cetacean (DKN1) and a porcine renal epithelial cell line (LLC-PK1), *Comp. Biochem. Physiol. A-Mol. Integr. Physiol.* 147 (2) (2007) 550–555.
- [30] V. Woshner, K. Knott, R. Wells, C. Willetto, R. Swor, T. O'Hara, Mercury and selenium in blood and epidermis of bottlenose dolphins (*Tursiops truncatus*) from Sarasota Bay, FL: interaction and relevance to life history and hematologic parameters, *Ecohealth* 5 (3) (2008) 360–370.
- [31] J.P. Vazquez-Medina, T. Zenteno-Savin, R. Elsner, Antioxidant enzymes in ringed seal tissues: potential protection against dive-associated ischemia/reperfusion, *Comp. Biochem. Physiol. C-Toxicol. Pharmacol.* 142 (3–4) (2006) 198–204.
- [32] Y. Ishii, M. Hatsumura, T. Ishida, N. Ariyoshi, K. Oguri, Significant induction of a 54-kDa protein in rat liver with homologous alignment to mouse selenium binding protein by a coplanar polychlorinated biphenyl, 3,4,5,3',4'-pentachlorobiphenyl and 3-methylcholanthrene, *Toxicol. Lett.* 87 (1) (1996) 1–9.
- [33] P.R. Becker, E.A. Mackey, R. Demiralp, M.M. Schantz, B.J. Koster, S.A. Wise, Concentrations of chlorinated hydrocarbons and trace elements in marine mammal tissues archived in the U.S. national biomonitoring specimen bank, *Chemosphere* 34 (9–10) (1997) 2067–2098.
- [34] C.S. Giometti, X.L. Liang, S.L. Tollaksen, D.B. Wall, D.M. Lubman, V. Subbarao, M.S. Rao, Mouse liver selenium-binding protein decreased in abundance by peroxisome proliferators, *Electrophoresis* 21 (11) (2000) 2162–2169.
- [35] A. Porat, Y. Sagiv, Z. Elazar, A 56-kDa selenium-binding protein participates in intra-golgi protein transport, *J. Biol. Chem.* 275 (19) (2000) 14457–14465.
- [36] Y.J. Kang, Metallothionein redox cycle and function, *Exp. Biol. Med.* 231 (9) (2006) 1459–1467.
- [37] A. Sarkar, D. Ray, A.N. Shrivastava, S. Sarker, Molecular biomarkers: their significance and application in marine pollution monitoring, *Ecotoxicology* 15 (4) (2006) 333–340.
- [38] Y.T. Kwohn, A. Okubo, H. Hirano, H. Kagawa, S. Yamazaki, S. Toda, Primary structure of striped dolphin renal metallothionein-II, *Agric. Biol. Chem.* 52 (3) (1988) 837–841.
- [39] Y.T. Kwohn, S. Yamazaki, A. Okubo, E. Yoshimura, R. Tatsukawa, S. Toda, Isolation and characterization of metallothionein from kidney of striped dolphin, *Stenella coeruleoalba*, *Agric. Biol. Chem.* 50 (11) (1986) 2881–2885.
- [40] A. Decataldo, A. Di Leo, S. Giandomenico, N. Cardellicchio, Association of metals (mercury, cadmium and zinc) with metallothionein-like proteins in storage organs of stranded dolphins from the Mediterranean sea (Southern Italy), *J. Environ. Monit.* 6 (4) (2004) 361–367.
- [41] K. Das, V. Debacker, J.M. Bouqueneau, Metallothioneins in marine mammals, *Cell. Mol. Biol.* 46 (2) (2000) 283–294.
- [42] R. Wagemann, R. Hunt, J.F. Klaverkamp, Subcellular distribution of heavy metals in liver and kidney of a narwhal whale (*Monodon monoceros*): an evaluation for the presence of metallothionein, *Comp. Biochem. Physiol. Part C: Comp. Pharmacol.* 78 (2) (1984) 301–307.
- [43] T. Ikemoto, T. Kunito, Y. Anan, H. Tanaka, N. Baba, N. Miyazaki, S. Tanabe, Association of heavy metals with metallothionein and other proteins in hepatic cytosol of marine mammals and seabirds, *Environ. Toxicol. Chem.* 23 (8) (2004) 2008–2016.
- [44] A.F. Ceylan-Isik, P. Zhao, B.F. Zhang, X.Y. Mao, G.H. Su, J. Ren, Cardiac over-expression of metallothionein rescues cardiac contractile dysfunction and endoplasmic reticulum stress but not autophagy in sepsis, *J. Mol. Cell. Cardiol.* 48 (2) (2010) 367–378.
- [45] Q.J. Liu, G.J. Wang, G.H. Zhou, Y. Tan, X.L. Wang, W. Wei, L.C. Liu, W.L. Xue, W.K. Feng, L. Cai, Angiotensin II-induced p53-dependent cardiac apoptotic cell death: its prevention by metallothionein, *Toxicol. Lett.* 191 (2–3) (2009) 314–320.
- [46] G. Ye, N.S. Metreveli, J. Ren, P.N. Epstein, Metallothionein prevents diabetes-induced deficits in cardiomyocytes by inhibiting reactive oxygen species production, *Diabetes* 52 (3) (2003) 777–783.
- [47] R. Giacconi, S. Kanoni, P. Mecocci, M. Malavolta, D. Richter, S. Pierpaoli, L. Costarelli, C. Cipriano, E. Muti, F. Mangialasche, F. Piacenza, S. Tessei, R. Galeazzi, E.V. Theodoraki, F. Lattanzio, G. Dedoussis, E. Mocchegiani, Association of MT1A haplotype with cardiovascular disease and antioxidant enzyme defense in elderly Greek population: comparison with an Italian cohort, *J. Nutr. Biochem.* 21 (10) (2010) 1008–1014.
- [48] I. Falnoga, M. Tusek-Znidaric, P. Stegnar, The influence of long-term mercury exposure on selenium availability in tissues: an evaluation of data, *Biomaterials* 19 (3) (2006) 283–294.
- [49] D. Behne, H. Hilmert, S. Scheid, H. Gessner, W. Elger, Evidence for specific selenium target tissues and new biologically important selenoproteins, *Biochim. Biophys. Acta* 966 (1) (1988) 12–21.
- [50] D. Behne, S. Scheid, A. Kyriakopoulos, H. Hilmert, Subcellular-distribution of selenoproteins in the liver of the rat, *Biochim. Biophys. Acta* 1033 (3) (1990) 219–225.
- [51] M. de Lorgeril, P. Salen, Selenium and antioxidant defenses as major mediators in the development of chronic heart failure, *Heart Fail. Rev.* 11 (1) (2006) 13–17.
- [52] N.A. McKeag, M.C. McKinley, J.V. Woodside, M.T. Harbinson, P.P. McKeown, The role of micronutrients in heart failure, *J. Acad. Nutr. Diet.* 112 (6) (2012) 870–886.
- [53] K.Y. Ge, G.Q. Yang, The epidemiology of selenium deficiency in the etiologic study of endemic diseases in China, *Am. J. Clin. Nutr.* 57 (2) (1993) 259–263.

Review

Naturally Occurring Food Toxins

Laurie C. Dolan *, Ray A. Matulka and George A. Burdock

Burdock Group, 801 N. Orange Ave., Suite 710, Orlando FL 32801, USA;

E-Mails: rmatulka@burdockgroup.com (R.A.M.); gburdock@burdockgroup.com (G.A.B.)

* Author to whom correspondence should be addressed; E-Mail: ldolan@burdockgroup.com;
Tel.: +1-407-802-1400; Fax: +1-407-802-1405.

Received: 12 August 2010; in revised form: 2 September 2010 / Accepted: 13 September 2010 /

Published: 20 September 2010

Abstract: Although many foods contain toxins as a naturally-occurring constituent or, are formed as the result of handling or processing, the incidence of adverse reactions to food is relatively low. The low incidence of adverse effects is the result of some pragmatic solutions by the US Food and Drug Administration (FDA) and other regulatory agencies through the creative use of specifications, action levels, tolerances, warning labels and prohibitions. Manufacturers have also played a role by setting limits on certain substances and developing mitigation procedures for process-induced toxins. Regardless of measures taken by regulators and food producers to protect consumers from natural food toxins, consumption of small levels of these materials is unavoidable. Although the risk for toxicity due to consumption of food toxins is fairly low, there is always the possibility of toxicity due to contamination, overconsumption, allergy or an unpredictable idiosyncratic response. The purpose of this review is to provide a toxicological and regulatory overview of some of the toxins present in some commonly consumed foods, and where possible, discuss the steps that have been taken to reduce consumer exposure, many of which are possible because of the unique process of food regulation in the United States.

Keywords: toxin; natural; environmental; exposure; processing; cooking; food

1. Introduction

Historically, we have learned that everything is toxic; it is only the dose that separates the toxic from the non-toxic. Even water is toxic if a large amount (4–5 liters) is consumed in a relatively short

time (2–3 hours). The pathogenesis of water intoxication includes hyponatremia, followed by cerebral edema, seizures and death.

Like water, too much of a good thing such as the antioxidant vitamin A, can have acute toxic effects leading to hepatotoxicity [1] or chronic high levels can have a pro-oxidant effect [2]. Something as innocent as licorice, when consumed in large amounts may be harmful. For example, Bannister and associates reported hypokalemia leading to cardiac arrest in a 58-year-old woman who had been eating about 1.8 kg of licorice per week [3]. This licorice intoxication (dubbed “glycyrrhizism” after glycyrrhizic acid, the active component of licorice), has an effect resembling that of aldosterone, which suppresses the renin-angiotensin-aldosterone axis, resulting in the loss of potassium. Clinically, hypokalemia with alkalosis, cardiac arrhythmias, muscular symptoms together with sodium retention and edema, and severe hypertension are observed. The syndrome may develop at a level of 100 g licorice per day but gradually abates upon withdrawal of the licorice [4].

Recently, public health and social agendas have become more proactive in food toxicology, such as regulating (or outright banning) trans fats or “endocrine disruptors” in foods on the basis of public safety, including a suggestion of removing the generally recognized as safe (GRAS) status for salt [5]. These agendas lose sight of the basic principle of toxicology that “the dose makes the poison” and that demanding “safety *per se*” or “safe at any dose”, for all foods and ingredients is a non-starter and as a concept, was abandoned with the adoption of the Federal Food and Drug Act (FFDCA) in 1958. For their part, the regulators can limit amounts of potentially toxic substances allowed in food and in those circumstances where setting limits is not effective and public health policy makers provide the public with sufficient information (e.g., label information), where possible, to protect the consumer from reasonably foreseeable problems. Labeling requirements by the FDA provide the consumer with helpful information about content of fats, carbohydrate, protein, potential allergens, caloric value, *etc.*, but do not provide information about toxins that may be inherent in the foods or formed during processing. Because some food toxins cannot be removed from foods and others may be created during processing or cooking, consumption of small quantities of food toxins is unavoidable. The purpose of this review is to illustrate the potential risks of these toxins when consumed at concentrations normally present in foods and the steps taken by regulators to mitigate exposure where possible. Although regulatory information from countries other than the United States is included, FDA legislation is emphasized. Readers from other countries are advised to consult regulations for their specific region, because regulations and regulatory practices in other countries may differ from those in the United States.

2. Regulatory Accommodation

Foods are regarded as such because they are edible—they cannot be unpalatable or toxic—and; foods must have nutritional, hedonic or satietal value—otherwise there would be no point in consuming them. Therefore, in the absence of a spontaneous change or contamination, the concept of a toxic food *per se* would seem to be an oxymoron. How then, could a food be toxic and still be considered a food—there are two principal means: (1) an ordinarily non-toxic food has become toxic, if even for a small subpopulation; and (2) over-consumption of an ordinarily non-toxic food. This shift between toxic and non-toxic or toxic only for a select group has the potential for creating headaches for regulatory agencies charged with protecting the health of the public, but as the reader will see in the

following pages, the FDA and other regulatory agencies have created some thoughtful and pragmatic solutions for achieving a balance of acceptable risk and unavoidable circumstances.

The large diversity of acceptable foods made it difficult for the framers of the Federal Food Drug and Cosmetic Act (FFDCA) to define what a food could be, so they settled on the pragmatic definition provided in §201(f) [6]:

The term “food” means (1) articles used for food or drink for man or other animals, (2) chewing gum, and (3) articles used for components of any such article.

The framers are to be congratulated on their realistic approach, but a little interpretation is required. In the first clause “...articles used for food...” includes what humans and animals will eat as such (including eggs, meat, kohlrabi, Velveeta[®] cheese and angel food cake). The third clause “articles used for components of any such article,” are simply those substances used to make food (defined in the first clause)—therefore, anything approved for addition to food, becomes a part of the food. The second clause was more of a political consideration than anything else, as there was some disagreement whether chewing gum was swallowed or expectorated; the swallowers prevailed and chewing gum is regulated as a food. Had the majority determined that chewing gum was expectorated (as is evident on a sidewalk outside of any theater or church), it would have been classified with breath mints (which are not swallowed) and are therefore regulated as a cosmetic, whose function is to “...promote attractiveness...” of the body [6]. It has also been ruled by the FDA that proposed dietary supplements (which are regulated as a subset of foods) meant to be held in the mouth, followed by expectoration, are not dietary supplements, because they are not swallowed.

The definition of food has generally held since the 1958 definition, although it was changed slightly in the 7th Circuit in 1983, to now indicate that a food is something consumed “...primarily, for [it’s] taste, aroma or nutritive value.” This court decision did not radically change the definition of food from the original context, but in this particular case, prohibited the use of a food extract for therapeutic intent (*i.e.*, amylase isolated from kidney beans as an inhibitor of carbohydrate breakdown and marketed for weight loss—so-called “starch blockers”).

In general, the law prohibits the sale of food “if it consists in whole or in part of any filthy, putrid, or decomposed substance, or if it is otherwise unfit for food” (in practice, “fitness” can be quite subjective). Also, some foods which are ordinarily safe to eat may become unsafe, as described in §402 of the FFDCA [7]:

§402. A food shall be deemed to be adulterated—(a) (1) If it bears or contains any poisonous or deleterious substance which may render it injurious to health; but in case the substance is not an added substance such food shall not be considered adulterated under this clause if the quantity of such substance in such food does not ordinarily render it injurious to health...

The first part of §402 is clear; if a food contains a poisonous or deleterious substance it cannot be used as a food—a fairly broad standard. The second part of the section “...but in case the substance is not an added...the quantity of such substance does not ordinarily render it injurious to health...” requires an explanation. This clause simply means that although toxic substances may be present in foods, the food is not adulterated if the amount present in the food is not ordinarily injurious to health. For example, tomatine in tomatoes, psoralens in celery or glycoalkaloids in potatoes are normally

present in concentrations that are not harmful; however, in the event these amounts are increased (through such processes as breeding, mishandling during harvesting, storage or transportation) and become harmful, these foods are then considered to be adulterated. This second and narrower part of the statute is followed up in §406 of the FFDCA [8]:

§406 Any poisonous or deleterious substance added to any food, except where such substance is required in the production thereof or cannot be avoided by good manufacturing practice shall be deemed to be unsafe for purposes of the application of clause (2) (A) of section 402(a); but when such substance is so required or cannot be so avoided, the Secretary shall promulgate regulations limiting the quantity therein or thereon to such extent as he finds necessary for the protection of public health, and any quantity exceeding the limits so fixed shall also be deemed to be unsafe for purposes of the application of clause (2) (A) of section 402(a).

§406 then, allows the FDA to establish tolerances for these unavoidable contaminants, that is, a food may contain a toxin (such as mercury), if the presence of that toxin is (a) unavoidable and (b) under the level tolerated, the food is not considered to be unsafe. Because establishing a “tolerance” requires an extensive rule-making process, the FDA has adopted the use of “action levels”, which are non-binding guidelines [9]. For food ingredients (e.g., additives), potentially harmful constituents or contaminants are addressed by limiting the amount present in the specifications; higher than allowed amounts render the ingredient and the food to which it has been added, adulterated.

A few potential foods are banned outright by regulation such as the slaughter of companion animals (cats, dogs and horses) for food, offal and colostrum or those foods whose preparation is regulated by guidelines other than current good manufacturing practices (e.g., pufferfish preparation). Some naturally sourced substances (while present in some foods) are banned for addition to food for reasons of safety and include safrole, calamus and coumarin (a full list of which may be seen in 21 Code of Federal Regulations (CFR) 189). Other foods which may contain toxic substances, such as prussic acid in peach leaves, β -thujone in wormwood, saxitoxin in seafood, *etc.*, are controlled by regulation through the use of tolerances, or more correctly, specifications for the product that limit the amount of toxin that may be present. For those foods or ingredients with potential for harm, but not addressed by a specific regulation, action level, *etc.*, the reference in the FFDCA to substances “unfit for food” and flowing from that provision, Sections 402 and 406 of the FFDCA, apply. That is, the lack of a specific action taken by the FDA (or any regulatory agency), for a potentially harmful substance is not a license to market that substance.

3. Factors Driving the Acceptance of Certain Foods

Beyond the basic requirements of nutritional or hedonic value, the concept of exactly what constitutes food is largely culturally based; that is, the consumption of pork, shellfish, eel, “rocky mountain oysters”, cracklings, chitlin’s (chitterlings), brain, monkey, guinea pig, dog, snake, insects and arachnids, *etc.*, may be prohibited by religious practices or a matter of personal taste and, in the case of brains (or neural tissue) at least from cattle, has recently become no longer acceptable. Interestingly, there are no fruits or vegetables on any theocratic forbidden list.

There are some personal prohibitions that are genetically driven, but may not be perceived as a “toxicity” concern. For example, a genetic variant has been described for cilantro, which is perceived

by some people as having an unpleasant soapy taste or rank smell [10]. Another, better known variant is the ability to taste phenylthiourea (also known as phenylthiocarbamide, PTU or PTC) [11]. The ability to taste and smell certain substances may be key to evolutionary survival, as while the alkaloids of many potentially poisonous plants confer a bitter flavor, Goff and Klee have indicated that certain flavors and odors may also provide sensory cues for nutritional value of some plants [12]. For example, the characteristic odor profile of tomato (e.g., “tomato”, “green”, or “grassy”) are derived from *cis*-3-hexenal, *cis*-3-hexenol and *trans*-hexenal along with visual cues, to promote repeated consumption of an enjoyable food. In the context of promoting consumption of a specific food anosmia (lack of odor perception) or “specific anosmia” (which may be genetically based), will put the individual at a competitive disadvantage in food selection. Persistent or total anosmia also represents a clear safety hazard as the individual could not detect the tell-tale signs of decay or putrefaction of unfit foods.

There are some food prohibitions that are medically driven, as the result of genetics or autoimmune disease, as shown in Table 1.

Table 1. Medically driven food prohibitions (compiled from NORD [13]).

Disease/Syndrome	Causative Food	Cause	Comment
Disaccharide intolerance	Sucrose, dextrins	Autosomal recessive trait characterized by the deficiency or absence of enzymes sucrase and isomaltase in the intestine.	Attacks characterized by bloating and diarrhea.
Favism	Broadbean (<i>Vicia faba</i>)	X-linked recessive trait resulting in low amounts of glucose-P-dehydrogenase. Several subtypes known.	Hemolytic anemia may result from consumption of offending foods.
Galactosemia	Galactose and lactose (dairy products)	Autosomal recessive trait with low levels of any one of three enzymes directly responsible for galactose metabolism.	High levels of galactose in the blood results in hepatomegaly, cirrhosis, and renal failure. Infant mortality is ~75%.
Gluten intolerance	Wheat, barley, gluten containing foods	Autoimmune disease	Sensitivity to storage protein (gliadin) in some grains.
Lactose intolerance	Dairy products	Inborn error of metabolism—low or no lactase enzyme in the intestine.	Lactase is required to cleave lactose (a disaccharide of galactose and glucose). Bloating and diarrhea may develop.
Ornithine transcarbamylase deficiency	Dietary nitrogen (primarily meat)	X-linked recessive disorder resulting in low production of hepatic ornithine transcarbamylase interrupting the urea cycle and leading to accumulation of ammonia.	Although usually first seen in neonates, there may be an adult onset. Citrullinemia is another genetic disease affecting the urea cycle.
Phenylketonuria (PKU disease)	Phenylalanine in foods	Autosomal recessive trait characterized by inadequate hepatic phenylalanine hydroxylase.	Leads to accumulation of phenylpyruvate which may accumulate in the brain and lead to seizures, mental retardation, etc. Products containing phenylalanine must be labeled.

Table 1. Cont.

Refractory sprue	Wheat, barley and rye	Autoimmune disorder triggered by gliadin, a gluten storage protein.	Unlike common celiac sprue, adherence to a gluten-free diet may not cause symptoms to abate.
Trimethylaminuria	Fish	Autosomal recessive resulting in low production of flavin containing monooxygenase enzyme 3 (FMO3).	Fish odor syndrome. Failure to breakdown trimethylamine, a build of which results in a fish odor.
Very long chain Acyl CoA dehydrogenase deficiency (LCAD)	Very long chain fatty acids	Autosomal recessive trait resulting from a mutation in the HADHA gene.	Prevents mitochondrial metabolism of very long chain fatty acids.

Other medically driven prohibitions include food allergies, the most common of which are to milk, egg, fish, crustacean shellfish, tree nuts, wheat, peanuts and soybeans which account for 90% of all food allergies in the US. The *Food Allergen Labeling and Consumer Protection Act of 2004* (FALCPA), effective January 1, 2006, requires labeling of any product containing these ingredients or a protein derived from one of these offending foods or incidental additives or flavors derived therefrom. Exceptions are limited to any highly refined oil derived from a major food allergen (e.g., peanut or soybean oil) or any food ingredient exempt from labeling under a *petition or notification process* specified in the law [14].

There are also a number of food-drug interactions, the consumption of one interfering with the metabolism of the other, which may result in an enhanced or abated effect of the drug (Table 2).

Table 2. Food drug interactions (used with permission from Kotsonis and Burdock [15]).

Enzyme or Transporter	Food	Drug
CYP1A2	Caffeine, theophylline, grapefruit juice (naringen and furanocoumarins bergmottin and dihydroxybergamotin), grape juice, cruciferous vegetables, apiaceous vegetables, cooked meat	Clozapine, fluvoxamine, imipramine
CYP2E1	Watercress and possibly other isothiocyanate-containing cruciferous vegetables; polyunsaturated fatty acids (corn oil, menhaden oil)	Ethanol, halothane, enflurane
CYP3A4	Grapefruit, orange juice, red wine, possibly other polyphenol-containing substances, St. Johns wort, garlic	Ketoconazole, cyclosporine, erythromycin, protease inhibitors, HMG-CoA reductase inhibitors
UGT and GST	Brussels sprouts, cabbage, watercress, broccoli	Acetaminophen, oxazepam, morphine, ibuprofen
P-glycopeptide and OATP	Vegetables, fruit juice, St. Johns wort	Digoxin, cyclosporine, parvastatin

UGT: uridine diphosphate glycuronosyltransferases; GST: glutathione-S-transferases; OATP: organic anion transporting polypeptides.

4. Toxin Incorporation during Growth, Storage or Processing

4.1. Environmental contaminants

4.1.1. Selenium in grain

Selenium (Se) enters the food chain via plant and microorganism conversion of inorganic selenium to organically bound forms [16]. Selenium toxicity (*i.e.*, selenosis), caused by excessive selenium intake, has occurred on a large scale in seleniferous regions in China as the result of increased consumption of selenium-containing foods (approximate daily intake of 3–6.5 mg Se/day) [17]. The most common symptoms of selenosis are loss of hair, deformity, and loss of nails. Other reported symptoms include increased blood selenium levels, diarrhea, fatigue, a garlic-like odor of the breath and bodily secretions, irritability, peripheral neuropathy, and skin lesions [18]. Selenium intake levels that cause selenosis have not yet been well defined. Studies in China suggest that approximately 3–5 mg/day (0.05–0.08 mg/kg/day) will cause selenosis. Residents of seleniferous regions in South Dakota who consumed approximately 700 µg selenium/day (0.01 mg/kg/day) showed no symptoms of selenosis. The EPA has proposed an oral reference dose (RfD) of 0.005 mg/kg bw/day, or 350 µg/day [19].

4.1.2. Methyl mercury in seafood

Exposure to elemental mercury is relatively rare, although was once an occupational disease of hat manufacturers as elemental mercury was used for the curing of animal pelts. Inhalation of the mercury fumes led to mental deterioration and subsequently named “mad hatter syndrome” [20].

Of interest to food toxicology, is the methyl derivative, methyl mercury, formed by bacterial action in an aquatic environment from anthropogenic and natural sources of elemental mercury. Anthropogenic sources include burning of coal (which contains mercury), chloralkali process and other sources of elemental mercury into aquatic environments. In the case of Minamata, Japan, there was a direct discharge of methyl mercury into the environment. Methyl mercury exposure may cause neurological paresthesias, ataxia, dysarthria, hearing defects and death. Developmental delays have been documented in children borne of mothers exposed to methyl mercury [21]. Other than direct exposure to methyl mercury, exposure usually comes about as the result of methyl mercury becoming incorporated into the food chain, moving up as each predator consumes the smaller and less fortunate animal. Near the peak of the food chain, methyl mercury becomes concentrated in fish including, bonito (*Sarda* spp.), halibut (*Hippoglossus* spp.), mackerel (*Scomberomorus* spp.), marlin (*Makaira* spp.), shark (all species), swordfish (*Xiphias gladius*), and bluefin tuna (*Thunnus* spp.). The selection of these species was based on historical data on levels of methyl mercury found in fish consumed in the U.S. The selection was also based on an FDA action level of 1.0 ppm in the edible portion of fish [22]. However, the allowable level of mercury depends on whether the mercury was “added”; that is, did the presence of mercury arise from an anthropogenic source (*i.e.*, was the fish caught in an area known for mercury discharge), or was not added and the result of mercury naturally present in the environment [23].

4.2. Naturally formed substances

4.2.1. β -Thujone

Thujone, a monoterpene ketone, is the primary constituent of essential oils derived from a variety of plants, including sage (*Salvia officinalis*), clary (*Salvia sclarea*), tansy (*Tanacetum vulgare*), wormwood (*Artemisia* spp. and white cedar (*Thuja occidentalis* L.) [24]. Essential oils from these plants are used in herbal medicines, as flavorings in alcoholic drinks and fragrances throughout the world. Thujone is potentially toxic and the presence of alpha- or beta-thujone in food and beverages is regulated by law in several countries. In the US, thujone as an isolated substance is banned as an ingredient to be added to food and many of the natural thujone-containing plant oils (e.g., wormwood, white cedar, oak moss (*Evernia prunastri*) and tansy) are used as flavorings in food under the condition that the finished food is thujone-free [25]. Absinthe (made from wormwood) contains significant levels of thujone and is available in Spain, Denmark and Portugal. Wormwood itself is a popular flavoring for vodka in Sweden, while vermouth, chartreuse, and Benedictine all contain small levels of thujone [26]. Sage oil is used to provide the characteristic flavor in sausages, meats, condiments and sauces, and contains approximately 20–30% thujone (alpha- and beta-) [27,28]. Both alpha- and beta-thujone act as noncompetitive blockers of the gamma-aminobutyric acid (GABA)-gated chloride channel [29]. The essential oils of sage, hyssop (*Hyssopus officinalis* L.), and cedar all contain thujone and have been cited to have caused central nervous system effects characterized by tonic-clonic or solely clonic convulsions [30]. Thujone is believed to be the toxic agent in absinthism, a syndrome produced by the chronic use of absinthe, made from the essence of wormwood. The syndrome is characterized by addiction, hyperexcitability and hallucinations. The debilitating illnesses suffered by Vincent Van Gogh and Henri de Toulouse-Lautrec have been linked to absinthism, while the toxicity of thujone was a major factor in banning absinthe in the early 1900s [31]. A published case report detailed a male subject that drank about 10 mL of essential oil of wormwood (believing it was absinthe) and became agitated, incoherent and disoriented, subsequently developing renal failure [32]. The no observable effect limit (NOEL) for convulsions in subchronic toxicity studies in female rats was 5 mg/kg bw/day [24]. Detoxification of thujone is thought to occur via CYP450-dependent oxidation and subsequent glucuronidation and excretion [33]. The FDA limits exposure to β -thujone from *Artemisia* spp., when used as a natural flavoring substance or natural substance used in conjunction with flavors (21 CFR 182.20).

4.2.2. Prussic acid in cherry, apple and peach pits

Prussic acid (also known as hydrocyanic acid, hydrogen cyanide, or cyanide) is formed when cyanogenic glycosides found in leaves, cherry, apple and peach pits, oak moss and other plant tissues are damaged and come into contact with *beta*-glycosidase or emulsion enzymes. The enzymes release the cyanide from the glycoside, and the cyanide prevents the body's cells from utilizing oxygen, resulting in cellular necrosis and tissue damage. The mucous membranes and blood are bright red as they are oxygenated, but the cells in the tissues cannot utilize the oxygen. Clinical signs of prussic acid poisoning include rapid breathing, trembling, incoordination and in extreme cases, respiratory and/or cardiac arrest [34]. Many fruit trees contain prussic acid glycosides in the leaves and seeds, but only


negligible levels are present in the fleshy parts of the fruit [35]. In the west African tropics, cassava is consumed as a dietary staple and inappropriate handling of the cassava prior to processing and consumption can result in a chronic form of cyanide poisoning termed “tropical ataxic neuropathy”, the result of demyelination of the optic, auditory, and peripheral nerve tracts [36].

Prussic acid as found in flavoring ingredients is limited to 25 ppm in cherry pits (*Prunus avium* L. or *P. cerasus* L.), cherry laurel leaves (*Prunus laurocerasus* L.), elder tree leaves (*Sambucus nigra* L.), and peach leaves (*Prunus persica* (L.) Batsch) (21 CFR 172.510); although the extract of bitter almond (*Prunus amygdalus* Batsch, *Prunus armeniaca* L., or *Prunus persica* (L.) Batsch) must be prussic acid free (21 CFR 182.20). There are no FDA regulations or guidelines restricting the presence of prussic acid in apple seed (*Malus* spp.), probably because extracts of these seeds have no economic value as flavor ingredients.

4.2.3. Hypericin in St. John’s wort

St. John’s wort (*Hypericum perforatum*; Figure 1) is an herbal thought to alleviate symptoms of depression, and standardized extracts of St. John’s wort are consumed typically in tablet or capsule form. The major active antidepressive constituents in St. John’s wort are thought to be hyperforin and hypericin [37,38]. The mechanism of action is not fully understood, but may involve inhibition of serotonin (5-HT) reuptake, similar to conventional antidepressive drugs. In this manner, hyperforin and hypericin taken in conjunction with other serotonin reuptake inhibitors may contribute to *serotonin syndrome*, a potentially life-threatening elevation of serotonin in the central nervous system. Hyperforin is also known to induce cytochrome P450 enzymes CYP3A4 and CYP2C9, which can lead to increased metabolism of certain drugs and decreased clinical response [39].

Figure 1. St. John’s wort (*Hypericum perforatum*) [40].



In large doses, St. John’s wort is poisonous to grazing animals, with published cases of livestock poisoning characterized by general restlessness and skin irritation, hindlimb weakness, panting, confusion, depression and in some instances, mania and hyperactivity resulting in the animal running in circles until exhausted [41]. In humans, consumption of St. John’s wort may result in

photosensitization, and at high continuous doses, some liver damage may occur [39]. The FDA limits exposure to St. Johns wort (*Hypericum perforatum*), including the leaves, flowers, and caulis, by mandating that only hypericin-free alcohol distillate form may be used and then, only in alcoholic beverages (21 CFR 172.510).

4.2.4. Goitrogens (glucosinolates) in *Brassica* spp.

Certain raw foods have been found to contain substances that suppress the function of the thyroid gland by interfering with the uptake of iodine, an essential nutrient in growth, cognitive function, and hormonal balance. A lack of functional iodine is known to result in cognitive deficiencies (e.g., Cretinism). The decrease in iodine uptake causes the thyroid gland to enlarge, forming a goiter. Foods that have been identified as goitrogenic include spinach, cassava, peanuts, soybeans, strawberries, sweet potatoes, peaches, pears, and vegetables in the *Brassica* genus, which include broccoli, brussels sprouts, cabbage, canola, cauliflower, mustard greens, radishes, and rapeseed [42]. Goiter has also been attributed to the consumption of large quantities of uncooked kale or cabbage.

High temperatures (*i.e.*, cooking) inactivate the goitrogenic substances, collectively termed glucosinolates. Cassava (*Manihot esculenta*) is an essential dietary source of energy in the tropics, but contains high levels of linamarin, a glucosinolate. Cassava must be properly processed-dried, soaked in water or baked to effectively reduce the linamarin content [43]. Glucosinolates are sulfur-containing substances that are metabolized in the body by thioglucosidase to form thiocyanate, isothiocyanate, nitriles and sulfur. Under certain conditions the isothiocyanates undergo cyclization to form goitrins, increasing their potent goitrogenic activity. The oils from rapeseed (*Brassica napus*) must be analyzed for potential goitrins to circumvent potential goitrogenic activity when consuming these oils [44]. No FDA regulations were located for permissible concentrations of glucosinolates in human food. Glucosinolates (calculated as epi-progoitrin) and goitrin are limited to not more than 4% and 0.1% (respectively) of the seed meal of *Crambe abyssinica* (Crambe meal) obtained after the removal of the oil and used as an animal feed ingredient (21 CFR 573.310).

4.2.5. Erucic acid in rape

Rape (*Brassica napus* L. or *Brassica campestris* L.) is an annual herb of the mustard family native to Europe and is grown in the United States because it produces oil-rich seeds for cooking oil [45]. Rapeseed oil had been used for hundreds of years as oil for lamps and more recently as machine oil lubricant. Widespread use of rapeseed oil as a food ingredient was not considered until the late 1940s and 50s. However, early studies found that feeding high levels of rapeseed oil to rats significantly increased cholesterol levels in the adrenal glands and lipidosis in the cardiac tissue [46,47]. This effect was also noted in chickens, ducks and turkeys fed high levels of rapeseed oil, resulting in growth retardation, mortality, and a thickening of the epicardium and increased fibrous tissue in different areas of the myocardium [48]. Erucic acid was identified as the causative agent of these effects of rapeseed oil. Erucic acid is a long-chain fatty acid with one unsaturated carbon-carbon bond (C22:1). High levels of erucic acid have been linked to fatty deposit formation in heart muscle in animals [49]. Erucic acid is poorly oxidized by the mitochondrial β -oxidation system, especially by the myocardial cells, which results in an accumulation of erucic acid, producing myocardial lipidosis which has been

reported to reduce the contractile force of the heart [50]. Although myocardial lipidosis due to erucic acid consumption has not been confirmed in humans, animal feeding studies confirmed the formation of myocardial lipidosis in a variety of animal species in a dose-dependent manner, which has been the standard assessment by government agencies of potential adverse effects in humans. Canola oil is obtained from Canola (Canadian oil, low acid), a rapeseed variety that was conventionally bred in the late 1970s in Canada to contain reduced levels of erucic acid and glucosinolates [51,52]. The FDA limits the amount of erucic acid in Canola oil to no more than 2% of the component fatty acids (21 CFR 184.1555).

4.2.6. Furocoumarins

Furocoumarins represent a family of natural food constituents with phototoxic and photomutagenic properties. They are found mainly in plants belonging to the *Rutaceae* (e.g., citrus fruits) and *Umbelliferae* (e.g., parsnip, parsley, celery, carrots) families. Furocoumarins are produced in response to stress, to aid plants in defense against viruses, bacteria, fungi, insects and animals, and are regarded as natural pesticides [53]. Concentrations may also increase after exposure to UV radiation, changes in temperature, prolonged storage, or treatment with hypochlorite or copper sulfate (Chaudhary et al., as cited in Wagstaff 1991 [53], p. 270 and Beier *et al.*, as cited in Ashwood-Smith [54], p. 916).

The three most active furocoumarins in producing photodermatitis are psoralen, 5-methoxypsoralen (5-MOP, bergapten), and 8-methoxypsoralen (8-MOP, xanthotoxin or methoxsalen) [55]. In the presence of near UV light (320–380 nm), these three linear furocoumarins can form adducts with DNA and DNA-crosslinks. The consequences of these photoadditions to cells are cell death, mutations and chromosome aberrations [54]. In the presence of ultraviolet A radiation, 5-MOP and 8-MOP produce skin tumors in experimental animals. At a chronic dose of 37.5 mg/kg bw/day in the diet, 8-MOP produces increased incidences of tubular cell hyperplasia, adenomas, and adenocarcinomas of the kidney and carcinomas of the Zymbal gland in rats [56]. Cases of skin cancer have been reported in patients treated with 8-MOP and long-wave ultraviolet light for treatment of psoriasis or mycosis fungoides [57,58]. IARC has classified 5-MOP and 8-MOP plus ultraviolet radiation in group A (probably carcinogenic in humans) and in group 1 (carcinogenic to humans), respectively [57,59].

Citrus fruits, especially grapefruit, produce a variety of chemicals in their peels that may have adverse interactions with drugs. Typically, citrus fruit juice is produced utilizing the whole fruit, including the peel. One chemical found in the peel is bergamottin (also known as bergamot), a natural furanocoumarin that is known to inhibit some isoforms of the cytochrome P450 enzyme (CYP) 3A4 [60]. Inhibition of this enzyme prevents oxidative metabolism of certain drugs, resulting in an elevated concentration of a drug in the bloodstream [61]. Bergamot and other chemicals in citrus (e.g., lime, grapefruit, orange, lemon) oils [62] are also phototoxic, causing significant toxicity to the skin when exposed to sunlight [63]. 5-Methoxypsoralen, the most phototoxic constituent of bergamot oil, showed mutagenic activity in bacterial assays and clastogenic effects in mammalian cells in culture when exposed to UV light [64].

Celery reportedly contains 100 ppb psoralens (100 micrograms/kg) and parsnips as much as 40 ppm (40 mg/kg) [65]. The estimated dietary intake of furocoumarins for people eating furocoumarin-containing foods (est. 80% of the population) is 1.31 mg/day [53], which is approximately 0.022 mg/kg bw/day

for a 60 kg human. This is approximately 1000-fold lower than the 13-week dietary no observable adverse effect level (NOAEL) for liver toxicity in the rat (25 mg 8-MOP/kg bw/day) and 1700-fold lower than the dietary dose that has been shown to induce cancer in rats (37.5 mg/kg). Therefore, the risk of developing liver toxicity or cancer due to ingestion of psoralens in the diet is low.

In humans, the phototoxic threshold dose of furocoumarin mixtures after dietary exposure is of the order of 10 mg 8-MOP plus 10 mg 5-MOP, which is equivalent to about 15 mg 8-MOP per person. This phototoxic threshold dose is not reached by the consumption of celery roots and other conventional vegetables under normal dietary habits, which result in intake of approximately 2–8 mg furocoumarins per person [66]. Therefore, ordinarily dietary exposure to psoralens is not considered to be a significant risk for development of photodermatitis, albeit the margin of safety is low [65]. There are no FDA regulations or guidelines specific to the presence of furocoumarins in food.

4.2.7. Amylase inhibitors

Naturally occurring inhibitors of α -amylase are found in aqueous extracts of wheat, rye and kidney beans. The physiological role of α -amylase inhibitors in plants is not well understood, but may protect them against insect infestation. In mammals, some amylase inhibitors have been shown to attenuate the normal increase in blood glucose that occurs after ingestion of starch. However, since α -amylase inhibitors have been shown to be inactivated by gastric acid, pepsin or pancreatic proteinases, their potential as “starch blockers” is limited [67]. α -Amylase inhibitors were once added to foods as “starch blockers” to limit carbohydrate absorption for the purpose of weight loss; however, the FDA later determined that at least this use of α -amylase inhibitors was as drug, and they were consequently taken off the market [68].

α -Amylase inhibitor protein is a major allergen (referred to as Asp o 2) that has been implicated in the development of occupational toxicity known as “baker’s asthma disease” [69]. Although α -amylase inhibitor protein is naturally found in wheat flour, it is also found in flour in which α -amylase from *Aspergillus oryzae* has been added to enhance carbohydrate fermentation by yeast [70]. Consequently, α -amylase inhibitor protein can be potentially found in baked products that are derived from sources other than wheat. Cases of food allergy have been reported in people ingesting bread containing α -amylase inhibitor protein. Symptoms of allergy include sneezing, rhinorrhea, oropharyngeal itching, hoarseness, cough and dyspnea [71].

High α -amylase inhibitor activity against human salivary α -amylase has been found in wheat flour (590 units/g), whole wheat flour (351 units/g) and whole rye flour (186 units/g). Bread baking reduces the activity by 80–100%, depending on type. The activity in uncooked spaghetti (248 units/g) is reduced more than 98% by 15 minutes of boiling. Boiling of red beans for 1.5 hours reduces activity to undetectable levels [71]. However, α -amylase has been shown to retain some allergenic activity when heated to 200 °C (Baur *et al.*, as cited in Phadia AB 2010 [72], p. 2).

4.2.8. Lectins in legumes

Lectins are a group of glycoproteins that are present in high levels in legumes (e.g., black beans, soybeans, lima beans, kidney beans and lentils) and grain products [73,74]. Lectins can reversibly bind to carbohydrates without altering their covalent structure [73]. The ability of lectins to bind to and

agglutinate red blood cells is well known and used for blood typing—hence the lectins are commonly called hemagglutinins. Lectins also can bind avidly to mucosal cells and interfere with nutrient absorption from the intestine [75]. Because the ability of the lectins to cause intestinal malabsorption is dependent on the presence of enteric bacteria, it has been hypothesized that lectins may also produce toxicity by facilitating bacterial growth in the GI tract [76].

Lectins isolated from black beans can produce growth retardation when fed to rats at 0.5% of the diet, and lectin from kidney beans causes death within two weeks when fed to rats at 0.5% of the diet. Soybean lectin produces growth retardation when fed to rats at 1% of the diet. The castor bean lectin ricin (one of the most toxic natural substances known) is notorious for causing deaths of children, and has been used as an instrument of bioterrorism [75].

Phytohaemagglutinin (PHA) is a lectin found in significant quantities (as much as 2.4–5% of total protein) in legumes such as red or white kidney beans, green beans and fava beans. PHA has a number of different properties, including the ability to induce mitosis, affect membrane transport and permeability to proteins, and agglutinate red blood cells. Rats fed a diet containing 6% PHA exhibit weight loss, associated with malabsorption of lipid, nitrogen and vitamin B12 [76]. PHA from red kidney beans inhibits sodium and chloride absorption in the rabbit ileum, indicating that PHA can affect electrolyte transport in the gut [77]. Symptoms of toxicity to PHA in humans such as nausea, vomiting, or diarrhea occur within three hours of ingestion. Recovery generally occurs within four or five hours of onset [78].

There are no FDA regulations or guidelines restricting the presence of lectins in food, but the FDA does provide recommended cooking practices prior to consuming legumes. Concentrations of PHA (and other lectins) are higher in uncooked than cooked beans. A raw, red kidney bean can contain up to 70,000 hemagglutinating units (hau). Most lectins are reduced by moist, but not dry heat. Therefore, steaming or boiling causes a significant reduction in concentrations of lectins in beans. Boiling for at least ten minutes has been shown to reduce hau in beans by 200-fold. Because cooking temperatures under 176 °F do not destroy lectin, use of slow cooking and/or a crockpot is not advised for cooking beans [79].

4.2.9. Anti-thiamine compounds

Substances that act on the availability of vitamins are commonly referred to as antivitamins. These include materials that can cause a deficiency of vitamins by competing with vitamins in various metabolic reactions as the result of similar chemical structure or destroying or decreasing the effects of a vitamin by modifying the molecular conformation or by forming a complex [67].

Thiaminase cleaves thiamine (vitamin B1) at the methylene linkage, rendering it biologically inactive. Activity of thiaminase requires a cosubstrate—usually an amine or sulfhydryl-containing protein such as proline or cysteine. Thiaminase is found in fish, crab, clams and in some fruits and vegetables such as blueberries, black currants, red beets, Brussels sprouts and red cabbage [67].

Thiamine is an essential vitamin involved in energy production. Thiamine deficiency is associated with impaired pyruvate utilization, resulting in a shortage of cellular ATP. In humans, thiamine deficiency may lead to weakness and weight loss. Severe thiamine deficiency produces “beri-beri”, a disease characterized by anorexia, cardiac enlargement, and muscular weakness leading to ataxia [80].

Cooking destroys thiaminases in fish and other sources. There are no FDA regulations or guidelines specific to the presence of thiaminase in food.

4.2.10. Pyrrolizidine alkaloids

Pyrrolizidine alkaloids (PAs) are found in some plants of the Apocyanaceae, Asteraceae, Boraginaceae, Compositae (*Senecionae* and *Eupatoriae*), Fabaceae, Leguminosae (*Crotalaria*), Ranunculaceae and Scrophulariaceae families. Herbs such as comfrey root and leaf (*Symphytum* spp.) (Figure 2), coltsfoot leaf and flower (*Tussilago farfara*) and borage leaf (*Borago officinale*), and several species of *Eupatorium* typically contain high levels of PAs. Humans are exposed to PAs through the accidental contamination of foodstuffs and intentional ingestion of PA-containing vegetables and herbal medicines. Serious incidences of illness have been reported in people consuming cereal grains that are contaminated with the seeds of PA-containing plants [81]. PAs are also present in milk from cows and goats and in honey [82].

Figure 2. Comfrey (*Symphytum officianale* L.) [83].



The pyrrolizidine structure is based on two fused, five-membered rings that share a bridgehead nitrogen atom, forming a tertiary alkaloid. The rings contain a hydroxymethylene group at the C-1 position and a hydroxyl group at the C-7 position, forming a necine base. Several PAs that contain unsaturated necine rings are hepatotoxic, mutagenic, teratogenic and/or carcinogenic. Toxicity is thought to be due to enzymatic conversion of PAs to pyrroles, which act as alkylating agents [67]. Pyrroles formed in the liver can travel to the lungs, causing thickening of the pulmonary vasculature and pulmonary hypertension [82].

The sale of comfrey products for internal use has been banned in the United States and Canada [82]. However, comfrey tea is still widely available. It is estimated that consumers of comfrey tea could be ingesting up to 5 mg of PAs per day (Speijers and Egmond, as cited in Deshpande 2002b [81], p. 368), or 0.083 mg/kg bw/day. The range of toxic doses in humans is thought to be 0.1–10 mg/kg per day [84], although the World Health Organization has reported a case of veno-occlusive disease in a subject ingesting 0.015 mg PAs/kg of body weight per day from comfrey.

4.2.11. Rhubarb and oxalic acid

Oxalic acid (oxalate) is generally found in rhubarb (0.2–1.3%), tea (0.3–2.0%), spinach (0.3–1.3%), parsley (1.7%) and purslane (1.3%), but may also be found in asparagus, broccoli, Brussels sprouts, collards, lettuce, celery, cabbage, cauliflower, turnips, beets, peas, coffee, cocoa, beans, potatoes, berries, and carrots [67,73,85].

Oxalic acid is an organic acid that can bind calcium and other minerals, making them insoluble and decreasing their bioavailability. Ingestion of foods containing high concentrations of oxalates may cause decreased bone growth, kidney stones, renal toxicity, vomiting, diarrhea, convulsions, coma and impaired blood clotting [73]. The significant role oxalate plays in kidney stone development is exemplified by the fact that approximately 65% of kidney stones consist of calcium oxalate [86].

Using the oral LD₅₀ value of 375 mg/kg in rats, it has been estimated that ingestion of approximately 22 g of oxalic acid could be lethal to a 59 kg human [85]. Because approximately 4.5 kg of rhubarb leaves would have to be ingested in order to achieve a lethal dose, it has been hypothesized that documented cases of fatal rhubarb poisoning in humans were due to consumption of some other substance than oxalic acid [67].

Because cooking does not remove oxalate, and mineral complexes with oxalate are insoluble in water, oxalates are somewhat difficult to remove from foods. Therefore, diets rich in oxalate-containing foods should be supplemented with minerals such as calcium or potassium to prevent deficiencies. Limits on oxalic acid have been cited in ferric ammonium ferrocyanide and ferric ferrocyanide when used as color additives (21 CFR 73.1298 and 21 CFR 73.1299) with oxalic acid or its salts at not more than 0.1% of the colorant.

4.2.12. Zucchini and cucurbitacins

Members of the *Cucurbitacea* family (zucchini, cucumbers, pumpkins, squash, melons and gourds) produce cucurbitacins (oxygenated tetracyclic terpenes) that act as movement arresters and compulsive feeding stimulants for Diabrotic beetle (corn rootworms and cucumber beetles). Cucurbitacins are among the most bitter compounds known, and in nanogram quantities they deter most non-Diabrotic herbivores [87].

Because cucurbitacins act as feeding stimulants, they are added to insecticidal baits to increase efficacy [88]. Therefore, dietary exposure to cucurbitacins could occur through ingesting plants that normally contain them or by ingesting plants to which cucurbitacin-containing pesticides have been applied.

Under normal circumstances, cucurbitacins are produced at low enough concentrations that are not perceived as being bitter by humans. In response to stresses such as high temperatures, drought, low soil fertility and low soil pH, concentrations in fruits such as cucumbers may increase and cause the fruits to have a bitter taste [89]. Occasional cases of stomach cramps and diarrhea have occurred in people ingesting bitter zucchini. Twenty-two cases of human poisoning from ingestion of as little as 3 grams of bitter zucchini were reported in Australia from 1981 to 1982, and in Alabama and California in 1984. The cultivar implicated in the Australia poisonings was “Blackjack” [90]. There are no FDA regulations or guidelines specific to the presence of cucurbitacins in food.

4.2.13. Coumarins (tonka bean, woodruff, clover)

Coumarin (2H-1-benzopyran-2-one) is found in herb teas made from tonka beans (*Dipteryx odorata*), melilot (*Melilotus officinalis* or *Melilotus arvensis*) and woodruff (*Asperula odorata*), the flavoring oil of bergamot (from *Citrus bergamia*) and the spice cassia (*Cinnamomum cassia*; sometimes sold as cinnamon) [91]. Coumarin is liberated from the glycoside melilotoside (an ether of glucose bonded with an ester bond to coumarin) on drying coumarin-containing herb material.

Molds present in spoiled sweet (Melilotus) clover and other hay products can metabolize coumarin to dicoumarol, which is similar in structure to vitamin K [92]. Vitamin K is necessary to activate prothrombin, which is converted to the blood clotting substance thrombin. By inhibiting vitamin K, dicoumarol promotes bleeding. Concentrations of dicoumarol in fodder >10 ppm have been responsible for fatalities by hemorrhaging in cattle [91].

The addition of coumarin to food in the United States was banned in 1954, based on reports of hepatotoxicity in rats. However, because a number of foods contain coumarin, humans ingest approximately 0.02 mg coumarin/kg bw/day. The chronic administration of high doses of coumarin causes liver tumors in the rat and liver and lung tumors in the mouse. Overall, available data indicate that coumarin is not genotoxic. It is thought that the carcinogenicity of coumarin is caused by metabolism to toxic epoxides. Because doses of coumarin that cause toxicity and carcinogenicity in the lung and liver of experimental animals are more than 100 times the maximum human intake, exposure to coumarin from food poses no health risk to humans [93].

The addition of coumarin is prohibited in 21 CFR 189.130. The regulation notes that coumarin is found in tonka beans and extract of tonka beans, among other natural sources, and is also synthesized. It has been used as a flavoring compound, therefore addressing not just natural products (which would include buffalo grass or sweetgrass (*Hierochloe odorata*) used in flavoring vodka and other natural sources (see above)), as well as synthesized coumarin. Further, according to the regulation, “(b) Food containing any added coumarin as such or as a constituent of tonka beans or tonka extract is deemed to be adulterated under the act, based upon an order published in the Federal Register of March 5, 1954 (19 Federal Register 1239).” An analytical method for detection of coumarin in foods is specified in 21 CFR 189.130.

4.2.14. Phytates and phytic acid

Phytic acid (also referred to as phytate) is found in bran and germ of many plant seeds and in grains, legumes and nuts. Phytic acid is a simple sugar (myo-inositol) containing six phosphate sidechains, and as such, is a dietary source of phosphorus and an effective chelator of divalent cations such as zinc, copper, iron, magnesium and calcium [67,94]. Studies indicate that phytate-mineral complexes are insoluble in the intestinal tract, reducing mineral bioavailability [73]. Phytate also has been shown to inhibit digestive enzymes such as trypsin, pepsin, α -amylase and β -glucosidase. Therefore, ingestion of foods containing high amounts of phytate could theoretically cause mineral deficiencies or decreased protein and starch digestibility. Vegetarians that consume large amounts of tofu and bean curd are particularly at risk of mineral deficiencies due to phytate consumption.

Because phytate-rich foods are digested at a slower rate and produce lower blood glucose responses than foods that do not contain phytate, it has been hypothesized that phytate could have a therapeutic role in management of diabetes [67]. It also may have utility as an antioxidant [95]. However, because the beneficial effects of phytate are outweighed by its ability to cause essential mineral deficiencies, consumption of a diet containing high amounts of phytate is not recommended. Food manufacturers are developing methods to reduce phytate in foods, such as addition of the microbial phytase, which releases phosphates from the inositol backbone of phytate [96].

Phytate is fairly heat stable, but can be removed by soaking or fermentation [67]. The soybean has one of the highest phytate levels of any grain or legume, and requires a long period of fermentation for reduction [94]. In people who consume large amounts of soy products, mineral deficiencies can be prevented by consumption of meat or dairy products or use of supplemental vitamins. There are no FDA regulations or guidelines restricting the presence of phytates in food.

4.2.15. Hypoglycin in Ackee

Ackee (*Blighia sapida*; Figure 3) is the national fruit of Jamaica and is also found in other Caribbean nations, Central America, South American and southern Florida [97]. Consumers of the unripe fruit sometimes suffer from “Jamaican vomiting sickness syndrome” allegedly caused by the alkaloids hypoglycin A (HGA) and B. Levels of HGA in the opened, ripe fruit are undetectable, making opened fruit safe for consumption [98].

The hypoglycin toxin (L-methylenecyclopropylalanine) inactivates several flavoprotein acyl-CoA dehydrogenases, causing disturbances of the oxidation of fatty acids and amino acids [99]. This leads to a secondary inhibition of gluconeogenesis which can precipitate an extreme, dangerous drop in blood-glucose levels (hypoglycemia) that can be fatal. Symptoms of poisoning from unripe ackee fruit occur within 6 to 48 hours of ingestion and include drowsiness, repeated vomiting, thirst, delirium, fever or loose bowels. Exhaustion of the muscular and nervous systems, collapse, coma, and death may ensue [100,101].

Figure 3. Unripe Ackee Fruit (left panel) and ripe Ackee Fruit (right panel) [100].



Dietary exposure to hypoglycin in Jamaicans ranges from 1.21–89.28 micrograms/gram ackee [102]. Ingestion of one 100 gram fruit could therefore result in a dose of approximately 300 micrograms/kg bw in a 30 kg child. This dose is approximately one-fifth of the maximum tolerated dose of HGA in male and female rats of 1500 micrograms/kg bw/day [103], indicating that normal use levels of ackee do not have a large margin of safety.

The importation of canned ackee fruit into the United States is restricted to certain manufacturers to insure that only properly ripened ackees are used for canning [104], and the FDA routinely analyzes incoming shipments of ackee for hypoglycin levels that could be a health concern, having issued a recall of canned ackee fruit for this very reason in 2005. If hypoglycin poisoning is expected, glucose, fluids and electrolytes should be administered. Antiemetics may be used to control vomiting and benzodiazepines to control seizures. Endotracheal intubation should be performed in people exhibiting seizures or coma [97].

4.2.16. Safrole

Safrole (1-allyl-3,4-methylenedioxybenzene) is found in aromatic oils of nutmeg (*Myristica fragrans*), cinnamon (*Cinnamomum verum*) and camphor (*Cinnamomum camphora*) and is a major constituent of oil of sassafras (*Sassafras albidum*) [105]. Prior to being banned as a food additive in the United States in 1960, safrole was commonly used to flavor root beer and other foods. Most commercial “sassafras teas” and root beers are now artificially flavored as a result of the FDA ban (21 CFR 189.180).


At a concentration of 1% in the diet, safrole produces weight loss, testicular atrophy, bone marrow depletion and malignant liver tumors in rats [106]. Based on sufficient evidence of carcinogenicity in experimental animals, safrole is reasonably anticipated to be a human carcinogen [107]. The mechanism of carcinogenicity is thought to involve cytochrome P450 catalyzed hydroxylation of safrole to 1'-hydroxysafrole, and its subsequent metabolism to highly reactive electrophiles that bind to DNA [108].

Despite the FDA ban, sassafras is still a popular ingredient in herb teas and preparations [73]. The hazardous dose of sassafras oil for humans (which typically contains 80% safrole) is considered to be 0.66 mg/kg [109]. This may be exceeded by ingesting sassafras tea, which has been estimated by Segelman and Bisset (as cited in Burfield 2009 [109], p. 3) to give a dose of 3 mg/kg for a 60 kg individual.

4.2.17. Myristicin

Myristicin (Figure 4) is a naturally occurring insecticide and acaricide that is found in nutmeg and mace (*Myristica* spp.) at concentrations of 1.3% and 2.7%, respectively [110]. It is also present in black pepper, carrot, celery parsley and dill [67]. It is estimated that the average total intake of myristicin from dietary sources is “in the order of a few mg per person per day” [110].

Figure 4. Structure of myristicin.



Myristicin is a weak inhibitor of monoamine oxidase, and is structurally related to mescaline. At a dose level of 6–7 mg/kg bw, it may cause psychotropic effects in man, such as increased alertness, and

a feeling of irresponsibility, freedom and euphoria. Unpleasant symptoms, such as nausea, tremor, tachycardia, anxiety and fear have also been reported in humans ingesting this dose. Although the metabolism of myristicin resembles that of safrole, there is no evidence to suggest that myristicin is carcinogenic [110]. There are no FDA regulations or guidelines specific to the presence of myristicin in food.

At the concentrations normally present in spices or food, the likelihood of toxicity arising from myristicin is low. However, ingestion of greater than 5 grams of nutmeg (corresponding to 1–2 mg/kg bw myristicin) has produced toxicological symptoms in humans that are similar to alcohol intoxication. Because the myristicin content of nutmeg is approximately 1–3%, it is likely that components of nutmeg in addition to myristicin contribute to nutmeg toxicity [110].

4.2.18. Tomatine in tomatoes

The leaves, stems and unripe fruit of the tomato plant contain α -tomatine, a steroidal alkaloid containing D-xylose, D-galactose, and two molecules of D-glucose. Tomatine is toxic to a number of different fungi, thereby acting as a natural fungicide. It has been hypothesized that the toxic effects of tomatine on fungi are due to the ability of tomatine to complex with membrane sterols, causing membrane disruption [111].

Currently, there is no evidence to suggest that tomatine is a substance of concern. There are no reports of acute toxicity in humans due to ingestion of green tomatoes and there are no FDA regulations or guidelines specific to the presence of tomatine in food. Ingestion of a rare variety of ripe tomato (*Lycopersicon esculentum* var. *cerasiforme*) that contains up to 5 mg tomatine/g of dry weight has no adverse effects on natives who commonly ingest them [112].

Concentrations of tomatine decrease as tomatoes ripen, and ripe fruit contains approximately 36 mg per a 100 gram tomato [73]. Microwaving or frying does not reduce content of tomatine, and delayed-ripening varieties of tomatoes contain similar concentrations of tomatine as other tomatoes [113]. At this time, there is no evidence to suggest that a diet high in green tomatoes would be injurious to human health. Tomatine forms strong, insoluble complexes with cholesterol *in vitro*, and has been shown to lower plasma LDL cholesterol in hamsters [114], suggesting that it may have beneficial effects on blood lipids of humans.

4.2.19. Japanese star anise

Chinese star anise (*Illicium verum*) is a common source of anethole, a popular flavoring ingredient. On the other hand, Japanese star anise (*Illicium anisatum*) is scientifically recognized as highly poisonous and not fit for human consumption. Japanese star anise contains the potent neurotoxins anisatin and neoanisatin, as well as the neurotoxic sesquiterpene lactone veranisatins that are normally found in other kinds of star anise, including Chinese star anise [115].

Brewed “teas” containing star anise have been associated with illnesses affecting about 40 individuals, including approximately 15 infants. The illnesses ranged from serious neurological effects, such as seizures, to vomiting, jitteriness and rapid eye movement. Due to the potential for adulteration, on September 10, 2003, the FDA issued an advisory to the public not to consume “teas”

brewed from star anise, until the FDA is able to differentiate between the Japanese star anise and Chinese star anise, which does not contain anisatin [116].

4.3. Substances formed as the result of product abuse

4.3.1. Glycoalkaloids (solanine and chaconine) in potatoes

The glycoalkaloids α -solanine and α -chaconine are natural pesticides that are produced in potatoes. α -Solanine is also found in eggplant, apples, bell peppers, cherries, sugar beets and tomatoes [74,117]. The only difference between α -solanine and α -chaconine is the sugars in the trisaccharide portion of the molecule, *i.e.*, glucose with two rhamnoses for α -solanine and a glucose, galactose and a rhamnose for α -chaconine [118].

Depending on variety and storage conditions, concentrations of α -chaconine and α -solanine in potato tubers vary between 0.5–635 ppm (0.0005–0.64 mg/g potato) and 5–125,100 molecule ppm (0.005–25.1 mg/g potato), respectively (Beckstrom-Sternberg, as cited in Tice 1998 [117], p. 9). Although glycoalkaloids are found throughout the potato tuber, the greatest concentrations are in the sprouts, peels and sun-greened areas [74]. The FDA considers the maximum acceptable glycoalkaloid content to be 20–25 mg/100 g fresh potato weight (or 200–250 ppm) (Crocco, as cited in FDA 2008 [119], p.1). Under current FDA regulations, 20 milligrams of solanine per 100 grams (a small potato) can render it unfit to eat.

Synthesis of α -chaconine and α -solanine is stimulated by light, mechanical injury, aging and potato beetle infestation [117,120]. Exposure of potatoes to light in the field or marketplace can lead to glycoalkaloid concentrations that are unsafe for human consumption. Concentrations of solanine in green or blighted potatoes have been shown to increase by seven fold [73].

The symptoms of acute toxicity to α -solanine and α -chaconine are due to their ability to act as inhibitors of acetylcholinesterase and disruptors of cell membranes. Glycoalkaloid doses of 1 to 5 mg/kg have been shown to be acutely toxic to humans, and doses of 3 to 6 mg/kg have resulted in death [117]. Symptoms of glycoalkaloid toxicity in humans include drowsiness, itchiness in the neck region, increased sensitivity (hyperesthesia), labored breathing and gastrointestinal symptoms (abdominal pain, nausea, vomiting and diarrhea) [74].

α -Solanine and α -chaconine are not mutagenic or only weakly mutagenic *in vitro*, are not genotoxic *in vivo*, and are embryotoxic and teratogenic to experimental animals. Teratogenic effects in mammals include central nervous system abnormalities (e.g., exencephaly, cranial bleb, encephalocele, and anophthalmia), mild hydronephrosis, hydroureter, and irregular or fused ribs. Although one human case study reported a correlation between the severity of potato late-blight and the incidence of spina bifida, no other studies in humans have found a correlation between the consumption of potatoes and birth defects [117]. There is no evidence that α -solanine and α -chaconine are carcinogenic in animals or humans.

In 1993, the National Institute of Environmental Health Sciences determined that the average consumption of glycoalkaloids from potatoes was 12.75 mg glycoalkaloids/person/day (0.18 mg/kg bw based on a bw of 70 kg) [117], which is approximately one-fifth of the lowest dose that has been shown to produce acute toxicity in humans (1 mg/kg bw).

4.3.2. Furocoumarin in parsnips

Ceska *et al.* reported that older 'spoiled' and diseased parsnips freely available in grocery stores may contain furocoumarin concentrations 2500% higher than fresh parsnips [121]. Microbial infection of parsnip roots can result in a dramatic increase in furocoumarin levels. Furocoumarin concentrations (the sum of five furocoumarins: angelicin, isopimpinellin, 5-MOP, 8-MOP and psoralen) in freshly harvested parsnips are generally lower than 2.5 mg/kg and do not increase after storage at $-18\text{ }^{\circ}\text{C}$ for up to 50 days. In contrast, storage of whole parsnips (but not cubes or homogenate) at $4\text{ }^{\circ}\text{C}$ resulted in a marked biphasic increase of furocoumarin concentrations (to approximately 40 mg/kg) after seven or 38 days of storage. A dramatic increase in furocoumarin concentrations (up to 566 mg/kg) was observed when whole parsnips were kept at room temperature over 53 days, resulting in a visible microbial (mold) infection [122].

In celery, infection with fungal pathogens has been shown to produce timethylpsoralen (which is absent from plants that are not infected) and increased concentrations of 8-MOP. The resulting "pink rot" has caused repeated outbreaks of photophytophthyodermatitis in commercial celery handlers [55]. Fungal infection also has been shown to stimulate a 155-fold increase in furocoumarin production by carrots (Ceska *et al.*, as cited in Wagstaff 1991 [53], p. 268). There are no FDA regulations or guidelines specific to the presence of furocoumarins in food.

4.4. Substances formed as the result of processing

4.4.1. Heterocyclic aromatic amines

There are two major classes of heterocyclic aromatic amines (HAAs). Pyrolytic HAAs are formed from the pyrolysis of amino acids or proteins at high temperature and aminoimidazoarenes (AIAs) are formed from creatine, free amino acids and monosaccharides, via the Maillard reaction. HAAs are present in many protein-rich foods of animal origin including cooked meat, fish, poultry and gravies and sauces derived from pan residues and scrapings of cooked meats. The formation and yield of HAAs are dependent on cooking temperature and time (concentrations increase with higher temperatures and longer cooking times), cooking technique and equipment (concentrations of HAAs in meat are generally higher after grilling and panfrying than broiling or roasting), and the ability of HAA precursors to migrate to the surface [123].

The AIAs 2-amino-3-methylimidazo-[4,5-f]quinoline (IQ), 2-amino-3,4-dimethylimidazo[4,5-f]quinoline (MeIQ) and 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) are among the most potent mutagens ever tested in the Ames assay. The pyrolytic AIA 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) and the HAAs 2-amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-P-1), 2-amino-4-methyl-5H-pyrido[4,3-b]indole (Trp-P-2), 2-amino-9H-pyrido[2,3-b]indole (A α C), 2-amino-3-methyl-9H-pyrido[2,3-b]indole (MeA α C) are also mutagenic. PhIP accounts for 75% of the mass of genotoxic material that has been attributed to HAAs in fried ground beef. Therefore, the potential for genotoxicity due to PhIP may be higher than that of more genotoxic HAAs in meat consumers [123].

Several HAAs are carcinogenic in rodents after long-term dietary administration. The doses required to induce tumors at a 50% rate (TD₅₀) vary for each HAA, and range from 0.1 to 64.6 mg/kg bw/day [123]. Four HAAs (IQ, MeIQ, MeIQx and PhIP) are "reasonably anticipated to be

human carcinogens” [124]. Due to the fact that exposure to HAAs in cooked meats is highly variable (concentrations in cooked meat may range from <1 to 500 ng/g), it has been estimated that the risk of developing cancer from exposure to HAAs in food is anywhere from 50 in one million to one in a thousand [123]. Currently, no tolerable upper limit of exposure to HAAs has been established.


4.4.2. Polycyclic aromatic hydrocarbons

Polycyclic aromatic hydrocarbons (PAHs) are known carcinogens that are formed from the incomplete combustion of fossil fuels such as wood, coal and oil. PAHs can enter the food chain from environmental contamination or from food processing. Foods containing the highest concentrations of PAHs include cooked or smoked meat or fish, smoked or cured cheese, tea and roasted coffee. Grilling or broiling of meat, fish or other foods over intense heat or direct contact with flames promotes production of PAHs. In general, concentrations of PAHs in meat are highest after charcoal grilling, followed by smoking, roasting and steaming. Concentrations of PAHs in smoked foods are influenced by temperature, type of wood, oxygen concentration and type of smoker. Concentrations of PAHs in tea dried over burning wood, oil or coal are generally higher than in tea dried over air, and coffee beans that are roasted over a direct fire contain higher concentrations than beans that do not come in contact with flames [125].

The European Commission’s (EC) Scientific Committee on Food and the Joint FAO/WHO Expert Committee on Food Additives (JECFA) has concluded that thirteen different PAHs are genotoxic and carcinogenic benz[a]anthracene, benzo[b]fluoranthene, benzo[j]fluoranthene, benzo[k]fluoranthene, benzo[a]pyrene, chrysene, dibenz[a,h]anthracene, dibenzo[a,e]pyrene, dibenzo[a,h]pyrene, dibenzo[a,i]pyrene, dibenzo[a,l]pyrene, indeno[1,2,3-cd]pyrene and 5-methylchrysene. Three of the four PAHs that have been tested for carcinogenicity in rats after oral exposure (benz[a]anthracene, benzo[a]pyrene and dibenz[a,h]anthracene) are carcinogenic. The estimated high and safe levels of intake of the benchmark PAH benzo[a]pyrene are 0.01 and 100 µg benzo[a]pyrene/kg bw/day, respectively, indicating that the estimated intake of PAHs in food is 10,000-fold lower than the level that is expected to cause toxicity in humans [126]. Currently, no tolerable upper limit of exposure to PAHs has been established by the FDA.

4.4.3. Acrylamide

Acrylamide (Figure 5) is found in a number of starch-based foods that are fried or baked at temperatures greater than 120 °C (248 °F), including bread, bakery products, breakfast cereal, and potato products (e.g., chips, french fries) [127]. It also is found in cocoa-based products and coffee. Acrylamide is formed via a Maillard reaction, a reaction between the carbonyl group of a reducing sugar and the nucleophilic group of an amino acid. Although a number of carbohydrates can be used as the source of the carbonyl group, the amino acid required for the formation of acrylamide is asparagine.

Figure 5. Structure of acrylamide.

Acrylamide is mutagenic and has been shown to be a neurotoxicant, reproductive toxicant and carcinogen in experimental animals and is classified by IARC as a probable human carcinogen. The main metabolite, glycidamide (an epoxide) is thought to be responsible for genotoxicity [127]. In humans, the only toxicological effect that has been linked to acrylamide is neurotoxicity in individuals occupationally exposed to high levels. Epidemiological studies have failed to show an increased risk of cancer from either occupational or dietary exposure to acrylamide and reproductive toxicity has not been reported in humans exposed to acrylamide [128]. Acrylamide is a unique substance that exemplifies the concept that the structure of the substance greatly influences the toxicity, as acrylamide is an animal feed ingredient (thickener and suspending agent) only when a part of a long-chain polymer having a minimum molecular weight of 3 million and a viscosity range of 3,000 to 6,000 centipoises at 77 °F. The residual acrylamide cannot be more than 0.05% (21 CFR 573.120).

In 2005, JECFA estimated that average and high intake consumers ingest 1 or 4 µg/kg bw/day acrylamide from food, respectively. Using a NOAEL for neurotoxicity of 200 µg/kg bw/day in animals, margins of safety of 200 and 50 for the average and high intake groups were derived, respectively. Utilizing a benchmark dose of 0.3 mg/kg bw/day and a NOAEL of 2 mg/kg bw/day for development of mammary tumors or reproductive in rats (respectively), higher margins of safety were calculated for carcinogenicity (300 and 75, respectively) and reproductive toxicity (200 and 50, respectively) [128].

Exposure to acrylamide can be reduced by avoiding deep-fried foods, soaking potato slices before cooking, cooking french fries at lower temperatures and to a lighter color, and toasting bread to a lighter color [127].

4.4.4. Chloropropanols

Chloropropanols are formed in hydrolyzed vegetable proteins (HVP) produced by hydrochloric acid (HCl) hydrolysis of proteinaceous by-products from edible oil extraction, such as soybean meal, rapeseed meal and maize gluten [129,130]. The chloropropanol most commonly found in food is 3-MCPD (3-monochloropropane-1,2-diol), although others may also be present, including 2-MCPD (2-monochloropropane-1,3-diol), 1,3-DCP (1,3-dichloro-2-propanol), and 2,3-DCP (2,3-dichloro-1-propanol) [130]. The two most widely studied chloropropanols are 3-MCPD and 1,3-DCP. It is thought that 3-MCPD is formed as a result of a reaction between a source of chlorine (chlorinated water or sodium chloride) in a food or a food contact material and a lipid. Two basic pathways have been proposed: thermally driven and enzyme-catalyzed (generally lipase) reactions. Direct precursors are thought to be glycerol and chloride. Recent work has also suggested glycidol (2,3-epoxy-1-propanol) as a precursor. 1,3-DCP is thought to arise from 3-MCPD.

High concentrations of 3-MCPD have been found in acid hydrolyzed HVP (acid-HVP), and soy or oyster sauce produced using an acid hydrolysis process. Other foods that may contain 3-MCPD are

cereal, toasted bread, coffee, cheese, licorice, baked goods, processed garlic, liquid smokes, malts, cured or smoked meat or fish or foods containing acid-HVP as a savory ingredient (soups, prepared meals, savory snacks, gravy mixes and stick cubes [129–132]. Foods containing 1,3-DCP include raw meat and soy sauce produced using an acid hydrolysis process [129].


In rats and mice, 3-MCPD is toxic to the kidney, producing renal tubule hyperplasia. It is also carcinogenic in rats when given in high doses over prolonged periods. Although 3-MCPD is genotoxic *in vitro*, it is not *in vivo*. The UK Committee on Carcinogenicity of Chemicals in Food, Consumer Products and the Environment (COC) has concluded that 3-MCPD is unlikely to present a carcinogenic risk to man, provided the exposure is 1000 times lower than the no observed effect level (NOEL) of 1.1 mg/kg bw/day for tumorigenicity. JECFA set a tolerable daily intake (TDI) of 2 µg 3-MCPD/kg of body weight in 2001 and a maximum allowable content of free 3-MCPD in liquid condiments at 0.4 mg/kg (400 µg/kg) in 2008 [130]. Assuming 400 µg/kg 3-MCPD is present in soy sauce, a 60 kg human would have to ingest 300 g of soy sauce (approximately two-thirds of a 444 mL bottle) per day to achieve the TDI. The FDA has provided a policy statement stating that acid-H[V]P or Asian sauces that contain 3-MCPD at levels greater than 1 ppm are not Generally Recognized As Safe (GRAS); therefore, these ingredients are unapproved food additives [133].

1,3-DCP is hepatotoxic, genotoxic and induces a variety of different types of tumors in rats. Therefore, 1,3-DCP is considered to be a potential carcinogen in humans. In 1993, FAO/WHO and JECFA concluded in that 1,3-DCP is an undesirable contaminant in food and that levels should be reduced to as low as “technologically achievable” [131].

4.4.5. Furan

Furan (Figure 6) is a by-product of high-energy and thermal treatment of carbohydrate. Meat and vegetable containing foods that are heat processed in cans and jars (such as soups, pastas, sauces, gravy and baby food) and brewed coffee, typically contain the highest concentrations. Concentrations of furan present in food and coffee range from undetectable to approximately 175 µg/kg [134]. Coffee powders may contain up to 5000 µg/kg on a dry weight basis. Although the mechanism of formation of furan in food is not completely understood, it can be synthesized from vitamin C, amino acids, reducing sugars, organic acids, carotenes and polyunsaturated fatty acids in the presence of heat [135].

Figure 6. Structure of Furan.



Furan is mutagenic and clastogenic in a number of *in vitro* mammalian cell assays, causes damage to chromosomes in mice, and is carcinogenic in both rats and mice after oral administration [134,136–138]. Furan is classified by IARC as possibly carcinogenic to humans [134].

In the United States and Europe, exposure to furan from food is estimated to be a maximum of 1.00 and 1.75 µg/kg bw/day, respectively [134]. The upper estimate of consumption is approximately

300 and 1000-fold lower than the NOAELs for cytotoxicity and hepatocarcinogenicity of 500 and 2000 µg/kg bw in female B6C3F1 mice, determined by Moser *et al.* [136].

Mitigation of furan in foods is difficult because the mechanism for its formation in food is unclear. Due to the fact that furan is volatile, it is thought that concentrations can be reduced by heating food in open containers or leaving ready-to-eat foods open to air after preparation. However, the effectiveness of this strategy in reducing exposure to furan has yet to be demonstrated [135]. Currently, there are no FDA regulations specific to the level of furan in food.

4.4.6. Trans fatty acids

Trans fatty acids (also known as trans fat) are the sum of all unsaturated fatty acids that contain one or more isolated double bonds in a trans configuration. Trans fatty acids more closely resemble saturated fatty acids than cis unsaturated fatty acids because their trans configuration makes them rigid. Trans fatty acids in the diet originate from two sources. The first is from bacterial hydrogenation in the forestomach of ruminants, which produces trans fatty acids that are found in beef and mutton fat, milk and butter. Trans fatty acids are also produced from the hydrogenation of liquid oils (mainly of vegetable origin). This produces solid fats and partially hydrogenated oils such as margarines, spreads, shortenings and frying oil, which are more stable than liquid oils [139].

Biochemically, trans-fatty acids act similarly to saturated fatty acids, raising low density lipoprotein (LDL) cholesterol and decreasing high-density lipoprotein (HDL) cholesterol levels [139]. High intakes of trans fatty acids have been associated with an increased risk of coronary heart disease (CHD) independent of other risk factors in large epidemiological studies [140]. A tolerable upper limit of trans fatty acids has not been set because any incremental increase in the intake of trans fatty acids increases the risk of coronary heart disease [141].

In the US, the main sources of intake of trans fatty acids are baked goods (28%), fried foods (25%), margarine, spreads and shortenings (25%), savory snacks (10%), milk and butter (9%) [139]. In 1996, processed foods and oils accounted for 80% of the trans fat in the diet [141]. In 1999, the FDA estimated that the average daily intake of trans fat in the United States is about 5.8 grams or 2.6% of calories per day [142]. It has been hypothesized that replacing 2% energy from trans fatty acids with 2% energy from oleic acid would reduce mean plasma LDL cholesterol concentration by 0.08 mmol/L, and increase plasma HDL concentration by 0.08 mmol/L. These changes could reduce the incidence of CHD by 5–15% [139].

Due to increased efforts by food manufacturers to reduce or eliminate the use of partially hydrogenated vegetable fat in food production, it is estimated that trans fatty acid content of processed foods has decreased over the last decade [143].

4.4.7. Nitrosamines formed during drying, curing and preserving

Nitrosamines are formed from the interaction of nitrites or other nitrosating agents with amines in food (or *in vivo*), under acidic conditions. Nitrites may be directly added to food or can be formed from bacterial reduction of nitrate. Nitrites and nitrates may occur naturally in water or foods such as leafy vegetables due to the use of fertilizer, or may be added to foods to prevent growth of *Clostridium botulinum*, or to add color or flavor [144].

Nitrosamines have been found in a variety of different foods such as cheese, soybean oil, canned fruit, meat products, cured or smoked meats, fish and fish products, spices used for meat curing, and beer and other alcoholic beverages [145]. Beer, meat products and fish are considered the main sources of exposure. Drying, kilning, salting, smoking or curing promotes formation of nitrosamines [146].

The nitrosamines most frequently found in food are nitrosodimethylamine (NDMA), *N*-nitrosopyrrolidine (NPYR), *N*-nitrosopiperidine (NPIP), and *N*-nitrosothiazolidine (NTHZ) [146]. NDMA, NPYR, NPIP are reasonably anticipated to be human carcinogens based on evidence of carcinogenicity in experimental animals [145,147,148]. Evidence from case-control studies supports an association between nitrosamine intake with gastric cancer, but not esophageal cancer in humans [149].

Levels of nitrosamines have been declining during the past three decades, concurrent with a lowering of the nitrite used in food, use of inhibitors such as ascorbic acid and use of lower operating temperatures and indirect heating during food processing. Based on an estimated exposure level of 3.3–5.0 ng/kg bw/day, the and the benchmark lower limit of 60 µg/kg bw/day, a margin of error associated with a low level of concern (12,000–18,2000) has been derived for NDMA, the most common nitrosamine in food [146].

Although current FDA regulations do not limit nitrosamine levels in foods, the FDA has provided an action level of 10 ppb for individual nitrosamines in both consumer and hospital rubber baby bottle nipples, while the FDA limits the approval of nitrites in curing mixes to the FDA-regulated food additive process (21 CFR 170.60), with the approval of sodium nitrite as a food additive (food preservative) (21 CFR 172.175). The USDA monitors finished meat products to insure that nitrite is not present in amounts exceeding 200 ppm (9 CFR 424.21).

4.4.8. Biogenic amines

Biogenic amines are normally formed in humans by normal cellular metabolism. In food, biogenic amines are mainly formed from microbial decarboxylation of amino acids. They are commonly found in fermented meat, beverages and dairy products, sauerkraut, and spoiled fish. The main biogenic amines in food are histamine, tyramine, cadaverine, putrescine, spermidine and spermine. The two biogenic amines that have been associated with acute toxicity are histamine and tyramine. Putrescine, spermine, sperimidine and cadaverine are not toxic in and of themselves, but may react with nitrite or nitrate to form nitrosamines (see Section 4.4.7 above) [150].

Scombrototoxicosis is a common seafood-borne disease associated with the consumption of toxic levels of histamine in spoiled scombroid fish such as tuna (*Thunnus* spp.), mackerel (*Scomber* spp.), saury (*Cololabis* saira) and bonito (*Sarda* spp.). Red wine may also contain relatively high levels of histamine. Symptoms of histamine intoxication from food are similar to allergies to other substances and include sneezing, nose congestion, breathing difficulties and urticaria [150].

Consumption of tyramine may precipitate migraine headache or a hypertensive crisis. The most serious case reports of tyramine toxicity have occurred in people consuming aged cheese. Because monoamine oxidase inhibitor (MAOI) drugs inhibit metabolism of amines, people taking these drugs may be particularly susceptible to tyramine toxicity. Whereas 200–800 mg of dietary tyramine induces only a mild rise in blood pressure in unmedicated adults, 10–25 mg may produce a serious adverse

event in those taking MAOI drugs. Other potentiating factors for tyramine toxicity include alcohol consumption, gastrointestinal distress and exposure to other amines [150].

Efforts taken by food manufacturers to reduce biogenic amine concentrations in fermented foods include using amine-negative starter cultures, adding probiotic bacterial strains alone or in combination with starter cultures, high pressure processing or low-dose gamma radiation [150]. FDA guidelines specify 50 mg/100 g as the toxic concentration of histamine in scombroid fish and the agency has published guidance on how to control levels [151].

5. Substances Passed from Animals to Humans

5.1. Toxins in seafood

5.1.1. Toxins involving algae

Consumption of seafood contaminated with algal toxins results in five different syndromes, paralytic, neurotoxic, amnesic, or diarrhetic shellfish poisoning and ciguatera fish poisoning [152].

5.1.1.1. Paralytic shellfish poisoning

Paralytic shellfish poisoning (PSP) is caused by the consumption of molluscan shellfish contaminated with heterocyclic guanidines called saxitoxins. Currently, over 21 known saxitoxins are produced by dinoflagellate species from three genera: *Alexandrium*, *Gymnodium* and *Pyrodinium*. Toxicity is caused by binding of saxitoxins to voltage-dependent sodium channels, which blocks neuronal activity. The primary site of action in humans is the peripheral nervous system. Symptoms of toxicity include tingling and numbness of the perioral area and extremities, loss of motor control, drowsiness, and incoherence. Ingestion of 1–4 mg saxitoxin has resulted in death from respiratory paralysis [152].

Outbreaks of PSP have occurred worldwide, due to the fact that saxitoxin-producing species of dinoflagellates can live in either temperate or tropical waters. Saxitoxins are not inactivated by cooking, and must be mitigated at their source to prevent ingestion. PSP is prevented by large-scale, proactive monitoring programs and rapid closures of harvest in areas containing dinoflagellate algal blooms [153]. In the United States, the permissible level of saxitoxin equivalents in shellfish is 80 micrograms/100 grams [154].

5.1.1.2. Neurotoxic shellfish poisoning

The dinoflagellate *Karenia brevis* produces brevetoxins that are lethal to fish, but not to mollusks such as oysters, clams and mussels. Consequently, they can accumulate in healthy-appearing mollusks to concentrations that are toxic to humans who ingest them. *Karenia brevis* brevetoxins cause the syndrome known as neurotoxic shellfish poisoning (NSP), which affects sodium transport in the autonomic nervous system and causes inhibition of neuromuscular transmission in skeletal muscle.

NSP is usually a relatively mild illness and should not be confused with the more serious condition of PSP. NSP symptoms usually occur within three hours of ingesting contaminated shellfish and may include abdominal pain, nausea and vomiting, vertigo, malaise, generalized muscle weakness, ataxia,

incoordination, chills, headache, myalgia, a reversal of hot/cold temperature sensation and progressive parasthesias. Dilated pupils, bradycardia and convulsions may occur in cases of severe poisoning [155]. Unlike PSP, no deaths have been reported from NSP [152].

K. brevis is the organism that is usually responsible for the red tides in the Gulf of Mexico and along the southern Atlantic coast of North America. Blooms along the west coast of Florida occur regularly [156]. Biotoxin control plans that are implemented during period of red tide are generally effective in preventing NSP, but have not eliminated NSP entirely.

The FDA has established an action level of 0.8 ppm (20 mouse units/100 g) brevetoxin-2 equivalents [154].

5.1.1.3. Amnesic shellfish poisoning (Domoic acid)

Amnesic shellfish poisoning (ASP) is caused by domoic acid produced by diatoms of the genus *Pseudo-nitzschia* (Figure 7), which are consumed by mussels, scallops, clams and crabs. Domoic acid is a water-soluble, tricarboxylic amino acid that is a structural analog of the neurotransmitter glutamate and is a glutamate receptor agonist. Persistent activation of the kainite glutamate receptor causes an increase in intracellular calcium, which can cause neuronal cell death and lesions of the brain where glutaminergic pathways are concentrated. Areas of the brain involved in learning and memory processing are particularly susceptible [152]. The symptoms of ASP are gastroenteritis, dizziness, disorientation, lethargy, seizures and loss of short term memory. Respiratory difficulty, coma and death may ensue [153]. Human toxicity has occurred after ingestion of 1–5 mg/kg domoic acid [152].

Figure 7. *Pseudo-nitzschia* [157].



In 1987, approximately 100 people became ill and died in Prince Edward Island, Canada, after eating contaminated mussels. In 1991, domoic acid poisoning caused the deaths of numerous pelicans and cormorants in Monterey Bay that ingested sardines and anchovies. Domoic acid also was responsible for a massive sea lion kill in Monterey Bay in 1998 [158]. *Pseudo-nitzschia* and domoic acid are now closely monitored throughout the world [159]. The FDA has established an action level of 20 ppm for domoic acid, except in the viscera of Dungeness crab, where 30 ppm is permitted [154]. Regulatory guidance has been effective in preventing ASP in humans, since no human outbreaks of ASP have occurred since 1987.

5.1.1.4. Diarrhetic shellfish poisoning

Diarrhetic shellfish poisoning (DSP) is caused by the production of okadaic acid and dinophysistoxins in the dinoflagellates *Dinophysis fortii* or *Prorocentrum lima*, which are consumed by mollusks. Okadaic acid and dinophysistoxins are inhibitors of serine/threonine phosphatases, critical components of signaling cascades that regulate a number of cellular processes involved in metabolism, ion balance, neurotransmission and cell cycle regulation [152].

Compared to other types of shellfish poisoning, symptoms of DSP are relatively mild, and generally consist of diarrhea, abdominal cramps, nausea, chills or vomiting within 30 minutes to a few hours after consumption of DSP toxins. Symptoms generally resolve within 2–3 days, with or without medical treatment [153]. Diarrhea is most likely due to the hyperphosphorylation of proteins (including ion channels) in the intestinal epithelia, resulting in impaired water balance and fluid loss. The long term consequences of low level exposure to DSP toxins may be more serious, as they have been shown to be tumor promoters [152]. The FDA has established an action level of 0.2 ppm okadaic acid plus 35-methyl okadaic acid (DXT 1) [154].

5.1.1.5. Ciguatera poisoning

Ciguatera fish poisoning (CFP) is caused by the dinoflagellate *Gambierdiscus toxicus*, which grows on filamentous macroalgae associated with coral reefs. The lipophilic precursors to ciguatoxin are biotransformed to ciguatoxins in herbivorous fish and invertebrates that consume the macroalgae, and bioaccumulate in large carnivorous fishes associated with coral reefs. High ciguatoxin concentrations may be found in barracuda, snapper, grouper and jacks [152].

Ciguatoxins are structurally related to the brevetoxins and compete with brevetoxin for binding to the same site on the voltage-dependent sodium channel. However, because ciguatoxin has a higher binding affinity for the site than brevetoxin, the toxic potency of ciguatoxin is higher than that of brevetoxin. The threshold level for toxicity in humans is estimated to be 0.5 ng/g [152].

CFP is estimated to affect over 50,000 people worldwide each year. The symptoms of CFP generally include gastrointestinal disturbances (nausea, vomiting and diarrhea) within 2–6 hours, followed by neurologic symptoms such as numbness of the perioral area and extremities, a reversal of hot/cold temperature sensation, muscle and joint aches, headache, itching, tachycardia, hypertension, blurred vision and paralysis. In rare cases, CFP is fatal [152].

Inasmuch as ciguatoxin is produced by organisms that live beneath the surface and is not routinely monitored for concentration in seafood, the only way to prevent consumption is to completely abstain from ingesting tropical reef fish, as the occurrence of toxic fish is sporadic, and not all fish of a given species or from a given locality will be toxic [153]. Currently, there are no FDA regulations limiting levels of ciguatoxins in fish, although a recent publication suggests an advisory level of 0.1 ppb pacific ciguatoxin equivalent (P-CTX-1) toxicity values in fish from the tropical Atlantic, Gulf of Mexico, Caribbean, and 0.01 ppb P-CTX-1 equivalent toxicity in fish from Pacific regions [160].

5.1.2. Toxins not involving algae

5.1.2.1. Gempylotoxin

There are naturally occurring toxins in some species that do not involve marine algae. Escolar (*Lepidocybium flavobrunneum*, Figure 8), and Oilfish or Cocco (*Ruvettus pretiosus*), a marine fish of the snake mackerel family, are sometimes sold under the category of “butterfish”, and contain a strong purgative oil, that when consumed can cause diarrhea known as Gempylid Fish Poisoning, Gempylotoxism or Keriorrhea [161]. The toxin consists of wax esters (C32, C34, C36 and C38 fatty acid esters), the primary component of which is C₃₄H₆₆O₂ [162]; these constitute a substantive portion of the lipid present in these fish (14–25% by weight). Escolar oil contains >90% wax esters [163]. Ingestion of fish containing wax esters in large amounts, coupled with their indigestibility and low melting point, results in diarrhea [164]. No tolerances have been established, and the FDA recommends avoidance of these fish [161].

Figure 8. Juvenile Oilfish (*Ruvettus pretiosus*) [165].



5.1.2.2. Tetramine in whelks

Tetramine is a toxin found in the salivary glands of *Buccinum*, *Busycon* or *Neptunia* spp., a type of whelk or sea snail that is distributed in temperate and tropic waters and has long been a food source for humans. Whelk are associated with a heat-stable neurotoxin, tetramine, which upon ingestion produces, among other symptoms, eyeball pain, headache, dizziness, abdominal pain, ataxia, tingling in the fingers, nausea and diarrhea [166,167]. Power *et al.* report that the highest concentration of tetramine is in the salivary gland (up to 6530 µg/g), but varies according to season [168]. Reid *et al.* reported levels of 37.5 µg tetramine/g of salivary gland tissue [166]. Because the whelk is a predator of bivalves, it is assumed the toxin is used for food procurement [168]. Although the FDA recommends removal of the salivary gland to avoid possible intoxication [154], tetramine is present in other tissues, albeit at lesser concentrations [169].

5.1.2.3. Trimethylamine oxide

The meat of the Greenland shark (*Somniosus microcephalus*) and the related member of the dogfish family, the pacific sleeper shark (*Somniosus pacificus*), is known to be poisonous to both man and dogs. The causative agent is trimethylamine oxide, which breaks down to trimethylamine in the gut, probably by enteric bacteria. The result is absorption of trimethylamine, which acts as a neurotoxin,


producing ataxia in both man and dogs. However, the flesh may be consumed if boiled several times with changes of water, or as the Inuit prepares it, by burying it in the ground and allowing the meat to go through several freezing and thawing cycles [170–172].

5.2. Toxins from animal, non-seafood sources passed on to humans

5.2.1. Grayanotoxins in honey and direct contact with food

Rhododendrons and azaleas (*Rhododendron* spp.), oleander (*Nerium oleander* or *Nerium indicum*), mountain laurel (*Kalmia latifolia*) and sheep laurel (*Kalmia angustifolia*), all produce grayanotoxins (Figure 9) whose action is to bind to sodium channels in muscle, including the heart. Although not all rhododendrons produce grayanotoxins (also known as oleander toxin, andromedotoxin, acetylandromedol or rhodotoxin), several species growing in the US are known to produce grayanotoxins and include *Rhododendron occidentale*, *Rhododendron macrophyllum* and *Rhododendron albiflorum*, all in the western US. Grayanotoxin is also found in the eastern US, within the botanical family Ericaceae, to which rhododendrons belong and are probably the most important sources of the toxin [173].

Figure 9. Grayanotoxins [173].



Grayanotoxin consists of a series of cardiac glycosides: thevetin, convallarin, steroidal, helleborein, ouabain, and digitoxin. At first, sympathetic nerves are paralyzed; the cardiotoxin stimulates the heart muscles similar to the action of digitalis, and gastric distress ensues. Symptoms start out as nausea, vomiting, abdominal pain and diarrhea; followed by tremor, drowsiness and ataxia. In severe cases, ectopic beats occur which may be followed by ventricular tachycardia and fibrillation. The origin of toxicity may be honey (made from the nectar of the flowers), milk from a cow having eaten the foliage and meat (e.g., hot dogs) roasted on oleander sticks [15,174]. The pooling of large quantities of grayanotoxin-containing honey or milk during commercial processing typically dilutes grayanotoxin to nontoxic levels. There are no FDA regulations specific to grayanotoxin levels in foods.

5.2.2. Tremetol contamination of milk from white snakeroot

“Milk sickness” also known as “puking fever”, “sick stomach”, “the slows” and “the trembles”, was a mysterious scourge of the Midwest United States in the 18th and 19th centuries. Thousands of people have been reported as dying, including Abraham Lincoln’s mother, Nancy Hanks Lincoln. In humans, milk sickness is characterized by loss of appetite, listlessness, weakness, vague pains, muscle stiffness, vomiting, abdominal discomfort, constipation, foul breath and finally, coma. For many years the origin of milk sickness was unknown, because there was nothing comparable in Europe (origin of most of the pioneers) and the outbreaks were sporadic. It was not recognized until the late 19th and early 20th century, that white snakeroot (*Ageratina altissima* née *Eupatorium rugosum*) and rayless goldenrod (*Bigelowia* spp., *Haplopappus heterophyllus* and *Isocoma pluriflora*) when eaten by cattle, was the source. The sporadic nature of outbreaks became clear when it was realized that cattle would consume these plants in over-grazed pasture or in years of drought; additionally, the toxin levels in plants can vary considerably, making identification of the source of poisonings difficult. Tremetol or tremetone is the toxic agent and consists of a mixture of sterols and derivatives of methyl ketone benzofuran. The three major benzofuran ketones are tremetone, dehydrotremetone and 3-oxyangeloyl-tremetone [173–177]. Currently, there is no USDA guidance specific to tremetol levels in dairy products.

6. Conclusions

Given the state of the science, the pressure on the food supply and the development of new products, the FDA has performed admirably in protecting the consumer from exposure to toxins in food with its judicious use of warning labels, action levels, tolerances, specifications, prohibitions and the ability conferred by Congress to declare substances “unsafe” or “unfit for food.” However, the FDA cannot protect consumers absolutely from exposure to toxins normally present in foods. At normal levels of food consumption, there is little potential for toxicity from natural food toxins. Nevertheless, there is always the possibility of an idiosyncratic response or undetected contamination.

References

1. Institute of Medicine (IOM). *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium and Zinc*; National Academies Press: Washington, DC, USA, 2001.
2. Burton, G.W.; Ingold, K.U. beta-Carotene: An unusual type of lipid antioxidant. *Science* **1984**, *224*, 569–573.
3. Bannister, B.; Gibsburg, G.; Shneerson, T. Cardiac arrest due to liquorice-induced hypokalaemia. *Br. Med. J.* **1977**, *2*, 738–739.
4. Isbrucker, R.A.; Burdock, G.A. Risk and safety assessment on the consumption of licorice root (*Glycyrrhiza* sp.), its extract and powder as a food ingredient, with emphasis on the pharmacology and toxicology of glycyrrhizin. *Regul. Toxicol. Pharmacol.* **2006**, *46*, 167–192.

5. United States Government Accountability Office (GAO). Food safety. FDA should strengthen its oversight of food ingredients determined to be generally recognized as safe (GRAS). GAO-10-246, February, 2010. Available online: <http://www.gao.gov/new.items/d10246.pdf> (accessed on 21 July 2010).
6. Food and Drug Law Institute. Sec. 201. [321] Definitions. In *FDCA Statutory Supplement Including FDA Amendments Act of 2007 and Related Sections of Additional Statutes*; Food and Drug Law Institute: Washington, DC, USA, 2008; pp. 1–2.
7. Food and Drug Law Institute. Sec. 402. [342] Adulterated Food. In *FDCA Statutory Supplement Including FDA Amendments Act of 2007 and Related Sections of Additional Statutes*; Food and Drug Law Institute: Washington, DC, USA, 2008; p. 31.
8. Food and Drug Law Institute. Sec. 406. [346] Tolerances for Poisonous Ingredients in Food. In *FDCA Statutory Supplement Including FDA Amendments Act of 2007 and Related Sections of Additional Statutes*; Food and Drug Law Institute: Washington, DC, USA, 2008; p. 31.
9. Kracov, D.A. The regulation of foods and food additives. In *A Practical Guide to Food and Drug Law Regulation*, 2nd ed.; Piña, K.R., Pines, W.L., Eds.; Food and Drug Law Institute: Washington, DC, USA, 2002; pp. 159–214.
10. Tullo, A. Newsprints: Vile weed or essential ingredient? *Chem. Eng. News* **2010**, *88*, 72.
11. Fischer, R.; Griffin, F.; Kaplan, A.R. Taste thresholds, cigarette smoking, and food dislikes. *Med. Exp. Int. J. Exp. Med.* **1963**, *9*, 151–167.
12. Goff, S.A.; Klee, H.J. Plant volatile compounds: Sensory clues for health and nutritional value? *Science* **2006**, *311*, 815–819.
13. National Organization for Rare Disorders (NORD), 2010. Available online: <http://www.rarediseases.org> (accessed on 21 July 2010).
14. Carabin, I.G.; Magnuson, B.A. New Labeling Requirements for Food Allergens, April, 2006. Nutritional Outlook. Available online: <http://www.nutritionaloutlook.com/article.php?ArticleID=2096> (accessed on 21 July 2010).
15. Kotsonis, F.N.; Burdock, G.A. Food Toxicology. In *Casarett and Doull's Toxicology: The Basic Science of Poisons*, 7th ed.; Klaassen, C.D., Ed.; McGraw-Hill: New York, NY, USA, 2008; pp. 1191–1236.
16. Sors, T.G.; Ellis, D.R.; Salt, D.E. Selenium uptake, translocation, assimilation and metabolic fate in plants. *Photosynth. Res.* **2005**, *86*, 373–389.
17. Yang, G.; Wang, S.; Zhou, R.; Sun, S. Endemic selenium intoxication of humans in China. *Am. J. Clin. Nutr.* **1983**, *37*, 872–881.
18. Reilly, C. Selenium: Physiology, dietary sources and requirements. In *Encyclopaedia of Human Nutrition*; Sadler, M.J., Ed.; Academic: San Diego, CA, USA, 1998; pp. 1752–1758.
19. United States Environmental Protection Agency (EPA). Selenium and compounds (CASRN 7782-49-2), March 1, 1991. Available online: <http://www.epa.gov/iris/subst/0472.htm> (accessed on 21 July 2010).
20. Waldron, H.A. Did the Mad Hatter have mercury poisoning? *Br. Med. J.* **1983**, *287*, 1961.

21. Carrington, C.; Bolger, M. An Exposure Assessment for Methylmercury from Seafood for Consumers in the United States. Available online: <http://www.fda.gov/downloads/Food/FoodSafety/Product-SpecificInformation/Seafood/FoodbornePathogensContaminants/Methylmercury/UCM114740.pdf> (accessed on 21 July 2010).
22. United States Food and Drug Administration (FDA). Chapter 10: Methyl Mercury. In *Fish and Fisheries Products Hazards and Controls Guidance*, 3rd ed, June, 2001. Available online: <http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Seafood/ucm092041.htm> (accessed on 21 July 2010).
23. Hutt, P.B.; Merrill, R.A.; Grossman, L.W. *Food and Drug Law*, 3rd ed.; Foundation Press: New York, NY, USA, 2007; p. 369.
24. European Commission. Scientific Committee on Food. Opinion of the scientific committee on food on thujone, February 6, 2003. Available online: http://ec.europa.eu/food/fs/sc/scf/out162_en.pdf (accessed on 21 July 2010).
25. United States Food and Drug Administration. *Code of Federal Regulations (CFR) 21 § 172.510*; U.S. Government Printing Office: Washington, DC, USA, 2006; pp. 55–57.
26. Galli, C.L.; Galli, G.; Tragni, E.; Caruso, D.; Fiecchi A. Quantitative analysis of alpha, beta-thujone, pulegone, safrole, coumarin and beta-asarone in alcoholic beverages by selected-ion monitoring. *J. Appl. Toxicol.* **1984**, *4*, 273–276.
27. Lawrence, B.M. Progress in essential oils. Sage oil. In *Essential Oils: 2001–2004*; Allured Publishing: Carol Stream, IL, USA, 2006; pp. 25–30.
28. Ben Farhat, M.; Jordán, M.J.; Chaouech-Hamada, R.; Landoulsi, A.; Sotomayor, J.A. Variations in essential oil, phenolic compounds, and antioxidant activity of tunisian cultivated *Salvia officinalis* L. *J. Agric. Food Chem.* **2009**, *57*, 10349–10356.
29. Patocka, J.; Plucar, B. Pharmacology and toxicology of absinthe. *J. Appl. Biomed.* **2003**, *1*, 199–205.
30. Millet, Y.; Jouglard, J.; Steinmetz, M.D.; Tognetti, P.; Joanny, P.; Arditti, J. Toxicity of some essential plant oils. Clinical and experimental study. *Clin. Toxicol.* **1981**, *18*, 1485–1498.
31. Bonkovsky, H.L.; Cable, E.E.; Cable, J.W.; Donohue, S.E.; White, E.C.; Greene, Y.J.; Lambrecht, R.W.; Srivastava, K.K.; Arnold, W.N. Porphyrinogenic properties of the terpenes camphor, pinene, and thujone. *Biochem. Pharmacol.* **1992**, *43*, 2359–2368.
32. United States National Toxicology Program (NTP). Alpha-Thujone, December 10, 1997. Available online: <http://ntp.niehs.nih.gov/index.cfm?objectid=03DB8C36-E7A1-9889-3BDF8436F2A8C51F> (accessed on 21 July 2010).
33. Hold, K.M.; Sirisoma, N.S.; Casida, J.E. Detoxification of alpha- and beta-thujones (the active ingredients of absinthe): Site specificity and species differences in cytochrome P450 oxidation *in vivo* and *in vivo*. *Chem. Res. Toxicol.* **2001**, *14*, 589–595.
34. Perdue University, Cooperative Extension Service (Perdue). Indiana plants poisonous to livestock and pets. Available online: <http://www.vet.purdue.edu/toxic/plant46.htm> (accessed on 21 July 2010).
35. Merck. Cyanide Poisoning: Introduction. In *The Merck Veterinary Manual*; 2008. Available online: <http://www.merckvetmanual.com/mvm/index.jsp?cfile=htm/bc/210800.htm&word=prussic%2cacid> (accessed on 21 July 2010).

36. Panter, K.E. Natural toxins of plant origin. In *Toxins in Food*; Dabrowski, W.M., Sikorski, Z.E., Eds.; CRC Press: Boca Raton, FL, USA, 2004; pp. 11–63.
37. Wentworth, J.M.; Agostini, M.; Love, J.; Schwabe, J.W.; Chatterjee, V.K. St John's wort, a herbal antidepressant, activates the steroid X receptor. *J. Endocrinol.* **2000**, *166*, R11–R16.
38. Karioti, A.; Bilia, A.R. Hypericins as potential leads for new therapeutics. *Int. J. Mol. Sci.* **2010**, *11*, 562–594.
39. Hammerness, P.; Basch, E.; Ulbricht, C.; Barrette, E.P.; Foppa, I.; Basch, S.; Bent, S.; Boon, H.; Ernst, E. St. John's Wort: A systematic review of adverse effects and drug interactions for the consultation psychiatrist. *Psychosomatics* **2003**, *44*, 271–282.
40. Britton, N.L.; Brown, A. *Hypericum perforatum* L. In *An illustrated Flora of the Northern United States, Canada and the British Possessions*; Charles Scribner's Sons: New York, NY, USA, 1913; Volume 2, p. 533. USDA-NRCS PLANTS Database. Available online: http://plants.usda.gov/java/profile?symbol=HYPE&photoID=hype_001_avd.tif (accessed on 31 August 2010).
41. State of Victoria Department of Primary Industries (Victoria). Landcare notes. St. John's wort, 2007. Available online: [http://www.dpi.vic.gov.au/dpi/nreninf.nsf/93a98744f6ec41bd4a256c8e00013aa9/9f65b9c41bbc7aa5ca25737500119160/\\$FILE/LC0177_Sep07.pdf](http://www.dpi.vic.gov.au/dpi/nreninf.nsf/93a98744f6ec41bd4a256c8e00013aa9/9f65b9c41bbc7aa5ca25737500119160/$FILE/LC0177_Sep07.pdf) (accessed on 21 July 2010).
42. Greer, M.A. Goitrogenic substances in food. *Am. J. Clin. Nutr.* **1957**, *5*, 440–444.
43. Conn, E.E. Cyanogenetic Glycosides. In *Toxicants Occurring Naturally in Foods*, 2nd ed.; Committee on Food Protection, Food and Nutrition Board, National Research Council, National Academy of Sciences: Washington, DC, USA, 1973; pp. 299–308.
44. VenEtten, C.H.; Wolff, I.A. Natural sulfur compounds. In *Toxicants Occurring Naturally in Foods*, 2nd ed.; Committee on Food Protection, Food and Nutrition Board, National Research Council, National Academy of Sciences: Washington, DC, USA, 1973; pp. 210–234.
45. United States Department of Agriculture (USDA). Plants Profile: *Brassica napus* L. Available online: <http://plants.usda.gov/java/profile?symbol=BRNA> (accessed on 21 July 2010).
46. Carroll, K.K. Erucic acid as the factor in rape oil affecting adrenal cholesterol in the rat. *J. Biol. Chem.* **1953**, *200*, 287–292.
47. Chien, K.R.; Bellary, A.; Nicar, M.; Mukherjee, A.; Buja, L.M. Induction of a reversible cardiac lipidosis by a dietary long-chain fatty acid (erucic acid). *Am. J. Pathol.* **1983**, *112*, 68–77.
48. Ratanasethkul, C.; Riddell, C.; Salmon, R.E.; O'Neil, J.B. Pathological changes in chickens, ducks and turkeys fed high levels of rapeseed oil. *Can. J. Comp. Med.* **1976**, *40*, 360–369.
49. Mattson, F.H. Potential toxicity of food lipids. In *Toxicants Occurring Naturally in Foods*, 2nd ed.; Committee on Food Protection, Food and Nutrition Board, National Research Council, National Academy of Sciences: Washington, DC, USA, 1973; pp. 189–209.
50. Mori, H.; Tanaka, T.; Hirono, I. Toxicants in Food: Naturally Occurring. In *Nutrition and Chemical Toxicity*; Ioannides, C., Ed.; John Wiley & Sons: West Sussex, England, UK, 1998; pp. 1–27.

51. Biotechnology Australia (Australian Government). "What is canola?" A problem with weeds—the canola story. Available online: <http://www.biotechnologyonline.gov.au/foodag/weeds.html> (accessed on 21 July 2010).
52. Health Canada. "Low Erucic Acid Rapeseed (Lear) Oil Derived From Canola-quality *Brassica juncea* (L.) CZERN. Lines PC 97-03, PC98-44 AND PC98-45", March 27, 2003. Available online: http://www.hc-sc.gc.ca/fn-an/gmf-agm/appro/low_erucic-faible_erucique-eng.php (accessed on 21 July 2010).
53. Wagstaff, D. Dietary exposure to furocoumarins. *Regul. Toxicol. Pharmacol.* **1991**, *14*, 261–272.
54. Ashwood-Smith, M.J.; Ceska, O.; Chaudhary, S.K.; Warrington, P.J.; Woodcock, P. Detection of furocoumarins in plants and plant products with an ultrasensitive biological photoassay employing a DNA-repair-deficient bacterium. *J. Chem. Ecol.* **1986**, *12*, 915–932.
55. Zobel, A.M.; Brown, S.A. Dermatitis-inducing psoralens on the surfaces of seven medicinal plant species. *J. Toxicol. Cutaneous Ocul. Toxicol.* **1991**, *10*, 223–231.
56. Dunnick, J.K. NTP Technical Report on the Toxicology and Carcinogenesis Studies of 8-Methoxypsoralen (CAS No. 298-81-7) in F344/N Rats. NIH Publication No. 89-2814. National Toxicology Program: Research Triangle Park, NC, USA, 1989.
57. International Agency for Research on Cancer (IARC). Summaries & Evaluations, 8-Methoxypsoralen (Methoxsalen) plus ultraviolet radiation. *IARC* **1987**, *7* (Suppl.), 261. Available online: <http://www.inchem.org/documents/iarc/suppl7/methoxypsoralen-8.html> (accessed on 21 July 2010).
58. Stern, R.S.; Nichols, K.T.; Vakeva, L.H. Malignant melanoma in patients treated for psoriasis with methoxsalen (psoralen) and ultraviolet A radiation (PUVA). The PUVA follow-up study. *N. Engl. J. Med.* **1997**, *336*, 1041–1045.
59. International Agency for Research on Cancer (IARC) Summaries & Evaluations, 5-Methoxypsoralen. *IARC* **1986**, *40*, 327. Available online: <http://www.inchem.org/documents/iarc/vol40/5-methoxypsoralen.html> (accessed on 21 July 2010).
60. Girenavar, B.; Poulouse, S.M.; Jayaprakasha, G.K.; Bhat, N.G.; Patil, B.S. Furocoumarins from grapefruit juice and their effect on human CYP3A4 and CYP1B1 isoenzymes. *Bioorg. Med. Chem.* **2006**, *14*, 2606–2612.
61. Bailey, D.G.; Malcom, J.; Arnold, O.; Spence, J.D. Grapefruit juice-drug interactions. *Br. J. Clin. Pharmacol.* **1998**, *46*, 101–110.
62. Duke, J.A. *Handbook of Phytochemical Constituents of GRAS Herbs and Other Economic Plants*; CRC Press: Boca Raton, FL, USA, 1992; pp. 171, 174, 180, 183.
63. Placzek, M.; Fromel, W.; Eberlein, B.; Gilbertz, K.P.; Przybilla, B. Evaluation of phototoxic properties of fragrances. *Acta Derm. Venereol.* **2007**, *87*, 312–316.
64. Marzulli, F.N.; Maibach, H.I. Perfume phototoxicity. *J. Soc. Cosmet. Chem.* **1970**, *21*, 695–715.
65. Coulumbe, R.A., Jr. Natural toxins and chemopreventives in plants. In *Food Toxicology*; Helferich, W., Winter, C.K., Eds.; CRC Press: Boca Raton, FL, USA, 2001; p. 152.
66. Schlatter, J.; Zimmerli, B.; Dick, R.; Panizzon, R.; Schlatter, C. Dietary intake and risk assessment of phototoxic furocoumarins in humans. *Food Chem. Toxicol.* **1991**, *29*, 523–530.

67. Deshpande, S.S. Food Additives. In *Handbook of Food Toxicology*; Marcel Dekker: New York, NY, USA, 2002a; pp. 219–284.
68. Nutrilab, Inc. v. S. Schweiker, 713 F.2d 335 (7th Cir. 1983). Available online: <http://openjurist.org/713/f2d/335> (accessed on 21 July 2010).
69. Franken, J.; Stephan, U.; Meyer, H.E.; Konig, W. Identification of alpha-amylase inhibitor as a major allergen of wheat flour. *Int. Arch. Allergy Appl. Immunol.* **1994**, *104*, 171–174.
70. Moreno-Ancillo, A.; Dominguez-Noche, C.; Gil-Arados, A.C.; Cosmes, P.M. Bread eating induced oral angioedema due to a-amylase allergy. *J. Investig. Allergol. Clin. Immunol.* **2004**, *14*, 346–347.
71. Granum, P.E. Studies on α -amylase in foods. *Food Chem.* **1979**, *4*, 173–178.
72. Phadia, A.B. http://www.immunocapinvitrosight.com/ImmunoCAPDefault___23027.aspx (accessed on 14 September 2010).
73. Jones, J.M.J. *Food Safety*; Eagan Press: St. Paul, MN, USA, 1995; pp. 71, 77, 84, 87.
74. Shibamoto, T.; Bjeldanes, L.F. Natural toxins in plant foodstuffs. In *Introduction to Food Toxicology*; Academic Press: San Diego, CA, USA, 1993; pp. 78–79, 82–84.
75. Omaye, S.T. Toxicity of Nutrients. In *Food and Nutritional Toxicology*; CRC Press: Boca Raton, FL, USA, 2004; pp. 205–213.
76. Banwell, J.G.; Boldt, D.H.; Meyers, J.; Weber, F.L., Jr. Phytohemagglutinin derived from red kidney bean (*Phaseolus vulgaris*): A cause for intestinal malabsorption associated with bacterial overgrowth in the rat. *Gastroenterology* **1983**, *84*, 506–515.
77. Dobbins, J.W.; Laurensen, J.P.; Gorelick, F.S.; Banwell, J.G. Phytohemagglutinin from red kidney bean (*Phaseolus vulgaris*) inhibits sodium and chloride absorption in the rabbit ileum. *Gastroenterology* **1986**, *90*, 1907–1913.
78. United States Food and Drug Administration (FDA). Bad Bug Book. Foodborne Pathogenic Microorganisms and Natural Toxins Handbook. Phytohaemagglutinin, May 14, 2009. Available online: <http://www.fda.gov/food/foodsafety/foodborneillness/foodborneillnessfoodborne pathogensnaturaltoxins/badbugbook/ucm071092.htm> (accessed on 21 July 2010).
79. Buhler, R. Eating raw, undercooked beans can be unpleasant. High Plains/Midwest AG Journal. Available online: <http://www.hpj.com/archives/2004/nov04/nov15/Eatingrawundercooke ddrybean.cfm> (accessed on 21 July 2010).
80. Cornell University. Plants poisonous to livestock. Thiaminases. Available online: <http://www.ansci.cornell.edu/plants/toxicagents/thiaminase.html> (accessed on 21 July 2010).
81. Deshpande, S.S. Toxicants and antinutrients in plant foods. In *Handbook of Food Toxicology*; Marcel Dekker: New York, NY, USA, 2002b; pp. 331–372.
82. Prakash, A.S.; Pereira, T.N.; Reilly, P.E.B.; Seawright, A.A. Pyrrolizidine alkaloids in human diet. *Mutat. Res.* **1999**, *443*, 53–67.
83. Britton, N.L.; Brown, A. *Symphytum officinale* L. In *An Illustrated Flora of the Northern United States, Canada and the British Possessions*; Charles Scribner's Sons: New York, NY, USA, 1913; Volume 2, p. 92. USDA-NRCS PLANTS Database. Available online: http://plants.usda.gov/java/profile?symbol=SYOF&photoID=syof_001_avd.tif (accessed on 31 August 2010).

84. Dharmananda, S. Safety issues affecting herbs: Pyrrolizidine alkaloids, November, 2001. Available online: <http://www.itmonline.org/arts/pas.htm> (accessed on 21 July 2010).
85. Lowry, N. Rhubarb and Oxalic Acid. Available online: <http://helios.hampshire.edu/~nlNS/mompdfs/oxalicacid.pdf> (accessed on 21 July 2010).
86. Finkelstein, V.A.; Goldfarb, D.S. Strategies for preventing calcium oxalate stones. *Can. Med. Assoc. J.* **2006**, *174* (10), 1407–1409, DOI:10.1503/cmaj.051517.
87. Subbiah, V. Method of isolating cucurbitacin, July 20, 1999. Available online: <http://www.freepatentsonline.com/5925356.html> (accessed on 21 July 2010).
88. Martin, P.A.W.; Blackburn, M.; Schroder, R.F.W.; Matsuo, K.; Li, B.W. Stabilization of cucurbitacin E-glycoside, a feeding stimulant for diabroticite beetles, extracted from bitter Hawkesbury watermelon. *J. Insect Sci.* **2002**, *2*, 1–6.
89. Feather, S. Growing zucchini. Why your garden zucchinis might taste bitter. Available online: <http://www.donnan.com/Zucchini.htm> (accessed on 21 July 2010).
90. Browning, S.; Hodges, L. Bitterness in Zucchini Squash and Cucumber, February 19, 2010. Available online: <http://cuke.hort.ncsu.edu/cucurbit/cuke/cukehndbk/cukebitterness.html> (accessed on 21 July 2010).
91. Burfield, T. Coumarin: The real story, January, 2008. Available online: <http://www.leffingwell.com/Coumarin%20-%20the%20real%20story%20update2.pdf> (accessed on 21 July 2010).
92. Cornell University. Plants poisonous to livestock. Coumarin Glycosides. Available online: <http://www.ansci.cornell.edu/plants/toxicagents/coumarin.html> (accessed on 21 July 2010).
93. Lake, B.G. Coumarin metabolism, toxicity and carcinogenicity: Relevance for human risk assessment. *Food Chem. Toxicol.* **1999**, *37*, 423–453.
94. Fallon, S.; Enig, M.G. Cinderella's dark side. Available online: http://www.mercola.com/article/soy/avoid_soy.htm (accessed on 21 July 2010).
95. Kumar, V.; Sinha, A.K.; Makkar, H.P.S.; Becker, K. Dietary roles of phytate and phytase in human nutrition: A review. *Food Chem.* **2010**, *120*, 945–959.
96. Baruah, K.; Sahu, N.P.; Pal, A.K.; Debnath, D. Dietary phytase: An ideal approach for a cost effective and low-polluting aquafeed. *NAGA, WorldFish Center Quarterly* **2004**, *27* (3 & 4), 15–19.
97. Schecter, J.C.; Wiener, S.W. Plant Poisoning, Hypoglycemics, December 16, 2009. Available online: <http://emedicine.medscape.com/article/817325-overview> (accessed on 21 July 2010).
98. Lancashire, R.J. Jamaican Ackee, November 21, 2008. Available online: <http://wwwchem.uwimona.edu.jm/lectures/ackee.html> (accessed on 21 July 2010).
99. Sherratt, H.S.A. Hypoglycin, the famous toxin of the unripe Jamaican ackee fruit. *Trends Pharmacol. Sci.* **1986**, *7*, 186–191.
100. United States Food and Drug Administration (FDA). Haitian ackee fruit, January, 2010. Available online: <http://www.fda.gov/Food/NewsEvents/WhatsNewinFood/ucm197850.htm> (accessed on 21 July 2010).
101. Henry, S.H.; Page, S.W.; Bolger, P.M. Hazard assessment of ackee fruit (*Blighia sapida*). *Hum. Ecol. Risk Assess.* **1998**, *4*, 1175–1187.

102. Blake, O.A.; Jackson, J.C.; Jackson, M.A.; Gordon, C.L.A. Assessment of dietary exposure to the natural toxin hypoglycin in ackee (*Blighia sapida*) by Jamaican consumers. *Food Res. Int.* **2004**, *37*, 833–838.
103. Blake, O.A.; Bennink, M.R.; Jackson, J.C. Ackee (*Blighia sapida*) hypoglycin A toxicity: Dose response assessment in laboratory rats. *Food Chem. Toxicol.* **2006**, *44*, 207–213.
104. United States Food and Drug Administration (FDA). Detention without Physical Examination of Ackees. Import Alert 21–11, June 3, 2010. Available online: http://www.accessdata.fda.gov/cms_ia/importalert_64.html (accessed on 21 July 2010).
105. McGuffin, M. *American Herbal Product Association's Botanical Safety Handbook*; CRC Press: Boca Raton, FL, USA, 1997; pp. 149–152.
106. Homburger, F.; Boger, E. The carcinogenicity of essential oils, flavors and spices: A review. *Cancer Res.* **1968**, *28*, 2372–2374.
107. United States National Institute of Environmental Health Sciences (NIEHS). Substance Profiles: Safrole (CAS No. 94-59-7). Report on Carcinogens, Eleventh Edition; January 31, 2005. Available online: <http://ntp.niehs.nih.gov/ntp/roc/eleventh/profiles/s159safa.pdf> (accessed on 21 July 2010).
108. Wislocki, P.G.; Miller, E.C.; Miller, J.A.; McCoy, E.C.; Rosenkranz, H.S. Carcinogenic and mutagenic activities of safrole, 1'-hydroxysafrole, and some known or possible metabolites. *Cancer Res.* **1977**, *37*, 1883–1891.
109. Burfield, T. Safrole: Human carcinogenicity risk over-stated? September, 2009. Available online: <http://www.cropwatch.org/Safrole%20human%20carcinogenicity.pdf> (accessed on 21 July 2010).
110. Hallstrom, H.; Thuvander, A. Toxicological evaluation of myristicin. *Nat. Toxins* **1997**, *5*, 186–192.
111. Arneson, P.A.; Drubin, R.D. Studies on the mode of action of tomatine as a fungitoxic agent. *Plant Physiol.* **1968**, *43*, 683–686.
112. Rick, C.M.; Uhlig, J.W.; Jones, A.D. High alpha-tomatine content in ripe fruit of Andean *Lycopersicon esculentum* var. *cerasiforme*: developmental and genetic aspects. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 12877–12881.
113. Friedman, M.; Levin, C.E. α -Tomatine content in tomato and tomato products determined HPLC with pulsed amperometric detection. *J. Agric. Food Chem.* **1995**, *43*, 1507–1511.
114. Friedman, M.; Fitch, T.E.; Yokayama, W.E. Lowering of plasma LDL cholesterol in hamsters by the tomato glycoalkaloid tomatine. *Food Chem. Toxicol.* **2000**, *38*, 549–553.
115. Ize-Ludlow, D.; Ragone, S.; Bruck, I.S.; Bernstein, J.N.; Duchowny, M.; Pena, M.G. Neurotoxicities in infants seen with consumption of star anise tea. *Pediatrics* **2004**, *114*, e653–e656.
116. United States Food and Drug Administration (FDA). Inspections, Compliance, Enforcement and Criminal Investigations. Available online: <http://www.fda.gov/ICECI/EnforcementActions/EnforcementStory/EnforcementStoryArchive/ucm095929.htm> (accessed on 21 July 2010).
117. Tice, R. α -Chaconine [20562-03-2] and α -Solanine [20562-02-1]. Review of toxicological literature. Prepared for Errol Zeiger, National Institute of Environmental Health Sciences, February, 1998. Available online: http://ntp.niehs.nih.gov/ntp/htdocs/Chem_Background/ExSumPdf/ChaconineSolanine.pdf (accessed on 21 July 2010).

118. Surak, J.G. Phytoalexins and human health—a review. *FSHS Proc.* **1978**, *91*, 256–258.
119. United States Food and Drug Administration (FDA). FDA Poisonous Database, January 1, 2008. Available online: <http://www.accessdata.fda.gov/scripts/Plantox/Detail.CFM?ID=6537> (accessed on 21 July 2010).
120. Dinkins, C.L.P.; Peterson, R.K.D. A human dietary risk assessment associated with glycoalkaloid response of potato to Colorado potato beetle defoliation. *Food Chem. Toxicol.* **2008**, *46*, 2837–2840.
121. Ceska, O.; Chaudhary, S.K.; Warrington, P.J.; Ashwood-Smith, M.J. Naturally-occurring crystals of phototoxic furocoumarins on the surface of parsnip roots sold as food. *Experientia* **1986**, *42*, 1302–1304.
122. Ostertag, E.; Becker, T.; Ammon, J.; Bauer-Aymanns, H.; Schrenk, D. Effects of storage conditions on furocoumarin levels in intact, chopped, or homogenized parsnips. *J. Agric. Food Chem.* **2002**, *50*, 2565–2570.
123. Turesky, R.J. Heterocyclic Aromatic Amines (Part 2.3). In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 75–115.
124. United States National Institute of Environmental Health Sciences (NIEHS). Selected Heterocyclic Amines. Report on Carcinogens, Eleventh Edition; January 31, 2005. Available online: <http://ntp.niehs.nih.gov/ntp/roc/eleventh/profiles/s092vhca.pdf> (accessed on 21 July 2010).
125. Park, J.-H.; Penning, T.M. Polyaromatic Hydrocarbons (Part 2.8). In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 243–282.
126. Joint FAO/WHO Expert Committee on Food Additives (JECFA). Sixty-fourth meeting (64/SC). Section 2.6. Available online: http://www.who.int/ipcs/food/jecfa/summaries/summary_report_64_final.pdf (accessed on 21 July 2010).
127. Mills, C.; Mottram, D.S.; Wedzicha, B.L. Acrylamide (Part 2.1). In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 23–50.
128. Exon, J.H. A review of the toxicology of acrylamide. *J. Toxicol. Environ. Health* **2006**, *9*, 397–412.
129. Hamlet, C.G.; Sadd, P.A. Chloropropanols and Chloroesters (Part 2.6). In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 175–214.
130. Watkins, C. Chloroesters in foods: An emerging issue; April, 2009. Available online: <http://www.aocs.org/Membership/FreeCover.cfm?itemnumber=1084> (accessed on 21 July 2010).
131. Directorate-General Health and Consumer Protection. Reports on tasks for scientific cooperation. Collection and collation of data on levels of 3-monochloropropanediol (3-MCPD) and related substances in foodstuffs; June, 2004. Available online: http://ec.europa.eu/food/food/chemicalsafety/contaminants/scoop_3-2-9_final_report_chloropropanols_en.pdf (accessed on 21 July 2010).

132. Food Standards Australia New Zealand (FSANZ). *Chloropropanols in Food—an Analysis of Public Health Risk*; Technical Report Series No. 15; Food Standards Australia New Zealand: Canberra, Australia, 2003.
133. United States Food and Drug Administration (FDA). Sec. 500.500 Guidance levels for 3-MCPD (3-chloro-1,2-propanediol) in acid-hydrolyzed protein and asian-style sauces; March 2008. Available online: <http://www.fda.gov/ICECI/ComplianceManuals/CompliancePolicyGuidanceManual/ucm074419.htm> (accessed on 31 August 2010).
134. Carthew, P.; DiNovi, M.; Setzer, R.W. Application of the margin of exposure (MoE) approach to substances in food that are genotoxic and carcinogenic. Example: Furan (CAS No. 110-00-9). *Food Chem. Toxicol.* **2010**, *48*, S69–S74.
135. Bolger, P.M.; Tao, S.; Dinovi, M. Hazards of Dietary Furan. In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 117–133.
136. Moser, G.J.; Foley, J.; Burnett, M.; Goldsworthy, T.L.; Maronpot, R. Furan-induced dose-response relationships for liver cytotoxicity, cell proliferation, and tumorigenicity (furan-induced liver tumorigenicity). *Exp. Toxicol. Pathol.* **2009**, *61*, 101–111.
137. Cordelli, E.; Leopardi, P.; Villani, P.; Marcon, F.; Macri, C.; Caiola, S.; Siniscalchi, E.; Conti, L.; Eleuteri, P.; Malchiodi-Albedi, F.; Crebelli, R. Toxic and genotoxic effects of oral administration of furan in mouse liver. *Mutagenesis* **2010**, *25*, 305–314.
138. Leopardi, P.; Cordelli, E.; Villani, P.; Cremona, T.P.; Conti, L.; DeLuca, G.; Crebelli, R. Assessment of *in vivo* genotoxicity of the rodent carcinogen furan: Evaluation of DNA damage and induction of micronuclei in mouse splenocytes. *Mutagenesis* **2010**, *25*, 57–62.
139. Sadler, M.J. Health effects of trans fatty acids. In *Encyclopedia of Human Nutrition*; Sadler, M.J., Strain, J.J., Caballero, B., Eds.; Academic: San Diego, CA, USA, 1999; Volume 2, pp. 769–776.
140. Ascherio, A.; Katan, M.B.; Zock, P.L.; Stampfer, M.J.; Willett, W.C. Trans fatty acids and coronary heart disease. *N. Eng. J. Med.* **1999**, *340*, 1994–1998.
141. Baxter, S.D. Nutrition for Healthy Children and Adolescents Aged 2 to 18 Years. In *Handbook of Nutrition and Food*, 2nd ed.; Berdanier, C.D., Dwyer, J., Feldman, E.B., Eds.; CRC Press: Boca Raton, FL, USA, 2008; p. 295.
142. United States Food and Drug Administration (FDA). Federal Register—68 FR 41433 July 11, 2003: Food Labeling; Trans Fatty Acids in Nutrition Labeling; Consumer Research to Consider Nutrient Content and Health Claims and Possible Footnote or Disclosure Statements; Final Rule and Proposed Rule. Available online: <http://www.fda.gov/food/labelingnutrition/labelclaims/nutrientcontentclaims/ucm110179.htm> (accessed on 21 July 2010).
143. United States Department of Agriculture (USDA). Dietary Guidelines for Americans 2005. Available online: <http://www.cnpp.usda.gov/publications/dietaryguidelines/2005/2005DGpolicydocument.pdf> (accessed on 21 July 2010).
144. Motarjemi, Y.; Stadler, R.H.; Studer, A.; Damiano, V. Application of the HACCP Approach for the Management of Processing Contaminants. In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; p. 573.

145. United States National Institute of Environmental Health Sciences (NIEHS). *N*-Nitrosodimethylamine. Report on Carcinogens, Eleventh Edition; January 31, 2005. Available online: <http://ntp.niehs.nih.gov/ntp/roc/eleventh/profiles/s128nitr.pdf> (accessed on 21 July 2010).
146. Habermeyer, M.; Eisenbrand, G. *N*-Nitrosamines, including *N*-Nitrosoaminoacids and potential further nonvolatiles (Part 4.1). In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 365–386.
147. United States National Institute of Environmental Health Sciences (NIEHS). *N*-Nitrosopyrrolidine. Report on Carcinogens, Eleventh Edition; January 31, 2005. Available online: <http://ntp.niehs.nih.gov/ntp/roc/eleventh/profiles/s137nsop.pdf> (accessed on 21 July 2010).
148. United States National Institute of Environmental Health Sciences (NIEHS). *N*-Nitrosopiperidine. Report on Carcinogens, Eleventh Edition; January 31, 2005. Available online: <http://ntp.niehs.nih.gov/ntp/roc/eleventh/profiles/s136nsop.pdf> (accessed on 21 July 2010).
149. Jakszyn, P.; Gonzalez, C.A. Nitrosamine and related food intake and gastric and oesophageal cancer risk: A systematic review of the epidemiological evidence. *World J. Gastroenterol.* **2006**, *12*, 4296–4303.
150. Sarkadi, L.S. Biogenic Amines (Part 3.2). In *Process-Induced Food Toxicants. Occurrence, Formation, Mitigation, and Health Risks*; Stadler, R.H., Lineback, D.R., Eds.; John Wiley & Sons: Hoboken, NJ, USA, 2009; pp. 321–361.
151. United States Food and Drug Administration (FDA). Fish and Fisheries Products Hazards and Controls Guidance, Third Edition. Chapter 7: Scombrototoxin (Histamine) Formation (A Chemical Hazard), June 2001. Available online: <http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Seafood/FishandFisheriesProductsHazardsandControlsGuide/ucm091910.htm> (accessed on 31 August 2010).
152. Van Dolah, F.M. Marine algal toxins: Origins, health effects, and their increased occurrence. *Environ. Health Perspect.* **2000**, *108*, 133–141.
153. Woods Hole Oceanographic Institution (WHOI). Human illness associated with harmful algae. Available online: <http://www.whoi.edu/science/B/redtide/illness/illness.html> (accessed on 21 July 2010).
154. United States Food and Drug Administration (FDA). Fish and Fisheries Products Hazards and Controls Guidance, Third Edition. Chapter 6: Natural Toxins (A Chemical Hazard). Available online: <http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Seafood/ucm091782.htm> (accessed on 29 August 2010).
155. Beauchamp, R.A.; Wiles, K.; Hendricks, K. Red Tide Information; May 20, 2008. Available online: <http://www.dshs.state.tx.us/seafood/redtide.shtm> (accessed on 21 July 2010).
156. University System of Maryland (USM). Harmful algal blooms. Available online: <http://aquaticpath.umd.edu/toxalg/nsp.html> (accessed on 21 July 2010).
157. United States Department of Commerce National Oceanic and Atmospheric Administration (NOAA). Microscopic image of Pseudo-nitzschia. Available online: http://www.noaanews.noaa.gov/stories2009/20091116_razor.html (accessed on 31 August 2010).

158. United States Department of Commerce National Oceanic and Atmospheric Administration (NOAA). Scientists report first remote, underwater detection of harmful algae, toxins; June 14, 2009. Available online: <http://www.physorg.com/news166807443.html> (accessed on 21 July 2010).
159. Kleivdal, H.; Kristiansen, S.; Nilsen, M.V. Single-laboratory validation of the Biosense Direct Competitive Enzyme-Linked Immunosorbent Assay (ELISA) for determination of domoic acid toxins in shellfish. *J. AOAC Int.* **2007**, *90*, 1000–1010.
160. Dickey, R.W.; Plakas, S.M. Ciguatera: A public health perspective. *Toxicon* **2009**, DOI:10.1016/j.toxicon.2009.09.008.
161. United States Food and Drug Administration (FDA). Bad Bug Book. Foodborne Pathogenic Microorganisms and Natural Toxins Handbook. BBB-Gemphylotoxin; May 20, 2010. Available online: <http://www.fda.gov/Food/FoodSafety/FoodborneIllness/FoodborneIllnessFoodbornePathogensNaturalToxins/BadBugBook/ucm071191.htm> (accessed on 21 July 2010).
162. Ukishima, Y.; Masui, T.; Masubara, S.; Goto, R.; Okada, S.; Tsuji, K.; Kosuge, T. Wax components of escolar (*Lepidocybium flavobrunneum*) and its application to base of medicine and cosmetics. *Yakugaku Zasshi* **1987**, *107*, 883–890.
163. Nicholas, P.D.; Mooney, B.D.; Elliott, N.G. Unusually high levels of non-saponifiable lipids in the fishes escolar and rudderfish identification by gas and thin-layer chromatography. *J. Chromatogr. A* **2001**, *936*, 183–191.
164. Berman, P.; Harley, E.H.; Spark, A.A. Keriorrhoea—the passage of oil per rectum—after ingestion of marine wax esters. *S. Afr. Med. J.* **1981**, *59*, 791–792.
165. SEFSC Pascagoula Laboratory; Collection of Brandi Noble. Photograph of Juvenile Oilfish (*Ruvettus pretiosus*), NOAA/NMFS/SEFSC. NOAA Photo Library. Available online: <http://www.photolib.noaa.gov/htmls/fish4425.htm> (accessed on 15 September 2010).
166. Reid, T.M.S.; Gould, I.M.; Mackie, I.M.; Ritchie, A.H.; Hobbs, G. Food poisoning due to the consumption of red whelks (*Neptunea antiqua*). *Epidemiol. Infect.* **1988**, *101*, 419–424.
167. Kim, J.H.; Lee, K.J.; Suzuki, T.; Kim, C.M.; Lee, J.Y.; Mok, J.S.; Lee, T.S. Identification of tetramine, a toxin in whelks, as the cause of a poisoning incident in Korea and the distribution of tetramine in fresh and boiled whelk (*Neptunea intersculpta*). *J. Food Prot.* **2009**, *72*, 1935–1940.
168. Power, A.J.; Keegan, B.G.; Nolan, K. The seasonality and role of the neurotoxin tetramine in the salivary glands of the red whelk *Neptunea antiqua* (L.). *Toxicon* **2002**, *40*, 419–425.
169. Anthoni, U.; Bohlin, L.; Larsen, C.; Nielsen, P.; Nielsen, N.H. The toxin tetramine from “edible” whelk *Neptunea antiqua*. *Toxicon* **1989**, *27*, 717–723.
170. Anthoni, U.; Christophersen, C.; Gram, L.; Nielsen, N.H.; Nielsen, P. Poisonings from flesh of the Greenland shark *Somniosus microcephalus* may be due to trimethylamine. *Toxicon* **1991**, *29*, 1205–1212.
171. Benz, G.W.; Hocking, R.; Kowunna, Sr.A.; Bullard, S.A.; George, J.C. A second species of Arctic shark: Pacific sleeper shark *Somniosus pacificus* from Point hope Alaska. *Polar Biol.* **2004**, *27*, 250–252.
172. Idboro, C.J. The pangnirtung inuit and the greenland shark. Masters Thesis. University of Manitoba, Canada, November, 2008. Available online: <http://www.umanitoba.ca/institutes/>

- natural_resources/canadaresearchchair/thesis/Idrobo.Masters%20Thesis.Feb%2009.pdf (accessed on 21 July 2010).
173. United States Food and Drug Administration (FDA). Bad Bug Book. Foodborne Pathogenic Microorganisms and Natural Toxins Handbook. BBB-Grayanotoxin. Available online: <http://www.fda.gov/Food/FoodSafety/FoodborneIllness/FoodbornIllnessFoodbornePathogensNaturalToxins/BadBugBook/ucm071128.htm> (accessed on 29 August 2010).
 174. Laborde, A. *Nerium oleader* L. Poisons Information Monograph 366. International Programme on Chemical Safety (INCHEM); November, 1989. Available online: <http://www.inchem.org/documents/pims/plant/pim366.htm> (accessed on 21 July 2010).
 175. Panter, K.E.; James, L.F. Natural plant toxicants in milk: A review. *J. Anim. Sci.* **1990**, *68*, 892–904.
 176. Lee, S.T.; Davis, T.Z.; Gardner, D.R.; Stegelmeier, B.L.; Evans, T.J. Quantitative method for the measurement of three benzofuran ketones in rayless goldenrod (*Isocoma pluriflora*) and white snakeroot (*Ageratina altissima*) by high-performance liquid chromatography (HPLC). *J. Agric. Food Chem.* **2009**, *57*, 5639–5643.
 177. National Park Service (NPS). Lincoln Boyhood National Memorial. The plant that killed Nancy Hanks Lincoln. Available online: http://www.nps.gov/archive/libo/white_snakeroot3.htm (accessed on 21 July 2010).

© 2010 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

From: acvimcardio <acvimcardio@vin.com>

To: (b) (6)

Subject: dilated cardiomyopathy and kangaroo and lentil diets (b) (6)

Date: Wed, Jan 17, 2018 10:51 am

ACVIM/ECVIM - Cardiologists ListServe



Email preferences

Password reminder

Board: ACVIM/ECVIM - Cardiologists
Folder: Myocardial Disease
Discussion: dilated cardiomyopathy and kangaroo and lentil diets

[Read the Message Board Discussion](#)

Reply By Email

Reply on Message Boards

Post New Thread By Email

New Messages (2)

12. Posted by (b) (6) on 01-16-18 07:50 ET

201C

Original Message

plasma taurine changes very quickly with diet so may not reflect very accurately blood taurine status

201D

This is definitely true in cats. We documented this in the late 80's. Fasting 24 hours can bring plasma taurine down below normal range, but not where we generally see in truly deficient cats. Still hard to interpret if fasted.

You also need to worry about false elevation from damaged WBCs and platelets that have high taurine concentrations.

201C

Original Message

snap frozen whole blood is better though a pain to collect and transport.

201D

As long as it's sterile, whole blood taurine should be pretty stable even if not frozen. We didn't note big degradation impacts of not freezing. Freezing and transporting likely is best still.

201C

Original Message

Equally blood taurine levels may not reflect cardiac muscle taurine

201D

These correlate pretty well in a steady state. There is a lag during depletion meaning that if you feed cats (I don't have data in dogs), a taurine depleting diet, the half-life of taurine in plasma is very short, whole blood relatively long and of skeletal muscle is longer. Cardiac muscle likely parallels skeletal, but we couldn't get as many data points since we could not sample cardiac muscle (by endomyocardial biopsy) as frequently as we sampled skeletal muscle.

>> (b) (6) <<

(b) (6)

13. Posted by (b) (6) on 01-17-18 10:45 ET

Hi All - I have been working with a few other cardiologists to collect a group of golden retrievers with taurine-deficiency and DCM. We have a total of 24 unrelated golden retrievers and the vast majority of these were eating diets that are labeled as grain free and use a large amount of peas or lentils in the ingredients.

Our nutritionists at UCD have been involved and analyzed some of the diets but certainly not all. We have reported the findings to the FDA. We are working on the (b) (4)

(b) (4). We are also interested in the trend that many of the designer diets include higher quality meats with relative little organ meat or byproducts. Perhaps this switch is also important as there is a fair amount of taurine to be found in the lower quality organ meats / byproducts that these designer diets try to avoid.

We have seen a fair number of other breeds as well but have not included them in our data gathering. A few of our cases were in CHF and have resolved with supplementation. Almost all have improved on supplementation. We have initial screening data and 3-4month follow-up on each of these cases now and I have a student working on writing this up. Many dogs had pretty low whole blood taurine levels, but a few had "low normal" values yet still responded favorable to supplementation.

Would love to chat with anyone interested.

(b) (6)

Reply By Email

Reply on Message Boards

Post New Thread By Email

Complete Discussions (13)

1. Created by (b) (6) on 01-10-18 02:15 ET

Hi,

I am currently seeing my fourth patient in a year with dilated cardiomyopathy that eats a

FDA-CVM-FOIA-2019-1704-000597

kangaroo and lentil diet. The first two were related Labrador retrievers (aunt and niece) in the same household. The second was a Cocker spaniel. Today's patient is a mixed breed 55 pound brindle lab mix. The first Labrador had a normal whole blood taurine, and we did not test the second housemate since she presented at a later date and we knew that her housemate fed the same diet was not taurine deficient. The Cocker Spaniel was taurine deficient based on a low whole blood taurine. The first Labrador is a year out from her initial heart rate episode, and still alive. We have changed diet, but not until her housemate was also found to have DCM (this patient was euthanized when she had a quick recurrence of CHF), just a few months ago, and an echo recently did not show significant change from baseline. We did look up carnitine levels, and supposedly kangaroo diets have high carnitine levels. I had just tacked the odd diet up to coincidence suspecting that the Labs may have had a heritable form of DCM, and the Cocker taurine-deficient DCM. Now, though, with number four, I thought I would extend the question to the group to see if anyone else has had any cases of DCM in dogs on a kangaroo diet.

Thanks for any input!

(b) (6)

2. Posted by (b) (6) on 01-10-18 06:03 ET

Hi (b) (6)

(b) (4)

Hopefully we will have more information soon.
Take care

(b) (6)

(b) (6), DVM, DACVIM (Cardiology)

3. Posted by (b) (6) on 01-11-18 11:38 ET

Hi,
When I was a resident I saw two American cockers with DCM that were being fed a kangaroo and lentil diet - they were taurine deficient and responded to diet change and supplementation. Both dogs belonged to the same owner but were unrelated. We consulted the nutrition team at UC Davis to see whether they thought there might be a problem with the diet - however, they thought it was more likely to be breed related, which seemed reasonable at the time.

I'm really intrigued by your cases though - were they all eating the same brand of food, or were they kangaroo and lentil diets from a variety of manufacturers?

Let me know if more information would help,

(b) (6)

4. Posted by (b) (6) on 01-12-18 09:54 ET

I have seen 3 dogs from one household on a lentil based diet with DCM. Mixed breeds. One died. Other 2 recovered with diet change and taurine supplementation.

Blood taurine levels were extremely low in all dogs. Owner was into vegan diets but fed nothing but lentils and a supplement for more than a year.

(b) (6)

5. Posted by (b) (6) on 01-12-18 11:32 ET

What good timing!

I too have had several atypical DCM breeds over the past few months, but the most recent one was this week - a 2 kg 5yo Pomeranian who is on Zignature Kangaroo diet and has been eating this for the past several years. I was highly suspicious for taurine deficiency based off the unusual breed, but it's interesting to know that others are seeing this as well.

(b) (6)

6. Posted by (b) (6) on 01-12-18 01:39 ET

We have also seen a few dogs recently. They were a 2 yo Boston eating kangaroo and lentil with a normal blood taurine, an 8 month old husky mix eating a grain free diet, and an 8 month old GSD mix also eating grain free (no blood taurines on either of the latter two).

(b) (6)

7. Posted by (b) (6) on 01-15-18 11:14 ET

Hi all,

I have three Cocker Spaniels, same family, all Taurine-Responsive systolic dysfunction receiving Zignature's Kangaroo & Lentil diet. One was in severe left sided congestive heart failure, hospitalized for 4 days with severely decreased systolic function and severe mitral regurgitation. Three months later, (on Taurine, pimobendan, furosemide and ACE inhibitor), his systolic function had resolved and his mitral regurgitation was significantly decreased (to trivial). Nine months out he has normal systolic function and persistently trivial mitral regurgitation (the previously noted severe MR is attributed to mitral valvular annular dilation and not primary valvular regurgitation). The other two have normal systolic function with Taurine supplementation, observed at six months following Taurine supplementation.

(b) (6)

(b) (6).com [mailto:(b) (6)]

(b) (6)

8. Posted by (b) (6) on 01-15-18 04:23 ET

Good afternoon,

I have had two dogs with DCM that were eating lentil and rice based diets. One was a mixed breed shepherd type dog that was eating a vegan diet (owner was vegan). The dog was not taurine deficient or responsive and died about six months after diagnosis. The second, a pit bull, developed after eating a lentil/rice diet for 12 weeks as part of a food allergy trial. The following three months the dog ate a few different proteins but went back to lentils/rice in between. This dog is low taurine and we are hoping will respond to supplementation. I am wondering if there is something inherently negative about the lentils and amino acid

processing...kind of a scary situation.

Best,

(b) (6)

9. Posted by (b) (6) on 01-15-18 10:36 ET

This compilation of reports related to a few food types is impressive, especially those demonstrating resolution after taurine supplementation. Someone needs to compile these and publish them.

Has anyone contacted the manufacturers of the foods? Response from them?

Has anyone analyzed the taurine content of the foods? I'm happy to arrange and pay for that if samples of the foods these dogs were eating are available.

>>(b) (6)<<

(b) (6)

10. Posted by (b) (6) on 01-16-18 03:11 ET

Hi All

Thankfully Kangaroo and lentils do not seem to have arrived en masse in the UK yet - I would have thought parasitic disease from kangaroos may also be a potential issue depending on how it has been prepared and stored as there is a reasonable prevalence of Toxoplasma - so just a thought on whether this could be related to cardiac disease.

Not sure how people are measuring taurine but plasma taurine changes very quickly with diet so may not reflect very accurately blood taurine status - snap frozen whole blood is better though a pain to collect and transport. Equally blood taurine levels may not reflect cardiac muscle taurine

(b) (6)

11. Posted by (b) (6) on 01-16-18 04:56 ET

Paul has forwarded an excellent idea. My understanding is that kangaroo has low levels of taurine and methionine and lentils are low in sulfur containing amino acids methionine and cysteine. The metabolism of these amino acids are complexly intertwined.

The dogs I saw were on lentils and begged oh supplement possibly before more taurine was added to the supplement. Certainly, one would not expect kangaroo meat to provide sufficient taurine to eliminate the risk of deficiency. Sounds like a good opportunity for a nutrition-oriented cardiologist or group to sort this out.

Until that time, consider supplementation of taurine and also possibly methionine.

(b) (6)

12. Posted by (b) (6) on 01-16-18 07:50 ET

201C

Original Message

plasma taurine changes very quickly with diet so may not reflect very accurately blood taurine status

201D

This is definitely true in cats. We documented this in the late 80's. Fasting 24 hours can bring plasma taurine down below normal range, but not where we generally see in truly deficient cats. Still hard to interpret if fasted.

You also need to worry about false elevation from damaged WBCs and platelets that have high taurine concentrations.

201C

Original Message

snap frozen whole blood is better though a pain to collect and transport.

201D

As long as it's sterile, whole blood taurine should be pretty stable even if not frozen. We didn't note big degradation impacts of not freezing. Freezing and transporting likely is best still.

201C

Original Message

Equally blood taurine levels may not reflect cardiac muscle taurine

201D

These correlate pretty well in a steady state. There is a lag during depletion meaning that if you feed cats (I don't have data in dogs), a taurine depleting diet, the half-life of taurine in plasma is very short, whole blood relatively long and of skeletal muscle is longer. Cardiac muscle likely parallels skeletal, but we couldn't get as many data points since we could not sample cardiac muscle (by endomyocardial biopsy) as frequently as we sampled skeletal muscle.

>>(b) (6)<<

(b) (6)

13. Posted by (b) (6) on 01-17-18 10:45 ET

Hi All - I have been working with a few other cardiologists to collect a group of golden retrievers with taurine-deficiency and DCM. We have a total of 24 unrelated golden retrievers and the vast majority of these were eating diets that are labeled as grain free and use a large amount of peas or lentils in the ingredients.

Our nutritionists at UCD have been involved and analyzed some of the diets but certainly not all. We have reported the findings to the FDA. (b) (4)

(b) (4)

We are also interested in the trend that many of the designer diets include higher quality meats with relative little organ meat or byproducts. Perhaps this switch is also important as there is a fair amount of taurine to be found in the lower quality organ meats / byproducts that these designer diets try to avoid.

We have seen a fair number of other breeds as well but have not included them in our data gathering. A few of our cases were in CHF and have resolved with supplementation. Almost all have improved on supplementation. We have initial screening data and 3-4month follow-up on each of these cases now and I have a student working on writing this up. Many dogs had pretty low whole blood taurine levels, but a few had "low normal" values yet still responded favorable to supplementation.

Would love to chat with anyone interested.

(b) (6)

Reply By Email

Reply on Message Boards

Post New Thread By Email

Email preferences

Password reminder

You are subscribed to the ACVIM/ECVIM Cardiology ListServe. This discussion is also posted on the "ACVIM/ECVIM Cardiology message board" hosted by VIN. All information posted is confidential and restricted to ListServe members. **Participation in the ListServe DOES NOT require VIN membership. VIN will never use or distribute any information about you or your participation in the ListServe in any way.**

To start new discussions, Select "Post New Thread By Email" or go to:
<http://WWW.vin.com/> (b) (6)

Please do not redistribute any of the discussions, images or other media posted on these Message Boards without prior permission from your colleagues. If you wish to use any of this information, please contact the appropriate colleague(s) directly or the list moderators at acvimcardio@vin.com and put "Permission to Distribute Information" in the Subject line.

To be removed from the ListServe entirely or to stop receiving email notifications but maintain access to the associated message board system, please email (b) (6)@vin.com and put REMOVE in the subject line.

800.700.4636 | VINGRAM@vin.com | 530.756.4881 | Fax: 530.756.6035

777 West Covell Blvd, Davis, CA 95616

Copyright 2018, Veterinary Information Network, Inc.

Unsubscribe from Cardiology ACVIM / ECVIM ListServe

Sample Submission Form

Amino Acid Laboratory
 University of California, Davis
 1020 Vet Med 3B
 1089 Veterinary Medicine Drive
 Davis, CA 95616
 Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
 Non-federal funds ID/Account Number
 to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/aal/aal.html>

Vet/Tech Contact: Account # _____ (b) (6) / Contact: (b) (6) Date: 9-5-17

Company Name: _____ (b) (6)

Address: _____ (b) (6)

Email: _____ (b) (6)

Tel: _____ (b) (6) Fax: _____ (b) (6)

Billing Contact: _____ (b) (6) TAX ID: _____

Email: _____ (b) (6) Tel: (b) (6)

Patient Name: _____ (b) (6)

Species: _____

Owner's Name: _____ (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
 Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 10 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

(b) (6)

(b) (6)

Patient Information

Patient: (b) (6) Age: 6 years Referring Veterinarian: (b) (6)
Patient Number (b) (6) Weight:(kg) 25.10 Cardiologist: (b) (6)
Breed: lab mix Sex: F Client Number: (b) (6)
Exam Date: (b) (6) 08:21 BSA: 0.87

History: (b) (6) was presented for evaluation of cough, labored breathing, multiple episodes of collapse, cardiomegaly, and suspected congestive heart failure. (b) (6) was initially seen by her rDVM on December 30th for cough. She was initially treated with a humidifier, hydrocodone, and a decrease in length of walks and did not improve. On January 4th she had a collapse episode during which she circled, fell over, and flopped for about a minute. She has a second less severe episode on January 5th. Labwork at this date showed mild elevation of ALT and AST with mild increase in CK. T4 was low normal. She has continued to be short of breath and tires easily. She chased a squirrel a couple of days ago and stood with a wide based stance afterward. She is on year-round heartworm, flea and tick preventative and was last tested negative for heartworm 9/9/17. She eats California Natural Kangaroo and Red Lentil dry food with vegetables and 2 tbsp of canned pumpkin. She is on Apoquel 8mg twice daily and sublingual immunotherapy. The hydrocodone was discontinued yesterday.

Physical Examination: T 102.7 P 208 R 150. Grade 3/6 left apical systolic murmur and gallop. Regular tachycardia. Quiet heart sounds. Localized fine crackles left cranial hilar region, dry cough. Poor femoral pulses. Unremarkable abdominal palpation. mm pale pink, normal refill. Hydration OK. Normal PLNs.

Diagnostic Tests:

(b) (6)
BP - 152mmHg 4 cm cuff LF
Echo - see below. Sinus tachycardia on ECG.
Taurine level (whole blood): pending, will call with results.
ECG - HR 189bpm, sinus tachycardia. Wide and tall P waves (0.06s, 0.5mV) consistent with atrial enlargement, tall R waves (3.7mV) consistent with LVE, normal PR (0.12s) and QT (0.2s) intervals.

(b) (6)
Thoracic radiographs: Mild decrease in severity of cardiomegaly (as compared to rDVM films from 1/9/18). Resolving cardiogenic edema.

Hospitalization:

An IV catheter was placed and (b) (6) was hospitalized in ICU with continuous ECG monitoring. She was started on IV Lasix (50 mg IV q8h) and pimobendan (5 mg in AM and 10 mg in PM). She did well overnight with an improvement in respiratory rate/effort. (b) (6) had occasional short paroxysms of "slow" ventricular tachycardia (160-270 bpm) that were noted noted to persist beyond ~7 pm.

(b) (6) was started on enalapril (10 mg q12h) the following morning. She continued to do well with a normal appetite and improved respiratory rate.

Echocardiographic Report

2D ECHO

LA Systolic Diameter LX 6.5 cm

DOPPLER

AV Peak Velocity	70.8 cm/s	PV Peak Gradient	1.3 mmHg
AV Peak Gradient	2 mmHg	TR Peak Velocity	255 cm/s
MR Peak Velocity	396 cm/s	TR Peak Gradient	26.1 mmHg
PV Peak Velocity	56.2 cm/s		

M-MODE

LV Diastolic Diameter MM	7.4 cm	LVPW Diastolic Thickness MM	0.92 cm
LV Systolic Diameter MM	6.5 cm	LVPW Systolic Thickness MM	0.96 cm
LV Fractional Shortening MM	12 %	LVPW Percent Thickening MM	0.048
LV Diastolic Volume Cube	398 cm ³	IVS to PW Ratio MM	1
LV Systolic Volume Cube	271 cm ³	LV Mass MM	324 g
LV Ejection Fraction Cube	0.32	LV Mass Normalized MM	374 g/m ²
IVS Diastolic Thickness MM	0.96 cm	RV Diastolic Diameter MM	0.86 cm
IVS Systolic Thickness MM	0.89 cm	LA Systolic Diameter MM	4.8 cm
IVS Percent Thickening MM	0.069	Aortic Root Diameter MM	2.1 cm

Left Ventricle: Severe dilation (normalized LVIDd 2.85) with severe myocardial dysfunction (normalized LVIDs 2.34). Increased sphericity.

Left Atrium: Severe dilation.

Right Ventricle: Mild dilation with subjective decrease in contractility.

Right Atrium: Mild dilation.

Mitral Valve: Normal valve morphology. 4+ central mitral regurgitation.

Aortic Valve: Normal.

Tricuspid Valve: Mildly thickened valve leaflets. 1+ tricuspid regurgitation. Normal regurgitant velocities.

Pulmonic Valve: Mildly thickened valve leaflets. Mild pulmonic insufficiency.

Aorta: Normal

Pericardium: Normal

Diagnosis

Dilated cardiomyopathy - This is a disease characterized by weakening of the heart muscle and dilation of the heart chambers. As the disease progresses, it can lead to congestive heart failure (fluid in the lungs causing shortness of breath and cough). Abnormal heart rhythms are common and can result in sudden death. Most commonly this is an inherited disease, though it can occur secondary to a deficiency in an amino acid called taurine.

Left sided congestive heart failure

Recommendations

Give all medications as directed:

Furosemide (Lasix, Salix) 40 mg tablets- Give 1 1/2 tablets by mouth every 12 hours. DUE: This evening.

This is a diuretic (water pill), that prevents the body from retaining excessive sodium and water. It will cause your pet to drink and urinate more frequently. It is important that fresh water is always available.

Benazepril 10 mg tablets- Give 1 tablet by mouth every 12 hours. DUE: ~9:00 PM

This medication is a strong drug that dilates blood vessels, permitting the heart to pump blood more efficiently. It can lower blood pressure (hypotension) and cause changes in kidney function and electrolyte values. If your pet develops weakness or depression, decrease the drug dose by 1/2 and call. A kidney panel and blood pressure should be reevaluated 7-10 days after beginning this medication.

Spirolactone (Aldactone) 50 mg tablets- Give 1 tablet by mouth every 24 hours. START: Tonight or tomorrow morning.

This is a diuretic (water pill) that also blocks a hormone that can injure the heart muscle. It works well in combination with the furosemide and enalapril.

Pimobendan (Vetmedin) 10 mg tablets- Give 1/2 of a tablet by mouth in the morning and 1 tablet in the evening. Give at 12 hour intervals. DUE: ~6:00 PM (1 tablet)

This is a drug that is approved for the treatment of congestive heart failure secondary to dilated cardiomyopathy or chronic valve disease (endocardiosis). Studies have shown improved quality of life and increase survival time when this drug is added to other standard cardiac medications. In our experience, side effects are uncommon, but it is important that you advise us if you feel your pet is having any potential adverse effects from this medication. The reported potential side effects listed for this medication are increased heart rate, vomiting, diarrhea, inappetence, uneasiness, incoordination, convulsions, increased drinking and increase urinating.

One thing that can be very helpful for home monitoring is checking sleeping or resting respiratory rates. A recent study showed that even pets with severe heart disease rarely have resting respiratory rates greater than 30 breaths per minute unless they are starting to decompensate for that disease. Elevated respiratory rates at home may be even more sensitive than chest radiographs at picking up early decompensation. Count your pet's respiratory rate when he/she is at rest or sleeping (not within 20 minutes of being active). If his/her respiratory rate is greater than 30 breaths per minute, recheck again in a couple of hours. If persistently elevated above this level, call.

With advanced heart disease, our biggest dietary concerns are adequate calorie content and low sodium content. We aim for less than 80mg sodium per 100 kilocalories (kcal) in patients that have developed congestive heart failure. We do not advise protein restriction unless there is concurrent kidney disease (i.e. kidney diets are not advised unless there is concurrent kidney disease). Please refer to our diet handouts with a list of currently adequate diets and treats, though this list is not exclusive. If you wish to feed a diet that is not on these lists, you will need to call the manufacturer of the diet to obtain a sodium content.

As we discussed, we have had three other cases of severe DCM where the dogs have been eating a kangaroo and lentil diet. There is no data that has shown an association with this diet and DCM but we are concerned there may be a connection there and are looking into it at this time. For this reason, we would consider changing (b) (6) diet.

We sent (b) (6) home with a few cans of Hill's Science Diet Canine Maintenance canned food. This food has an appropriate level of sodium for dogs in congestive heart failure and is available at most pet stores. Lamb should be avoided as a protein source but any other protein is appropriate (with the exception of kangaroo).

The very best diet for dogs with DCM/heart failure is probably Hill's Science Diet Prescription j/d. This food has a good source of taurine, carnitine and fatty acids. However, this diet is rather costly.

We have submitted a taurine level and will call you with the results when they are available.

Exercise is also a concern in advanced heart disease. While cage rest is ideal with active heart failure, some exercise is permissible in asymptomatic disease. However, vigorous or extended exercise should be avoided.

***As long as (b) (6) does well at home we would like to re-evaluate her in 7-10 days. At this time we will recheck her kidney values/electrolytes and blood pressure as well as repeat chest x-rays.

(b) (6)

(b) (6) 08:21

(b) (6)

(Electronically Signed)

Final Date:

Like us on Facebook!

(b) (6)

Notes to our clients

-Please bring all medications to your pet's scheduled appointments.

-We require a 48 hour notice for all refills. When you call to request a refill, please leave the pharmacy phone number or clearly indicate if you plan on picking up the medication at our facility. PRESCRIPTION REFILLS ARE NOT AVAILABLE AFTER (b) (6) (b) (6) REGULAR BUSINESS HOURS (Evenings, Fridays, holidays and weekends).

-Check out WWW.GOODRX.COM and enter your local zip code to search for the best prices on your medications at your local pharmacies.

-If an emergency arises with your pet, (b) (6) is a 24 hour facility.

From: [Conway, Charlotte](#)
To: [Edwards, David](#)
Subject: FW: DCM investigation - Pet Food Institute
Date: Monday, June 24, 2019 12:38:31 PM

FYI

From: Ask CVM <AskCVM@fda.hhs.gov>
Date: June 24, 2019 at 10:20:06 AM CDT
To: Forfa, Tracey <Tracey.Forfa@fda.hhs.gov>, Norris, Anne <Anne.Norris@fda.hhs.gov>, Steinberg, Nadine <Nadine.Steinberg@fda.hhs.gov>, DeLancey, Siobhan <Siobhan.Delancey@fda.hhs.gov>, Hartogensis, Martine <Martine.Hartogensis@fda.hhs.gov>, Conway, Charlotte <Charlotte.Conway@fda.hhs.gov>
Subject: FW: DCM investigation - Pet Food Institute

From: Susan Thixton <susan@truthaboutpetfood.com>
Sent: Monday, June 24, 2019 9:37 AM
To: Ask CVM <AskCVM@fda.hhs.gov>
Subject: DCM investigation - Pet Food Institute

We were alerted that the FDA has provided the Pet Food Institute details of a soon to be released update of the Agency's investigation of pet food related DCM. It is more than concerning that FDA shared details of their investigation with industry well in advance of sharing those details with pet owners.

Pet owners – not industry – are the ones suffering the most from this nutritional failure of pet food. Pet owners are reporting to us that veterinarians are telling clients *“they are literally seeing thousands of new cases (of DCM) everyday”* and veterinarians are pushing pet owners to provide their pets ONLY a grain-based pet food made by Purina, Mars, or Hill's. Needless to say, the focus of this situation is completely out of control.

The focus should be on the actual problem – a nutritional failure of 'Complete and Balanced' pet foods. We ask the FDA to keep their focus on the Complete and Balanced nutritional failure and to update pet owners in the same timely manner as they do industry.

Susan Thixton

--

Susan Thixton
Pet Food Consumer Advocate
TruthaboutPetFood.com
AssociationforTruthinPetFood.com

Compendium

Cardiovascular Effects of Thyroid Disease

Jodi K. Sangster, DVM

David L. Panciera, DVM, MS, DACVIM (Small Animal Internal Medicine)

Jonathan A. Abbott, DVM, DACVIM (Cardiology)

Virginia Tech

Blacksburg, Virginia


Abstract: *Thyroid hormones have many effects on cardiovascular function, and deficiency or excess of thyroid hormones can result in cardiac dysfunction. Abnormalities of the cardiovascular system are often identified during examination of hyperthyroid and hypothyroid patients. This article addresses the effects of thyroid hormones on the cardiovascular system and the clinical relevance of the cardiovascular response to thyroid dysfunction. In addition, treatment recommendations are presented.*

(b) (4)

(b) (4)




(b) (6)





(b) (6)






(b) (6)




(b) (6)





(b) (6)





(b) (6)



(b) (6)



(b) (6)





(b) (6)

Vet-LIRN Case Summary Document

Vet-LIRN Case Number:	800.218
EON/CC #:	EON-323515-323519
Vet-LIRN Initiation Date:	7/13/2017
MedRec: Requested:	7/12/2017
MedRec: Received:	7/13/2017
MedRec: Significant finding:	Cardiogenic heart failure clinically but less supported histologically
Vet-LIRN Tests (planned):	(b) (4), (b) (5)
Vet-LIRN Test Results:	TAMU: <ul style="list-style-type: none"> Fumonisin-Negative (b) (4) Taurine, Carnitine-<i>pending</i>
Result Interpretation:	
IF NFA, justification:	

COMPLAINT Narrative:

(b) (6) housemate, (b) (6) (separate report submitted) was diagnosed with DCM and CHF 2/17 and was euthanized after aggressive treatment of CHF. At that time (b) (6) had 2 syncopal events closely related to each other. His appetite for dog food declined but he would eat it if tempted with treats mixed in. He was presented (b) (6) for more syncopal events and was similarly diagnosed with severe DCM and CHF. He was able to be successfully treated however and is clinically doing well on CHF medications as of 7/10/17. A re-review of the myocardial histopathology for (b) (6) housemate (b) (6) was requested at this time because of the unusual diagnosis of DCM in a small breed dog living in the same house as another dog similarly diagnosed a few months ago. This re-review by one of our pathologists showed myofiber vacuoles reminiscent of the changes seen in doxorubicin toxicity. Since the dog had not received doxorubicin, the pathologist recommended culturing the food for *Streptomyces peucetius* - the bacterium which produces doxorubicin. He also recommended testing for *Fusarium* spp. a fungus which produces Fumonisin B1, a toxin that produces heart failure in pigs. Like (b) (6) (unrelated, younger miniature schnauzer), (b) (6) had been fed California Naturals Adult - both kangaroo with lentils and venison with lentils along with Milo's kitchen treats. We have samples of these foods from 6/17 but not the original bags from when he was presented 2/17. These samples were provided at the time (b) (6) also presented with severe DCM and CHF. Like (b) (6), (b) (6) had extensive infectious disease testing which was negative and nutritional amino acid deficiencies were ruled out. Because of this, their unrelated lineages (although the same breed, they were from different lines), different ages but similar time of presentation, we are considering common environmental factors which could precipitate DCM, including food contamination or toxin exposure. We have plasma, serum, urine and myocardial tissue samples (latter only for (b) (6)) stored at -80 Celsius in addition to food and treat samples.

3 week history of cough treated unsuccessfully with doxycycline and prednisone. 3 day history of inappetence and vomiting prior to presentation to (b) (6) emergency service for dyspnea. Radiographs showed severe pulmonary edema and echocardiogram showed severe Dilated Cardiomyopathy. There was an initial response to diuretic therapy however, he declined and was placed on the ventilator for respiratory support and continued CHF treatment. Attempts to wean off the ventilator were unsuccessful and aquaphoresis was performed. He continued to decline despite aggressive therapy and was euthanized. Infectious disease testing was negative and taurine and carnitine analysis showed adequate levels. Necropsy initially did not reveal a cause for DCM and supported alveolar injury (possibly ventilator related). A rereview of the myocardial histopathology by one of our pathologist showed myofiber vacuoles reminiscent of the changes seen in doxorubicin toxicity. Since the dog had not received doxorubicin, the pathologist recommended culturing the food for Streptomyces peucetius - the bacterium which produces doxorubicin. He also recommended testing for Fusarium spp. a fungus which produces Fumonisin B1, a toxin that produces heart failure in pigs. (b) (6) had been fed California Naturals Adult - both kangaroo with lentils and venison with lentils along with Milo's kitchen treats. We have samples of these foods from 6/17 but not the original bags from when he was presented 2/17. These samples were provided at the time his housemate, (b) (6) (unrelated, older miniature schnauzer) also presented with severe DCM and CHF. I will enter this dog as a separate affected patient. Both dogs had extensive infectious disease testing which was negative and nutritional amino acid deficiencies were ruled out. Because of this, their unrelated lineages (although the same breed, they were from different lines), different ages but similar time of presentation ((b) (6) had clinical signs at the time (b) (6) was treated, but didn't present with CHF for several months), we are considering common environmental factors which could precipitate DCM, including food contamination or toxin exposure.

Signalment:

- (b) (6): 7 yr MC Miniature Schnauzer
- (b) (6): 2 yr MC Miniature Schnauzer-deceased

Signs: syncopal episodes, dyspnea, cough, heart failure

Food: Alternated feedings between:

- California Natural Adult Limited Ingredient Grain Free Venison & Green Lentils Recipe Dog Food
- "" Kangaroo and Lentils

Vet-LIRN PLAN OF ACTION:

- (b) (5)
- (b) (5)

Medical Record Review:

(b) (6):

Presenting complaint (b) (6) : dyspnea, cough of 3 week duration-wheezing type more frequent at night → rDVM, treated w/ prednisone and doxycycline for kennel cough → (b) (6) inappetance, vomiting → (b) (6) dyspneic and recheck, hospitalized and treated for pneumonia, regurgitated → (b) (6) treated as outpatient, (b) (6) as syring feeding, dog regurgitated and had marked dyspnea → ER → refer to NCSU → (b) (6) put on mechanical ventilator → (b) (6) euthanized

PE (b) (6): P 160 bpm, R 64 rpm, pale pink mm, Gr I-II/VI left apical systolic murmur, femoral pulse hypokinetic but synchronous, jugular venous distention

Labwork:

(b) (6)

Labs: unremarkable (unclear what was done)

(b) (6)

Big 4: Glu 135, Azo 15-20

- (b) (6) vBGA: Lact 2.4, rest wnl
- Chem: P 6.2, K 4.9, Na 140, TP 4.2
- Chem: BG 225, BUN 29, P 11.7, K 3.3, Cl 95, Na 144
- Chem: BG 136, P 4.6, CK 13,621, K 4.3, Na 151, Cl 109, AST 577
- Chem: BG 165, BUN 37, P 8.1, ALT 147, AST 1006, CK 35,930, Na 135, K 3.8, Cl 90

(b) (6)

CBC: WBC 9.4, NP 7.9, Band .18, Plt 157

- (b) (6): WBC 9.9, NP 8, Band .7
- WBC 6.8, TP 6.9, NP 4.2, Ban .54, Toxic NP-mild, Plt 97

(b) (6)

BP sys: 90

(b) (6)

UA post Lasix: 1.011**Cardiac troponin** 0.79**BAP GM**-pending**Vector borne panel:** pending**Taurine/Carnitine:** pending

(b) (6)

Coag: PT 9.1, PTT 14, Dimer 189, Fib 539, INR 1.09**Urine Creat:** 27.9**Urine Na:** pending**ECG:** suspected atrial tachycardia**Rads** (b) (6): concern for aspiration pneumonia

- (b) (6): cardiomegaly, severe diffuse mixed interstitial to alveolar pattern most severe caudo-dorsally, hepatomegaly, dec abdominal serosal contrast
- (b) (6): severe generalized cardiomegaly with biventricular heart failure, improved vs rDVM rads
- (b) (6): worsening cardiogenic pulmonary edema, cannot exclude lung induced injury +/- pneumonia
- (b) (6): post ultrafiltration, improved cardiogenic edema, hypovolemia, residual interstitial to patchy alveolar
- (b) (6): improved CHF with possible concern for bronchopneumonia, suspected hiatal hernia
- (b) (6): markedly progressive alveolar pattern with significantly worse cardiogenic edema

tFAST (b) (6): severe cardiomegaly with ventricular hypocontractility**Echo** (b) (6): dcm vs. myocarditis vs pacing induce vs. other (severely dilated & hypocontractile left & right ventricles, severely dilated left and right atria)**Necropsy:** Lung-severe diffuse alveolar injury with marked fibrin deposition (hyaline) and marked alveolar histiocytosis and multifocal type II pneumocyte hyperplasia; mod to marked diffuse pulmonary edema; mild cardiomegaly with mild mitral valve endocardiosis and mild left ventricular hypertrophy and left atrial dilation; thorax with mild pleural effusion; Suspect primary non-cardiogenic etiology but if clinical cardiac dysfunction then functional cardiac abnormalities cannot be ruled out**Prior MHx:** coffee brown urine including clumping after strenuous activity when it is hot outside and resolves with 24-36 hours; also Crystalluria

(b) (6)

Presented (b) (6): episodes of collapse, first occurred mid February, fall 6 seconds without losing consciousness → immediately return to normal → 2 weeks later again collapse, then on → 6/3 post 2 hour hike collapsed again; panting more than usual; good appetite for treats but reluctant to eat food since February; → recheck (b) (6), doing better, no collapsing episodes except a stumbling moment when excited, respiratory rate normal, diet changed to Hill's

(b) (6) **PE:** P 130 bpm, R pant, mild increased breath sounds in all lung fields

(b) (6): T 99.7F, P 136 bpm, R 36 rpm, equivocal mild dehydration <5%, Gr II/VI left apical systolic murmur

Labs: (b) (6) **Big 4:** BG 64 (recheck 79), BUN 15-26

BP-sys: 130 mmHg

-7/10: 110 mmHg

ECG: left ventricular enlargement suggested

UA: 1.019

Taurine & Carnitine: normal (no values)

Vector borne panel (PCR and IFA): normal

BAP GM

Troponin 1

T4

Toxoplasma/Neospora

Chagas

Complete AA: no significant abnormalities, consulting with UC Davis

-7/10 **Renal Panel:** K 4.0

(b) (6) **Rads:** left sided congestive heart failure

-7/10: moderate left sided cardiomegaly without heart failure, moderate hepatomegaly

(b) (6) **Echo:** mitral valve endocardiosis with left atrial enlargement and heart failure, decreased left ventricular systolic function, suspected DCM

Thoughts: possible fumonisin induced cardiomyopathy, vs. other non-food related toxic compound (lily of valley, digitalis, ionophores, sicklepod, gossypol, white snake root), ethyl alcohol, foxglove, buttercups

7/18/2017

JJ-DR agrees fumonisin worth testing and suggests asking about fish/sardines/anchovies being fed to dogs in February (re: domoic acid toxicity which can appear doxorubicin-like).

Will ask vet re: fish/sardines/anchovies and also how much bag of food weighs. Emailed TAMU requesting fumonisin testing.

7/19/2017

JJ-TAMU can do the testing at (b) (6). Vet sent product weight and dimensions. Vet confirms, neither dog received anchovies, sardines, or seafood in February or Chronically.

7/22/2017

JJ-Vet did not receive a return shipping label. Will make PO for shipping to Vet-LIRN.

7/28/2017

JJ-Vet sent info-she's had 2 other cases of DCM with dogs eating the California Naturals diet. Info below:

I don't know if there is anything to this but I have treated 2 other dogs in the last 2 weeks with DCM and CHF that are being fed California natural food (one kangaroo and lentil, and we are trying to find out about the other one which is in our ER now). One is mixed breed and we'll recommend testing for taurine deficiency. The other was a golden and taurine was a bit low but not super low so although we supplemented, I wasn't totally convinced it was the cause. Unfortunately she died a week later.

I don't have a sense for how widely fed this diet is but I don't see it on the top selling lists I can find by google, so seeing 4 DCM dogs recently eating this particular food is interesting to say the least. I thought this might be of interest to you as you start to look at the California natural food sample from the Ramsammy dogs.

Vet also sent journal article about Acrolein chronic oral ingestion causing idiopathic DCM in rats.

Found (b) (6) can test for Acrolein in pet food.

Sent info to group, and DR thinks testing for taurine, carnitine, and acrolein (if not too expensive) worth it. I can send an email to (b) (6) to request testing cost/method info/shipping info/amount needed for testing.

8/3/2017

JJ-Prepared pet food on 8/2 and lab submission forms. Make boxes today and send for testing.

JG – Shipped samples to (b) (4)

8/4/2017

JJ-(b) (6) responded. Acrolein testing would require method development and be quite costly. Will not pursue at this time.

8/7/2017

JJ-(b) (4) sent final results. Fumonisin negative. Will send all product test results to vet when received.

8/9/2017

JJ-DR suggested-maybe vet at (b) (6) has funds to do acrolein.

8/22/2017

JJ-Received (b) (4) results for Taurine and Carnitine. Filed. Vet asked about results- Send Fumonisin results and taurine/carnitine in progress until we interpret. Plan to send those after interpretation finalized. Asked vet which dry food we were sent.

Flavor	Moisture Max (label)	Ingredient lists:	Label claims	Nutrient Analysis (web)
Kangaroo	10%			(b) (5)

1. Taurine 231 mg/100g food → 0.23% As Is basis → assume max 10% moisture (label) for either food:

Kangaroo → 0.26% (on DMB calculated using label moisture max)

(b) (5)

Interpretation: No AAFCO minimum for Taurine in dogs. (b) (5)

(b) (5) - If the food is either flavor, it is compliant for minimums of cat taurine levels.

2. L-Carnitine 69,900 ppb = 69 ppm = 0.0069% As Is basis → assume max 10% moisture (label):

Kangaroo → 0.0077% (on DMB calculated using label moisture max)

(b) (5)

Interpretation: No carnitine minimum for dogs or cats. Unclear whether or not this is low, normal, or high.

Final Conclusion: The cause of the two dogs' DCM is unclear. The bloodwork for these dogs showed normal taurine and carnitine levels. Based on the dogs' blood taurine/carnitine levels and the dry dog food test results, it is unlikely that Fumonisin, taurine, or carnitine caused the dogs' illness.

Vet responded-flavor is Kangaroo. Let her know acrolein not available as a test unless she wants to pursue method development with (b) (6) .

DR agrees-sent final interpretation to vet. NFA.

10/16/2017

JJ-From: Garland T. Overview of Gossypol Poisoning. Merck Veterinary Manual. Found at:

<http://www.merckvetmanual.com/toxicology/gossypol-poisoning/overview-of-gossypol-poisoning>

“Differential diagnoses (for gossypol) include poisonings by cardiotoxic ionophoric antibiotics (eg, monensin, lasalocid, salinomycin, narasin) and ammonia, nutritional or metabolic disorders (eg, selenium, vitamin E, or copper deficiency), infectious diseases, noninfectious diseases (eg, pulmonary adenomatosis, emphysema), mycotoxicoses caused by Fusarium-contaminated grain, and toxicoses caused by plants with cardiotoxic and other effects. Cardiotoxic plants (see Poisonous Plants), which may cause confusing or similar clinical signs and postmortem lesions, include English yew (*Taxus baccata*), Japanese yew (*T. cuspidata*), laurel (*Kalmia* spp), azalea (*Rhododendron* spp), oleander (*Nerium oleander*), yellow oleander or yellow-be-still tree (*Thevetia peruviana*), purple foxglove (*Digitalis purpurea*), lily-of-the-valley (*Convallaria majalis*), dogbane (*Apocynum* spp), coffee senna (*Senna occidentalis*), bracken fern (*Pteridium aquilinum*), white snakeroot (*Eupatorium rugosum*), death camas (*Zygadenus* spp), lantana (*Lantana camara*), monkshood (*Aconitum napellum*), and milkweed (*Asclepias* spp).”

14768

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/index.cfm>

Vet/Tech Contact: (b) (6)
Company Name: (b) (6)
Address: (b) (6)

Email: (b) (6)
Tel: (b) (6) Fax: (b) (6)

Billing Contact: Finance Dept. TAX ID: (b) (6)
Email: (b) (6) Tel: ext

Patient Name: (b) (6)
Species: Feline
Owner's Name: (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 196 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
6/6/2016	C	(b) (6)	RECEPTION ACTIONS NOTE Sympathy card sent-AG
6/6/2016	C	(b) (6)	MEDICAL COMMENTS 6/6/2016 11:15 FDA complaint submitted: Pet Food Safety Report, ID 53897, was successfully submitted on 6/6/2016 11:15:17 AM EST to the FDA, and it was issued an Individual Case Safety Report Number (ICSR) of 1053335.
(b) (6)	C	(b) (6)	MEDICAL COMMENTS (b) (6) 16:26 (b) (6) ReplyReply AllForwardActions To: (b) (6) Sent Items(b) (6) Hi (b) (6) , Sorry for the delay in getting back with you, I needed to get permission from the owner's before providing you with their contact information. Below is the their information as well as the names of the individual cats. The cat, (b) (6) , with dilated cardiomyopathy was euthanized yesterday. Owners: (b) (6) and (b) (6) Cats: -(b) (6) : (b) (6) Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016, euthanized on (b) (6) 5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016 -(b) (6) : 9yr male neutered domestic long hair: 196 nmol/ml -(b) (6) : 8y female spayed domestic short hair: 368 nmol/ml -(b) (6) : 9yr male neutered domestic long hair: 124 nmol/ml -(b) (6) : 9yr male neutered domestic long hair: 536 nmol/ml Please let me know if you have any other questions.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Sincerely,

(b) (6)
(b) (6)

(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)

(b) (6) @merrickpetcare.com]Reply All

(b) (6) 12:56 PM

Thank You for providing me this information (b) (6) . Could you provide us the pet parents information as well. We would like to reach out to the pet parent as well and speak with her. Thanks.

(b) (6) C

(b) (6)

COMMUNICATIONS WITH CLIENT

(b) (6) 15:55

SWO - expressed my condolences. asked for permission to provide contact info to company and the FDA - owner consented. Discussed what to expect when talking to company. Owner thankful for call.

(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

TO: (b) (6)
(b) (6)
(b) (6)
(b) (6)

FAX #: (b) (6)
FROM: (b) (6)
DATE: (b) (6)

RE:
Client: (b) (6)
Patient: (b) (6) / ID: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 5 Mos. **Sex:** Spayed Female

~ EUTHANASIA NOTIFICATION ~

Dear (b) (6) :

This letter is to inform you that your patient, (b) (6) was visited by (b) (6) At Home house call service today for end-of-life care.

If you have any questions, please feel free to contact me at the location noted above.

Thank you,

(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)	C	(b) (6)	<p>MEDICAL COMMENTS - Closed Jun 04/2016 (b) (6) 18:03 Seen on emergency today, embolic event secondary to DCM. Discussed necropsy and advised that nutritionist and cardiologist agreed that prior test results were sufficient; necropsy would not reveal anything not already documented. Owner had already admin 0.2ml buprenex sublingually, requested I admin the remaining 0.3ml dispensed today which I did. They then spent time privately with the patient prior to euthanasia.</p> <p>Flushed cephalic catheter in right front leg; patent. Admin 20mg (2ml) expired propofol IV, apneic and unresponsive Admin 975mg (2.5ml) beuthanasia IV, 3 exhalation spasms followed Confirmed deceased by prolonged thoracic auscultation Removed IVC, placed (b) (6) in coffin, nested in owner's blanket</p>
---------	---	---------	--

(b) (6)	D	(b) (6)	Pleural Effusion Final
(b) (6)	D	(b) (6)	Feline Arterial Thromboembolic Disease Final

(b) (6)	R	(b) (6)	Referral Letter - Cardio Resident Eval and labs - FINAL (b) (6) - REF fxd
---------	---	---------	---

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

TO: (b) (6)
(b) (6)
(b) (6)
(b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: (b) (6)
RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 5 Mos. **Sex:** Spayed Female
Current Weight: 5.3 kilograms as of 5/25/2016

Thank you for referring (b) (6). The following is a case summary.

Date of evaluation: (b) (6)

Date of previous cardiac evaluation: Wednesday, May 25, 2016

CHIEF COMPLAINT: heavy breathing, dragging RH limb

HISTORY: (b) (6) previously doing well. Appetite had improved, eating 2/3 can max cal per day. Normal breathing. Owner noted acute onset of dragging RH limb this morning and heavy breathing. No interest in food this morning. Brought in to ER immediately. Received 9mg lasix IV total and 0.075mg buprenorphine IV on presentation. Previous hx: Diagnosed with DCM (b) (6). Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4. Taurine level confirmed taurine deficiency

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was tachypnic with moderate increased effort, RR 48. Normal BV sounds, no crackles on auscultation. Unable to bear weight on RH limb, dragging. Femoral pulses were fair in left hind, absent in left hind limb. Right paw pads cold to the touch. Heart rate was 160 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals (b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, age cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, entative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Cursory Ultrasound: small volume pleural effusion. No pericardial effusion. Large thrombus in LV.

Brief Echo 5/25/16: moderate volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle unchanged from previous.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM (b) (6)/2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), taurine deficiency, moderate left atrial enlargement, pleural effusion, hx azotemia, LV thrombus, FATE (partial)

THORACOCENTESIS 4/10/16: a total of 180ml of yellow fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed scant effusion remained in the patient.

THORACOCENTESIS (b) (6)/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: (b) (6) clinical signs are due to a partial aortic thromboembolus and recurrent left sided congestive heart failure. Thrombus previously noted in LV is still present. There is a very guarded prognosis of recovery with ATE. Discussed with owners, if they wants to go forward with treatment recommend hospitalization to manage CHF and pain. Discussed risk of reperfusion injury, main concern hyperkalemia which can be fatal. At home care will consist primarily of pain medication and nursing care. She may or may not gain function back to hind limb, and this could take up to 1-2 months to see improvement if any. Owner elected to take home to spend a few hours to spend time and have (b) (6) come for euthanasia this evening. IVC left in place (right cephalic). Sent home with additional dose of buprenorphine to keep her comfortable until euthanasia. Owners understand she may pass on her own. Gave owners my condolences.

MEDICATIONS:

Buprenorphine 0.3mg/ml- give entire contents of syringe (0.5ml) at 3pm today for pain control

Thank you for the courtesy of this interesting referral. Please feel free to contact me

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

with any questions or comments.

Sincerely,

(b) (6)

(b) (6)

Sent electronically - no signature required

(b) (6)

(b) (6)

Client ID: (b) (6) Patient ID: (b) (6) Patient Name: (b) (6)

Pain Condition Scores

Have you noticed the "PCS" aka Pain Condition Score in our medical records? As part of our AAHA specialty accreditation, we evaluate the level of pain or discomfort for each and every patient we see. You will see this reflected in the physical exam recordings as a "PCS". We follow the guidelines created and published by Colorado State University. If you have questions about interpreting pain or treatment options, or you need to schedule a pain management consult, please speak with our Anesthesiology & Pain Management Department by calling (b) (6).

Clinical Studies

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

We are recruiting for a variety of clinical studies across multiple specialties and are looking for eligible patients. For a complete list, please visit our website at [\[redacted\]](#) and click on Veterinary Professionals and then Current Clinical Studies.

Avian & Exotics

(b) (6) is happy to re-introduce our Avian & Exotics Department and welcome (b) (6) to our team! We offer advanced diagnostics and treatment methods to complement the care provided by referring veterinarians. In addition to referral services, we provide primary care and medical boarding. Our facility maintains a dedicated exotics ward and a 24-hour monitored critical care unit. The Avian & Exotics Department accepts appointments 5 days a week (Tuesday-Saturday) and our emergency service receives patients, including exotic species, 24 hours a day. You can contact (b) (6) at (b) (6)

(b) (6)	C	(b) (6)	EMERGENCY PHYSICAL EXAM - Closed Jun 04/2016 (b) (6)
---------	---	---------	---

Chief Complaint: Respiratory distress

History: (b) (6) presented for STAT evaluation of respiratory distress. Owner noticed progressive tachypnea this AM and difficulty using right hindlimb. She did not want to eat this AM so she did not receive her AM medication. She is currently under the care of our Cardiology Service for Dilated Cardiomyopathy (DCM) (suspect secondary to taurine deficiency), moderate left atrial enlargement, pleural effusion, azotemia, and LV thrombus.

Other Medical Problems: None

Medications/Supplements: Pimobendan, Lasix, taurine supplementation, appetite stimulant

Environment: indoors only, several other cats

Vaccination Status: UTD

Current Diet (Type): Tempting to eat
- Frequency:

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

- Amount:

Physical Examination:

S(ubjective): BAR/distressed, hydration WNL, BCS 7/9, pain score: 1/4

O(bjective):

Weight: 5.3 kilograms

TPR: T: 94.8 HR: 188, RR/RE: 60/rapid/shallow

EENT: clear AU/OU, no nasal discharge, normal cervical palpation, mm pink, moist/CRT<2 sec

INTEG: Hair coat ok

PLN: WNL

CV: NSR, no murmur ausculted, left femoral pulse moderate/synchronous, right very difficult to feel to absent

RESP: tachypneic, sl. dull ventrally, no crackles/wheezes

GI: soft, nonpainful, no masses

UG: FS, NSF

M/S: laterally recumbent, a

Neuro: alert/appropriate, cranial nerves intact, no placing deficits or spinal/neck pain

Problems/Differential Diagnoses:

Respiratory distress, decreased motor/absent femoral pulse RHL

Diagnostics:

None performed

Assessment:

12yo FS DSH

- absent to faint femoral pulse RHL, decreased motor, hypothermia, hx: DCM with LV thrombus- r/o saddle thrombus vs. other

- respiratory distress, mild amount pleural effusion on TFAST- r/o secondary to CHF secondary to DCM (suspect taurine deficiency)

- hx: Dilated Cardiomyopathy (DCM) (suspect secondary to taurine deficiency), moderate left atrial enlargement, pleural effusion, azotemia, and LV thrombus.

Treatment:

Placed in oxygen. IVC placed. 4mg Lasix IV, followed by additional 5mg IV. 0.015mg/kg Buprenorphine IV. Improved rr/re with above.

Plan/Recommendations:

Discussed PE at length with owner. Concerned for partial vs. full saddle thrombus RHL secondary to LV thrombus we know she has. Discussed options- point to

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

severity of underlying disease- ATH, repeat Echo, supportive care, vs. euthanasia. Owner elected to continue supportive care until they could speak with (b) (6), considering euthanasia. Elected RED code, transferred to cardiology.

(b) (6)	P	(b) (6)	0.50 mg of Buprenex (Buprenorphine) 0.3mg/mL MG C3 (MOBHL0) Rx #: 2579780 0 Of 0 Refills Give the entire contents of the syringe (0.5ml) under the tongue at 3pm.
---------	---	---------	---

(b) (6)	C	(b) (6)	CARDIAC EVALUTION - CLOSED 06/04/2016 - Cardiac Evaluation
---------	---	---------	--

Date of evaluation: (b) (6)

Date of previous cardiac evaluation: Wednesday, May 25, 2016

CHIEF COMPLAINT: heavy breathing, dragging RH limb

HISTORY: (b) (6) previously doing well. Appetite had improved, eating 2/3 can max cal per day. Normal breathing. Owner noted acute onset of dragging RH limb this morning and heavy breathing. No interest in food this morning. Brought in to ER immediately. Received 9mg lasix IV total and 0.075mg buprenohpine IV on presentation.

Previous hx: Diagnosed with DCM (b) (6)/16. Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4. Taurine level confirmed taurine deficiency

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was tachypnic with moderate increased effort, RR 48. Normal BV sounds, no crackles on auscultation. Unable to bear weight on RH limb, dragging. Femoral pulses were fair in left hind, absent in left hind limb. Right paw pads cold to the touch. Heart rate was 160 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) (b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Cursory Ultrasound: small volume pleural effusion. No pericardial effusion. Large thrombus in LV.

Brief Echo 5/25/16: moderate volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle unchanged from previous.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM (b) (6)/2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), taurine deficiency, moderate left atrial enlargement, pleural effusion, hx azotemia, LV thrombus, FATE (partial)

THORACOCENTESIS 4/10/16: a total of 180ml of yellow fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed scant effusion remained in the patient.

THORACOCENTESIS (b) (6)/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

ER thoracocentesis (b) (6) 16: 25ml yellow tinged fluid from the right side.

SUMMARY: (b) (6) clinical signs are due to a partial aortic thromboembolus and recurrent left sided congestive heart failure. Thrombus previously noted in LV is still present. There is a very guarded prognosis of recovery with ATE. Discussed with owners, if they want to go forward with treatment recommend hospitalization to manage CHF and pain. Discussed risk of reperfusion injury, main concern hyperkalemia which can be fatal. At home care will consist primarily of pain medication and nursing care. She may or may not gain function back to hind limb, and this could take up to 1-2 months to see improvement if any. Owner elected to take home to spend a few hours to spend time and have (b) (6) come for euthanasia this evening. IVC left in place (right cephalic). Sent home with additional dose of buprenorphine to keep her comfortable until euthanasia. Owners understand she may pass on her own. Gave owners my condolences.

MEDICATIONS:

Buprenorphine 0.3mg/ml- give entire contents of syringe (0.5ml) at 3pm today for pain control

(b) (6) CK (b) (6) STAT
Reason for Visit: Emergency
Date Patient Checked Out: (b) (6) Practice TF

(b) (6) TC (b) (6) MEDICAL COMMENTS - TENTATIVE
(b) (6) 10:40
SW (b) (6) at Merrick - updated company that we have documented taurine deficiency in 2 other cats in house hold. The quality assurance team indicated that the level of taurine in the lot # I gave them was sufficient - discussed that this likely takes 3-6 month to develop and likely to be related to earlier lot and they need to

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

investigate further. Asked if the level they gave me was from them retesting the food after I called or from an earlier test - not sure. Asked for information from the taurine tests - told I can send summary of lab results. Also indicated that I am reporting this to the FDA. She give new info to her manager and quality assurance. Told them I expect them to follow up with me. Below email sent to Merrick:

Taurine Levels

(b) (6)
To:
(b) (6) @merrickpetcare.com
Hi (b) (6) ,

Thank you for your help with these cases. Here is the summary of the lab results:

12yr female spayed domestic short hair diagnosed and clinical for dilated cardiomyopathy

(b) (6)/2016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016

5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016

-9yr male neutered domestic long hair: 196 nmol/ml
-8y female spayed domestic short hair: 368 nmol/ml
-9yr male neutered domestic long hair: 124 nmol/ml
-9yr male neutered domestic long hair: 536 nmol/ml

Please let me know if you have any other questions.

Sincerely,

(b) (6)

(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)
(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE</p> <p>(b) (6) 08:42</p> <p>O Imom- was previously doing well. Eating ~2/3 can of max cal per day, sRR 6-7breaths/15sec. Now, this morning, dragging RH limb and breathing heavier.</p> <p>SWO- recommended to come in as soon as possible since (b) (6) breathing heavy. Unfortunately, cardiology will be in surgery this morning. Should go through emergency and I will consult.</p>
---------	----	---------	---

(b) (6)	B	(b) (6)	.08 mg of Buprenex (Buprenorphine) 0.3mg/mL MG C3 (MOBHL0) by (b) (6)
	B		1.00 Specialty/Referral Exam Level 3 (REF03) by (b) (6)
	B		Echo Guided Thoracocentesis Group (EGT) by (b) (6)
	B		1.00 EGT Procedure (USSC50) by (b) (6)
	B		1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
	B		1.00 Cared for by (b) (6) (b) (6) by (b) (6)
	B		.50 mg of Buprenex (Buprenorphine) 0.3mg/mL MG C3 (MOBHL0) by (b) (6)
	B		At Home Euthanasia Group (HCEUTH) by (b) (6)
	B		1.00 At Home Euthanasia Service (HC08) by (b) (6)
	B		1.00 At Home Burial (HC10) by (b) (6)
	B		100.00 mg of Telazol 100mg/mL inj per mg (C3-N) 2103 (MLTZL1) by (b) (6)
	B		10.00 mg of Acepromazine 10mg/mL Inj per mg (MLA2L1) by (b) (6)
	B		1.00 mg of Butorphanol 10mg/mL Inj per mg (C4) (MOB2L10) by (b) (6)
	B		100.00 mg of Beuthanasia Soln 390 mg/mL (C-3N) / MG (MOB2L1) by (b) (6)
	B		1.00 IV Catheter Placement (CATH) by (b) (6)
	B		1.00 each of Tx Catheter IV 22g x 1" Surflo (BLUE) (H113) by (b) (6)
	B		1.00 each of Tx IV Ext T Set Hospira 1265028 (H027) by (b) (6)
	B		At Home Euthanasia Group (HCEUTH) by (b) (6)
	B		1.00 At Home Euthanasia Service (HC08) by (b) (6)
	B		1.00 At Home Burial (HC10) by (b) (6)
	B		-100.00 mg of Telazol 100mg/mL inj per mg (C3-N) 2103 (MLTZL1) by (b) (6)
	B		-10.00 mg of Acepromazine 10mg/mL Inj per mg (MLA2L1) by (b) (6)
	B		-1.00 mg of Butorphanol 10mg/mL Inj per mg (C4) (MOB2L10) by (b) (6)
	B		875.00 mg of Beuthanasia Soln 390 mg/mL (C-3N) / MG (MOB2L1) by (b) (6)
	B		1.00 Cared for by (b) (6) (b) (6) by (b) (6)
	B		1.00 each of Tx Injection Cap/Plug Termo 007110 (H118) by (b) (6)
	B		Oxygen Therapy (Caged)/Hour (Group) (O2CAGE) by (b) (6)
	B		1.00 O2 Therapy Per Hour (T044) by (b) (6)
	B		1.00 Oxygen-related Patient Care / Hour (O2CARE) by (b) (6)
	B		1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
	B		4.00 mg of Furosemide (Lasix) 5% Injection per mg (MMF2L5) by (b) (6)
	B		5.00 mg of Furosemide (Lasix) 5% Injection per mg (MMF2L5) by (b) (6)

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6) (b) (6) 5/29/2016	B B P	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6) 1.00 Emergency Exam Level 4 (EE04) by (b) (6) 7.00 can of lams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) Rx #: 2578487 0 Of 6 Refills Feed up to 1 can daily.
5/29/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/29/2016 13:52 SWO- (b) (6) doing ok. sRR7breaths/15 sec. Ate ~3/4 can of the lams max cal last night. Had normal BM yesterday. Hind limbs are very weak, one is worse than the other, but able to take a few steps on it before needing a rest. Does not see painful or distressed. Rec continue lasix 1/4 tab SID for now until appetite is consistent, then may consider increasasing. Continue pimo and taurine. Will put refill through for max cal. Emailed Client: I put through a prescription for 7 cans of food for (b) (6) . She needs just under 1 can per day (although if she eats a whole can per day that is fine). There are also refills on the prescription if you need more. I will call to check in on her in a few days. Please call me with any concerns.
5/29/2016	B	(b) (6)	7.00 can of lams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) by (b) (6)
5/28/2016	C	(b) (6)	PHARMACY NOTE Returned O call, left voice message that medication is ready for pick up
5/28/2016	P	(b) (6)	21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472) Rx #: 2569385 1 Of 12 Refills Filled by: (b) (6) Give 1 tablet by mouth twice daily with food.
5/28/2016	B	(b) (6)	21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472) by (b) (6)
5/26/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/26/2016 10:12 SWO- (b) (6) back to licking gravy, not eating a lot of solid food. Hind legs are weak. Owner not able to get sRR yet but seems comfortable. A/o to continue with current meds. If stops eating, then stop lasix. Otherwise will touch base in a few days. Gave owner my cell phone number if they need anything.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates,
 I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended,
 R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/25/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/25/2016 18:38 SWO- BW wnl, kidney values have decreased to normal. A/o to give lasix 12.5mg tabs- 1/4 tab SID. Will adjust based on appetite and breathing. Continue with other meds (pimo, taurine and app stimulant). Owner thankful.
5/25/2016	D	(b) (6)	Pleural Effusion Final
5/25/2016	C	(b) (6)	CARDIAC EVALUTION - CLOSED 05/28/2016 - Cardiac Evaluation

Date of evaluation: Wednesday, May 25, 2016

Date of previous evaluation: Sunday, May 15, 2016

CHIEF COMPLAINT: heavy breathing

HISTORY: Owners noted heavy breathing yesterday. Decreased appetite yesterday and today. Prior to that her appetite was improving. Owners transitioned her to royal canin and she started eating small amounts of solid food, previously only licking gravy.

Previous hx: Diagnosed with DCM (b) (6) 16. Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4. Taurine level confirmed taurine deficiency

PHYSICAL EXAM: The patient was bright, alert and responsive. No murmur on auscultation. The patient was tachypnic with mild increased effort, RR 48. Normal BV sounds, no crackles on auscultation. Faint referred upper airway noise. Femoral pulses were fair and synchronous. Heart rate was 180 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) (b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Brief Echo 5/25/16: moderate volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle unchanged from previous.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM (b) (6) 2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia, LV thrombus

THORACOCENTESIS 4/10/16: a total of 180ml of yellow fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed scant effusion remained in the patient.

THORACOCENTESIS (b) (6) 6: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

ER thoracocentesis 5/8/16: 25ml yellow tinged fluid from the right side.

SUMMARY: (b) (6) had 180ml fluid removed from her chest today. A renal panel showed normal renal values (BUN 31, creatinine 1.3)- previous azotemia. Will start with very low dose of lasix since decreased appetite right now (decreased appetite seemed to correlate with onset of heavy breathing). If appetite improves, can consider increasing lasix dose. Continue other medications as below. Recheck in 2 weeks, sooner if concerns.

MEDICATIONS:

START:

Lasix 12.5mg tablets- give 1/4 tablet by mouth once daily

CONTINUE:

Taurine 250mg by mouth twice daily

Mirtazepine 15mg tablets: Give 1/4 tablet by mouth every 3 days as needed.

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

5/25/2016 | (b) (6) Cardiology Discharge Instructions

(b) (6)
(b) (6)
(b) (6)
5/25/2016

(b) (6) had 160ml of fluid removed from her chest tonight. The clot in her heart appears similar to previous. I will call you with the bloodwork results and we can determine what to do with her lasix dose.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Recheck in 2 weeks, sooner if concerns.

MEDICATIONS:
CONTINUE:

Taurine 250mg by mouth twice daily
 Mirtazepine 15mg tablets: Give ¼ tablet by mouth every 3 days as needed.
 Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

WAIT TO BE INSTRUCTED FURTHER ON LASIX DOSE

Watch (b) (6) for the following clinical signs and call a veterinarian if you see any of these:

- Initiation of or increase in cough
- Excessive panting or wheezing
- Restlessness, unable to get comfortable
- Decreased appetite
- Lethargy/weakness
- Collapse or fainting

It has been a pleasure caring for (b) (6). Thank you for entrusting us with his care. If you have any further questions or problems, please don't hesitate to call.

5/25/2016 L	(b) (6)	(b) (6)	Cardiac Panel #10 results from (b) (4) Requisition ID: (b) (4) 0 Posted Final <table border="0" style="width: 100%;"> <thead> <tr> <th style="text-align: left;">Test</th> <th style="text-align: left;">Result</th> <th style="text-align: left;">Reference Range</th> </tr> </thead> <tbody> <tr> <td>HCT =</td> <td>34 %</td> <td></td> </tr> <tr> <td>NA+ =</td> <td>147.3 mmol/L</td> <td>146.2 - 156.2</td> </tr> <tr> <td>K+ =</td> <td>6.65 mmol/L H</td> <td>3.41 - 4.71</td> </tr> <tr> <td>CL- =</td> <td>115.1 mmol/L L</td> <td>117.0 - 125.3</td> </tr> <tr> <td>BUN =</td> <td>31 mg/dL</td> <td>22 - 33</td> </tr> <tr> <td>CREA =</td> <td>1.3 mg/dL</td> <td>0.07 - 1.9</td> </tr> </tbody> </table> Manually entered. PCV = 32% TS = 6.0g/dL	Test	Result	Reference Range	HCT =	34 %		NA+ =	147.3 mmol/L	146.2 - 156.2	K+ =	6.65 mmol/L H	3.41 - 4.71	CL- =	115.1 mmol/L L	117.0 - 125.3	BUN =	31 mg/dL	22 - 33	CREA =	1.3 mg/dL	0.07 - 1.9
Test	Result	Reference Range																						
HCT =	34 %																							
NA+ =	147.3 mmol/L	146.2 - 156.2																						
K+ =	6.65 mmol/L H	3.41 - 4.71																						
CL- =	115.1 mmol/L L	117.0 - 125.3																						
BUN =	31 mg/dL	22 - 33																						
CREA =	1.3 mg/dL	0.07 - 1.9																						

5/25/2016 P	(b) (6)	(b) (6)	30.00 ml of DNULsix 10mg/ml/ML (M0568) Rx #: 2576809 0 Of 12 Refills Give 0.5ml by mouth once daily or as directed by your veterinarian.
-------------	---------	---------	--

5/25/2016 V	(b) (6)	(b) (6)	May 25, 2016 04:26 PM Staff: (b) (6) ----- Weight : 5.30 kilograms Rm. 14
-------------	---------	---------	--

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/25/2016	CK	(b) (6)	breathing heavy Reason for Visit: Recheck Date Patient Checked Out: 05/25/16 Practice (b) (6)
5/25/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/25/2016 14:40 SWO- (b) (6) breathing heavy today. No interest in food yesterday or today. Owner to bring in this afternoon.
5/25/2016	B	(b) (6)	1.00 Specialty/Referral Exam Level 2 (REF02) by (b) (6)
5/25/2016	B	(b) (6)	Echo Guided Thoracocentesis Group (EGT) by (b) (6)
5/25/2016	B	(b) (6)	1.00 EGT Procedure (USSC50) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Equipment Service & Preparation (USEOPT) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Thoracocentesis Therapeutic (R33) by (b) (6)
5/25/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/25/2016	B	(b) (6)	1.00 In-house lab (XNBALIX) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Cardiac (b) (6) Panel #10 ((b) (6)) by (b) (6)
5/25/2016	B	(b) (6)	1.00 Cared for by (b) (6) ((b) (6)) by (b) (6)
5/25/2016	B	(b) (6)	30.00 ml of DNULsix 10mg/ml/ML (M0568) by (b) (6)
5/25/2016	B	(b) (6)	-30.00 ml of DNULsix 10mg/ml/ML (M0568) b (b) (6)
5/22/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/22/2016 16:23 SWO- Appetite is still the same, but now (b) (6) will go to the food on her own instead of owners bringing it to her. Still only eating gravy, no solid food yet. Owner has not tried Max Cal, recommended trying that. Cats who go prolonged period without eating at risk for hepatic lipidosis. Personality wise, she is much improved, almost back to normal self. Ambulating around the house as before. Very social. Owner bought Royal Canin as new diet. Gets taurine and pimo BID now. Told owner to continue appetite stimulant for now (had stopped this). Urinating and defecating outside of the litter box, not a SE of meds, likley behavioral. Soft stools. A/o let me know if soft stools continue.
5/19/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT 5/19/2016 16:18

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/19/2016	C	(b) (6)	<p>SWO - updating that I did talk with Merrick customer service today to take my complaint and are filing it with their quality assurance. I am not sure when they will get back with me, but I will let them know as soon as I hear anything. Owner thankful for call.</p>
5/19/2016	C	(b) (6)	<p>MEDICAL COMMENTS ***ADDENDUM 5/19/2016 5/19/2016 11:49 Called Merrick at 1(800)664-7387 to report taurine deficiency possibly related to consumption of their product, Merrick Purrfect Bistro Grain Free Real Chicken Recipe feline dry (best by 7/26/2017, lot #16025 DL1 38310 14131 - lost # difficult to read), USB# 22808 38310). Owner has been feeding this food for approximately 3 years, 5 cats total in household, product has been purchased from the (b) (6) in (b) (6) in (b) (6). Requesting that the company investigate this possible deficiency, also discussed that I would like for the other cats in the household to be tested. SW (b) (6) @ Merrick - said I could expect call back in 2 weeks, let her know I would like to know when to expect a call. She will submit complaint and let me know. ADDENDUM on 5/19/2016 at 15:28:19 from (b) (6) Merrick called back - additional questions of how long the cat has been sick - presented to ER (b) (6) and sick day before; also wanted to know if bag was new - yes bag was purchased about 2 weeks prior per owner. My concern however is that it takes several months for this to develop and I do not believe this is a single bag/lot issue.</p>
5/19/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT 5/19/2016 10:10 SWO - introduce3d myself, asked owner about diet history, has been feeding Merrick Purrfect Bistro Grain Free Real Chicken Recipe for approximately 3 years and purchasing from (b) (6) in (b) (6). Prior to this feeding Dick Van Pattons Indoor Formula Dry, chicken and salmon flavor. Discussed with owner that I will contact the company and also report to the FDA. Will let owner know of communication. In my experience sometimes the company will also want to reach out to the client. Owner thankful for call.</p>
5/18/2016	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE 5/18/2016 18:16 SWO- (b) (6) still not eating, had a little gravy this morning. Drank a lot of water today. Breathing is normal. Owner dropping food off tonight.</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/18/2016	TC	(b) (6)	COMMUNICATIONS WITH DOCTOR - TENTATIVE 5/18/2016 18:15 Imom for (b) (6) regarding appt on Saturday with other cats in house. Would like whole blood taurine levels sent to the UC Davis amino acid lab. Call me or speak with nutrition regarding any questions.
5/17/2016	P	(b) (6)	3.00 can of Iams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) Rx #: 2573240 0 Of 3 Refills Feed as directed
5/17/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/17/2016 15:58 SWO- did not eat much this morning. O gave appetite stimulant this morning and then left her alone with some food. Has not checked on her yet. Normal BM last night. sRR6brs/15sec last night. Vet coming for house call Saturday morning to take taurine sample for other cats. A/o to transition after that- recommended Hills Science Diet, Purina, Royal Canin. Will also rx Iams max cal for her to pick up here and offer (b) (6) .
5/17/2016	B	(b) (6)	3.00 can of Iams K9/Fel Max-Cal Plus 6oz cn 041484 (FI791) by (b) (6)
5/16/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/16/2016 18:04 SWO- (b) (6) seemed to be doing better last night. A little brighter when they got home. Started taurine supplementation last night. Not eating solid foods yet, but licking gravy- had the gravy from almost 3 cans last night. Owner put solid food in blender, but (b) (6) not interested (may have been too thick still). Drinking water. No BM, not a concern because she is not eating. Asked owner to bring in the food in original package as soon as possible, owner was planning on dropping off tomorrow. Also discussed to get other 4 cats tested for taurine levels this week, as since we are changing their diet we would like to know levels on current diet. Owner will call to either have mobile vet come to house or schedule here with GP this week. Told owner I will talk to nutrition about recommended diets.
5/15/2016	R	(b) (6)	Referral Letter - Cardio Resident Eval and labs - FINAL 05/15/2016 - REF

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

TO: (b) (6)
(b) (6)
(b) (6)
(b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: Sunday, May 15, 2016
RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 4 Mos. **Sex:** Spayed Female
Current Weight: 5.2 kilograms as of 5/15/2016

Thank you for referring (b) (6). The following is a case summary.

Date of evaluation: Sunday, May 15, 2016

Date of previous cardiac evaluation: Monday, (b) (6) 2016

CHIEF COMPLAINT: Recheck, not eating

HISTORY: (b) (6) has not eaten since discharge on (b) (6). Owner gave mirtazapine yesterday. Today licking some of the liquid off the food and very polydyspic, but no interest in eating any solid food. Will take a few steps and then lay down, very weak. Owner not able to get sRR, awake RR 6breaths/15sec. No heavy breathing noted. Previous hx: Diagnosed with DCM (b) (6) 16. Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4.

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was eupnic, RR 32. Normal BV sounds, no crackles on auscultation. Femoral pulses were fair and synchronous. Heart rate was 180 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) (b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted

, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, age cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, entative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

in left ventricle.

ECHOCARDIOGRAM (b) (6)/2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM). Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia, LV thrombus

THORACOCENTESIS (b) (6) 6: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the pati (b) (6) 6: 25ml yellow tinged fluid from the right side.

SUMMARY: The small amount of pleural effusion was not enough to warrant thoracocentesis, especially since patient eupnic. (b) (6) taurine level came back low (24nmol/ml). Recommended supplementing taurine at below dose. Discussed with owner it can take up to 2-3 weeks to see an effect of this and even longer (3 months) to see changes on echocardiogram. There is a chance (b) (6) will succumb to congestive heart failure before we see the positive effects of the taurine. A renal panel revealed azotemia. Discussed with owner the challenge going forward of managing the heart failure with azotemia. If (b) (6) develops trouble breathing, may not have options to treat. Euthanasia may be the most humane option for (b) (6) at that time.

The echocardiogram showed a large mass in the left ventricle, consistent with a thrombus. There is a risk that this clot, or a piece of it, leaves the heart and causes a thromboembolic event. The most common place for thromboembolic disease in cats is the aortic bifurcation, but can occur anywhere including the lungs which can be fatal. Advised owners, if FATE suspected at home, bring (b) (6) to closest emergency facility. At that time, euthanasia is recommended. We discussed holding off on an aspirin or Plavix for now, as it will not affect the current clot, and (b) (6) is not yet eating.

Owners willing to start taurine supplementation and take it day by day. Owners understand the possible outcomes we discussed today. I will call to check in tomorrow.

MEDICATIONS:

START:

Taurine 250mg by mouth twice daily

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

CONTINUE:

Mirtazepine 15mg tablets: Give 1/4 tablet by mouth every 3 days as needed.

HOLD FOR NOW:

Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

Thank you for the courtesy of this interesting referral. Please feel free to contact me with any questions or comments.

Sincerely,

(b) (6)

(b) (6)

Sent electronically - no signature required

(b) (6)

(b) (6)

Client ID: (b) (6) Patient ID: (b) (6) Patient Name: (b) (6)

DATE/TIME TEST	RESULT	REFERENCE RANGE
5/15/2016 BUN	= 61 mg/dL (H)	22 - 33
5/15/2016 CL-	= 104.5 mmol/L (L)	117.0 - 125.3
5/15/2016 CREA	= 3.1 mg/dL (H)	0.07 - 1.9
5/15/2016 HCT	= 43 %	
5/15/2016 K+	= 3.33 mmol/L (L)	3.41 - 4.71
5/15/2016 NA+	= 145.3 mmol/L (L)	146.2 - 156.2

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Lab Comments: Manually entered.

Additional Comments: BUN/Cre=19.7mg/mg PCV=39% TS=7.8g/dL Normal

Pain Condition Scores

Have you noticed the "PCS" aka Pain Condition Score in our medical records? As part of our AAHA specialty accreditation, we evaluate the level of pain or discomfort for each and every patient we see. You will see this reflected in the physical exam recordings as a "PCS". We follow the guidelines created and published by Colorado State University. If you have questions about interpreting pain or treatment options, or you need to schedule a pain management consult, please speak with our Anesthesiology & Pain Management Department by calling (b) (6).

Clinical Studies

We are recruiting for a variety of clinical studies across multiple specialties and are looking for eligible patients. For a complete list, please visit our website at [and click on Veterinary Professionals and then Current Clinical Studies.](#)

Avian & Exotics

(b) (6) is happy to re-introduce our Avian & Exotics Department and welcome (b) (6) to our team! We offer advanced diagnostics and treatment methods to complement the care provided by referring veterinarians. In addition to referral services, we provide primary care and medical boarding. Our facility maintains a dedicated exotics ward and a 24-hour monitored critical care unit. The Avian & Exotics Department accepts appointments 5 days a week (Tuesday-Saturday) and our emergency service receives patients, including exotic species, 24 hours a day. You can contact (b) (6) at (b) (6).

5/15/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/15/2016 16:34 SWO- discussed azotemia. Since (b) (6) is eupnic, would hold off on lasix for now. Hope would be that she may be able to breathe comfortably without lasix for enough time that taurine may start to help. Otherwise may give low dose of lasix, but going to be a big challenge with azotemia. Owners are to start taurine tonight. Discussed case with nutrition. Will file a complaint about the food. Will have more
-----------	----	---------	---

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			information on this tomorrow. Will call to check in tomorrow.
5/15/2016	D	(b) (6)	Taurine Deficiency Final
5/15/2016	D	(b) (6)	Azotemia Tentative
5/15/2016	D	(b) (6)	Pleural Effusion Final

5/15/2016 C (b) (6) CARDIAC EVALUTION - CLOSED 05/18/2016 - Cardiac Evaluation

Date of evaluation: Sunday, May 15, 2016

Date of previous cardiac evaluation: Monday, (b) (6) 2016

CHIEF COMPLAINT: Recheck, not eating

HISTORY: (b) (6) has not eaten since discharge on (b) (6) Owner gave mirtazapine yesterday. Today licking some of the liquid off the food and very polydyspic, but no interest in eating any solid food. Will take a few steps and then lay down, very weak. Owner not able to get sRR, awake RR 6breaths/15sec. No heavy breathing noted.

Previous hx: Diagnosed with DCM (b) (6) 16. Initially presented to ER for lethargy and ADR. Pleural effusion present. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3). Prior to discharge BUN was 74, creatinine was 1.4.

PHYSICAL EXAM: The patient was quiet, alert and responsive. No murmur on auscultation. The patient was eupnic, RR 32. Normal BV sounds, no crackles on auscultation. Femoral pulses were fair and synchronous. Heart rate was 180 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals)(b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Brief Echo 5/15/16: small volume pleural effusion. No pericardial effusion. Large mass noted in left ventricle.

ECHOCARDIOGRAM (b) (6) 2016:

IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm
IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia, LV thrombus

THORACOCENTESIS (b) (6) 16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.
ER thoracocentesis (b) (6) /16: 25ml yellow tinged fluid from the right side.

SUMMARY: The small amount of pleural effusion was not enough to warrant thoracocentesis, especially since patient eupnic. (b) (6) taurine level came back low (24nmol/ml). Recommended supplementing taurine at below dose. Discussed with owner it can take up to 2-3 weeks to see an effect of this and even longer (3 months) to see changes on echocardiogram. There is a chance (b) (6) will succumb to congestive heart failure before we see the positive effects of the taurine. A renal panel revealed azotemia. Discussed with owner the challenge going forward of managing the heart failure with azotemia. If (b) (6) develops trouble breathing, may not have options to treat. Euthanasia may be the most humane option for (b) (6) at that time. The echocardiogram showed a large mass in the left ventricle, consistent with a thrombus. There is a risk that this clot, or a piece of it, leaves the heart and causes a thromboembolic event. The most common place for thromboembolic disease in cats is the aortic bifurcation, but can occur anywhere including the lungs which can be fatal. Advised owners, if FATE suspected at home, bring (b) (6) to closest emergency facility. At that time, euthanasia is recommended. We discussed holding off on an aspirin or Plavix for now, as it will not affect the current clot, and (b) (6) is not yet eating. Owners willing to start taurine supplementation and take it day by day. Owners understand the possible outcomes we discussed today. I will call to check in tomorrow.

MEDICATIONS:

START:

Taurine 250mg by mouth twice daily

CONTINUE:

Mirtazepine 15mg tablets: Give 1/4 tablet by mouth every 3 days as needed.

HOLD FOR NOW:

Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

5/15/2016	I	CDS	Cardiology Discharge Instructions (b) (6) (b) (6) (b) (6) 5/15/2016
-----------	---	-----	---

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6) has a small amount of fluid in her chest today. It was not enough to warrant draining today.

Her taurine levels came back low. Please start supplementing taurine as below. It can take up to 2-3 weeks to see an effect of this.

The echocardiogram showed a large mass in one of the chambers of her heart (the left ventricle). There is a risk that this clot, or a piece of it, leaves the heart. If that happens, it can travel to any part of the body (lungs, hind legs, etc) and this can be fatal. We discussed holding off on an aspirin or Plavix medication for now, as it will not do anything for the current clot, and (b) (6) is not yet eating. I will call you with her bloodwork results this afternoon.

MEDICATIONS:

START:

Taurine 250mg by mouth twice daily

Watch (b) (6) for the following clinical signs and call a veterinarian if you see any of these:

- Initiation of or increase in cough
- Excessive panting or wheezing
- Restlessness, unable to get comfortable
- Decreased appetite
- Lethargy/weakness
- Collapse or fainting

It has been a pleasure caring for (b) (6). Thank you for entrusting us with his care. If you have any further questions or problems, please don't hesitate to call.

5/15/2016 L	(b) (6)	(b) (6)	Cardiac Panel #10 results from (b) (4) Requisition ID: <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">0</th> <th style="width: 10%;">Posted</th> <th style="width: 10%;">Final</th> <th style="width: 10%;"></th> <th style="width: 10%;"></th> <th style="width: 10%;"></th> <th style="width: 10%;"></th> <th style="width: 10%;"></th> </tr> </thead> <tbody> <tr> <td>Test</td> <td></td> <td></td> <td>Result</td> <td></td> <td></td> <td></td> <td>Reference Range</td> </tr> <tr> <td>HCT =</td> <td></td> <td></td> <td>43 %</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>NA+ =</td> <td></td> <td></td> <td>145.3 mmol/L L</td> <td></td> <td></td> <td></td> <td>146.2 - 156.2</td> </tr> <tr> <td>K+ =</td> <td></td> <td></td> <td>3.33 mmol/L L</td> <td></td> <td></td> <td></td> <td>3.41 - 4.71</td> </tr> <tr> <td>CL- =</td> <td></td> <td></td> <td>104.5 mmol/L L</td> <td></td> <td></td> <td></td> <td>117.0 - 125.3</td> </tr> <tr> <td>BUN =</td> <td></td> <td></td> <td>61 mg/dL H</td> <td></td> <td></td> <td></td> <td>22 - 33</td> </tr> <tr> <td>CREA =</td> <td></td> <td></td> <td>3.1 mg/dL H</td> <td></td> <td></td> <td></td> <td>0.07 - 1.9</td> </tr> <tr> <td colspan="8">Manually entered.</td> </tr> <tr> <td colspan="2">BUN/Cre=19.7mg/mg</td> <td colspan="2">PCV=39%</td> <td colspan="2">TS=7.8g/dL</td> <td colspan="2">Normal</td> </tr> </tbody> </table>	0	Posted	Final						Test			Result				Reference Range	HCT =			43 %					NA+ =			145.3 mmol/L L				146.2 - 156.2	K+ =			3.33 mmol/L L				3.41 - 4.71	CL- =			104.5 mmol/L L				117.0 - 125.3	BUN =			61 mg/dL H				22 - 33	CREA =			3.1 mg/dL H				0.07 - 1.9	Manually entered.								BUN/Cre=19.7mg/mg		PCV=39%		TS=7.8g/dL		Normal	
0	Posted	Final																																																																																	
Test			Result				Reference Range																																																																												
HCT =			43 %																																																																																
NA+ =			145.3 mmol/L L				146.2 - 156.2																																																																												
K+ =			3.33 mmol/L L				3.41 - 4.71																																																																												
CL- =			104.5 mmol/L L				117.0 - 125.3																																																																												
BUN =			61 mg/dL H				22 - 33																																																																												
CREA =			3.1 mg/dL H				0.07 - 1.9																																																																												
Manually entered.																																																																																			
BUN/Cre=19.7mg/mg		PCV=39%		TS=7.8g/dL		Normal																																																																													

5/15/2016 V (b) (6) May 15, 2016 03:24 PM Staff: (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

			----- Weight : 5.20 kilograms cardio baby scale
5/15/2016	CK	(b) (6)	Reason for Visit: Recheck Date Patient Checked Out: 05/15/16 Practice (b)

5/15/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/15/2016 13:29 SWO (mrs)- owner gave mirtazapine, no improvement in appetite. Drinking excessively. Having a hard time walking, very weak. Owner not able to get sRR, awake breathing 6breaths/15sec. Offered to see (b) (6) today. Made appt for 3pm
-----------	----	---------	---

5/15/2016	B	(b) (6)	1.00 Specialty/Referral Exam Level 2 (REF02) by (b) (6)
5/15/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/15/2016	B	(b) (6)	1.00 In-house lab (XNBALIX) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Cardiac (b) (6) Panel #10 (b) (6) by (b) (6)
5/15/2016	B	(b) (6)	Echo Guided Thoracocentesis Group (EGT) by (b) (6)
5/15/2016	B	(b) (6)	1.00 EGT Procedure (USSC50) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
5/15/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)

5/14/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/14/2016 17:42 Mrs called and Imovm that (b) (6) hasn't been eating well. I called back and sw Mr. He said she is eating only very tiny amounts and not improving, wanted to know if I had suggestions. Owners feel breathing is still ok, 6 breaths/15 seconds but coughed a little today. I told Mr she could have poor app due to fluid reforming or azotemia or her heart disease in general. She may need to be rechecked sooner than later to evaluate this and r/o fluid and worsening azo. Owners plan to discuss w/(b) (6) but wanted to know if there is something they could give her before morning. I offered to prescribe appetite stimulant, explained that this may not work b/c it doesn't override what is causing the inappetance in the 1st place but it's fine to try. Mr was thankful, said he may or may not pick it up tonight but is glad to have the option.
-----------	----	---------	--

5/14/2016	P	(b) (6)	2.00 tablet of Mirtazapine 15mg Tablet (M1052)
-----------	---	---------	--

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/14/2016	B	(b) (6)	Rx #: 2571986 0 Of 3 Refills Give 1/4 tablet once every 3 days as needed to stimulate appetite. 2.00 tablet of Mirtazapine 15mg Tablet (M1052) by (b) (6)
5/12/2016	C	(b) (6)	TRIAGE CALL 5/12/2016 21:23 Per owner, (b) (6) appetite has been decreasing over the last couple days. Yesterday only ate about 2 tablespoons, today less. Let owner know that if the appetite has been decreasing recommend a recheck. Owner wants to talk to cardio first to see about an appetite stimulant. She will call tomorrow to speak with cardio department.
5/11/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/11/2016 13:16 SWO (mrs)- still weak and unstable, but up and walking around short distances. sRR was 5breaths/15sec this morning. Ate 2 teaspoons canned food last night, so owner gave 1/4 tab lasix. Has not eaten yet this morning. Advised owner since sRR wnl, hold off on lasix for now. Will restart when either (b) (6) has a good appetite or if sRR >8br/15esc. Owner understands. Also hold off on pimo and taurine supplement. Should have taurine level back by recheck in 2 weeks. Owner asked about starting aspirin. Can consider aspirin/plavix at recheck if appetite is good. Discussed that they may lower risk, but do not prevent risk of clot formation. Will call to check on appetite in a few days. Owner to call sooner with concerns.
5/10/2016	TC	(b) (6)	COMMUNICATIONS WITH CLIENT - TENTATIVE 5/10/2016 10:13 SWO (mrs)- (b) (6) was drinking a lot last night. Has not eaten anything yet. sRR 7breaths/15sec. Has gotten up and walked around, otherwise sleeping. Advised owner to hold off on meds today, would like her eating before restarting them. Will call tomorrow to check on appetite and advised what to do with lasix.
5/10/2016	L	(b) (6)	Miscellaneous results from (b) (6) ID: 189206 Posted Final Asc: (b) (6) Profile: Taurine RE: 16758 Sample: PLASMA, HEPARIN RE: 16759 Taurine 24 NMOL/ML nmol/ml Feline taurine ranges: normal plasma 60-120 nmol/mL critical level <40 nmol/mL; whole blood normal 300-600 nmol/mL

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			<p>critical level <200 nmol/mL. TEST PERFORMED AT THE UNIVERSITY OF WISCONSIN</p> <p>FELINE PLASMA NORMALS: 60-120 nmol/ml WHOLE BLOOD NORMALS: 300-600 nmol/ml PLASMA CRITICAL: LESS THAN 40 nmol/ml WHOLE BLOOD CRITICAL: LESS THAN 200 nmol/ml</p> <p>K9 PLASMA NORMALS: 60-120 nmol/ml WHOLE BLOOD NORMALS: 200-350 nmol/ml PLASMA CRITICAL: LESS THAN 40 nmol/ml WHOLE BLOOD CRITICAL: LESS THAN 150 nmol/ml</p>
(b) (6) 2016	R	(b) (6)	Referral Letter - Cardio Resident Eval and labs - FINAL 05/09/2016 - REF

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

TO: (b) (6)
(b) (6)
(b) (6)
(b) (6)
FAX #: (b) (6)
FROM: (b) (6)
DATE: Monday, May 09, 2016
RE:
Client: (b) (6)
Patient: (b) (6)
Breed: Shorthair, Domestic
Age: 12 Yrs. 4 Mos. **Sex:** Spayed Female
Current Weight: 15.6 pounds as of 12/28/2009

Thank you for referring (b) (6). The following is a case summary.

Date of evaluation: Monday, May 09, 2016

CHIEF COMPLAINT: pleural effusion

HISTORY: Presented to ER last night for lethargy and ADR. Cursorsy ultrasound revealed pleural effusion. Thoracocentesis yielded 25ml yellow tinged fluid from the right side. Patient received 12mg lasix IV and was placed in oxygen overnight. Her RR was wnl, with slight effort noted overnight. She ate a small amount. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3).

PHYSICAL EXAM: The patient was bright, alert and responsive. No murmur on auscultation, but heart sound slightly muffled. The patient was eupnic, RR 30 with slight effort. Normal BV sounds, no crackles on auscultation. Pulses were fair and synchronous. Heart rate was 170 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals) (b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM). Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial

, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, age cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6) **Patient:** (b) (6)
Phone: (b) (6) **Species:** Feline **Breed:** Shorthair, Domestic
Address: (b) (6) **Age:** 12 Yrs. 5 Mos. **Sex:** Spayed Female
(b) (6) **Color:** Black

Date	Type	Staff	History
------	------	-------	---------

effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia

THORACOCENTESIS (b) (6)/16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

SUMMARY: (b) (6) has dilated cardiomyopathy (DCM) and has developed pleural effusion due to congestive heart failure. The pleural effusion was removed today and we are starting the medications as described below. Although taurine deficiency is a rare cause for dilated cardiomyopathy in cats, we submitted a taurine level to the lab today to look for this. The test results can take up to 2 weeks; we will fax results when they are available. Unfortunately the prognosis for DCM (with no taurine deficiency) in cats is poor, with survival time 3-6 weeks or in rare cases 4-6 months. A renal panel was recheck today and showed elevated BUN (74) and normal creatinine (1.4). Discussed with owner the azotemia may make treating the congestive heart failure challenging.

We discussed that with heart enlargement there is a risk for developing a blood clot, or stroke. Although aspirin +/- plavix can be given in hopes of reducing clot formation, they have not been definitively proven to prevent blood clot formation in cats. If the owners elect to start this medication I would recommend waiting until we know she is eating and feeling well on the below medications as aspirin/plavix can cause GI side effects (vomiting, inappetance) in some cats. A recheck with cardiology is recommended in 2 weeks, or sooner if clinical signs.

Advised owner congestive heart failure likely contributing to lethargy, but concern that weakness may be secondary to the DCM and low cardiac output. If that is the case, may not improve once we get her out of failure. Owner comfortable taking (b) (6) home to see how she does. (b) (6) was discharged on the below medications. If she does not improve at home over the next few days, owner to consider euthanasia.

MEDICATIONS:

START: Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

START in 3 DAYS IF EATING

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.
If eating and easy to medicate, start: Taurine 250 mg by mouth twice a day with food.

Thank you for the courtesy of this interesting referral. Please feel free to contact me

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

with any questions or comments.

Sincerely,

(b) (6)

(b) (6)

Sent electronically - no signature required

(b) (6)

(b) (6)

Client ID: (b) (6) Patient ID: (b) (6) Patient Name: (b) (6)

DATE/TIME	TEST	RESULT	REFERENCE RANGE
(b) (6)	CREA	= 1.4 mg/dL	0.8 - 2.4

Lab Comments: CREA: Test results for the latest analyzer run have been multiplied by the dilution factor for a dilution of 1 in 4 total.

DATE/TIME	TEST	RESULT	REFERENCE RANGE
(b) (6)	ALB	= 2.8 g/dL	2.3 - 3.9
(b) (6)	ALB/GLOB	= 0.8	
(b) (6)	ALKP	= 11 U/L (L)	14 - 111
(b) (6)	ALT	= 140 U/L (H)	12 - 130
(b) (6)	BUN/UREA	= 74 mg/dL (H)	16 - 36
(b) (6)	Chloride	= 100 mmol/L (L)	112 - 129
(b) (6)	CREA	= --- mg/dL	0.8 - 2.4
(b) (6)	GLOB	= 3.3 g/dL	2.8 - 5.1
(b) (6)	GLU	= 105 mg/dL	71 - 159
(b) (6)	Na/K	= 29	
(b) (6)	OSM calc	= 298 mmol/kg	
(b) (6)	PHOS	= 7.0 mg/dL	3.1 - 7.5
(b) (6)	Potassium	= 4.7 mmol/L	3.5 - 5.8
(b) (6)	Sodium	= 138 mmol/L (L)	150 - 165

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
		(b) (6)	= 6.1 g/dL 5.7 - 8.9

Pain Condition Scores

Have you noticed the "PCS" aka Pain Condition Score in our medical records? As part of our AAHA specialty accreditation, we evaluate the level of pain or discomfort for each and every patient we see. You will see this reflected in the physical exam recordings as a "PCS". We follow the guidelines created and published by Colorado State University. If you have questions about interpreting pain or treatment options, or you need to schedule a pain management consult, please speak with our Anesthesiology & Pain Management Department by calling (b) (6).

Clinical Studies

We are recruiting for a variety of clinical studies across multiple specialties and are looking for eligible patients. For a complete list, please visit our website at [and click on Veterinary Professionals and then Current Clinical Studies.](#)

Avian & Exotics

(b) (6) is happy to re-introduce our Avian & Exotics Department and welcome (b) (6) to our team! We offer advanced diagnostics and treatment methods to complement the care provided by referring veterinarians. In addition to referral services, we provide primary care and medical boarding. Our facility maintains a dedicated exotics ward and a 24-hour monitored critical care unit. The Avian & Exotics Department accepts appointments 5 days a week (Tuesday-Saturday) and our emergency service receives patients, including exotic species, 24 hours a day. You can contact (b) (6) at (b) (6).

(b) (6) 2016 | (b) (6) Cardiology Discharge Instructions
(b) (6)
(b) (6) 2016

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

A cardiologist has evaluated (b) (6) and has diagnosed her with Dilated Cardiomyopathy (DCM). DCM means she has poor muscle contraction of the heart and she has developed significant heart enlargement over time. Her clinical signs were due to congestive heart failure (fluid buildup around the lungs called pleural effusion), which developed secondary to the enlarged heart. We removed all the pleural effusion today. The fluid will reform but how fast this occurs is unpredictable. Please start the medications as below to help clear fluid and slow the fluid formation.

Although taurine deficiency is a rare cause for cardiomyopathy in cats, we submitted a taurine level to the lab today to look for this. The test results can take up to 2 weeks; we will call you when the results are available.

As we discussed, (b) (6) has elevations in her kidney values. This can make treating her heart disease challenging because she may not tolerate the lasix. If her kidney values become elevated to a certain degree, it will make her feel sick and she will likely have a decreased appetite or stop eating. We will monitor her kidney values with bloodwork over time.

Cats with heart enlargement are at risk for developing a blood clot, or stroke. Although aspirin and/or plavix can be given in hopes of reducing clot formation, they have not been proven to prevent blood clot formation in cats. If you elect to start this medication I would recommend waiting until she is eating and feeling well at home as these medications can cause GI side effects (vomiting, inappetance) in some cats.

Please periodically take a sleeping respiratory rate (sRR) at home. WHILE (b) (6) IS SLEEPING, count the number of times she breathes in over 15 seconds. She should breathe 8 or fewer breaths in 15 seconds.

A recheck with cardiology is recommended in 2 weeks, or sooner if you see any of the below signs.

MEDICATIONS:

START TODAY: Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet by mouth once a day

Furosemide: Also called Salix or Lasix. This is a diuretic and will help clear the fluid from your pet's lungs. Side effects include electrolyte abnormalities (if they stop eating), dehydration and kidney enzyme elevations. Blood work can be done to monitor these. This medication will be probably given for the life of your pet.

START IN 3 DAYS IF EATING:

Pimobendan 1.5 mg tiny tabs: Give 1 tablet by mouth two times a day WITH

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

FOOD.

Pimobendan is a phosphodiesterase inhibitor that gives increased contractility and arterial vasodilation. This will help the heart function better, allow your cat to feel better and live longer. Any medication can upset the stomach. This drug does not typically cause this, but if you see any changes, please stop the drug till you talk to a doctor here at (b) (6). Please give this with (b) (6) meals. Giving on empty stomach is more likely to make her nauseous.

We have called this medication into (b) (6). Please call them to order it and they will mail it to you.

If eating, start: Taurine 250 mg by mouth twice a day with food. I have submitted blood for a taurine level. The result may not return for 2 weeks. In the meantime, please start Taurine at home, 250 mg two times a day with food. This can be purchased at any health food store. If she is not eating well or if it is difficult to give her this medication, you can skip this until we get the taurine result from the blood work.

Watch (b) (6) for the following clinical signs and call a veterinarian if you see any of these:

- Initiation of or increase in cough
- Excessive panting or wheezing
- Restlessness, unable to get comfortable
- Decreased appetite
- Lethargy/weakness
- Collapse or fainting

It has been a pleasure caring for (b) (6). Thank you for entrusting us with his care. If you have any further questions or problems, please don't hesitate to call.

(b) (6) L

(b) (6)
Chemistry results from (b) (6) In-clinic Laboratory Requisition ID: 197

Test	Result	Posted	Final Reference Range
CREA =	1.4 mg/dL		0.8 - 2.4

CREA: Test results for the latest analyzer run have been multiplied by the dilution factor for a dilution of 1 in 4 total.

(b) (6) C

(b) (6)

PHARMACY NOTE

Called (b) (6) and spoke to (b) (6). Ordered Pimobendan

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			1.5mg tiny tabs. Give 1 tablet by mouth twice daily with food. #100, 12 refills

(b) (6) L	Chemistry results from (b) (6) Laboratory Requisition ID: 197	In-clinic Posted	Final
Test	Result	Reference	Range
ALB =	2.8 g/dL	2.3 - 3.9	
ALKP =	11 U/L L	14 - 111	
ALT =	140 U/L H	12 - 130	
BUN/UREA =	74 mg/dL H	16 - 36	
Chloride =	100 mmol/L L	112 - 129	
CREA -	--.-- mg/dL	0.8 - 2.4	
GLU =	105 mg/dL	71 - 159	
PHOS =	7.0 mg/dL	3.1 - 7.5	
Potassium =	4.7 mmol/L	3.5 - 5.8	
Sodium =	138 mmol/L L	150 - 165	
TP =	6.1 g/dL	5.7 - 8.9	
GLOB =	3.3 g/dL	2.8 - 5.1	
ALB/GLOB =	0.8		
Na/K =	29		
OSM calc =	298 mmol/kg		

(b) (6) TC (b) (6) COMMUNICATIONS WITH CLIENT - TENTATIVE
 (b) (6) 2016 12:35
 10am- SWO (mrs)- Discussed echo confirmed heart disease, DCM. Reviewed causes of DCM (unlikley taurine def, but will submit for levels) and prognosis with owner. Risk of future episodes of CHF, when is unpredictable. Oenwer consented to thoracocentesis. If continues to breath comfortably out of oxygen can go home this afternoon.

 12:30pm- SWO- (b) (6) breathing is stable out of oxygen. Very weak and lethargic. Ate a small amount of food this morning. Discussed since breathing is comfortable, can try at home. If energy level does not improve at home over the next few days, may consider euthanasia. Discussed elevated kidney values and how that is giong to make treating CHF with lasix challenging. Owner is comfortable with trying (b) (6) at home to see how she does. Will have husband call back to set up a time.

(b) (6) P (b) (6)
 21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472)
 Rx #: 2569385 0 Of 12 Refills
 Give 1 tablet by mouth twice daily with food.
 60.00 tablet of Lasix (Salix / Furosemide) 12.5mg Tablet (M568)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)	D	(b) (6)	Rx #: 2569382 0 Of 0 Refills Give 1/2 tablet by mouth twice daily. Pleural Effusion Final Left Atrial Enlargement Final Dilated Cardiomyopathy Final
---------	---	---------	--

(b) (6) C (b) (6) RADIOLOGY REVIEW - FINAL 05/09/2016
The DV view of the thorax obtained on May 8, 2016 has been reviewed and there is a moderate to large amount of pleural effusion that obscures visualization of the cardiac silhouette. There is also an area of increased opacity in the left hemithorax in the region of the caudal segment of left cranial lung lobe. The remaining lung parenchyma appears to be within normal limits and the pulmonary vessels appear normal. This combination of findings may be the result of heart failure or neoplasia and a cardiac consult is recommended for further evaluation.

This review was written by: (b) (6)

(b) (6) C (b) (6) CARDIAC EVALUTION - CLOSED 05/12/2016 - Cardiac Evaluation

Date of evaluation: Monday, May 09, 2016

CHIEF COMPLAINT: pleural effusion

HISTORY: Presented to ER last night for lethargy and ADR. cursory ultrasound revealed pleural effusion. Thoracocentesis yielded 25ml yellow tinged fluid from the right side. Patient received 12mg lasix IV and was placed in oxygen overnight. Her RR was wnl, with slight effort noted overnight. She ate a small amount. A renal panel performed overnight revealed azotemia (BUN 67, Creat 5.3).

PHYSICAL EXAM: The patient was bright, alert and responsive. No murmur on auscultation, but heart sound slightly muffled. The patient was eupnic, RR 30 with slight effort. Normal BV sounds, no crackles on auscultation. Pulses were fair and synchronous. Heart rate was 170 bpm, regular rhythm. PCS 0/4. BCS 8/9.

RADIOGRAPHS (DV, both laterals (b) (6) 16: Diffusely increased opacity obscuring the cardiac silhouette. Area of moderate increased opacity in the region of the caudal segment of the left cranial lobe. Pulmonary vasculature appears wnl.

ECHOCARDIOGRAM (b) (6) 2016:
IVSd: 0.37 cm LVIDd: 1.94 cm LVPWd: 0.48 cm

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (5)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (5)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

IVSs: 0.35 cm LVDs: 1.86 cm LVPWs: 0.48 cm %FS: 4 %
Ao: 0.8 cm LAD: 1.6 cm LA:Ao ratio 2 LA max: 1.5 cm LLAD: 1.57 cm

Comments: The left atrium is moderately enlarged. The left ventricle is enlarged in systole and diastole with poor left ventricular function. Mild right atrial and ventricular enlargement. Trivial MR and TR noted. No evidence for systolic anterior motion of the mitral valve (SAM).

Aortic and pulmonic flows were reduced. Moderate volume pleural effusion, no pericardial effusion. ECG on echo: HR 160, sinus rhythm.

DIAGNOSIS/PROBLEM LIST: Dilated Cardiomyopathy (DCM), moderate left atrial enlargement, pleural effusion, azotemia

THORACOCENTESIS (b) (6) 16: a total of 120 cc of yellow tinged fluid was removed with an 18 gauge needle from the right side of the chest in right lateral recumbency. No sedation. A post-tap echo showed mild effusion remained in the patient.

SUMMARY: (b) (6) has dilated cardiomyopathy (DCM) and has developed pleural effusion due to congestive heart failure. The pleural effusion was removed today and we are starting the medications as described below. Although taurine deficiency is a rare cause for dilated cardiomyopathy in cats, we submitted a taurine level to the lab today to look for this. The test results can take up to 2 weeks; we will fax results when they are available. Unfortunately the prognosis for DCM (with no taurine deficiency) in cats is poor, with survival time 3-6 weeks or in rare cases 4-6 months. A renal panel was recheck today and showed elevated BUN (74) and normal creatinine (1.4). Discussed with owner the azotemia may make treating the congestive heart failure challenging.

We discussed that with heart enlargement there is a risk for developing a blood clot, or stroke. Although aspirin +/- plavix can be given in hopes of reducing clot formation, they have not been definitively proven to prevent blood clot formation in cats. If the owners elect to start this medication I would recommend waiting until we know she is eating and feeling well on the below medications as aspirin/plavix can cause GI side effects (vomiting, inappetance) in some cats. A recheck with cardiology is recommended in 2 weeks, or sooner if clinical signs.

Advised owner congestive heart failure likely contributing to lethargy, but concern that weakness may be secondary to the DCM and low cardiac output. If that is the case, may not improve once we get her out of failure. Owner comfortable taking (b) (6) home to see how she does. (b) (6) was discharged on the below medications. If she does not improve at home over the next few days, owner to consider euthanasia.

MEDICATIONS:

START: Furosemide (Salix, Lasix) 12.5 mg tablets: Give 1/4 tablet once daily.

START in 3 DAYS IF EATING

Pimobendan 1.5mg tiny tabs: Give 1 tablet by mouth two times a day WITH FOOD.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
If eating and easy to medicate, start: Taurine 250 mg by mouth twice a day with food.			

(b) (6) 2016 L (b) (6)

Basic Metabolic Profile, (b) (6) results from (b) (6)
In-Clinic Requisition ID: 0 **Posted** **Final**

Test	Result	Reference Range
HCT =	40 %	12 - 70
HB =	13.1 g/dL	9.9 - 14.9
NA+ =	146.3 mmol/L	146.2 - 156.2
K+ =	4.99 mmol/L H	3.41 - 4.71
CL+ =	107.8 mmol/L L	117.0 - 125.3
CA++ =	1.17 mmol/L	1.16 - 1.35
MG++ =	1.08 mmol/L H	0.33 - 0.49
GLU =	156 mg/dL H	72 - 132
LAC =	9.7 mg/dL H	0.7 - 1.9
BUN =	67 mg/dL H	22 - 33
CREAT =	5.3 mmol/L H	1.1 - 3.5
O2CAP =	18.2 mL/dL	
TCO2 =	19.9 mmol/L	
GAP =	20.3 mmol/L	
CA/MG =	1.1 mol/mol	
OSM =	313.5 mOsm/kg	
BUN/CREA =	12.7 mg/mg	

Manually entered.
 PCV: 43% T.S: 6.6mg/dl

(b) (6) TC (b) (6)

LAB RESULTS - NOTES - TENTATIVE
 (b) (6) 00:00
 Lab Results: PCV: 42% TS g/dl: 6.8 Serum: Normal
 Original Lab Date:

B (b) (6)
 B
 B
 B
 B
 B
 B
 B

Laboratory Request / Sample Handling (LABS) by (b) (6)
 1.00 Sample Handling & Disposal (LFEE) by (b) (6)
 1.00 Basic Metabolic (b) (6) Panel # 2 (b) (6) by (b) (6)
 1.00 Specialty/Referral Exam Level 3 (REF03) by (b) (6)
 Echocardiogram Level 3 Group (USSC19) by (b) (6)
 1.00 VRC04 Procedure (VRC04) by (b) (6)
 1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
 1.00 Thoracocentesis Therapeutic (R33) by (b) (6)

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
(b) (6)	2016 B	(b) (6)	60.00 tablet of Lasix (Salix / Furosemide) 12.5mg Tablet (M568) by (b) (6)
	2016 B		21.00 tablet of Pimobendan 1.5mg Tab (Cpd) (M00472) by (b) (6)
	2016 B		Oxygen Therapy (Caged)/Hour (Group) (O2CAGE) by (b) (6)
	2016 B		7.00 O2 Therapy Per Hour (T044) by (b) (6)
	2016 B		7.00 Oxygen-related Patient Care / Hour (O2CARE) by (b) (6)
	2016 B		1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
	2016 B		Hospitalization Hours Smart Group (HOSPIT) by (b) (6)
	2016 B		17.00 Hospitalization Hours- Feline (H01) by (b) (6)
	2016 B		1.00 In-House Nutrition Assessment Level 1 (NTR012) by (b) (6)
	2016 B		17.00 Critical Care Level 2- Hours (CCU2) by (b) (6)
	2016 B		.20 ml of DNULsix 50mg/ml/ML (T106) by (b) (6)
	2016 B		1.00 Cared for by (b) (6) (b) (6) by (b) (6)
	2016 B		1.00 Cared by (b) (6) - Cardiology (CDS) by (b) (6)
	2016 B		Laboratory Request / Sample Handling (LABS) by (b) (6)
	2016 B		1.00 Taurine U of Wisc (via (b) (6) S16755) (L0245) by (b) (6)
	2016 B		1.00 Taurine U of Wisc (via (b) (6) S16755) (L0245) by (b) (6)
	2016 B		Laboratory Request / Sample Handling (LABS) by (b) (6)
	2016 B		1.00 Chemistry IV Renal Panel RBVH (CH25) by (b) (6)
	2016 B		Laboratory Request / Sample Handling (LABS) by (b) (6)
	2016 B		1.00 Dilution Verification Catalyst CREA (CH11DV) by (b) (6)

(b) (6) C (b) (6)

EMERGENCY PHYSICAL EXAM - Closed May 10/2016
 (b) (6)

Chief Complaint: Lethargic

History: Starting yesterday patient was noted to be lethargic and not herself. 4 other cats so difficult to say if she was eating but they think she was. Not sure about U/BM. Indoor only. Did not notice she was having issues breathing.

Other Medical Problems: None

Medications/Supplements: None

Environment: Indoor only

Vaccination Status:

Current Diet (Type):
 - Frequency:
 - Amount:

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

Physical Examination:

S(ubjective): QAR, hydration WNL, BCS 7/9, pain score: 0/4

O(b)jective:

Weight: 15.6 pounds

TPR: [temp - 93]F, [HR - 150] bpm, [RR - 60] bpm

EENT: clear AU/OU, no nasal discharge, normal cervical palpation, mm pink, moist/CRT < 2 sec

INTEG: Hair coat ok

PLN: WNL

CV: Heart sounds muffled

RESP: Increased RE, dull lung sounds

GI: soft, nonpainful, no masses

UG: SF, WNL

M/S: amb x 4

Neuro: alert/appropriate, cranial nerves intact

Problems/Differential Diagnoses:

Dyspnea

Lethargy

Diagnostics:

Cursory ultrasound - mild to moderate amount of pleural effusion R>L

DV thoracic radiograph - cardiac silhouette difficult to visualize, pleural effusion

(b) (6) 2

Assessment:

12 yr SF DSH

1. Pleural effusion, dyspnea - r/o cardiac (HCM) vs neoplasia (lymphoma vs other)

Treatment:

12 mg Lasix IM at 10 PM

Place in O2 cage

Thoracocentesis - 25 mL clear to yellow fluid removed from the right side

Place IVC, 12 mg Lasix IV at 2 AM

Plan/Recommendations:

Discussed differentials for pleural effusion - cardiac vs neoplasia. Due to pleural effusion cannot tell on radiographs if this is cardiac over neoplastic. Rec thoracocentesis to make (b) (6) breathe more comfortably - o consents. Rec echocardiogram in the morning to see if this is heart disease. If this is (b) (6) spoke about disease process and prognosis. If this is neoplasia owner's may decide to

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
 Phone: (b) (6)
 Address: (b) (6)
 (b) (6)

Patient: (b) (6)
 Species: Feline
 Age: 12 Yrs. 5 Mos.
 Color: Black

Breed: Shorthair, Domestic
 Sex: Spayed Female

Date Type Staff History

stop. Discussed possibility of taking repeat radiographs tomorrow if needed.

(b) (6) R

(b) (6)

Tx Template (blank)- Old WTS - TENTATIVE

WARD TREATMENT SHEET

DATE: (b) (6)		WARD: ccu	
LAST NAME: (b) (6)		CLINICIAN: (b) (6)	
PATIENT NAME: (b) (6)		TRANSFER DOCTOR:	
BREED: Shorthair, Domestic	COLOR: Black	LEGEND: O = scheduled X = performed D/C = discontinued inc = increase dec = decrease ⊖ = not given	
AGE: 12 Yrs. 4 Mos.	SEX: Spayed Female		
WEIGHT: 15.6 pounds as of: 12/28/2009			
PROBLEM/WORKING DIAGNOSIS:		SURGERY:	
Pleural effusion		IV CATHETER:	
ALERT: increased RR		CODE: RED	

Technician STAFF ID ==>			8	9	10	11	N	1	2	3	4	5	6	7	8	9	10	11	M	1	2	3	4	5	6	7
DR Staff ID	TREATMENTS	Call Parameters																								
	Temperature	<99, >102.5																								
	HR/MM/CRT	<140, >240																								
	RR/RE	>40, increased RE	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O
	Pulse Quality	wk																								
	Assess Pain & Note Score	2																								
	WEIGH																									
	RER: 262 kcal/day Diet: 115 g EN BID - any -				O											O										
	Quantify Food Intake							O									O									
	Lasix 12 mg IV - ask		O																			O				
	O2 cage																									

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6) _____
Phone: (b) (6) _____
Address: (b) (6) _____
 (b) (6) _____

Patient: (b) (6) _____
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

	AM SHIFT
	MID SHIFT
	OVER

TIME	Init. TECH	Temp	HR	Pulse qual.	RR/effort	MM color	CRT	PCV/ ts	BG	Turn	Blood Pressure	Pain Score	BM	Urine	Vomit	Grams Consumed	Init. AA
8 a																	
9 a																	
10 a																	
11a																	
N																	
1 p																	
2 p																	
3 p																	
4 p																	
5 p																	
6 p																	
7 p																	
8 p																	
9 p																	
10 p																	
11 p																	
M																	
1 a																	
2 a																	
3 a																	
4 a																	
5 a																	
6 a																	
7 a																	

TIME	ADDITIONAL COMMENTS

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 12 Yrs. 5 Mos.
Color: Black

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6)

3.00 O2 Therapy Per Hour (T044) by (b) (6)
3.00 Oxygen-related Patient Care / Hour (O2CARE) by (b) (6)
1.00 Equipment Service & Preparation (USEQPT) by (b) (6)
IV Catheter with Injection Cap (IVCATCP) by (b) (6)
1.00 IV Catheter Placement (CATH) by (b) (6)
1.00 each of Tx Catheter IV 20g x 2" Surflo (PINK) (H0112) by (b) (6)
1.00 each of Tx IV Ext T Set Hospira 1265028 (H027) by (b) (6)
1.00 each of Tx Injection Cap/Plug Termo 007110 (H118) by (b) (6)
1.00 Cared for by (b) (6) (b) (6) by (b) (6)
Thorax Radiographic Study Group (RADTH) by (b) (6)
1.00 Radiograph Preparation (XFEE) by (b) (6)
1.00 One view rdgh stdy (RAD1V) by (b) (6)
1.00 Radiologist Review Fee (RADGN) by (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
6/7/2016	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE 6/7/2016 12:33 On the phone with client discussing (b) (6), also discussed (b) (6). T4 and liver values need to be rechecked 6 weeks after his meds began to test for any needed dose adjustments. We can come to the home or he can schedule with a GP in the hospital. We discussed that he's enjoyed working with (b) (6) before.</p>
6/6/2016	C	(b) (6)	<p>MEDICAL COMMENTS 6/6/2016 11:47 FDA complaint submitted: Pet Food Safety Report, ID 54405, was successfully submitted on 6/6/2016 11:44:41 AM EST to the FDA, and it was issued an Individual Case Safety Report Number (ICSR) of 1053339</p>
(b) (6)	TC	(b) (6)	<p>MEDICAL COMMENTS - TENTATIVE (b) (6) 10:46 SW (b) (6) at Merrick - updated company that we have documented taurine deficiency in 2 other cats in house hold. The quality assurance team indicated that the level of taurine in the lot # I gave them was sufficient - discussed that this likely takes 3-6 month to develop and likely to be related to earlier lot and they need to investigate further. Asked if the level they gave me was from them retesting the food after I called or from an earlier test - not sure. Asked for information from the taurine tests - told I can send summary of lab results. Also indicated that I am reporting this to the FDA. She give new info to her manager and quality assurance. Told them I expect them to follow up with me. Below email sent to Merrick:</p> <p>Taurine Levels (b) (6) To: (b) (6) @merrickpetcare.com Hi (b) (6),</p> <p>Thank you for your help with these cases. Here is the summary of the lab results:</p> <p>12yr female spayed domestic short hair diagnosed and clinical for dilated cardiomyopathy (b) (6) 016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016</p> <p>5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016

-9yr male neutered domestic long hair: 196 nmol/ml

-8y female spayed domestic short hair: 368 nmol/ml

-9yr male neutered domestic long hair: 124 nmol/ml

-9yr male neutered domestic long hair: 536 nmol/ml

Please let me know if you have any other questions.

Sincerely,

(b) (6)

(b) (6)

5/31/2016 C

(b) (6)

COMMUNICATIONS WITH CLIENT - Closed (b) (6)

5/31/2016 16:14

Spoke with husband, he confirmed email was received and (b) (6) and (b) (6) have begun taurine supplementation. Discussed results are a different normal range for the whole blood testing I did vs the plasma testing the cardiologist did on (b) (6). He notes (b) (6) had been getting some Fancy Feast so that likely explains why his values are top of the normal range. Discussed (b) (6) elevated globulins, need protein electrophoresis to better define the issue, can be chronic inflammatory response or a few types of cancer. If cancer, because they are RB clients they can consult with oncology at no charge to hear what the treatment options would be. They can bring (b) (6) into the office with a GP doctor for the bloodwork or I can do the bloodwork at the house. If I am doing it, I would not charge any recheck exam, just the travel and diagnostic test costs. If a GP in the hospital, they shouldn't have to charge for an exam either because she was just checked in late May. Owner likes (b) (6); advised he could schedule that with her. Discussed rechecking (b) (6) T4 and liver values 6 weeks after starting meds to check if dose needs adjusting; can be done at the house or in the office as well. Advised Nutrition is contacting Merrick about the taurine results and she feels the owner shouldn't have to pay for the taurine testing; she wants Merrick to have to pay for it directly. So I

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

told the owner that first and foremost, they are responsible for payment of the testing to (b) (6) and once we advise them of a charge, they would be required to pay it. If the Nutritionist is able to circumvent that by having Merrick pay us directly, that would be a nice advantage for the client. He understands.

5/31/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed Jun 02/2016 5/31/2016 15:39 LMOM on husband cell making sure they received my treatment advice in the email from over the weekend. Please call back or reply to email so I can be certain the treatment guidelines were received.
-----------	---	---------	---

5/28/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 30/2016 5/28/2016 15:46 Email to client: Hi Mr and Mrs (b) (6) – I left a (long-winded) message on Mr (b) (6) voicemail earlier today. The nutritionist has since been in contact with me and advised that both (b) (6) and (b) (6) should be started on taurine supplementation. She recommends 250mg taurine twice daily for 2-3 weeks. Because you've already switched them to another diet, after 2-3 weeks, supplementation can be discontinued. (b) (6) and (b) (6) tested safely within the normal range for taurine, so they do not require any supplementation. I presume since you have already been treating (b) (6), you likely have a supply of taurine supplement. If not, feel free to contact me (b) (6) or the nutritionist or cardiologist to get a larger supply in order to treat the brothers. The nutritionist also advised she'll be contacting Merrick again now that the data has been received. Once she has heard more from them, she'll be in contact with you, as well. I also mentioned in the voicemail that (b) (6) blood test was repeated and verified that she does have elevated globulins. The most harmless reason would be chronic inflammation, but since she's been otherwise healthy, it is valuable to pursue further diagnostic inquiry. Unfortunately, elevated globulins can also indicate cancer, so we want to determine precisely what is happening with her. We can collect another blood sample from her at any time in order to perform a test called protein electrophoresis which further defines which specific immunoglobulins are elevated. You may choose to bring her into the office or have us out to the home again. (b) (6) will need repeat bloodwork after he's been on his thyroid supplement for 6 weeks, we could collect her second sample at that time as well, if
-----------	---	---------	---

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

you choose.

Feel free to contact me with any questions. I will next be in the office on Tuesday May 31st.

Happy Memorial Day weekend –

(b) (6)

5/28/2016 C

(b) (6)

COMMUNICATIONS WITH DOCTOR - Closed May 30/2016

5/28/2016 15:45

Email response from (b) (6) to (b) (6)

I'm also going to talk with (b) (6) this week about the cost of the taurine test. In my opinion this should be paid for by the company. I don't want the owner to pay the cost yet until I talk with (b) (6) and the company again.

(b) (6)

From: (b) (6)

Sent: Saturday, May 28, 2016 2:16 PM

To: (b) (6)

Subject: RE: price for taurine test

There is a risk of deficiency with anything <200, so that's why I would go ahead and supplement both cats...and it's harmless:-)

From: (b) (6)

Sent: Saturday, May 28, 2016 2:13 PM

To: (b) (6)

Subject: RE: price for taurine test

It is really interesting...probably the same reason some puppies raised on an unbalanced home cooked diet never have issues and other do.

Great the diet has been changed. We should get the cats that tested low on some supplementation for 2-3 weeks just to cover our bases. 250mg taurine PO BID...if she needs to use a powder form and mix with the cats food that's fine

Let the owner know I will touch base with the company after Memorial day...I have not heard back from them yet. This will also give me much more to go on when reporting to the FDA...who know this might turn into a pet food recall (it should turn into a recall)!

Thank you so much for the update!

(b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

(b) (6)

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
			<p>From: (b) (6) Sent: Saturday, May 28, 2016 1:55 PM To: (b) (6) Subject: RE: price for taurine test Hi (b) (6) – Thanks for providing this justification for the lab decision; I really appreciate it! I left you a voicemail earlier today – the results are in. 2 cats tested within the normal range [(b) (6) 368, (b) (6) 536 (300-600)]. (b) (6) was 196 and (b) (6) was way down at 124. All 4 cats were switched to Royal Canin food about 7 days ago. I left a voicemail for the client advising of the results, but told him I wanted your input before devising a treatment strategy. I would think of these 4, only (b) (6) would benefit substantially from taurine supplementation. I presume (b) (6) levels are sufficient now that he's been put on a properly formulated diet. Do you agree? This case is so interesting... how the cats fall all along the clinical spectrum, including some that have sufficient taurine, despite all eating the same presumably flawed diet. Thanks for your input, (b) (6)</p>
5/28/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 30/2016 5/28/2016 13:44 LMOM - advised client taurine results have been received, I have notified nutrition dept who will weigh in on treatment decision-making. Normal range is 300-600 and (b) (6) and (b) (6) tested within that range. Clinical signs are unlikely above 200, (b) (6) was 196, so it is likely he wouldn't show any issues. (b) (6) tested at 124 so he might be the one to benefit from additional supplementation, aside from just the diet change to the Royal Canin food. We will wait to initiate any therapy until the nutritionist has a chance to comment; we are working as a team on this. Since we have results, we likely have an invoice from the lab as well, so we should be able to advise of the cost of this testing in the short-term. I had spoken with his wife about (b) (6) having elevated globulins and on the re-test that status persists, was verified. Recommend additional blood testing for further work-up, could be collected when we visit (b) (6) for bloodwork 6 weeks after starting his thyroid meds. Please call back to discuss these results.</p>
5/27/2016	TC	(b) (6)	<p>LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:32 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
5/24/2016	P	(b) (6)	100.00 tablet of Tapazole (Methimazole) 5mg Tablet (M066) Rx #: 2576298 0 Of 0 Refills Give 1 tablet by mouth twice daily. Check bloodwork for dose adjustment 6 weeks after starting medication.
5/24/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:41 Spoke with Mrs; advised hyperthyroid with some liver elevations. Reviewed life-long treatment, bloodwork 6 weeks after med started and then twice yearly if stable. If dose is changed after first bloodwork, we repeat bloodwork again 6 weeks later until properly regulated. Med can be tablet, liquid or transdermal. Owner wants to crush tablet into canned food; advised this is fine as long as we're certain he's the only one who might consume the medicated food within their group-housing situation. Owner feels she can guarantee that. Meds will be at 197 pharmacy. Taurine pending, will call. Advised final pricing on taurine at (b) (6) lab not yet determined, will be in touch with that info as soon as finalized. Owner asked why use a diff lab; advised nutritionists recommended this lab, specialized testing at university, two labs finding low levels strengthens case against food company.
5/24/2016	B	(b) (6)	100.00 tablet of Tapazole (Methimazole) 5mg Tablet (M066) by (b) (6)
5/24/2016	B	(b) (6)	1.00 Cared for by (b) (6) (RHS) by (b) (6)
5/21/2016	C	(b) (6)	GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Weight loss/Taurine check History: Owner notes chronic weight loss across the recent months. Was losing hair for over a year, but was told it was related to anxiety. Eats with voracious appetite. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; he is one of 4 cats that live together in a room above the garage. Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household. Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food Meds: none S: BAR, PCS 0/4 - slightly feisty O: MM / ORPH: Pink, moist, crt <2 sec, mild tartar E/E: mild black debris in outer cartilages of left ear, deep canal WNL, right ear WNL. ophtho WNL.

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

INT: alopecia caudal dorsum, ventrum, lateral thighs. no ectoparasites observed.
 PLN: WNL
 CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic
 GI/UG: Compliant, no masses
 MS/NS: Normal amb x4
 BCS: 1.5-2/5 4.4kg
 A: 9yr9mo MN DLH
 1) weight loss - r/o hyperthyroid, diabetes mellitus, organ dz (kidney, liver), other endocrinopathy, neoplasia, nutritional problem
 2) alopecia - r/o FAD, other derm issue, psychogenic (stress, pain-related)
 3) dental disease
 4) otic debris - r/o infection vs inadequate grooming
 P: PE
 Taurine level
 CBC/Superchem/T4

Advised client of marked weight loss from last documented weight. Systemic bloodwork may illuminate the reason; will call with results next week. Taurine level will take 7-10 days.

Advised client we are sending taurine test to a different lab than the one that tested (b) (6) sample, at the advice of the nutrition service. We do not have a price in our computer system for this test through this lab, so the client will be invoiced for the taurine level (for all 4 cats) once that is established. Client paid today's services during the visit and is aware of the pending charge; advised the (b) (6) charge was \$214 and the charge at the other lab will likely be within \$50 under/over that fee. He commits to paying taurine test fees once advised of final fee. Stated we want to submit samples for testing ASAP and he understands fee structure will not be set until after tests are underway.

5/21/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 23/2016 5/21/2016 16:25 mailed welcome card, magnet, Rabies certificates (b) (6), (b) (6)) and feedback postcard
-----------	---	---------	---

5/21/2016	V	(b) (6)	May 21, 2016 11:21 AM Staff: (b) (6) ----- Weight : 4.40 kilograms HC-RS scale
-----------	---	---------	---

5/21/2016	L	(b) (6)	Hematology results from (b) (6) ID: 209396 Posted Final
-----------	---	---------	--

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

Test	Result	Reference Range
HCT	42 %	29 - 48
HGB	15.0 g/dL	9.3 - 15.9
MCHC	35.7 g/dL	30 - 38
WBC	14.2 10 ³ /uL	3.5 - 16.0
Bands	0 %	0 - 3
RBC	9.5 10 ⁶ /uL	5.92 - 9.93
MCV	44 fL	37 - 61
MCH	15.8 pg	11 - 21
ABS BASO	0 /uL	0 - 150
ABS NEUTB	0 /uL	0 - 150
Platelet C	254 10 ³ /uL	200 - 500
Platelet E	ADEQUATE	ADEQUATE -
Neutrophil	53 %	35 - 75
Lymphocyte	41 %	20 - 45
Monocytes	2 %	1 - 4
Eosinophil	4 %	2 - 12
Basophils	0 %	0 - 1
Absolute N	7526 /uL	2500 - 8500
Absolute L	5822 /uL	1200 - 8000
Absolute M	284 /uL	0 - 600
Absolute E	568 /uL	0 - 1000

Ascn: (b) (6) Profile: Complete Blood Count

5/21/2016 L

Chemistry results from (b) (6)

ID: 209396	Posted	Final	
Test	Result		Reference Range
ALB	3.4 g/dL		2.5 - 3.9
ALKP	174 U/L H		6 - 102
ALT	243 U/L H		10 - 100
AMYL	882 U/L		100 - 1200
AST	46 U/L		10 - 100
BUN/UREA	17 mg/dL		14 - 36
Ca	9.2 mg/dL		8.2 - 10.8
Chloride	111 mEq/L		104 - 128
CHOL	181 mg/dL		75 - 220
CK	124 U/L		56 - 529
CREA	0.6 mg/dL		0.6 - 2.4
GGT	3 U/L		1 - 10
GLU	80 mg/dL		64 - 170
Mg	1.7 mEq/L		1.5 - 2.5
PHOS	5.9 mg/dL		2.4 - 8.2
Potassium	4.5 mEq/L		3.4 - 5.6
Sodium	150 mEq/L		145 - 158
TBIL	0.1 mg/dL		0.1 - 0.4
TP	5.9 g/dL		5.2 - 8.8
TRIG	58 mg/dL		25 - 160
GLOB	2.5 g/dL		2.3 - 5.3

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates,
 I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended,
 R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)	Patient: (b) (6)	
Phone: (b) (6)	Species: Feline	Breed: Longhair, Domestic
Address: (b) (6)	Age: 9 Yrs. 10 Mos.	Sex: Neutered Male
(b) (6)	Color: Calico	

Date	Type	Staff	History																
			<table style="width: 100%; border: none;"> <tr> <td style="width: 33%;">A/G Ratio</td> <td style="width: 33%;">1.4 Ratio</td> <td style="width: 33%;">0.35 - 1.5</td> </tr> <tr> <td>B/C Ratio</td> <td>28 Ratio</td> <td>4 - 33</td> </tr> <tr> <td>Na/K Ratio</td> <td>33</td> <td></td> </tr> </table>	A/G Ratio	1.4 Ratio	0.35 - 1.5	B/C Ratio	28 Ratio	4 - 33	Na/K Ratio	33								
A/G Ratio	1.4 Ratio	0.35 - 1.5																	
B/C Ratio	28 Ratio	4 - 33																	
Na/K Ratio	33																		
5/21/2016	L		<p>Endocrinology results from (b) (6)</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 33%;"></td> <td style="width: 33%;">ID: 209396</td> <td style="width: 33%;">Posted</td> <td style="width: 33%;">Final</td> </tr> <tr> <td>Test</td> <td>Result</td> <td>Reference</td> <td>Range</td> </tr> <tr> <td>T4</td> <td>20.2 ug/dL H</td> <td>0.8 - 4.0</td> <td></td> </tr> <tr> <td>Ascن:</td> <td>(b) (6)</td> <td>Profile: Total T4</td> <td></td> </tr> </table> <p>Result verified.</p>		ID: 209396	Posted	Final	Test	Result	Reference	Range	T4	20.2 ug/dL H	0.8 - 4.0		Ascن:	(b) (6)	Profile: Total T4	
	ID: 209396	Posted	Final																
Test	Result	Reference	Range																
T4	20.2 ug/dL H	0.8 - 4.0																	
Ascن:	(b) (6)	Profile: Total T4																	
5/21/2016	L		<p>Miscellaneous results from (b) (6)</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 33%;"></td> <td style="width: 33%;">ID: 209396</td> <td style="width: 33%;">Posted</td> <td style="width: 33%;">Final</td> </tr> <tr> <td>Ascن:</td> <td>(b) (6)</td> <td>Profile: Superchem</td> <td></td> </tr> </table> <p>RE: 1045 PrecisionP 28 U/L 8 - 26 PrecisionPSL elevations correlate closely with abnormal PLI concentrations. In cats with appropriate clinical signs, this PrecisionPSL is supportive of, but not definitive, for a diagnosis of pancreatitis. In cats without clinical signs of pancreatitis, a mild elevation is an insignificant finding. RE: 11067 Comment Hemolysis 1+ No significant interference.</p>		ID: 209396	Posted	Final	Ascن:	(b) (6)	Profile: Superchem									
	ID: 209396	Posted	Final																
Ascن:	(b) (6)	Profile: Superchem																	
5/21/2016	B	(b) (6)	1.00 Superchem Cbc T4 (b) (6) Sa120 (L85) by (b) (6)																
5/21/2016	B	(b) (6)	1.00 House Call Travel Level 2 (HC06) by (b) (6)																
5/21/2016	B	(b) (6)	1.00 At Home Appointment (HC04) by (b) (6)																
5/21/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)																
5/21/2016	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)																
5/21/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)																
5/21/2016	B	(b) (6)	1.00 Lab Sample Label (b) (6) by (b) (6)																
5/21/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)																
5/20/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 21/2016 5/20/2016 15:00 Called to confirm tomorrow's appointment fro (b) (6) , (b) (6) , (b) (6) and (b) (6) at 9 am. I also mentioned in my message that we should use the address (b) (6) in the GPS. If any questions please call (b) (6)</p>																

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

5/17/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 19/2016 5/17/2016 12:29 Responding to owner's message, booked (b) (6), (b) (6), (b) (6), (b) (6) for House Call on Saturday 5/21. (b) (6) was seen on emergency and diagnosed with low taurine, so all cats need to be screened. Had been eating Merrick dry food. Cats are kept in a finished room above the garage; he thinks they won't need to be confined/isolated more than that in order to work on them. Discussed senior bloodwork as well. He notes this emergency with (b) (6) was a wake-up call and he'd like to thoroughly have everyone checked out. (b) (6) is losing weight. (b) (6) and (b) (6) have been to another vet and have a current Rabies; (b) (6) and (b) (6) haven't been to the vet in a long time. Discussed PureVax 1yr vs 3yr vs standard RabVac, vaccine-associated sarcoma issue - owner wants the purified vaccine, prefers the one year since they should be examined annually anyway. Advised if (b) (6) status progresses and we need to be checking her as well, please call to inform us in case we need special items/supplies for her care. Owner notes we should use (b) (6) with the GPS; his home address was renamed/renumbered a few years ago, but GPS cannot often find (b) (6)</p>
-----------	---	---------	--

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

6/7/2016 TC (b) (6) COMMUNICATIONS WITH CLIENT - TENTATIVE
6/7/2016 12:25
Spoke with owner, advised of specialist's comments about protein electrophoresis. Can conduct further testing through our Internal Medicine service of ultrasounds of chest and abdomen, essentially searching for origin of chronic inflammation. It is possible that testing will come back normal, despite the bloodwork indicating the elevated globulins. Another option is to track her body weight at home once monthly and report any weight loss promptly; if none noted, recheck bloodwork in 3 months to see if globulins are resolved. Specific cancers cause specific spikes and her results do not show those elevations, so that is good news. But we don't have a definite reason for her chronic inflammation. Owner understands and will relate to wife.

6/7/2016 TC (b) (6) COMMUNICATIONS WITH DOCTOR - TENTATIVE
6/7/2016 12:11
Spoke with (b) (6) at (b) (6); she notes monoclonal globulin spikes are most concerning because they define lymphoma, myeloma. Polyclonal spikes can be FIP, but also any cause of chronic inflammation. There is a chance chronic inflammation can be associated with neoplastic process though. She notes the degree of elevation is mild. If the owners want to work this up aggressively, full body imaging, best with ultrasound, to search for cancer. If they would like to monitor, recheck chemistry panel in 3 months and assess globulin count at that time. Only repeat electrophoresis if significantly higher elevation. If pet is losing weight, neoplasia moves up the list of differentials. Advised chronic otitis externa in this otherwise healthy patient; doctor concurs that wouldn't be sufficient to cause systemic response.

(b) (6) L

Chemistry results from (b) (6)

ID:	209396	Posted	Final	
Test	Result		Reference	Range
ALB	2.9 g/dL		2.5	3.9
TP	8.2 g/dL		5.2	8.8
GLOB	5.3 g/dL		2.3	5.3
ALPHA 1	0.3 g/dL		0.2	1.1
ALPHA 2	0.7 g/dL		0.4	0.9
BETA	0.6 g/dL		0.3	0.9
GAMMA	3.6 g/dL H		0.3	2.5

(b) (6) L **Miscellaneous results from (b) (6)**

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

(b) (6) ID: 209396 Posted Final
 Asc: (b) (6) Profile: Protein Electrophoresis, Serum
 RE: 1140 Interpretation
 The gamma globulin fraction is elevated, characterized by a broad polyclonal band, resulting from a mixture of increased immunoglobulins associated with an immune response. Potential causes include suppurative disease, chronic infectious disease (bacterial; protozoal; viral; rickettsial; fungal), connective tissue disease, chronic granulomatous disease, etc. Correlate with clinical findings.
PATHOLOGIST:

(b) (6), BVSc (Hons 1), DACVP
 (b) (6)

Due to difference in method of analysis, there may be slight differences in the quantitative albumin and calculated globulin results between serum electrophoresis results compared to a general chemistry panel.

(b) (6)	C	(b) (6)	MEDICAL COMMENTS - Closed Jun 04/2016 (b) (6) 18:35 Drew sample for protein electrophoresis while at the home for EOL care for (b) (6).
---------	---	---------	---

(b) (6)	B	(b) (6)	1.00 House Call Travel Level 2 (HC06) by (b) (6) Laboratory Request / Sample Handling (LABS) by (b) (6) 1.00 Outside Lab (XTBALUO) by (b) (6) 1.00 Protein Electrophor. Serum (b) (6) T240 (L018) by (b) (6) 1.00 Sample Handling & Disposal (LFEE) by (b) (6) 1.00 Lab Sample Label (TL) by (b) (6) 1.00 Cared for by (b) (6) (RHS) by (b) (6)
---------	---	---------	---

(b) (6)	C	RHS	COMMUNICATIONS WITH CLIENT - Closed Jun 02/2016 (b) (6) 16:21 (See full phone call under (b) (6) record)
---------	---	-----	--

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/27/2016	TC	(b) (6)	<p>Discussed (b) (6) elevated globulins, need protein electrophoresis to better define the issue, can be chronic inflammatory response or a few types of cancer. If cancer, because they are (b) (6) clients they can consult with oncology at no charge to hear what the treatment options would be. They can bring (b) (6) into the office with a GP doctor for the bloodwork or I can do the bloodwork at the house. If I am doing it, I would not charge any recheck exam, just the travel and diagnostic test costs. If a GP in the hospital, they shouldn't have to charge for an exam either because she was just checked in late May. Owner likes (b) (6); advised he could schedule that with her.</p>
5/27/2016			<p>LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:36 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)</p>
5/24/2016	C	(b) (6)	<p>COMMUNICATIONS WITH DOCTOR - Closed May 26/2016 5/24/2016 16:48 Spoke with doctor at (b) (6) consult line - she opted to rerun the full chemistry profile to validate the results since (b) (6) remaining profile is so normal. If globulins are truly elevated, protein electrophoresis is the next step. Ddx: myeloma, lymphoma, FIP, other neoplasia, chronic inflammatory condition. Asked specifically about taurine based on (b) (6) and current investigation into whole household's taurine status; not aware of any relationship between globulins and taurine.</p>
5/24/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:46 Spoke with Mrs; (b) (6) has elevated globulins which can indicate cancer or a chronic inflammatory condition. Spoke with specialist and no correlation with taurine deficiency. Lab is going to re-run her full profile to validate the results. Expect an update in 1-2 days. If verified, we may need to collect additional blood for the next level of testing which tells us which specific pattern of globulins is elevated. Taurine pending, will call.</p>
5/21/2016	C	(b) (6)	<p>GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Wellness/Taurine check History: Doing well. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; she is one of 4 cats that live together in a room above the</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
------	------	-------	---------

garage. Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household.
 Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food
 Meds: none
 S: BAR, PCS 0/4
 O:
 MM / ORPH: Pink, moist, crt <2 sec, small suspect FORL right upper PM3, mild tartar overall.
 E/E: copious black debris AU, mildly pruritic while cleaning. ophtho WNL.
 INT: WNL; no evidence of ectoparasites observed
 PLN: WNL
 CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic
 GI/UG: Compliant, no masses
 MS/NS: Normal amb x4
 BCS: 3/5 4.15kg
 A: 8yr FS DSH
 1) otitis externa - r/o bacterial/fungal vs ear mites
 2) dental disease
 P: PE
 Taurine level
 CBC/Vetscreen
 Disp Tresaderm 7.5ml - apply 2-3 drops in each ear twice daily for 7-10 days, keep in fridge
 ear cleaning
 PureVax Rabies 1yr SQ right hind (lot#17390B, exp 12/11/2016)

Discussed ear infection and treatment. Will call with lab results; systemic early next week, taurine in 7-10 days.

5/21/2016	I	(b) (6)	An animal is not considered immunized for at least 28 days after the initial or primary vaccination is administered. For this reason, pets receiving their first rabies vaccine should not be left outdoors unattended.						
5/21/2016	P	(b) (6)	1.00 bottle of Tresaderm 7.5ml (Merial] (M225) Rx #: 2574865 0 Of 0 Refills Apply 2-3 drops in each ear twice daily for 7-10 days.						
5/21/2016	V	(b) (6)	May 21, 2016 11:15 AM Staff: (b) (6) ----- Weight : 4.15 kilograms HC-RS scale						
5/21/2016	L	(b) (6)	Hematology results from (b) (6) <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%;">ID: 209396</th> <th style="width: 30%;">Posted</th> <th style="width: 40%;">Final</th> </tr> <tr> <th>Test</th> <th>Result</th> <th>Reference Range</th> </tr> </thead> </table>	ID: 209396	Posted	Final	Test	Result	Reference Range
ID: 209396	Posted	Final							
Test	Result	Reference Range							

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
			HCT 31 % 29 - 48 HGB 11.0 g/dL 9.3 - 15.9 MCHC 35.5 g/dL 30 - 38 WBC 13.4 10 ³ /uL 3.5 - 16.0 Bands 0 % 0 - 3 RBC 6.7 10 ⁶ /uL 5.92 - 9.93 MCV 46 fL 37 - 61 MCH 16.4 pg 11 - 21 ABS BASO 0 /uL 0 - 150 ABS NEUTB 0 /uL 0 - 150 Platelet C 375 10 ³ /uL 200 - 500 Platelet E ADEQUATE ADEQUATE - Neutrophil 55 % 35 - 75 Lymphocyte 37 % 20 - 45 Monocytes 2 % 1 - 4 Eosinophil 6 % 2 - 12 Basophils 0 % 0 - 1 Absolute N 7370 /uL 2500 - 8500 Absolute L 4958 /uL 1200 - 8000 Absolute M 268 /uL 0 - 600 Absolute E 804 /uL 0 - 1000 Ascn: (b) (6) Profile: Complete Blood Count

5/21/2016 L

Chemistry results from (b) (6)

ID:	209396	Posted	Final	Reference Range
Test	Result			
ALB	2.6 g/dL			2.5 - 3.9
ALKP	18 U/L			6 - 102
ALT	14 U/L			10 - 100
AST	11 U/L			10 - 100
BUN/UREA	29 mg/dL			14 - 36
Ca	8.8 mg/dL			8.2 - 10.8
Chloride	113 mEq/L			104 - 128
CHOL	94 mg/dL			75 - 220
CK	62 U/L			56 - 529
CREA	1.2 mg/dL			0.6 - 2.4
GLU	79 mg/dL			64 - 170
PHOS	5.7 mg/dL			2.4 - 8.2
Potassium	4.8 mEq/L			3.4 - 5.6
Sodium	148 mEq/L			145 - 158
TBIL	0.1 mg/dL			0.1 - 0.4
TP	9.1 g/dL H			5.2 - 8.8
GLOB	6.5 g/dL H			2.3 - 5.3
A/G Ratio	0.4 Ratio			0.35 - 1.5
B/C Ratio	24 Ratio			4 - 33
Na/K Ratio	31			

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 8 Yrs. 0 Mos.
Color: brown tabby

Breed: Shorthair, Domestic
Sex: Spayed Female

Date	Type	Staff	History
5/21/2016	L		<p>Miscellaneous results from (b) (6) ID: 209396 Posted Final Asc: (b) (6) Profile: Vet Screen RE: 11067 Comment Hemolysis 1+ No significant interference.</p>
5/21/2016	B	(b) (6)	1.00 At Home Additional Pet Appointment (HC03) by (b) (6)
5/21/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Vetscreen Cbc Antec SA030 (L00030) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Lab Sample Label (TL) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Fel Vax Rabies 1 Year Purevax (Merial) (V21) by (b) (6)
5/21/2016	B	(b) (6)	1.00 bottle of Tresaderm 7.5ml (Merial) (M225) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)

5/20/2016 C (b) (6)

COMMUNICATIONS WITH CLIENT - Closed May 21/2016
 5/20/2016 15:04
 Called to confirm tomorrow's appointment from (b) (6), (b) (6), (b) (6) and (b) (6) at 9 am. I also mentioned in my message that we should use the address (b) (6) in the GPS. If any questions please call (b) (6)

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

(b) (6) TC (b) (6)

MEDICAL COMMENTS - TENTATIVE

(b) (6) 10:46

SW (b) (6) at Merrick - updated company that we have documented taurine deficiency in 2 other cats in house hold. The quality assurance team indicated that the level of taurine in the lot # I gave them was sufficient - discussed that this likely takes 3-6 month to develop and likely to be related to earlier lot and they need to investigate further. Asked if the level they gave me was from them retesting the food after I called or from an earlier test - not sure. Asked for information from the taurine tests - told I can send summary of lab results. Also indicated that I am reporting this to the FDA. She give new info to her manager and quality assurance. Told them I expect them to follow up with me. Below email sent to Merrick:

Taurine Levels

(b) (6)

To:

(b) (6) @merrickpetcare.com

Hi (b) (6) ,

Thank you for your help with these cases. Here is the summary of the lab results:

12yr female spayed domestic short hair diagnosed and clinical for dilated cardiomyopathy

(b) (6) 2016 Plasma Taurine 24nmol/ml (normal 60-120, critical level <40) - test performed at University of Wisconsin, results were received on 5/15/2016

5/21/2016 - Whole Blood Taurine submitted at the University of California Davis on remaining 4 cats consuming this food (normal 300-600 nmol/ml, no known risk for deficiency >200), results were received on 5/27/2016

-9yr male neutered domestic long hair: 196 nmol/ml

-8y female spayed domestic short hair: 368 nmol/ml

-9yr male neutered domestic long hair: 124 nmol/ml

-9yr male neutered domestic long hair: 536 nmol/ml

Please let me know if you have any other questions.

Sincerely,

(b) (6)

(b) (6)

Clinical Nutrition Department

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
			(b) (6)
5/27/2016	TC	(b) (6)	LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:34 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)
5/24/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:50 Spoke with Mrs; systemic blood results WNL for (b) (6). Taurine pending.
5/21/2016	C	(b) (6)	GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Wellness/Taurine check History: Doing well. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; he is one of 4 cats that live together in a room above the garage. Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household. Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food Meds: none S: BAR, PCS 0/4 O: MM / ORPH: Pink, moist, crt <2 sec, moderate tartar overall E/E: ophtho/otoscopic exams WNL INT: no evidence of ectoparasites observed. matted hair present. PLN: WNL CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic GI/UG: Compliant, no masses MS/NS: Normal amb x4 BCS: 4/5 9.5kg A: 9yr9mo MN DLH 1) overweight 2) dental disease

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

3) matted hair
 P: PE
 Taurine level
 CBC/Vetscreen

Recommend weight loss, frequent brushing +/- clippers to remove mats or using a groomer. Dental condition warrants treatment. Will call with systemic blood results early next week, taurine level in 7-10 days.

5/21/2016	V	(b) (6)	May 21, 2016 11:21 AM Staff: (b) (6) ----- Weight : 9.50 kilograms HC-RS scale
-----------	---	---------	---

5/21/2016	L	(b) (6)	Hematology results from (b) (6) <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>ID: 209396</th> <th>Posted</th> <th>Final</th> <th></th> </tr> </thead> <tbody> <tr> <td>Test</td> <td>Result</td> <td></td> <td>Reference Range</td> </tr> <tr> <td>HCT</td> <td>40 %</td> <td></td> <td>29 - 48</td> </tr> <tr> <td>HGB</td> <td>12.3 g/dL</td> <td></td> <td>9.3 - 15.9</td> </tr> <tr> <td>MCHC</td> <td>30.8 g/dL</td> <td></td> <td>30 - 38</td> </tr> <tr> <td>WBC</td> <td>11.6 10³/uL</td> <td></td> <td>3.5 - 16.0</td> </tr> <tr> <td>Bands</td> <td>0 %</td> <td></td> <td>0 - 3</td> </tr> <tr> <td>RBC</td> <td>7.9 10⁶/uL</td> <td></td> <td>5.92 - 9.93</td> </tr> <tr> <td>MCV</td> <td>51 fL</td> <td></td> <td>37 - 61</td> </tr> <tr> <td>MCH</td> <td>15.6 pg</td> <td></td> <td>11 - 21</td> </tr> <tr> <td>ABS BASO</td> <td>0 /uL</td> <td></td> <td>0 - 150</td> </tr> <tr> <td>ABS NEUTB</td> <td>0 /uL</td> <td></td> <td>0 - 150</td> </tr> <tr> <td>Platelet C</td> <td>188 10³/uL L</td> <td></td> <td>200 - 500</td> </tr> <tr> <td>Platelet E</td> <td>ADEQUATE</td> <td></td> <td>ADEQUATE -</td> </tr> <tr> <td>Neutrophil</td> <td>72 %</td> <td></td> <td>35 - 75</td> </tr> <tr> <td>Lymphocyte</td> <td>21 %</td> <td></td> <td>20 - 45</td> </tr> <tr> <td>Monocytes</td> <td>3 %</td> <td></td> <td>1 - 4</td> </tr> <tr> <td>Eosinophil</td> <td>4 %</td> <td></td> <td>2 - 12</td> </tr> <tr> <td>Basophils</td> <td>0 %</td> <td></td> <td>0 - 1</td> </tr> <tr> <td>Absolute N</td> <td>8352 /uL</td> <td></td> <td>2500 - 8500</td> </tr> <tr> <td>Absolute L</td> <td>2436 /uL</td> <td></td> <td>1200 - 8000</td> </tr> <tr> <td>Absolute M</td> <td>348 /uL</td> <td></td> <td>0 - 600</td> </tr> <tr> <td>Absolute E</td> <td>464 /uL</td> <td></td> <td>0 - 1000</td> </tr> </tbody> </table> <p> Ascn: (b) (6) Profile: Complete Blood Count Ascn: (b) (6) Profile: Complete Blood Count </p> <p> Platelet count reflects the minimum number due to platelet clumping. </p>	ID: 209396	Posted	Final		Test	Result		Reference Range	HCT	40 %		29 - 48	HGB	12.3 g/dL		9.3 - 15.9	MCHC	30.8 g/dL		30 - 38	WBC	11.6 10 ³ /uL		3.5 - 16.0	Bands	0 %		0 - 3	RBC	7.9 10 ⁶ /uL		5.92 - 9.93	MCV	51 fL		37 - 61	MCH	15.6 pg		11 - 21	ABS BASO	0 /uL		0 - 150	ABS NEUTB	0 /uL		0 - 150	Platelet C	188 10 ³ /uL L		200 - 500	Platelet E	ADEQUATE		ADEQUATE -	Neutrophil	72 %		35 - 75	Lymphocyte	21 %		20 - 45	Monocytes	3 %		1 - 4	Eosinophil	4 %		2 - 12	Basophils	0 %		0 - 1	Absolute N	8352 /uL		2500 - 8500	Absolute L	2436 /uL		1200 - 8000	Absolute M	348 /uL		0 - 600	Absolute E	464 /uL		0 - 1000
ID: 209396	Posted	Final																																																																																													
Test	Result		Reference Range																																																																																												
HCT	40 %		29 - 48																																																																																												
HGB	12.3 g/dL		9.3 - 15.9																																																																																												
MCHC	30.8 g/dL		30 - 38																																																																																												
WBC	11.6 10 ³ /uL		3.5 - 16.0																																																																																												
Bands	0 %		0 - 3																																																																																												
RBC	7.9 10 ⁶ /uL		5.92 - 9.93																																																																																												
MCV	51 fL		37 - 61																																																																																												
MCH	15.6 pg		11 - 21																																																																																												
ABS BASO	0 /uL		0 - 150																																																																																												
ABS NEUTB	0 /uL		0 - 150																																																																																												
Platelet C	188 10 ³ /uL L		200 - 500																																																																																												
Platelet E	ADEQUATE		ADEQUATE -																																																																																												
Neutrophil	72 %		35 - 75																																																																																												
Lymphocyte	21 %		20 - 45																																																																																												
Monocytes	3 %		1 - 4																																																																																												
Eosinophil	4 %		2 - 12																																																																																												
Basophils	0 %		0 - 1																																																																																												
Absolute N	8352 /uL		2500 - 8500																																																																																												
Absolute L	2436 /uL		1200 - 8000																																																																																												
Absolute M	348 /uL		0 - 600																																																																																												
Absolute E	464 /uL		0 - 1000																																																																																												

5/21/2016	L	(b) (6)	Chemistry results from (b) (6)
-----------	---	---------	---------------------------------------

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative med note, V: Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

ID: 209396	Posted	Final	
Test	Result		Reference Range
ALB	3.1 g/dL		2.5 - 3.9
ALKP	27 U/L		6 - 102
ALT	64 U/L		10 - 100
AST	44 U/L		10 - 100
BUN/UREA	26 mg/dL		14 - 36
Ca	9.3 mg/dL		8.2 - 10.8
Chloride	112 mEq/L		104 - 128
CHOL	98 mg/dL		75 - 220
CK	157 U/L		56 - 529
CREA	1.2 mg/dL		0.6 - 2.4
GLU	90 mg/dL		64 - 170
PHOS	5.9 mg/dL		2.4 - 8.2
Potassium	5.1 mEq/L		3.4 - 5.6
Sodium	150 mEq/L		145 - 158
TBIL	0.1 mg/dL		0.1 - 0.4
TP	8.4 g/dL		5.2 - 8.8
GLOB	5.3 g/dL		2.3 - 5.3
A/G Ratio	0.6 Ratio		0.35 - 1.5
B/C Ratio	22 Ratio		4 - 33
Na/K Ratio	29		

5/21/2016 L (b) (6) **Miscellaneous results from (b) (6)**
 ID: 209396 Posted Final
 Asc: (b) (6) Profile: Vet Screen
 RE: 11067 Comment
 Hemolysis 1+ No significant interference.
 Asc: (b) (6) Profile: Vet Screen
 RE: 11067 Comment
 Hemolysis 1+ No significant interference.

5/21/2016 B (b) (6) 1.00 At Home Additional Pet Appointment (HC03) by (b) (6)
 5/21/2016 B (b) (6) Laboratory Request / Sample Handling (LABS) by (b) (6)
 5/21/2016 B (b) (6) 1.00 Outside Lab (XTBALUO) by (b) (6)
 5/21/2016 B (b) (6) 1.00 Vetscreen Cbc Antec SA030 (L00030) by (b) (6)
 5/21/2016 B (b) (6) 1.00 Sample Handling & Disposal (LFEE) by (b) (6)
 5/21/2016 B (b) (6) 1.00 Lab Sample Label (b) (6) by (b) (6)
 5/21/2016 B (b) (6) 1.00 Cared for by (b) (6) (b) (6) by (b) (6)

5/20/2016 C (b) (6) **COMMUNICATIONS WITH CLIENT - Closed May 21/2016**
 5/20/2016 15:04
 Called to confirm tomorrow's appointment fro (b) (6) , (b) (6) , (b) (6) and (b) (6) at 9

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
(b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 10 Mos.
Color: Calico

Breed: Longhair, Domestic
Sex: Male

Date	Type	Staff	History
------	------	-------	---------

am. I also mentioned in my message that we should use the address (b) (6) in the GPS. If any questions please call (b) (6)

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6) **Patient:** (b) (6)
Phone: (b) (6) **Species:** Feline **Breed:** Longhair, Domestic
Address: (b) (6) **Age:** 9 Yrs. 7 Mos. **Sex:** Neutered Male
(b) (6) **Color:**

Date	Type	Staff	History
6/7/2016	TC	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - TENTATIVE 6/7/2016 12:32 While speaking with owner about (b) (6), discussed (b) (6) dental. Spends the day at 197, but most often home that same night after procedure. Bloodwork is good for 2 months. Can schedule with GP or dentistry according to owner's preference.</p>
5/27/2016	TC	(b) (6)	<p>LAB RESULTS - NOTES - TENTATIVE 5/27/2016 15:38 Lab Results: Fax from U of Cal Davis. Original Lab Date: - Attachment(s)</p>
5/24/2016	C	(b) (6)	<p>COMMUNICATIONS WITH CLIENT - Closed May 26/2016 5/24/2016 16:51 Spoke with Mrs; bloodwork WNL, excellent news for planning anesthesia and dental work. Important that taurine status is addressed prior to anesthesia, but dental work should be planned for the next 4-8 weeks. Taurine pending, will call.</p>
5/21/2016	C	(b) (6)	<p>GP PHYSICAL EXAM - Closed May 26/2016 Date Presented: 5/21/2016 Chief Complaint: Wellness/Taurine check History: Doing well. Normal activity level. Normal ur/def. No known V/D/C/S. 5 cat, 2 dog household; he is the one cat who lives in the house (b) (6) is aggressive toward (b) (6), so he lives away from other cats). Indoor only. (b) (6) has DCM from taurine deficiency, checking status of other cats in household. Diet (type, freq, amt): Merrick Purrfect Bistro Grain Free chicken dry food Meds: none S: BAR, PCS 0/4 O: MM / ORPH: Pink, moist, crt <2 sec; right upper canine tooth loose, significant gingivitis locally. heavy tartar on PM3s bilaterally. missing incisors. E/E: brown debris in outer ear cartilages bilaterally, but canals clean/free of debris. ophtho exam WNL. INT: matted hair. no evidence of ectoparasites observed. PLN: WNL CV / Resp: Reg rhythm, no murmur, SSP, clear and eupneic GI/UG: Compliant, no masses</p>

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative med note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 7 Mos.
Color:

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

MS/NS: Normal amb x4
 BCS: 3-3.5/5 6.7kg
 A: 9yr7mo MN DLH
 1) dental disease
 2) matted hair
 P: PE
 Taurine level
 CBC/Superchem
 PureVax Rabies 1yr SQ right hind (lot# 17390B, exp 12/11/2016)

Advised dental status is poor and likely painful; recommend prompt dental cleaning under general anesthesia with extraction of canine +/- other teeth. Can use dental specialists or general practitioner depending on owner's preference. Recommend waiting for taurine level and any management pertaining to that issue before scheduling anesthetic procedure. Can be shaved down during anesthesia; frequent brushing +/- intermittent clipping or taking to a groomer is needed. Will call with systemic blood results early next week; taurine level will take 7-10 days.

5/21/2016	I	(b) (6)	An animal is not considered immunized for at least 28 days after the initial or primary vaccination is administered. For this reason, pets receiving their first rabies vaccine should not be left outdoors unattended.
-----------	---	---------	---

5/21/2016	V	(b) (6)	May 21, 2016 11:24 AM Staff: (b) (6) ----- Weight : 6.70 kilograms HC-RS scale
-----------	---	---------	---

5/21/2016	L	(b) (6)	Hematology results from (b) (6) <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">ID: 209396</th> <th style="text-align: left;">Posted</th> <th style="text-align: left;">Final</th> <th style="text-align: left;">Reference Range</th> </tr> </thead> <tbody> <tr> <td>Test</td> <td>Result</td> <td></td> <td></td> </tr> <tr> <td>HCT</td> <td>36 %</td> <td></td> <td>29 - 48</td> </tr> <tr> <td>HGB</td> <td>11.8 g/dL</td> <td></td> <td>9.3 - 15.9</td> </tr> <tr> <td>MCHC</td> <td>32.8 g/dL</td> <td></td> <td>30 - 38</td> </tr> <tr> <td>WBC</td> <td>9.8 10³/uL</td> <td></td> <td>3.5 - 16.0</td> </tr> <tr> <td>Bands</td> <td>0 %</td> <td></td> <td>0 - 3</td> </tr> <tr> <td>RBC</td> <td>7.9 10⁶/uL</td> <td></td> <td>5.92 - 9.93</td> </tr> <tr> <td>MCV</td> <td>46 fL</td> <td></td> <td>37 - 61</td> </tr> <tr> <td>MCH</td> <td>14.9 pg</td> <td></td> <td>11 - 21</td> </tr> <tr> <td>ABS BASO</td> <td>0 /uL</td> <td></td> <td>0 - 150</td> </tr> <tr> <td>ABS NEUTB</td> <td>0 /uL</td> <td></td> <td>0 - 150</td> </tr> <tr> <td>Platelet C</td> <td>490 10³/uL</td> <td></td> <td>200 - 500</td> </tr> <tr> <td>Platelet E</td> <td>ADEQUATE</td> <td></td> <td>ADEQUATE -</td> </tr> <tr> <td>Neutrophil</td> <td>59 %</td> <td></td> <td>35 - 75</td> </tr> <tr> <td>Lymphocyte</td> <td>33 %</td> <td></td> <td>20 - 45</td> </tr> <tr> <td>Monocytes</td> <td>2 %</td> <td></td> <td>1 - 4</td> </tr> <tr> <td>Eosinophil</td> <td>6 %</td> <td></td> <td>2 - 12</td> </tr> </tbody> </table>	ID: 209396	Posted	Final	Reference Range	Test	Result			HCT	36 %		29 - 48	HGB	11.8 g/dL		9.3 - 15.9	MCHC	32.8 g/dL		30 - 38	WBC	9.8 10 ³ /uL		3.5 - 16.0	Bands	0 %		0 - 3	RBC	7.9 10 ⁶ /uL		5.92 - 9.93	MCV	46 fL		37 - 61	MCH	14.9 pg		11 - 21	ABS BASO	0 /uL		0 - 150	ABS NEUTB	0 /uL		0 - 150	Platelet C	490 10 ³ /uL		200 - 500	Platelet E	ADEQUATE		ADEQUATE -	Neutrophil	59 %		35 - 75	Lymphocyte	33 %		20 - 45	Monocytes	2 %		1 - 4	Eosinophil	6 %		2 - 12
ID: 209396	Posted	Final	Reference Range																																																																								
Test	Result																																																																										
HCT	36 %		29 - 48																																																																								
HGB	11.8 g/dL		9.3 - 15.9																																																																								
MCHC	32.8 g/dL		30 - 38																																																																								
WBC	9.8 10 ³ /uL		3.5 - 16.0																																																																								
Bands	0 %		0 - 3																																																																								
RBC	7.9 10 ⁶ /uL		5.92 - 9.93																																																																								
MCV	46 fL		37 - 61																																																																								
MCH	14.9 pg		11 - 21																																																																								
ABS BASO	0 /uL		0 - 150																																																																								
ABS NEUTB	0 /uL		0 - 150																																																																								
Platelet C	490 10 ³ /uL		200 - 500																																																																								
Platelet E	ADEQUATE		ADEQUATE -																																																																								
Neutrophil	59 %		35 - 75																																																																								
Lymphocyte	33 %		20 - 45																																																																								
Monocytes	2 %		1 - 4																																																																								
Eosinophil	6 %		2 - 12																																																																								

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 7 Mos.
Color:

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
------	------	-------	---------

Basophils 0 % 0 - 1
Absolute N 5782 /uL 2500 - 8500
Absolute L 3234 /uL 1200 - 8000
Absolute M 196 /uL 0 - 600
Absolute E 588 /uL 0 - 1000
Ascن: (b) (6) **Profile:** Complete Blood Count

5/21/2016 L

Chemistry results from (b) (6)
 ID: 209396 Posted Final

Test	Result	Reference Range
ALB	3.8 g/dL	2.5 - 3.9
ALKP	25 U/L	6 - 102
ALT	32 U/L	10 - 100
AMYL	1067 U/L	100 - 1200
AST	14 U/L	10 - 100
BUN/UREA	32 mg/dL	14 - 36
Ca	9.9 mg/dL	8.2 - 10.8
Chloride	112 mEq/L	104 - 128
CHOL	125 mg/dL	75 - 220
CK	76 U/L	56 - 529
CREA	1.3 mg/dL	0.6 - 2.4
GGT	1 U/L	1 - 10
GLU	99 mg/dL	64 - 170
Mg	2.2 mEq/L	1.5 - 2.5
PHOS	5.5 mg/dL	2.4 - 8.2
Potassium	5.1 mEq/L	3.4 - 5.6
Sodium	150 mEq/L	145 - 158
TBIL	0.1 mg/dL	0.1 - 0.4
TP	7.8 g/dL	5.2 - 8.8
TRIG	97 mg/dL	25 - 160
GLOB	4.0 g/dL	2.3 - 5.3
A/G Ratio	1.0 Ratio	0.35 - 1.5
B/C Ratio	25 Ratio	4 - 33
Na/K Ratio	29	

5/21/2016 L

Miscellaneous results from (b) (6)
 ID: 209396 Posted Final
 (b) (6) **Profile:** Superchem
RE: 1045 PrecisionP 17 U/L 8 - 26
Acute pancreatitis is unlikely. Chronic pancreatitis is not excluded by a normal PrecisionPSL.
RE: 11067 Comment
Hemolysis 1+ No significant interference.

B:Billing, C:Med note, CB:Call back, CK:Check-in, CM:Communications, D:Diagnosis, DH:Declined to history, E:Examination, ES:Estimates, I:Departing instr, L:Lab result, M:Image cases, P:Prescription, PA:PVL Accepted, PB:problems, PP:PVL Performed, PR:PVL Recommended, R:Correspondence, T:Images, TC:Tentative medl note, V:Vital signs

Patient History Report

Client: (b) (6)
Phone: (b) (6)
Address: (b) (6)
 (b) (6)

Patient: (b) (6)
Species: Feline
Age: 9 Yrs. 7 Mos.
Color:

Breed: Longhair, Domestic
Sex: Neutered Male

Date	Type	Staff	History
5/21/2016	B	(b) (6)	1.00 At Home Additional Pet Appointment (HC03) by (b) (6)
5/21/2016	B	(b) (6)	Laboratory Request / Sample Handling (LABS) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Outside Lab (XTBALUO) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Superchem Cbc (b) (6) Sa020 (L07) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Sample Handling & Disposal (LFEE) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Lab Sample Label (b) (6) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Fel Vax Rabies 1 Year Purevax (Merial) (V21) by (b) (6)
5/21/2016	B	(b) (6)	1.00 Cared for by (b) (6) (b) (6) by (b) (6)
<hr/>			
5/20/2016	C	(b) (6)	COMMUNICATIONS WITH CLIENT - Closed May 21/2016 5/20/2016 15:05 Called to confirm tomorrow's appointment from (b) (6), (b) (6), (b) (6) and (b) (6) at 9 am. I also mentioned in my message that we should use the address (b) (6) in the GPS. If any questions please call (b) (6)

B: Billing, C: Med note, CB: Call back, CK: Check-in, CM: Communications, D: Diagnosis, DH: Declined to history, E: Examination, ES: Estimates, I: Departing instr, L: Lab result, M: Image cases, P: Prescription, PA: PVL Accepted, PB: problems, PP: PVL Performed, PR: PVL Recommended, R: Correspondence, T: Images, TC: Tentative medl note, V: Vital signs

14770

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/index.cfm>

Vet/Tech Contact: (b) (6)
Company Name: (b) (6)
Address: (b) (6)

Email: (b) (6)
Tel: (b) (6) Fax: (b) (6)

Billing Contact: (b) (6) TAX ID: (b) (6)
Email: (b) (6) Tel: (b) (6)

Patient Name: (b) (6)
Species: Feline
Owner's Name: (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 368 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

14769

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aa/index.cfm>

Vet/Tech Contact: (b) (6)
Company Name: _____
Address: _____

Email: (b) (6)
Tel: _____ Fax: (b) (6)

Billing (b) (6)
Email: _____ TAX ID: _____
Tel: (b) (6)

Patient Name: (b) (6)
Species: Feline
Owner's Name: (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)

Plasma: _____ Whole Blood: 124 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150

14771

Sample Submission Form

Amino Acid Laboratory
University of California, Davis
1020 Vet Med 3B
1089 Veterinary Medicine Drive
Davis, CA 95616
Tel: (530)752-5058, Fax: (530)752-4698

UC CUSTOMERS ONLY:
Non-federal funds ID/Account Number
to bill: _____

<http://www.vetmed.ucdavis.edu/vmb/labs/aal/index.cfm>

Vet/Tech Contact: (b) (6)
Company Name: (b) (6)
Address: (b) (6)

Email: (b) (6)
Tel: (b) (6) Fax: (b) (6)

Billing Contact: (b) (6) TAX ID: (b) (6)
Email: (b) (6) Tel: (b) (6)

Patient Name: (b) (6)
Species: Feline
Owner's Name: (b) (6)

Sample Type: Plasma Whole Blood Urine Food Other: _____
Test Items: Taurine Complete Amino Acid Other: _____

Taurine Results (nmol/ml)
Plasma: _____ Whole Blood: 536 Urine: _____ Food: _____

Reference Ranges (nmol/ml)

	Plasma		Whole Blood	
	Normal Range	No Known Risk for Taurine Deficiency	Normal Range	No Known Risk for Taurine Deficiency
Cat	80-120	>40	300-600	>200
Dog	60-120	>40	200-350	>150